$$1 \mid \int \frac{\sqrt{x-1}}{x} dx$$

Let
$$u = \sqrt{x-1}$$
, $du = \frac{1}{2\sqrt{x-1}}$

$$\int \frac{\sqrt{x-1}}{x} dx = \int \frac{u}{TODO} dx$$

- 2 | 2
- 3 | 3
- 4 | 4
- 5 | 5

$$6 \mid \int \tan^2 x + 1 dx$$

$$\int \tan^2 x + 1 dx = \int \sec^2 x - 1 + 1 dx$$
$$= \int \sec^2 x dx$$

$$\mathsf{Let} u = x, du = 1$$

$$= \int \sec^2 u du$$
$$= \tan u + C$$
$$= \boxed{\tan x + C}$$

- 7 | 7
- 8 | 8
- $9 \mid \int \frac{\sec^2 x}{\csc x} sinx dx$

$$\int \frac{\sec^2 x}{\csc x} \sin x dx = \int \tan^2 x dx$$
$$= \int \sec^2 x - 1 dx$$
$$= \int \sec^2 x dx - \int 1 dx$$
$$= \tan x - x$$

- 10 | 10
- 11 | $\int \frac{e^{2\ln\sin x} + e^{2\ln\cos x}}{e^{2\ln\tan x} + e^{2\ln 1}} dx$

$$\int \frac{e^{2\ln\sin x} + e^{2\ln\cos x}}{e^{2\ln\tan x} + e^{2\ln 1}} dx = \int \frac{\sin^2 x + \cos^2 x}{\tan^2 x + 1} dx$$

- 12 | **12**
- 13 | **13**
- 14 | **14**