

Source:

1 | eigenvalues

eigenvalue: multiplied by a scalar? a subspace that, when put through a linear map, only gets scaled.

$$Tv = \lambda v$$

Where $v \neq 0$. (we ignore it because its no fun to send zero to zero, and bc the span is empty).

T must be an operator! Otherwise the matrix sizes don't work out when subtracting λI .

where v is the eigenvector and λ is the eigenvalue. The equation is often rewritten as:

$$Tv - \lambda v = 0Tv - \lambda Iv = 0(T - \lambda I)v = 0$$

now this can be factored and roots can be found. also it's an operator.

1.1 | Axler 5.6 equivalent conditions

Only when V is finite dimensional!

1.1.1 $|T - \lambda I$ is not injective, because both $v, 0$ are in the null space.

1.1.2 $|T - \lambda I$ is also not surjective or invertible bc finite dim operator.

2 | an example

Given the matrix $\begin{pmatrix} 3 & 1 \\ 0 & 2 \end{pmatrix}$, find the eigenvalues and eigenvectors.

Now that we have that other fomulation, we can just subtract λI from T to get

$$\begin{pmatrix} 3 - \lambda & 1 \\ 0 & 2 - \lambda \end{pmatrix}$$

Then, we just need to find whether it is non-invertible aka singular aka determinant.

$$(3 - \lambda)(2 - \lambda) = 0$$

The solutions are $\lambda = 2$ or 3 . These are the eigenvalues.

3 | depends on

3.1 | finding roots is helpful