

Source: [KBIO101CentralDogma](#)

1 | Protein Synthesis

Let's make a protein together!

1.1 | Before we begin, some background

Genetic Code => "nucleotide code" found in the DNA that helps make protein. There are two parts of this: translation and transcription.

- The process of **Transcription** involves taking the DNA, separating it, and copying its corresponding pairs to RNA
- The process of **Translation** involves taking the RNA and making proteins.
- *Non-coding sequence*: metadata for DNA for the processors
- *Coding sequence*: DNA content for amino-acid production

Occasionally, the RNA is what we want to end up with, so then obviously we no longer need the process of Translation.

1.2 | Transcription => converting DNA to mRNA

The process of transcription is the process by which DNA is converted to messenger RNA, a type of RNA that travels to the ribosome to create a protein. This process is dependent on the enzyme **RNA Polymerase**, which is the primary driver that handles DNA transcription.

See [KBhBIO101DNATranscription](#)

1.3 | mRNA processing => splicing mRNA

After the transcribed mRNA is finished, Eukaryotes only will need to go through one additional process called "mRNA processing" that both remove the non-protein-synthesizing Introns of the mRNA sequence, and mark the mRNA for maturity.

Notably, **Prokaryotes does not do this!** Prokaryotes' coding sequence always makes a full protein, so we just start at promoter and end at terminator and make a protein!

See [KBhBIO101mRNAPreprocessing](#)

1.3.1 | Translation => RNA-directed polypeptide synthesis

And now, this is what we are here for. Now that we have a constructed and mature mRNA, let's make a protein!

See [KBhBIO101Translation](#)

Mature mRNA sent to ribosome. mRNA must travel to the cytoplasm in the Eukaryotes to catch the RNA, whereas in prokaryotes they don't have to go anywhere.

Ribosomes has two units: 50S unit + 30S unit => they come together whenever a mRNA needs it. Each contained specialized rRNA + tRNA to catalyze attachment of and carry amino acids + adapt the incoming mRNA respectively.

Note! The beginning of mRNA is not translated. There a portion on the 5' end of the mRNA (starts with AGGAGG) — about 170 nucleotides in humans, and shorter in bacteria — that's called UTR (untranslated region.) This region helps ribosomes bind to it + stabilize the binds.

- 3 protein factors IF1, IF2, IF3 forms a complex for transcription by binding to a subunit on the ribosome
- Methionine-carrying tRNA binds to the start of the mRNA, which forms the initiation complex. This is typically removed after translation if not coded for (if M-A amino acid pair coded for, methionine removed; but if M-L pairs coded for, methionine not removed.)
- A-site: translates mRNA to tRNA — anti-codon pairs
- P-site: amino acid dumped from tRNA to the actual chain being built
- Spent tRNA ejected to the E-site, which is then recycled
- Catalyst tRNA combines with rRNA to catalyze amino acid peptide bond
- Each codon (group of 3 units in tRNA), matches a specific [KBhBIO101AminoAcids](#)

Smaller ribosome unit grabs, larger attaches + forms amino acid

After the amino acids are assembled, it's time for [KBe2020bio101refProteinFolding](#). See also [KBhBIO101Proteins](#).

=> Shaperones fold proteins, and if its finds proteins impossible to fold, it flags it using ubiquitin to send to the garbage