Source: [KBBiologyMasterIndex]

## 0.1 | **#flo #ret**

## 1 | Notes

## 1.0.1 | **Overview**

- · Organisms turn genes on and off which is called Gene Expression
  - This can be done in response to external and internal signals
    - · These signals are based off of environmental factors
  - · This is also be done in order to specialize cells
    - Certain cells need certain genes to preform their specific role

### Differential Gene Expression - Human Cells can express about 20% of it's protein coded genes at any given time - Most cells contain the same genome - Each cell type must use specific parts of this genome - This is called Differential gene expression - Exception would be cells of the immune system - Due to the importance of gene expression when it has issues it can affect the organism significantly - Process of Gene expression in a Eukaryotic cell - Chromatin (DNA unpacking) -> - RNA processing -> - Transport to cytoplasm -> - Translation -> - Protein processing -> - Transport to cellular destination-> - This process can often be equated to transcription for Prokaryote cells

## 1.0.2 | Regulation of Chromatin Structure

- The chromatin structure itself allows for the regulation of gene expression
  - This is partially due to the location of the promoter
- · Chemical modifications to the histone proteins can affect the structure
  - · This in turn can affect gene expression
  - · Histone proteins are the proteins in which the DNA is wrapped
  - There are many types of modifications that can take place
    - · Histone acetylation can tend to promote transcriptions by opening up the chromatin
    - Additional methyl groups tend to close up the chromatin and decrease transcription
  - · DNA methylation occurs in most plants and animals as well as fungi
  - · Methylated DNA will stay methalated through cell divisions
    - · This accounts for genomic imprinting
    - These epigenetic markers can be inherited
      - There is continually more evidence for the importance of epigentics in gene expression ### Regulation of Transcription
- Chromatin changes are not permanent and can be reversed
- The next step of gene expression regulation is in the transcription factors
  - These either allow for or inhibit transcription
- · These factors usually bind to proteins, but some of them bind to DNA
- High levels of transcription factors created for specific genes are associated with another protein thought creatively of as specific transcription factors
- Gene expression is dramatically increased or decreased by the binding of specific transcription factors

- · These are either activators or repressors
- · There are many transcription factors
  - Repressors act in many different ways, but some bind directly to control element DNA blocking activator binding
  - · Others interfere with the activator itself
- Coordinated control of genes can need to happen when multiple genes need to be expressed at the same time for something to function
  - These can often be signaled from the outside with something like a hormone
- The activation of receptors on the surface of the cell can release specific repressors and activators

### Mechanisms of Post-Transcriptional Regulation - Transcription is not the only thing that regulates gene expression - How much of the protein is created once the RNA is received is also a factor - There is also mu