



—title: Nuclear Physics source: [KBPhysicsMasterIndexcourse](#): PHYS201 author: Houjun Liu — # Nuclear Physics First of all, recall [KBhPHYS201CoulombsLaw](#). Given the force between two particles is  $\frac{kQ^2}{R^2}$ , we could hand-wavily calculate the *work* between two particles if we know how much they travel near/far from each other. Through this, we could show that nuclear forces (through nuclear distance, proton=>electron) are much larger than that of the chemical forces (atom/atom, electron=>electron). #compilefromnote Remember:  $A_{nucl} = \frac{1}{10^{10}} A_{atom}$  ## Radioactivity Radiation is the emission of waves — lights, heat, etc. etc. We call something “radioactive” if it emits ionizing radiation: that it has enough energy to liberate an electron from an atom. ### Geiger Counter #insert how geiger counters work Because of the fact that Geiger counters require time to discharge, there is a certain rate called “dead time” during which Geiger counters simply sit and do nothing. As such, we have to account for this lossy “deadtime” of Geiger counters by relating the two values with the following equation  $T = \frac{M}{1-(M/L)}$ , where  $M$  is the measured rate of radiation and  $L$  is the “dead time” — the upper limit of the Geiger counter in question. ### Radio Charge Types-  $\alpha$ : positively charged + relatively massive (low  $\frac{q}{m}$ )-  $\beta$ : negatively charged + relatively high charge (high  $\frac{q}{m}$ )-  $\gamma$ : neutral This could be seen by how these three




types of charge curve into a magnetic field.





Apply right hand rule 1.5. ### Creating a ray “Split a nucleus, somehow” **Alpha Decay:**

alpha decay, a massive nucleolus spits out a Helium—resulting part of itself to get rid of 2 protons and 2 neu-




trons. So, formally...

**Gamma Decay** Instead of splitting part of the nucleus, gamma decay spits an electrically excited (so... chemistry, charged, energy level, that stuff) atom




into a normal, non-excited atom and also emits a photon.

now, the most confusing one...**Beta Decay** There's two types of beta decay: "beta-minus" decay and "beta-plus" decay. When folks talk about just "beta-decay", they are talking about beta-minus decay. An el-



ement decays from the parent element into a different nucleus. Beta minus decay

this case, the nucleus gained a proton and lost a neutron. What happened? A neutron in the nucleus turned into a positive proton and a negative electron. The newly-formed electron comes flying out as a "beta-minus" particle. Also, this process creates an "antineutrino", which is a tiny, charge-less element that will



become important later. Beta plus decay

This is the opposite of beta-minus decay. The element takes one of its protons, splits it to a positron (a positive elec-

isolated




$$m_n > m_{p^+}$$




tron, this is antimatter), a neutrino, and a neutron. Wait... But *how*?

minus decay makes sense, because it would be energetically favorable as a neutron is *slightly* more massive and hence will lose some of the mass during beta-minus decay. But, during beta-plus decay, the reac-



tants are less massive than the product (!) — so thermally it won't really work out.




in order to actually create beta-plus decay, you have to shove the protons, antimatter, etc. together really fast with some kinetic energy. Or, this could happen spontaneously as long as the mass of the daughter atom is smaller than the mass of the parent (as in... as long as mass of daughter + mass of anti electron + mass of neutrino < mass of parent, this should work.) To make sense of this, stop thinking that atoms' masses could be deducted by just counting the number of neutrons+protons. Electron cap-



ture\_

this case, you will gain one neutron by simply capturing an electron out of thin air (the electron cloud) and

In

 *$\beta$  decay:*

merge it with a proton with the help of an antineutrino to form.