

Source: [KBe2020math530floIndex](#)

#flo

1 | Polynomials

- See [KBrefPolynomial](#)

1.1 | 0 polynomial

- Has degree $-\infty$
- Degrees are usually positive, except for the 0 degree
- “that’s too hard, and we’re not going to do it here”

1.2 | Identically zero

- Like 0 or $0x^0$
- Most polynomials are sometimes zero, but polynomials that are “identically zero” means that it’s always zero (instead of just sometimes zero)

1.3 | $\mathcal{P}_m(F)$

- Polynomials with coefficients in F whose highest degree is m
- It can’t be “whose degree is exactly m ” because otherwise you won’t have the identity and it won’t be closed under addition (in the case where coefficient sum $a_m + b_m = 0$)

1.3.1 | It’s a finite dimensional vector space

$$\bullet \quad a_0z^0 + \dots + a_mz^m + b_0z^0 + \dots + b_mz^m = (a_0 + b_0)z^0 + \dots + (a_m + b_m)z^m$$

1.4 | Proof of 2.16

- Structure: proof by contradiction

2 | Linear Independence

- “non-trivial” means “simplest possible”, which has usually got the most zeros
- See [KB20math530refLinearIndependence](#)

2.1 | 2.21 Linear Dependence Lemma 2.21

- it’s saying that any linearly independent list has a vector inside that doesn’t “contribute anything”, and that if you remove it you’ll have the same span. Implicitly, maybe through induction?) if you remove a dependent vector enough times then you get a linearly independent list.
- The list $(1, 1, 1), (2, 2, 2), (3, 3, 3)$ is really dependent, but $(0), (0), (0)$ is the most dependent (you have to remove all to get independence).

- how
- to
- pagebreak

3 | Exercise 2.A.1

3.1 | Lemma

If vectors v_1, v_2, v_3, v_4 span V , then the list

$$v_1 - v_2, v_2 - v_3, v_3 - v_4, v_4$$

also spans V .

3.2 | Proof

We prove the lemma by showing that any vector $v \in V$ can be written in the form $a_1 v_1 + a_2 v_2 + a_3 v_3 + a_4 v_4$ can also be written as a linear combination of the form

$$b_1(v_1 - v_2) + b_2(v_2 - v_3) + b_3(v_3 - v_4) + b_4 v_4$$

If we set

$$b_1 = a_1$$

$$b_2 = a_1 + a_2$$

$$b_3 = a_1 + a_2 + a_3$$

$$b_4 = a_1 + a_2 + a_3 + a_4$$

then the two combinations will be equivalent:


$$\begin{aligned} & a_1(v_1 - v_2) + (a_1 + a_2)(v_2 - v_3) + (a_1 + a_2 + a_3)(v_3 - v_4) + (a_1 + a_2 + a_3 + a_4)v_4 \\ &= \cancel{a_1 v_1 - a_1 v_2 + a_1 v_2} + \cancel{a_2 v_2 - (a_1 + a_2)v_3 + (a_1 + a_2)v_3} + \cancel{a_3 v_3 - (a_1 + a_2 + a_3)v_4 + (a_1 + a_2 + a_3)v_4} \\ &= a_1 v_1 + a_2 v_2 + a_3 v_3 + a_4 v_4 \end{aligned}$$

4 | Rock

And here's a rock for good measure:



Look at these colors!



```
4 BUILDNUMBER_FILE='buildID.txt'
5
6 while true; do
7     printf "working..." \r
8     echo "\n\nLog for attempt at $(date)" >> log.txt
9     git pull >> log.txt
10
11     changes="$(git status --porcelain | cut -c4-)"
12
13     if [[ -n "$changes" ]]; then
14         #find . -type f -name '*.md' -exec sed -i'' -e 's/\Users/houliu/Documents/School Work/2020-2021/KnowledgeBase/./g' {} +
15
16         ./compile.sh
17
18         #if make >> log.txt 2>>recent_errors.txt; then
19         # [[ -f $BUILDNUMBER_FILE ]] || echo 0 > $BUILDNUMBER_FILE;
20         # echo $(( $(cat $BUILDNUMBER_FILE) + 1 )) > $BUILDNUMBER_FILE
21         #fi
22         [[ -f $BUILDNUMBER_FILE ]] || echo 0 > $BUILDNUMBER_FILE;
23         echo $(( $(cat $BUILDNUMBER_FILE) + 1 )) > $BUILDNUMBER_FILE
24
25         (git add -A >> log.txt && \
26         git reset -- log.txt >> log.txt && \
27         git commit -m "b$(cat buildID.txt): $(echo $changes | paste -sd ' ' -) $(printf "\n\n")automated compilation $(cat buildID.txt))" >> log.txt && \
28         git push --quiet && \
29         printf "Snapshoted and committed at $(date)\r" || printf "Sync Failed!!! $(date)"
30     else
31         printf "Nothing to sync at $(date)" \r
32     fi
33
34     sleep 30
35 done
36
```

