

Source: [KBiologyMasterIndex](#)

0.1 | #flo #ret

1 | Notes

1.0.1 | Overview

- Organisms turn genes on and off which is called Gene Expression
 - This can be done in response to external and internal signals
 - These signals are based off of environmental factors
 - This is also be done in order to specialize cells
 - Certain cells need certain genes to preform their specific role

Differential Gene Expression - Human Cells can express about 20% of it's protein coded genes at any given time - Most cells contain the same genome - Each cell type must use specific parts of this genome - This is called Differential gene expression - Exception would be cells of the immune system - Due to the importance of gene expression when it has issues it can affect the organism significantly - Process of Gene expression in a Eukaryotic cell - Chromatin (DNA unpacking) -> - RNA processing -> - Transport to cytoplasm -> - Translation -> - Protein processing -> - Transport to cellular destination-> - This process can often be equated to transcription for Prokaryote cells

1.0.2 | Regulation of Chromatin Structure

- The chromatin structure itself allows for the regulation of gene expression
 - This is partially due to the location of the promoter
- Chemical modifications to the histone proteins can affect the structure
 - This in turn can affect gene expression
 - Histone proteins are the proteins in which the DNA is wrapped
 - There are many types of modifications that can take place
 - Histone acetylation can tend to promote transcriptions by opening up the chromatin
 - Additional methyl groups tend to close up the chromatin and decrease transcription
 - DNA methylation occurs in most plants and animals as well as fungi
 - Methylated DNA will stay methalated through cell divisions
 - This accounts for genomic imprinting
 - These epigenetic markers can be inherited
 - There is continually more evidence for the importance of epigenetics in gene expression

Regulation of Transcription

- Chromatin changes are not permanent and can be reversed
- The next step of gene expression regulation is in the transcription factors
 - These either allow for or inhibit transcription
- These factors usually bind to proteins, but some of them bind to DNA
- High levels of transcription factors created for specific genes are associated with another protein thought creatively of as specific transcription factors
- Gene expression is dramatically increased or decreased by the binding of specific transcription factors

- These are either activators or repressors
- There are many transcription factors
 - Repressors act in many different ways, but some bind directly to control element DNA blocking activator binding
 - Others interfere with the activator itself

Coordinately Controlled Genes in Eukaryotes - Coordinated control of genes can need to happen when multiple genes need to be expressed at the same time for something to function - These can often be signaled from the outside with something like a hormone - The activation of receptors on the surface of the cell can release specific repressors and activators -