Axler 6.A exercise 9 April 27, 2021

## 1 | Axler 6.A exercise 9

Suppose  $u, v \in V$  and  $||u|| \le 1$  and  $||v|| \le 1$ . Prove that

$$\sqrt{1 - \|u\|^2} \sqrt{1 - \|u\|^2} \le 1 - |\langle u, v \rangle|$$

## 2 | **Proof**

We will prove this by showing that the left hand side is less than or equal to an intermediate term, which is less than or equal to the right hand side.

$$|\langle u, v \rangle| \le ||u|| ||v||$$
$$1 - ||u|| ||v|| \le 1 - |\langle u, v \rangle|$$

This intermediate value is obtained using the Cauchy-Schwarz inequality.

Now, to show that the square of the left hand side is less than or equal to the square of the right hand side,

$$\begin{split} &= (1 - \|u\|^2)(1 - \|v\|^2) \\ &= 1 - \|u\|^2 - \|v\|^2 + \|u\|^2\|v\|^2 \\ &= 1 - \|u\|^2 - \|v\|^2 + \|u\|^2\|v\|^2 \\ &= 1 - \|u\|^2 - \|v\|^2 + \|u\|^2\|v\|^2 \\ &= 1 - 2\|u\|\|v\| + \|u\|^2\|v\|^2 \end{split}$$

Taproot · 2020-2021 Page 1 of 1