$$1 \mid \int \frac{\sqrt{x-1}}{x} dx$$

Let
$$u = \sqrt{x-1}$$
, $du = \frac{1}{2\sqrt{x-1}}dx$

$$\int \frac{\sqrt{x-1}}{x} dx = \int \frac{u}{u^2+1} 2u du$$
$$= 2 \int \frac{(u^2+1)-1}{u^2+1} du$$
$$= 2 \int \frac{-1}{1} du$$

- 2 | 2
- 3 | 3
- 4 | 4
- 5 | 5

$$6 \mid \int \tan^2 x + 1 dx$$

$$\int \tan^2 x + 1 dx = \int \sec^2 x - 1 + 1 dx$$

$$= \int \sec^2 x dx$$
Let $u = x, du = 1$

$$= \int \sec^2 u du$$

$$= \tan u + C$$

$$= \boxed{\tan x + C}$$

- 7 | 7
- 8 | 8
- $9 \mid \int \frac{\sec^2 x}{\csc x} sinx dx$

$$\int \frac{\sec^2 x}{\csc x} \sin x dx = \int \tan^2 x dx$$
$$= \int \sec^2 x - 1 dx$$
$$= \int \sec^2 x dx - \int 1 dx$$
$$= \tan x - x$$

10 | 10

11 | TODO
$$\int \frac{e^{2\ln\sin x} + e^{2\ln\cos x}}{e^{2\ln\tan x} + e^{2\ln 1}} dx$$

$$\int \frac{e^{2\ln\sin x} + e^{2\ln\cos x}}{e^{2\ln\tan x} + e^{2\ln 1}} dx = \int \frac{\sin^2 x + \cos^2 x}{\tan^2 x + 1} dx$$
$$= \int \frac{1}{\tan^2 x + 1} dx$$
$$= \int \frac{1}{\sec^2 x} dx$$
$$= \int \cos^2 x dx$$

- 12 | **12**
- 13 | 13
- 14 | **14**