


Function	1 st four terms of expansion	General Term/Sigma Notation
$\frac{1}{1-2x}$		
	$1 - \frac{9x^2}{2!} + \frac{81x^4}{4!} - \frac{729x^6}{6!} + \dots$	
		$\sum_{n=0}^{\infty} \frac{x^n}{e^{2n!}}$
	$x^2 - \frac{x^6}{3!} + \frac{x^{10}}{5!} - \frac{x^{14}}{7!} + \dots$	
$\frac{x}{1+x^4}$		
	$1 + (x-1)^2 + \frac{(x-1)^4}{2} + \frac{(x-1)^6}{6} + \dots$	
$\frac{\cos(x) - 1}{x^2}$		
$2x \ln(1+2x)$		
	$2x - 2x^3 + 2x^5 - 2x^7 + \dots$	

Lagrange Error in Taylor Polynomials= remainder of a Taylor Polynomial



Error = $|R_n(x)| = |f(x) - P_n(x)|$.

THEOREM 9.19 Taylor's Theorem

If a function f is differentiable through order $n + 1$ in an interval I containing c , then, for each x in I , there exists z between x and c such that

$$f(x) = f(c) + f'(c)(x - c) + \frac{f''(c)}{2!}(x - c)^2 + \cdots + \frac{f^{(n)}(c)}{n!}(x - c)^n + R_n(x)$$

where

$$R_n(x) = \frac{f^{(n+1)}(z)}{(n+1)!}(x - c)^{n+1}.$$

A proof of this theorem is given in Appendix A.

See LarsonCalculus.com for Bruce Edwards's video of this proof.

6. The Maclaurin series for the function f is given by

$$f(x) = \sum_{n=0}^{\infty} \frac{(2x)^{n+1}}{n+1} = 2x + \frac{4x^2}{2} + \frac{8x^3}{3} + \frac{16x^4}{4} + \cdots + \frac{(2x)^{n+1}}{n+1} + \cdots$$

on its interval of convergence.

- (a) Find the interval of convergence of the Maclaurin series for f . Justify your answer.
- (b) Find the first four terms and the general term for the Maclaurin series for $f'(x)$.
- (c) Use the Maclaurin series you found in part (b) to find the value of $f'\left(-\frac{1}{3}\right)$.

6. The function f is defined by the power series

$$f(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n+1)!} = 1 - \frac{x^2}{3!} + \frac{x^4}{5!} - \frac{x^6}{7!} + \cdots + \frac{(-1)^n x^{2n}}{(2n+1)!} + \cdots$$

for all real numbers x .

- (a) Find $f'(0)$ and $f''(0)$. Determine whether f has a local maximum, a local minimum, or neither at $x = 0$.
Give a reason for your answer.
- (b) Show that $1 - \frac{1}{3!}$ approximates $f(1)$ with error less than $\frac{1}{100}$.
- (c) Show that $y = f(x)$ is a solution to the differential equation $xy' + y = \cos x$.

6. Let f be the function given by $f(x) = \sin\left(5x + \frac{\pi}{4}\right)$, and let $P(x)$ be the third-degree Taylor polynomial for f about $x = 0$.
- (a) Find $P(x)$.
 - (b) Find the coefficient of x^{22} in the Taylor series for f about $x = 0$.
 - (c) Use the Lagrange error bound to show that $\left|f\left(\frac{1}{10}\right) - P\left(\frac{1}{10}\right)\right| < \frac{1}{100}$.
 - (d) Let G be the function given by $G(x) = \int_0^x f(t) dt$. Write the third-degree Taylor polynomial for G about $x = 0$.

6. Let f be a function with derivatives of all orders and for which $f(2) = 7$. When n is odd, the n th derivative of f at $x = 2$ is 0. When n is even and $n \geq 2$, the n th derivative of f at $x = 2$ is given by $f^{(n)}(2) = \frac{(n-1)!}{3^n}$.
- (a) Write the sixth-degree Taylor polynomial for f about $x = 2$.
 - (b) In the Taylor series for f about $x = 2$, what is the coefficient of $(x - 2)^{2n}$ for $n \geq 1$?
 - (c) Find the interval of convergence of the Taylor series for f about $x = 2$. Show the work that leads to your answer.

6. The function f is defined by the power series

$$f(x) = -\frac{x}{2} + \frac{2x^2}{3} - \frac{3x^3}{4} + \cdots + \frac{(-1)^n nx^n}{n+1} + \cdots$$

for all real numbers x for which the series converges. The function g is defined by the power series

$$g(x) = 1 - \frac{x}{2!} + \frac{x^2}{4!} - \frac{x^3}{6!} + \cdots + \frac{(-1)^n x^n}{(2n)!} + \cdots$$

for all real numbers x for which the series converges.

- (a) Find the interval of convergence of the power series for f . Justify your answer.
- (b) The graph of $y = f(x) - g(x)$ passes through the point $(0, -1)$. Find $y'(0)$ and $y''(0)$. Determine whether y has a relative minimum, a relative maximum, or neither at $x = 0$. Give a reason for your answer.

$$f(x) = \begin{cases} \frac{\cos x - 1}{x^2} & \text{for } x \neq 0 \\ -\frac{1}{2} & \text{for } x = 0 \end{cases}$$

6. The function f , defined above, has derivatives of all orders. Let g be the function defined by

$$g(x) = 1 + \int_0^x f(t) dt.$$

- (a) Write the first three nonzero terms and the general term of the Taylor series for $\cos x$ about $x = 0$. Use this series to write the first three nonzero terms and the general term of the Taylor series for f about $x = 0$.
- (b) Use the Taylor series for f about $x = 0$ found in part (a) to determine whether f has a relative maximum, relative minimum, or neither at $x = 0$. Give a reason for your answer.
- (c) Write the fifth-degree Taylor polynomial for g about $x = 0$.
- (d) The Taylor series for g about $x = 0$, evaluated at $x = 1$, is an alternating series with individual terms that decrease in absolute value to 0. Use the third-degree Taylor polynomial for g about $x = 0$ to estimate the value of $g(1)$. Explain why this estimate differs from the actual value of $g(1)$ by less than $\frac{1}{6!}$.

6. The function g has derivatives of all orders, and the Maclaurin series for g is

$$\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+3} = \frac{x}{3} - \frac{x^3}{5} + \frac{x^5}{7} - \dots.$$

- (a) Using the ratio test, determine the interval of convergence of the Maclaurin series for g .
- (b) The Maclaurin series for g evaluated at $x = \frac{1}{2}$ is an alternating series whose terms decrease in absolute value to 0. The approximation for $g\left(\frac{1}{2}\right)$ using the first two nonzero terms of this series is $\frac{17}{120}$. Show that this approximation differs from $g\left(\frac{1}{2}\right)$ by less than $\frac{1}{200}$.
- (c) Write the first three nonzero terms and the general term of the Maclaurin series for $g'(x)$.