Page 1

Source:

1 | linear approximations

1.1 | cube root

1.1.1 | approximation

$$(1+x)^{\frac{1}{3}} \to \frac{1}{3}(1+x)^{\frac{-2}{3}}$$

at x = 0 is

$$\frac{1}{3}(1+0)^{...} = \frac{1}{3}$$

so the linear approximation is

$$y \approx m(x-0) + f(0) = \frac{1}{3}x + 1$$

1.1.2 | estimations

value	estimate
0.05	1.016666
-0.25	0.916666

These will be overestimates because the graph is concave down in this reigon.

1.2 | sin(x)

1.2.1 | approximation

$$y \approx \frac{d}{dx}\sin x\Big|_{0}(x-0) + \sin 0 = x$$

1.2.2 | estimates

value	estimate
-0.1	-0.1
0.1	0.1

The first estimate will be an underestimate because $\sin x$ is concave up in that reigon. The opposite is true for the second estimate.

Exr0n · **2020-2021**

1.3 unknown function (only some points known

1.3.1 | approximation

$$y \approx \frac{d}{dx} f(x) \Big|_{c} (x - c) + f(c)$$

plugging in c = 1,

$$y \approx 5(x-1) - 4$$

1.3.2 | estimations

value	estimate
1.2	-3

This will be an underestimate because the second derivative is positive and the graph is thus concave up.

2 | differentials

For a function y = f(x), dy and dx are differentials and the relationship is dy = f'(x)dx.

For a function written f(x) = (something), the differentials are df and dx and the relationship is the same: df = f'(x)dx.

2.1 | cube error

2.1.1 | differential

$$df = f'(x)dx$$
$$= 3x^2 dx$$

2.1.2 | volume error

If I understand the use of differentials corretly, then x is the measured value (2) and dx is the uncertainty (delta x), or 0.2ft. Then, the change in the volume (change in fuction or df) would be $3(2)^2(0.2)=2.4$

2.1.3 | max error for some ϵ

$$df = 3x^{2}dx$$

$$dx = \frac{df}{3x^{2}}$$

$$= \frac{0.1}{3(2)^{2}}$$

$$= 8.\overline{3}$$

2.2 | sphere measuring

$$f(r) = 4\pi r^{2}$$

$$\frac{d}{dr}f(r) = 8\pi r$$

$$df = 8\pi r (dr)$$

$$= 8\pi 21(0.05) \approx$$

Exr0n · 2020-2021 Page 3