Axler 6.A exercise 9 April 27, 2021

1 | Axler 6.A exercise 9


Suppose $u,v\in V$ and $\|u\|\leq 1$ and $\|v\|\leq 1$. Prove that

$$\sqrt{1 - \|u\|^2} \sqrt{1 - \|u\|^2} \le 1 - |\langle u, v \rangle|$$

2 | **Proof**

2.1 | Useful Lemma

$$||u||^2 + ||v||^2 \ge 2||u||||v||$$



This proof is only valid for inner product spaces over \mathbb{F}^n and the Euclidean norm. An algebraic proof would be better.

2.2 | Cauchy-Schwarz Corollary

$$|\langle u, v \rangle| \le ||u|| ||v||$$
$$1 - ||u|| ||v|| \le 1 - |\langle u, v \rangle|$$

This intermediate value is obtained using the Cauchy-Schwarz inequality.

2.3 | Main Proof

Now, to show that the square of the left hand side is less than or equal to the square of the right hand side,

$$\begin{split} (1-\|u\|^2)(1-\|v\|^2) = &1-\|u\|^2-\|v\|^2+\|u\|^2\|v\|^2\\ = &1-(\|u\|^2+\|v\|^2)+\|u\|^2\|v\|^2\\ \leq &1-2\|u\|\|v\|+\|u\|^2\|v\|^2 \qquad \text{by the earlier lemma}\\ = &(1-\|u\|\|v\|)^2\\ \leq &(1-|\langle u,v\rangle|)^2 \qquad \text{by the Cauchy-Schwarz corollary} \end{split}$$

Taproot · 2020-2021 Page 1 of 2

Axler 6.A exercise 9 April 27, 2021

Taking square roots of both sides proves the desired result.

Taproot · **2020-2021** Page 2 of 2