

# CS172: COMPUTER SYSTEMS II

## Lecture 17

# Natural Deduction

*- rules for or, not*

James Power



# Proofs using 'or'

The rules for dealing with disjunction (i.e. 'or'):

- Introduction:

To prove  $A \vee B$ :

you must prove  $A$ , or you must prove  $B$  (your choice)

# Proofs using 'or'

The rules for dealing with disjunction (i.e. 'or'):

- Introduction:

To prove  $A \vee B$ :

you must prove  $A$ , or you must prove  $B$  (your choice)

- Elimination:

If you know  $A \vee B$ :

Unfortunately you don't know *which* one of them is true,  
so you must prove the theorem both

- for the case where  $A$  is true
- and also for the case where  $B$  is true

(Proof by cases).

# Formal proof rules for 'or'

$$\frac{A}{A \vee B} \vee_{I1} \quad \frac{B}{A \vee B} \vee_{I2}$$

$$\frac{A \vee B \quad \begin{array}{|l} \text{Suppose } A : \\ \vdots \\ C \end{array} \quad \begin{array}{|l} \text{Suppose } B : \\ \vdots \\ C \end{array}}{C} \vee_E$$

## Example of a maths proof using 'or'

### Theorem

*If  $A \subseteq C$  and  $B \subseteq C$  then  $(A \cup B) \subseteq C$*

## Example of a maths proof using 'or'

### Theorem

If  $A \subseteq C$  and  $B \subseteq C$  then  $(A \cup B) \subseteq C$

### Proof.

Suppose that  $A \subseteq C$  and  $B \subseteq C$ ,  
and suppose also that  $x$  is some element of  $(A \cup B)$ .

Thus we know that  $x \in A$  or  $x \in B$ .

Case 1: If  $x \in A$ , then since  $A \subseteq C$  we know  $x \in C$ .

Case 2: If  $x \in B$ , then since  $B \subseteq C$  we know  $x \in C$ .

Since we know that at least one case is true, we conclude  $x \in C$ .

But  $x$  was an *arbitrary* element of  $(A \cup B)$ , so we conclude that  
 $(A \cup B) \subseteq C$ . □

## A more formal version of that proof

### Theorem

*If  $A \subseteq C$  and  $B \subseteq C$  then  $(A \cup B) \subseteq C$*

## A more formal version of that proof

### Theorem

*If  $A \subseteq C$  and  $B \subseteq C$  then  $(A \cup B) \subseteq C$*

$$\text{11 } (A \subseteq C) \wedge (B \subseteq C) \rightarrow ((A \cup B) \subseteq C)$$



## A more formal version of that proof

### Theorem

*If  $A \subseteq C$  and  $B \subseteq C$  then  $(A \cup B) \subseteq C$*

① Suppose  $A \subseteq C$  and  $B \subseteq C$

⑩  $(A \cup B) \subseteq C$

⑪  $(A \subseteq C) \wedge (B \subseteq C) \rightarrow ((A \cup B) \subseteq C)$

$\rightarrow_I$ , lines 1-10

# A more formal version of that proof

## Theorem

*If  $A \subseteq C$  and  $B \subseteq C$  then  $(A \cup B) \subseteq C$*

① Suppose  $(x \in A) \rightarrow (x \in C)$  and  $(x \in B) \rightarrow (x \in C)$  :

⑩  $(x \in (A \cup B)) \rightarrow (x \in C)$

⑪  $(A \subseteq C) \wedge (B \subseteq C) \rightarrow ((A \cup B) \subseteq C)$

$\rightarrow_I$ , lines 1-10

# A more formal version of that proof

## Theorem

If  $A \subseteq C$  and  $B \subseteq C$  then  $(A \cup B) \subseteq C$

- 1 Suppose  $(x \in A) \rightarrow (x \in C)$  and  $(x \in B) \rightarrow (x \in C)$  :
- 2     Suppose  $x \in (A \cup B)$  :

9           Thus  $x \in C$

10           $(x \in (A \cup B)) \rightarrow (x \in C)$   $\rightarrow_I$ , lines 2-9

11          $(A \subseteq C) \wedge (B \subseteq C) \rightarrow ((A \cup B) \subseteq C)$   $\rightarrow_I$ , lines 1-10

# A more formal version of that proof

## Theorem

If  $A \subseteq C$  and  $B \subseteq C$  then  $(A \cup B) \subseteq C$

- ① Suppose  $(x \in A) \rightarrow (x \in C)$  and  $(x \in B) \rightarrow (x \in C)$  :
- ②     Suppose  $(x \in A) \vee (x \in B)$  :

⑨           Thus  $x \in C$

⑩      $(x \in (A \cup B)) \rightarrow (x \in C)$   $\rightarrow_I$ , lines 2-9

⑪    $(A \subseteq C) \wedge (B \subseteq C) \rightarrow ((A \cup B) \subseteq C)$   $\rightarrow_I$ , lines 1-10

# A more formal version of that proof

## Theorem

If  $A \subseteq C$  and  $B \subseteq C$  then  $(A \cup B) \subseteq C$

① Suppose  $(x \in A) \rightarrow (x \in C)$  and  $(x \in B) \rightarrow (x \in C)$  :

②       Suppose  $(x \in A) \vee (x \in B)$  :

③           Suppose  $(x \in A)$  :

⑥           Suppose  $(x \in B)$  :

⑨           Thus  $x \in C$

$\vee_{\mathcal{E}}$ , lines 2,3-5,6-8

⑩        $(x \in (A \cup B)) \rightarrow (x \in C)$

$\rightarrow_{\mathcal{I}}$ , lines 2-9

⑪    $(A \subseteq C) \wedge (B \subseteq C) \rightarrow ((A \cup B) \subseteq C)$

$\rightarrow_{\mathcal{I}}$ , lines 1-10

# A more formal version of that proof

## Theorem

If  $A \subseteq C$  and  $B \subseteq C$  then  $(A \cup B) \subseteq C$

- 1 Suppose  $(x \in A) \rightarrow (x \in C)$  and  $(x \in B) \rightarrow (x \in C)$  :
- 2     Suppose  $(x \in A) \vee (x \in B)$  :
- 3         Suppose  $(x \in A)$  :
- 4             But  $(x \in A) \rightarrow (x \in C)$   $\wedge_{\mathcal{E}1}$ , line 1
- 5             So  $(x \in C)$   $\rightarrow_{\mathcal{E}}$ , lines 3,4
- 6         Suppose  $(x \in B)$  :
- 7             But  $(x \in B) \rightarrow (x \in C)$   $\wedge_{\mathcal{E}2}$ , line 1
- 8             So  $(x \in C)$   $\rightarrow_{\mathcal{E}}$ , lines 6,7
- 9         Thus  $x \in C$   $\vee_{\mathcal{E}}$ , lines 2,3-5,6-8
- 10      $(x \in (A \cup B)) \rightarrow (x \in C)$   $\rightarrow_{\mathcal{I}}$ , lines 2-9
- 11  $(A \subseteq C) \wedge (B \subseteq C) \rightarrow ((A \cup B) \subseteq C)$   $\rightarrow_{\mathcal{I}}$ , lines 1-10

## Duality between 'and' and 'or'

Note that the introduction/elimination rules for 'and' and 'or' seem to *reflect* each other:

## Duality between 'and' and 'or'

Note that the introduction/elimination rules for 'and' and 'or' seem to *reflect* each other:

- The introduction rule for 'or' and the elimination rule for 'and' give you the choice (which to prove/use).
- The elimination rule for 'or' and the introduction rule for 'and' insist you deal with both possibilities.



## Duality between 'and' and 'or'

Note that the introduction/elimination rules for 'and' and 'or' seem to *reflect* each other:

- The introduction rule for 'or' and the elimination rule for 'and' give you the choice (which to prove/use).
- The elimination rule for 'or' and the introduction rule for 'and' inist you deal with both possibilities.
- ... a little bit similar to the way they interact in de Morgan's laws

## Duality between 'and' and 'or'


Note that the introduction/elimination rules for 'and' and 'or' seem to *reflect* each other:

- The introduction rule for 'or' and the elimination rule for 'and' give you the choice (which to prove/use).
- The elimination rule for 'or' and the introduction rule for 'and' inist you deal with both possibilities.
- ... a little bit similar to the way they interact in de Morgan's laws

The connectives 'and' and 'or' are not exactly logical opposites: instead we say that are **dual** to each other.

Other examples of dual concepts:

- for-all and there-exists
- satisfiability and validity
- set theory: union and intersection
- platonic solids: cube and octahedron



# Negation

- Dealing with negation is a little awkward...
  - To assert “ $\neg A$ ” must be the same as saying “ $A$  is false”, but we only have rules to prove things that are *true*.

# Negation

- Dealing with negation is a little awkward...
  - To assert “ $\neg A$ ” must be the same as saying “ $A$  is false”, but we only have rules to prove things that are *true*.
- The usual approach is to work by means of a **contradiction**:
  - if we assume  $A$  and derive a contradiction,  
then it must have been wrong to assume  $A$ ,  
so deduce  $\neg A$
- We use the symbol  $\perp$  to denote “contradiction”.

# Formal proof rule for 'not'

For the moment we will adopt just an *introduction rule* for negation:

$$\frac{\boxed{\begin{array}{l} \text{Suppose } A : \\ \vdots \\ \perp \end{array}}}{\neg A} \neg I$$

- To prove  $\neg A$ :

Assume  $A$ , and then derive a contradiction.

# Formal proof rule for 'not'

For the moment we will adopt just an *introduction rule* for negation:

$$\frac{\boxed{\begin{array}{l} \text{Suppose } A : \\ \vdots \\ \perp \end{array}}}{\neg A} \neg I$$

- To prove  $\neg A$ :  
Assume  $A$ , and then derive a contradiction.
- So how do we derive a contradiction?

# Formal proof rules for 'contradiction'

$$\frac{B \quad \neg B}{\perp} \perp_{\mathcal{I}}$$

$$\frac{\perp}{B} \perp_{\mathcal{E}}$$

Explanation:

- Introduction:

You get a contradiction by  
proving both  $B$  and  $\neg B$ , for some formula  $B$ .

- Elimination:

If you have a contradiction in your proof,  
then anything can be deduced.

# Formal proof rules for ‘contradiction’

$$\frac{B \quad \neg B}{\perp} \perp_{\mathcal{I}}$$

$$\frac{\perp}{B} \perp_{\mathcal{E}}$$

## Notes

- Introduction:

The introduction rule for contradiction is also a sort of *elimination rule* for negation:

If you have a premise of the form  $\neg B$ ,  
try to prove  $B$ , and then deduce a contradiction.

- Elimination:

It's rarely useful to deduce just “anything” from  $\perp$ .

Typically this rule is used to tidy up impossible cases in a proof.



# A formal(-ish) example of a proof using negation

## Theorem

*If  $x^2 + y = 13$  and  $y \neq 4$  then  $x \neq 3$*

# A formal(-ish) example of a proof using negation

## Theorem

*If  $x^2 + y = 13$  and  $y \neq 4$  then  $x \neq 3$*

## Proof.

⑨  $(x^2 + y = 13) \wedge (y \neq 4) \rightarrow (x \neq 3)$

# A formal(-ish) example of a proof using negation

## Theorem

*If  $x^2 + y = 13$  and  $y \neq 4$  then  $x \neq 3$*

## Proof.

9  $(x^2 + y = 13) \wedge \neg(y = 4) \rightarrow \neg(x = 3)$

# A formal(-ish) example of a proof using negation

## Theorem

*If  $x^2 + y = 13$  and  $y \neq 4$  then  $x \neq 3$*

## Proof.

① Suppose  $(x^2 + y = 13) \wedge \neg(y = 4)$  :

⑧  $\neg(x = 3)$

⑨  $(x^2 + y = 13) \wedge \neg(y = 4) \rightarrow \neg(x = 3)$   $\rightarrow_I$ , lines 1-8

# A formal(-ish) example of a proof using negation

## Theorem

If  $x^2 + y = 13$  and  $y \neq 4$  then  $x \neq 3$

## Proof.

① Suppose  $(x^2 + y = 13) \wedge \neg(y = 4)$  :

② Suppose  $x = 3$  :

⑦  $\perp$

⑧  $\neg(x = 3)$

$\neg_I$ , lines 2-7

⑨  $(x^2 + y = 13) \wedge \neg(y = 4) \rightarrow \neg(x = 3)$

$\rightarrow_I$ , lines 1-8

# A formal(-ish) example of a proof using negation

## Theorem

If  $x^2 + y = 13$  and  $y \neq 4$  then  $x \neq 3$

## Proof.

- ① Suppose  $(x^2 + y = 13) \wedge \neg(y = 4)$  :
- ②     Suppose  $x = 3$  :
- ③          $x^2 + y = 13$   $\wedge_{\mathcal{E}1}$ , line 1
- ⑦          $\perp$
- ⑧          $\neg(x = 3)$   $\neg_{\mathcal{I}}$ , lines 2-7
- ⑨  $(x^2 + y = 13) \wedge \neg(y = 4) \rightarrow \neg(x = 3)$   $\rightarrow_{\mathcal{I}}$ , lines 1-8

# A formal(-ish) example of a proof using negation

## Theorem

*If  $x^2 + y = 13$  and  $y \neq 4$  then  $x \neq 3$*

## Proof.

- ① Suppose  $(x^2 + y = 13) \wedge \neg(y = 4)$  :
- ②     Suppose  $x = 3$  :
- ③          $x^2 + y = 13$   $\wedge_{\mathcal{E}1}$ , line 1
- ④          $3^2 + y = 13$  “substitution”, lines 2,3
- ⑦          $\perp$
- ⑧          $\neg(x = 3)$   $\neg_{\mathcal{I}}$ , lines 2-7
- ⑨  $(x^2 + y = 13) \wedge \neg(y = 4) \rightarrow \neg(x = 3)$   $\rightarrow_{\mathcal{I}}$ , lines 1-8

# A formal(-ish) example of a proof using negation

## Theorem

If  $x^2 + y = 13$  and  $y \neq 4$  then  $x \neq 3$

## Proof.

- ① Suppose  $(x^2 + y = 13) \wedge \neg(y = 4)$  :
- ②     Suppose  $x = 3$  :
- ③          $x^2 + y = 13$   $\wedge_{\mathcal{E}1}$ , line 1
- ④          $3^2 + y = 13$  “substitution”, lines 2,3
- ⑤          $y = 4$  “arithmetic”, line 4
- ⑦          $\perp$
- ⑧          $\neg(x = 3)$   $\neg_{\mathcal{I}}$ , lines 2-7
- ⑨  $(x^2 + y = 13) \wedge \neg(y = 4) \rightarrow \neg(x = 3)$   $\rightarrow_{\mathcal{I}}$ , lines 1-8



# A formal(-ish) example of a proof using negation

## Theorem

If  $x^2 + y = 13$  and  $y \neq 4$  then  $x \neq 3$

## Proof.

- ① Suppose  $(x^2 + y = 13) \wedge \neg(y = 4)$  :
- ②     Suppose  $x = 3$  :
- ③          $x^2 + y = 13$   $\wedge_{\mathcal{E}1}$ , line 1
- ④          $3^2 + y = 13$  “substitution”, lines 2,3
- ⑤          $y = 4$  “arithmetic”, line 4
- ⑥          $\neg(y = 4)$   $\wedge_{\mathcal{E}2}$ , line 1
- ⑦          $\perp$
- ⑧      $\neg(x = 3)$   $\neg_{\mathcal{I}}$ , lines 2-7
- ⑨  $(x^2 + y = 13) \wedge \neg(y = 4) \rightarrow \neg(x = 3)$   $\rightarrow_{\mathcal{I}}$ , lines 1-8

# A formal(-ish) example of a proof using negation

## Theorem

*If  $x^2 + y = 13$  and  $y \neq 4$  then  $x \neq 3$*

## Proof.

- ① Suppose  $(x^2 + y = 13) \wedge \neg(y = 4)$  :
- ②     Suppose  $x = 3$  :
- ③          $x^2 + y = 13$   $\wedge_{\mathcal{E}1}$ , line 1
- ④          $3^2 + y = 13$  “substitution”, lines 2,3
- ⑤          $y = 4$  “arithmetic”, line 4
- ⑥          $\neg(y = 4)$   $\wedge_{\mathcal{E}2}$ , line 1
- ⑦          $\perp$   $\perp_{\mathcal{I}}$ , lines 5,6
- ⑧      $\neg(x = 3)$   $\neg_{\mathcal{I}}$ , lines 2-7
- ⑨  $(x^2 + y = 13) \wedge \neg(y = 4) \rightarrow \neg(x = 3)$   $\rightarrow_{\mathcal{I}}$ , lines 1-8

## A less formal example of a proof using negation

### Theorem

*The square root of 2 is irrational.*

## A less formal example of a proof using negation

### Theorem

*The square root of 2 is irrational.*

### Proof.

- So  $\sqrt{2}$  is not rational.

# A less formal example of a proof using negation

## Theorem

*The square root of 2 is irrational.*

## Proof.

- Suppose that  $\sqrt{2}$  is rational.

- Contradiction!
- So  $\sqrt{2}$  is not rational.

# A less formal example of a proof using negation

## Theorem

*The square root of 2 is irrational.*

## Proof.

- Suppose that  $\sqrt{2}$  is rational.
- Then it can be written as  $a/b$  for some integers  $a$  and  $b$ ,  
and we can assume that  $a$  and  $b$  have no common factors.
- But if  $\sqrt{2} = a/b$ , then  $a^2 = 2b^2$ .
- Therefore  $a^2$  must be even. Therefore  $a$  must be even.
- This means that  $b$  must be odd.
- However if  $a$  is even, then  $a^2$  must be a multiple of 4.
- Since  $a^2 = 2b^2$ ,  $2b^2$  is also a multiple of 4.
- Therefore  $b^2$  is even, and thus  $b$  is even.
- Contradiction!
- So  $\sqrt{2}$  is not rational.

# A less formal example of a proof using negation

## Theorem

*The square root of 2 is irrational.*

## Proof.

- Suppose that  $\sqrt{2}$  is rational.
- Then it can be written as  $a/b$  for some integers  $a$  and  $b$ ,  
and we can assume that  $a$  and  $b$  have no common factors.
- But if  $\sqrt{2} = a/b$ , then  $a^2 = 2b^2$ .
- Therefore  $a^2$  must be even. Therefore  $a$  must be even.
- This means that  $b$  must be odd.
- However if  $a$  is even, then  $a^2$  must be a multiple of 4.
- Since  $a^2 = 2b^2$ ,  $2b^2$  is also a multiple of 4.
- Therefore  $b^2$  is even, and thus  $b$  is even.
- **Contradiction!**
- So  $\sqrt{2}$  is not rational.

## Elimination rule for 'not' (Classical Logic Only)

- We deferred discussing the elimination rule for negation as it is not accepted in all kinds of logic.
- However, it is accepted in *classical logic*.
- Elimination rule:

$$\frac{\neg\neg A}{A} \neg\mathcal{E}$$



## Elimination rule for 'not' (Classical Logic Only)

- We deferred discussing the elimination rule for negation as it is not accepted in all kinds of logic.
- However, it is accepted in *classical logic*.

- Elimination rule:

$$\frac{\neg\neg A}{A} \neg\mathcal{E}$$

- Do you believe this rule is valid?

# Brouwer's Intuitionism



L. E. J. Brouwer  
(1881-1966)

- Developed **intuitionism** (philosophical view of the foundation of mathematics)
- “On the significance of the principle of excluded middle in mathematics, especially in function theory.”, 1923.
- Ideas developed into *constructive logic* by Kolmogorov and Heyting (“BHK”).
- Close links with the *lambda-calculus* (via the Curry-Howard isomorphism)

# Some rules that are equivalent to $\neg_{\mathcal{E}}$

“Law of the Excluded Middle”

$$\frac{\cdot}{A \vee \neg A} \text{LEM}$$

This is a direct consequence of  $\neg_{\mathcal{I}}$  (tricky proof), and is the most famous rule that is rejected by constructive logic.

(Full) “Proof by contradiction”

$$\frac{\boxed{\begin{array}{l} \text{Suppose } \neg A : \\ \vdots \\ \perp \end{array}}}{A} \text{PBC}$$

This *PBC* rule is just  $\neg_{\mathcal{I}}$  followed immediately by  $\neg_{\mathcal{E}}$ .

## Example of a proof using LEM

### Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

## Example of a proof using LEM

### Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

### Proof.

- So  $a^b$  is rational

## Example of a proof using LEM

### Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

### Proof.

- Let  $b = \sqrt{2}$  (so  $b$  is not rational).

- So  $a^b$  is rational

## Example of a proof using LEM

### Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

### Proof.

- Let  $b = \sqrt{2}$  (so  $b$  is not rational).
- Either  $b^b$  is rational or it is not.

*LEM.*

- So  $a^b$  is rational

## Example of a proof using LEM

### Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

### Proof.

- Let  $b = \sqrt{2}$  (so  $b$  is not rational).
- Either  $b^b$  is rational or it is not.
- - Suppose  $b^b$  is rational:

*LEM.*

- - Suppose  $b^b$  is not rational:

- So  $a^b$  is rational either way.

$\forall \mathcal{E}$



## Example of a proof using LEM

### Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

### Proof.

- Let  $b = \sqrt{2}$  (so  $b$  is not rational).
- Either  $b^b$  is rational or it is not.
- - Suppose  $b^b$  is rational:
  - Easy: choose  $a = b$ , (so  $a$  is not rational).
  - and thus  $a^b = b^b$  is rational
- - Suppose  $b^b$  is not rational:

LEM.

- So  $a^b$  is rational either way.

$\forall \mathcal{E}$

# Example of a proof using LEM

## Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

## Proof.

- Let  $b = \sqrt{2}$  (so  $b$  is not rational).
- Either  $b^b$  is rational or it is not. LEM.
- - Suppose  $b^b$  is rational:
  - Easy: choose  $a = b$ , (so  $a$  is not rational).
  - and thus  $a^b = b^b$  is rational
- - Suppose  $b^b$  is not rational:
  - Then choose  $a = b^b = (\sqrt{2}^{\sqrt{2}})$  (so  $a$  is not rational).
- So  $a^b$  is rational either way.  $\forall \mathcal{E}$

# Example of a proof using LEM

## Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

## Proof.

- Let  $b = \sqrt{2}$  (so  $b$  is not rational).
- Either  $b^b$  is rational or it is not. LEM.
- - Suppose  $b^b$  is rational:
  - Easy: choose  $a = b$ , (so  $a$  is not rational).
  - and thus  $a^b = b^b$  is rational
- - Suppose  $b^b$  is not rational:
  - Then choose  $a = b^b = (\sqrt{2}^{\sqrt{2}})$  (so  $a$  is not rational).
  - But now both  $a$  and  $b$  are irrational,
  - and  $a^b = (\sqrt{2}^{\sqrt{2}})^{\sqrt{2}} = \sqrt{2}^{(\sqrt{2}\sqrt{2})} = 2$ , which is rational.
- So  $a^b$  is rational either way.  $\forall \mathcal{E}$

## Example of a proof using LEM

### Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

Q: Is this proof OK?

## Example of a proof using LEM

### Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

Q: Is this proof OK?

A: (classical)

Yes, it follows the rules of natural deduction.

A: (constructive)

No, you haven't given me *actual* irrationals  $a$  and  $b$  such that  $a^b$  is rational.