

# CS172: COMPUTER SYSTEMS II

## Lecture 22

# Cardinality *(Infinite Sets)*

James Power



# Galileo's paradox

Consider the function  $(\lambda x \in \mathbb{N} \cdot x^2)$  over the natural numbers.

# Galileo's paradox

Consider the function  $(\lambda x \in \mathbb{N} \cdot x^2)$  over the natural numbers.

- For every number there is *exactly one* perfect square, and for every perfect square there is *exactly one* number that is its (positive) square root.
- The function forms a **bijection** between the natural numbers,  $\{0, 1, 2, 3, \dots\}$ , and the perfect squares,  $\{0, 1, 4, 9, \dots\}$ .

# Galileo's paradox

Consider the function  $(\lambda x \in \mathbb{N} \cdot x^2)$  over the natural numbers.

- For every number there is *exactly one* perfect square, and for every perfect square there is *exactly one* number that is its (positive) square root.
- The function forms a **bijection** between the natural numbers,  $\{0, 1, 2, 3, \dots\}$ , and the perfect squares,  $\{0, 1, 4, 9, \dots\}$ .
- So the set  $\{0, 1, 2, 3, \dots\}$  must have the *same number of elements* as the set  $\{0, 1, 4, 9, \dots\}$ .

# Galileo's paradox

Consider the function  $(\lambda x \in \mathbb{N} \cdot x^2)$  over the natural numbers.

- For every number there is *exactly one* perfect square, and for every perfect square there is *exactly one* number that is its (positive) square root.
- The function forms a **bijection** between the natural numbers,  $\{0, 1, 2, 3, \dots\}$ , and the perfect squares,  $\{0, 1, 4, 9, \dots\}$ .
- So the set  $\{0, 1, 2, 3, \dots\}$  must have the *same number of elements* as the set  $\{0, 1, 4, 9, \dots\}$ .



“we cannot speak of infinite quantities as being the one greater or less than or equal to another”

Galileo Galilei

*Dialogues concerning two new sciences*, 1638

# Defining “infinite sets”

# Defining “infinite sets”

Q: How can we define what is mean by an “*infinite set*” in terms of the concepts we’ve defined so far?

Some set  $S$  is infinite if...

# Defining “infinite sets”

Q: How can we define what is mean by an “*infinite set*” in terms of the concepts we’ve defined so far?

Some set  $S$  is infinite if...

A: ... there is no  $n \in \mathbb{N}$  such that  $\#S = n$ .



# Defining “infinite sets”

Q: How can we define what is mean by an “*infinite set*” in terms of the concepts we’ve defined so far?

Some set  $S$  is infinite if...

A: ... there is no  $n \in \mathbb{N}$  such that  $\#S = n$ . ❌  
- *we haven't defined cardinality for infinite sets.*

# Defining “infinite sets”

Q: How can we define what is mean by an “*infinite set*” in terms of the concepts we’ve defined so far?

Some set  $S$  is infinite if...

A: ... there is no  $n \in \mathbb{N}$  such that  $\#S = n$ . ❌  
- *we haven't defined cardinality for infinite sets.*

A: ... we can remove an element and it stays infinite.

# Defining “infinite sets”

Q: How can we define what is mean by an “*infinite set*” in terms of the concepts we’ve defined so far?

Some set  $S$  is infinite if...

A: ... there is no  $n \in \mathbb{N}$  such that  $\#S = n$ . ❌  
- *we haven't defined cardinality for infinite sets.*

A: ... we can remove an element and it stays infinite. ❌  
- *we still haven't defined “infinite” yet.*

# Defining “Infinite sets”



Richard  
Dedekind  
(1831-1916)

- A set is **infinite** if there exists a bijection between it and one of its own proper subsets.

*Was sind und was sollen die Zahlen?*, 1888  
(“What are numbers and what should they be?”)

# Cardinality of infinite sets

Two (finite or infinite) sets **have the same cardinality** iff there exists a bijection between them.

## Cardinality of infinite sets

Two (finite or infinite) sets **have the same cardinality** iff there exists a bijection between them.

So, the set of natural numbers has the same cardinality as:

- The set of numbers greater than zero

# Cardinality of infinite sets

Two (finite or infinite) sets **have the same cardinality** iff there exists a bijection between them.

So, the set of natural numbers has the same cardinality as:

- The set of numbers greater than zero

Proof:  $(\lambda x \in \mathbb{N} \cdot x + 1)$  is a bijection.

## Cardinality of infinite sets

Two (finite or infinite) sets **have the same cardinality** iff there exists a bijection between them.

So, the set of natural numbers has the same cardinality as:

- The set of numbers greater than zero

Proof:  $(\lambda x \in \mathbb{N} \cdot x + 1)$  is a bijection.

- The set of numbers greater than  $k$  (for any given  $k \in \mathbb{N}$ )



# Cardinality of infinite sets

Two (finite or infinite) sets **have the same cardinality** iff there exists a bijection between them.

So, the set of natural numbers has the same cardinality as:

- The set of numbers greater than zero

Proof:  $(\lambda x \in \mathbb{N} \cdot x + 1)$  is a bijection.

- The set of numbers greater than  $k$  (for any given  $k \in \mathbb{N}$ )

Proof:  $(\lambda x \in \mathbb{N} \cdot x + k)$  is a bijection.

# Cardinality of infinite sets

Two (finite or infinite) sets **have the same cardinality** iff there exists a bijection between them.

So, the set of natural numbers has the same cardinality as:

- The set of numbers greater than zero

Proof:  $(\lambda x \in \mathbb{N} \cdot x + 1)$  is a bijection.

- The set of numbers greater than  $k$  (for any given  $k \in \mathbb{N}$ )

Proof:  $(\lambda x \in \mathbb{N} \cdot x + k)$  is a bijection.

- The set of even numbers

# Cardinality of infinite sets

Two (finite or infinite) sets **have the same cardinality** iff there exists a bijection between them.

So, the set of natural numbers has the same cardinality as:

- The set of numbers greater than zero

Proof:  $(\lambda x \in \mathbb{N} \cdot x + 1)$  is a bijection.

- The set of numbers greater than  $k$  (for any given  $k \in \mathbb{N}$ )

Proof:  $(\lambda x \in \mathbb{N} \cdot x + k)$  is a bijection.

- The set of even numbers

Proof:  $(\lambda x \in \mathbb{N} \cdot 2 * x)$  is a bijection.

# Cardinality of infinite sets

Two (finite or infinite) sets **have the same cardinality** iff there exists a bijection between them.

So, the set of natural numbers has the same cardinality as:

- The set of numbers greater than zero

Proof:  $(\lambda x \in \mathbb{N} \cdot x + 1)$  is a bijection.

- The set of numbers greater than  $k$  (for any given  $k \in \mathbb{N}$ )

Proof:  $(\lambda x \in \mathbb{N} \cdot x + k)$  is a bijection.

- The set of even numbers

Proof:  $(\lambda x \in \mathbb{N} \cdot 2 * x)$  is a bijection.

- The set of odd numbers

# Cardinality of infinite sets

Two (finite or infinite) sets **have the same cardinality** iff there exists a bijection between them.

So, the set of natural numbers has the same cardinality as:

- The set of numbers greater than zero

Proof:  $(\lambda x \in \mathbb{N} \cdot x + 1)$  is a bijection.

- The set of numbers greater than  $k$  (for any given  $k \in \mathbb{N}$ )

Proof:  $(\lambda x \in \mathbb{N} \cdot x + k)$  is a bijection.

- The set of even numbers

Proof:  $(\lambda x \in \mathbb{N} \cdot 2 * x)$  is a bijection.

- The set of odd numbers

Proof:  $(\lambda x \in \mathbb{N} \cdot (2 * x) + 1))$  is a bijection.

## Cardinality of infinite sets

Two (finite or infinite) sets **have the same cardinality** iff there exists a bijection between them.

So, the set of natural numbers has the same cardinality as:

- The set of numbers greater than zero  
Proof:  $(\lambda x \in \mathbb{N} \cdot x + 1)$  is a bijection.
- The set of numbers greater than  $k$  (for any given  $k \in \mathbb{N}$ )  
Proof:  $(\lambda x \in \mathbb{N} \cdot x + k)$  is a bijection.
- The set of even numbers  
Proof:  $(\lambda x \in \mathbb{N} \cdot 2 * x)$  is a bijection.
- The set of odd numbers  
Proof:  $(\lambda x \in \mathbb{N} \cdot (2 * x) + 1)$  is a bijection.
- The set of numbers divisible by  $k$  (for any given  $k > 0$ )

# Cardinality of infinite sets

Two (finite or infinite) sets **have the same cardinality** iff there exists a bijection between them.

So, the set of natural numbers has the same cardinality as:

- The set of numbers greater than zero  
Proof:  $(\lambda x \in \mathbb{N} \cdot x + 1)$  is a bijection.
- The set of numbers greater than  $k$  (for any given  $k \in \mathbb{N}$ )  
Proof:  $(\lambda x \in \mathbb{N} \cdot x + k)$  is a bijection.
- The set of even numbers  
Proof:  $(\lambda x \in \mathbb{N} \cdot 2 * x)$  is a bijection.
- The set of odd numbers  
Proof:  $(\lambda x \in \mathbb{N} \cdot (2 * x) + 1)$  is a bijection.
- The set of numbers divisible by  $k$  (for any given  $k > 0$ )  
Proof:  $(\lambda x \in \mathbb{N} \cdot k * x)$  is a bijection.

# Countable

Any set that has the same cardinality as the natural numbers is called a **countable set**.

**Notation:** It is usual to denote the cardinality of the natural numbers by the symbol  $\aleph_0$  (pronounced “aleph null”)



# Countable

Any set that has the same cardinality as the natural numbers is called a **countable set**.

**Notation:** It is usual to denote the cardinality of the natural numbers by the symbol  $\aleph_0$  (pronounced “aleph null”)

Thus the following *subsets* of  $\mathbb{N}$  are all countable sets:

- $\{n \in \mathbb{N} \mid n + 1\}$
- $\{n \in \mathbb{N} \mid n + k\}$ , for any fixed  $k \in \mathbb{N}$
- $\{n \in \mathbb{N} \mid 2 * x\}$
- $\{n \in \mathbb{N} \mid (2 * x) + 1\}$
- $\{n \in \mathbb{N} \mid n * k\}$ , for any fixed  $k \in \mathbb{N}$

# Countable

Any set that has the same cardinality as the natural numbers is called a **countable set**.

**Notation:** It is usual to denote the cardinality of the natural numbers by the symbol  $\aleph_0$  (pronounced “aleph null”)

Thus the following operations do not change the cardinality of  $\mathbb{N}$

- Deleting an element of  $\mathbb{N}$
- Deleting  $k$  elements of  $\mathbb{N}$ , for any fixed  $k \in \mathbb{N}$
- Deleting every second element in  $\mathbb{N}$
- Deleting every  $k^{\text{th}}$  element in  $\mathbb{N}$ , for any fixed  $k > 0$

# Cardinality of the integers

Suppose we consider a *superset* of  $\mathbb{N}$ ...

What is the cardinality of the integers?

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

# Cardinality of the integers

Suppose we consider a *superset* of  $\mathbb{N}$ ...

What is the cardinality of the integers?

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

Surprise: The cardinality of  $\mathbb{Z}$  is  $\aleph_0$

# Cardinality of the integers

Suppose we consider a *superset* of  $\mathbb{N}$ ...

What is the cardinality of the integers?

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

Surprise: The cardinality of  $\mathbb{Z}$  is  $\aleph_0$

**Proof:** There is a bijective function  $f$  from  $\mathbb{Z}$  to  $\mathbb{N}$  defined as:

$$f(x) = \begin{cases} -2 * x, & \text{if } x \leq 0 \\ (2 * x) - 1, & \text{if } x > 0. \end{cases}$$

# Cardinality of the integers

Suppose we consider a *superset* of  $\mathbb{N}$ ...

What is the cardinality of the integers?

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

Surprise: The cardinality of  $\mathbb{Z}$  is  $\aleph_0$

**Proof:** There is a bijective function  $f$  from  $\mathbb{Z}$  to  $\mathbb{N}$  defined as:

$$f(x) = \begin{cases} -2 * x, & \text{if } x \leq 0 \\ (2 * x) - 1, & \text{if } x > 0. \end{cases}$$

That is:

- map the negative integers to the even natural numbers
- and the positive integers to the odd natural numbers

# Cardinality of the integers

Suppose we consider a *superset* of  $\mathbb{N}$ ...

What is the cardinality of the integers?

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

Surprise: The cardinality of  $\mathbb{Z}$  is  $\aleph_0$

**Proof:** There is a bijective function  $f$  from  $\mathbb{Z}$  to  $\mathbb{N}$  defined as:

$$f(x) = \begin{cases} -2 * x, & \text{if } x \leq 0 \\ (2 * x) - 1, & \text{if } x > 0. \end{cases}$$

This gives the mapping:

$$\{\dots (-3, 6), (-2, 4), (-1, 2), (0, 0), (1, 1), (2, 3), (3, 5), \dots\}$$

## Another even 'bigger' set?

What is the cardinality of the set  $\mathbb{N} \times \mathbb{N}$ ?




## Another even 'bigger' set?

What is the cardinality of the set  $\mathbb{N} \times \mathbb{N}$ ?

(0,0)	(1,0)	(2,0)	(3,0)	...
(0,1)	(1,1)	(2,1)	(3,1)	...
(0,2)	(1,2)	(2,2)	(3,2)	...
(0,3)	(1,3)	(2,3)	(3,3)	...
⋮	⋮	⋮	⋮	


# Another even 'bigger' set?

What is the cardinality of the set  $\mathbb{N} \times \mathbb{N}$ ?



# Another even 'bigger' set?


What is the cardinality of the set  $\mathbb{N} \times \mathbb{N}$ ?



$$= \{(0,0), (1,0), (0,1), (0,2), (1,1), (2,0), (3,0), (2,1), (1,2), (0,3), \dots\}$$

## Another even 'bigger' set?

What is the cardinality of the set  $\mathbb{N} \times \mathbb{N}$ ?



Note each diagonal is just the (finite) set  $\{i, j \in \mathbb{N} \mid i + j = k \bullet (i, j)\}$  for some fixed value of  $k \in \mathbb{N}$

# Defining the mapping

We can define this more formally...

- Define an *injective* function  $f$  from  $(\mathbb{N} \times \mathbb{N})$  to  $\mathbb{N}$  as:

$$f((i,j)) = 2^i * 3^j$$

# Defining the mapping

We can define this more formally...

- Define an *injective* function  $f$  from  $(\mathbb{N} \times \mathbb{N})$  to  $\mathbb{N}$  as:

$$f((i,j)) = 2^i * 3^j$$

This uses an arithmetical trick:

# Defining the mapping

We can define this more formally...

- Define an *injective* function  $f$  from  $(\mathbb{N} \times \mathbb{N})$  to  $\mathbb{N}$  as:

$$f((i,j)) = 2^i * 3^j$$

This uses an arithmetical trick:

- The **fundamental theorem of arithmetic** tells us that any natural number can be factored uniquely into the product of powers of primes.

# Defining the mapping

We can define this more formally...

- Define an *injective* function  $f$  from  $(\mathbb{N} \times \mathbb{N})$  to  $\mathbb{N}$  as:

$$f((i,j)) = 2^i * 3^j$$

This uses an arithmetical trick:

- The **fundamental theorem of arithmetic** tells us that any natural number can be factored uniquely into the product of powers of primes. Since 2 and 3 are prime,  $2^i * 3^j$  is unique for any given  $i$  and  $j$ .



## Defining the mapping

We can define this more formally...

- Define an *injective* function  $f$  from  $(\mathbb{N} \times \mathbb{N})$  to  $\mathbb{N}$  as:

$$f((i,j)) = 2^i * 3^j$$

This uses an arithmetical trick:

- The **fundamental theorem of arithmetic** tells us that any natural number can be factored uniquely into the product of powers of primes. Since 2 and 3 are prime,  $2^i * 3^j$  is unique for any given  $i$  and  $j$ .
- Since  $f$  is an *injection* into  $\mathbb{N}$ , it must be a *bijection* into some infinite subset of  $\mathbb{N}$ , hence:

## Defining the mapping

We can define this more formally...

- Define an *injective* function  $f$  from  $(\mathbb{N} \times \mathbb{N})$  to  $\mathbb{N}$  as:

$$f((i, j)) = 2^i * 3^j$$

This uses an arithmetical trick:

- The **fundamental theorem of arithmetic** tells us that any natural number can be factored uniquely into the product of powers of primes. Since 2 and 3 are prime,  $2^i * 3^j$  is unique for any given  $i$  and  $j$ .
- Since  $f$  is an *injection* into  $\mathbb{N}$ , it must be a *bijection* into some infinite subset of  $\mathbb{N}$ , hence:

The cardinality of  $(\mathbb{N} \times \mathbb{N})$  is  $\aleph_0$

## Extending this to any size tuple

- The same trick - based on prime factorisation - will work for tuples of any (finite) size.
- Example: suppose we have 6-tuples of numbers, that is, elements of the set  $(\mathbb{N} \times \mathbb{N} \times \mathbb{N} \times \mathbb{N} \times \mathbb{N} \times \mathbb{N}) \dots$
- Then we can map any 6-tuple of the form  $(a_1, a_2, a_3, a_4, a_5, a_6)$  to the number

$$2^{a_1+1} * 3^{a_2+1} * 5^{a_3+1} * 7^{a_4+1} * 11^{a_5+1} * 13^{a_6+1}$$

- Example:  $(3, 4, 2, 0, 2, 1)$  maps to

$$2^4 * 3^5 * 5^3 * 7^1 * 11^3 * 13^2 = 765,242,478,000$$

- We know that this will work for  $k$ -tuples for any given  $k \in \mathbb{N}$ , since there will always be enough prime numbers.

## Other countable sets

The previous work implies that some other infinite sets are also countable:

- The rational numbers,  $\mathbb{Q}$ , since any fraction of the form  $\frac{a}{b}$  can be mapped to the tuple  $(a, b)$  and then into the natural numbers.

## Other countable sets

The previous work implies that some other infinite sets are also countable:

- The rational numbers,  $\mathbb{Q}$ , since any fraction of the form  $\frac{a}{b}$  can be mapped to the tuple  $(a, b)$  and then into the natural numbers.
- Any formal language (e.g. defined by a regular expression or a context-free grammar), since a finite-length string over any given alphabet is really just a tuple of characters.

## Other countable sets

The previous work implies that some other infinite sets are also countable:


- The rational numbers,  $\mathbb{Q}$ , since any fraction of the form  $\frac{a}{b}$  can be mapped to the tuple  $(a, b)$  and then into the natural numbers.
- Any formal language (e.g. defined by a regular expression or a context-free grammar), since a finite-length string over any given alphabet is really just a tuple of characters.
- The set of all valid Java programs (or C#, or ...), since these can be arranged in order: shortest first, and then alphabetically for programs of the same length.

## Hilbert's Hotel: a *Gedankenexperiment*

Imagine a hotel with infinitely many rooms, numbered  $\{0, 1, 2, 3, 4, \dots\}$ , and suppose the hotel is currently full.

## Hilbert's Hotel: a *Gedankenexperiment*

Imagine a hotel with infinitely many rooms, numbered  $\{0, 1, 2, 3, 4, \dots\}$ , and suppose the hotel is currently full.



60-second video from the **Open University**:

<http://www.youtube.com/watch?v=faQBrAQ8714>



# Hilbert's Hotel: a *Gedankenexperiment*

Imagine a hotel with infinitely many rooms, numbered  $\{0, 1, 2, 3, 4, \dots\}$ , and suppose the hotel is currently full.



Let's Begin...

The Infinite Hotel, a thought experiment created by German mathematician David Hilbert, is a hotel with an infinite number of rooms but one person wants to check in? What about 407? Or an infinitely full bus of people? Jeff Dekofsky solves that.



6-minute video by Jeff Dekofsky at **TedEd**:

[http://ed.ted.com/lessons/  
the-infinite-hotel-paradox-jeff-dekofsky/](http://ed.ted.com/lessons/the-infinite-hotel-paradox-jeff-dekofsky/)

## Hilbert's Hotel: a *Gedankenexperiment*

Imagine a hotel with infinitely many rooms, numbered  $\{0, 1, 2, 3, 4, \dots\}$ , and suppose the hotel is currently full.

How would Hilbert accommodate:

- one new guest
- $k$  new guests, for some given  $k \in \mathbb{N}$
- a bus containing infinitely many new guests
- $k$  buses, with each bus containing infinitely many new guests, for some given  $k \in \mathbb{N}$
- infinitely many buses, with each bus containing infinitely many new guests
- $k$  aircraft carriers, each of which contains infinitely many buses, with each bus containing infinitely many new guests, for some given  $k \in \mathbb{N}$

Welcome to...

# Hilbert's Hotel

- never any vacancy -
- always room for more -