

COMP9020 19T1

Week 3

Functions and Relations

- Textbook (R & W) - Ch. 1, Sec. 1.7
Ch. 3, Sec. 3.1, 3.3
- Problem set 3
- Supplementary Exercises Ch. 1, Ch. 3 (R & W)
- Quiz 3 (due Monday week 4)

Functions

Reminder:

We deal with functions as a set-theoretic concept, it being a special kind of correspondence (between two sets)

$f : S \rightarrow T$ describes pairing of the sets: it means that f assigns to every element $s \in S$ a unique element $t \in T$.

S — **domain** of f , symbol: $\text{Dom}(f)$

T — **codomain** of f , symbol: $\text{Codom}(f)$

$\{f(x) : x \in \text{Dom}(f)\}$ — **image** of f , symbol: $\text{Im}(f)$

$$\text{Im}(f) \subseteq \text{Codom}(f)$$

We observe that every function maps its domain **into** its codomain, but only **onto** its image.

1

Navigation icons

2

Navigation icons

Exercise

1.5.3 Regarding $\text{length} : \{a, b\}^* \rightarrow \mathbb{N}$

- (c) $\text{length}(\lambda) = ?$
- (d) $\text{Im}(\text{length}) = ?$

1.5.4 Σ^* as above and $g(n) \stackrel{\text{def}}{=} \{\omega \in \Sigma^* : \text{length}(\omega) \leq n\}$, $n \in \mathbb{N}$
Here $g(n)$ is a function that has a complex object as its value for any given argument — it maps \mathbb{N} into $\text{Pow}(\Sigma^*)$

- (a) $g(0) = ?$
- (b) $g(1) = ?$
- (c) $g(2) = ?$
- (d) Are all $g(n)$ finite?

Exercise

1.5.3 Regarding $\text{length} : \{a, b\}^* \rightarrow \mathbb{N}$

- (c) $\text{length}(\lambda) = 0$
- (d) $\text{Im}(\text{length}) = \mathbb{N}$

1.5.4 Σ^* as above and $g(n) \stackrel{\text{def}}{=} \{\omega \in \Sigma^* : \text{length}(\omega) \leq n\}$, $n \in \mathbb{N}$
Here $g(n)$ is a function that has a complex object as its value for any given argument — it maps \mathbb{N} into $\text{Pow}(\Sigma^*)$

- (a) $g(0) = \{\lambda\}$
- (b) $g(1) = \{\lambda, a, b\}$
- (c) $g(2) = \{\lambda, a, b, aa, ab, ba, bb\}$

In general $g(n) = \bigcup_{i=0}^n \Sigma^i = \Sigma^{\leq n}$

- (d) Are all $g(n)$ finite?

Yes; $|g(n)| = 2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$

3

Navigation icons

4

Navigation icons

Exercise (cont'd)

(e) Give an example of a set in $\text{Pow}(\Sigma^*)$ that is not in $\text{Im}(g)$

Exercise (cont'd)

(e) Give an example of a set in $\text{Pow}(\Sigma^*)$ that is not in $\text{Im}(g)$

- any infinite subset of Σ^* (infinite language)
- any finite language that excludes some intermediate length words, e.g. $\{\lambda, a\}, \{a, b\}, \{\lambda, a, aa\}, \dots$

5

6

Exercise (cont'd)

1.5.6 Regarding $\gcd : \mathbb{P} \times \mathbb{P} \longrightarrow \mathbb{P}$

(c) $\text{Im}(\text{gcd}) \stackrel{?}{=}$

1.5.7 Define $f : \mathbb{R} \rightarrow \mathbb{R}$ by

$$f(x) = \begin{cases} x^3 & x \geq 1 \\ x & 0 \leq x < 1 \\ -x^3 & x < 0 \end{cases}$$

(c) $\text{Im}(f) \stackrel{?}{=}$

Exercise (cont'd)

1.5.6 Regarding $\gcd : \mathbb{P} \times \mathbb{P} \longrightarrow \mathbb{P}$

(c) $\text{Im}(\gcd) = \mathbb{P}$ since $\gcd(n, n) = n$

1.5.7 Define $f : \mathbb{R} \rightarrow \mathbb{R}$ by

$$f(x) = \begin{cases} x^3 & x \geq 1 \\ x & 0 \leq x < 1 \\ -x^3 & x < 0 \end{cases}$$

(c) $\text{Im}(f) = \mathbb{R}_{\geq 0}$

7

8

Composition of Functions

Auxiliary notation

$$f : x \mapsto y, \quad f : A \mapsto B$$

The former means that x is mapped to y ; the latter means that B is the image of A under f .

NB

Observe the difference between \longrightarrow and \mapsto

Composition of functions is described as

$$g \circ f : x \mapsto g(f(x)), \quad \text{requiring } \text{Im}(f) \subseteq \text{Dom}(g)$$

If a function maps a set into itself, i.e. when $\text{Dom}(f) = \text{Codom}(f)$ (and thus $\text{Im}(f) \subseteq \text{Dom}(f)$), the function can be composed with itself — **iterated**

$$f \circ f, f \circ f \circ f, \dots, \quad \text{also written } f^2, f^3, \dots$$

Composition is associative

$$h \circ (g \circ f) = (h \circ g) \circ f, \quad \text{can write } h \circ g \circ f$$

Identity function on S

$$\text{Id}_S(x) = x, x \in S; \text{Dom}(i) = \text{Codom}(i) = \text{Im}(i) = S$$

For $g : S \longrightarrow T$ $g \circ \text{Id}_S = g, \text{Id}_T \circ g = g$

gcd Example

Reconsider gcd as a **higher-order function**, defined by

$$\text{gcd}(f)(m, n) = \begin{cases} m & \text{if } m = n \\ f(m - n, n) & \text{if } m > n \\ f(m, n - m) & \text{if } m < n \end{cases}$$

Its type is now $\text{gcd} : (\mathbb{P}^2 \rightharpoonup \mathbb{P}) \longrightarrow (\mathbb{P}^2 \rightharpoonup \mathbb{P})$
that is, it maps each partial function (from pairs of positive integers to a positive integer) to a (partial) function of the same type. The worst such function is the “nowhere defined” function

$$f_{\perp}(m, n) = \perp.$$

NB

A **partial function** $f : S \rightharpoonup T$ is a function $f : S' \longrightarrow T$ for $S' \subseteq S$

gcd Example cont'd

Consider the sequence

$$f_{\perp}, \text{gcd}(f_{\perp}), \text{gcd}(\text{gcd}(f_{\perp})), \dots, \text{gcd}(\text{gcd}(\dots(f_{\perp})\dots)), \dots$$

and observe that the i 'th element of this sequence is an approximation of the gcd function that works as long as the depth of the recursion is less than $i - 1$. Since we proved that the original gcd function terminates, we can deduce that the limit of this sequence exists, and is the original gcd. It also is the **least fixpoint** of gcd i.e. the “simplest” solution f to the equation $f = \text{gcd}(f)$. This, in a nutshell, explains how the semantics of recursive procedures is defined in CS. How all this works is somewhat beyond the scope of COMP9020 but still serves the purpose of motivating why we discuss functions and their composition, iteration.

Properties of Functions

Function is called **onto** (or **surjective**) if every element of the codomain is mapped to by at least one x in the domain, i.e.

$$\text{Im}(f) = T$$

Examples (of functions that are not onto)

- $f : \mathbb{N} \rightarrow \mathbb{N}$ with $f(x) \mapsto x^2$
- $f : \{a, \dots, z\}^* \rightarrow \{a, \dots, z\}^*$ with $f(w) \mapsto awe$

1-1 Functions

Function is called **1-1** (**one-to-one**) or **injective** if different x implies different $f(x)$, i.e.

$$f(x) = f(y) \Rightarrow x = y$$

Examples (of functions that are not 1-1)

- absolute value
- floor, ceiling
- length of a word

13

Navigation icons

14

Inverse Functions

Definition

Inverse function for a given $f : S \rightarrow T$

$$f^{-1} : T \rightarrow S \quad \text{s.t.} \quad f^{-1}(f(x)) = x \text{ for all } x \in S$$

exists exactly when f is both 1-1 and onto

Image of a subdomain A under a function

$$f(A) = \{ f(s) : s \in A \} = \{ t \in T : t = f(s) \text{ for some } s \in A \}$$

Inverse image — $f^{\leftarrow}(B) = \{ s \in S : f(s) \in B \} \subseteq S$;

it is defined for every f

If f^{-1} exists then $f^{\leftarrow}(B) = f^{-1}(B)$

$$f(\emptyset) = \emptyset, f^{\leftarrow}(\emptyset) = \emptyset$$

15

Navigation icons

16

Exercise

1.7.5 f and g are 'shift' functions $\mathbb{N} \rightarrow \mathbb{N}$ defined by $f(n) = n + 1$, and $g(n) = \max(0, n - 1)$

- (c) Is f 1-1? onto?
- (d) Is g 1-1? onto?
- (e) Do f and g commute, i.e. $\forall n ((f \circ g)(n) = (g \circ f)(n))$?

Navigation icons

Exercise

1.7.5 f and g are 'shift' functions $\mathbb{N} \rightarrow \mathbb{N}$ defined by $f(n) = n + 1$, and $g(n) = \max(0, n - 1)$

(c) f is 1-1, not onto: $f(\mathbb{N}) = \mathbb{N} \setminus \{0\} = \mathbb{P}$

(d) g is onto, not 1-1: $g(0) = g(1)$

(e) f and g do not commute:

$g \circ f : n \mapsto (n + 1) - 1 = n$, thus $g \circ f = \text{Id}_{\mathbb{N}}$

$f \circ g : 0 \mapsto 1$, hence $f \circ g \neq \text{Id}_{\mathbb{N}}$

NB

For a **finite** set S and $f : S \rightarrow S$ the properties

- ① onto, and
- ② 1-1

are equivalent. (Proof suggestion?)

NB

$f \circ g$ is the identity when restricted to \mathbb{P}

17



18



Exercise

1.7.6 $\Sigma = \{a, b, c\}$

(c) Is $\text{length} : \Sigma^* \rightarrow \mathbb{N}$ onto? Yes: $\text{length}^{\leftarrow}(\{n\}) = \Sigma^n \neq \emptyset$

(d) $\text{length}^{\leftarrow}(2) = \{aa, ab, ac, bb, \dots, cc\}$

Exercise

1.7.6 $\Sigma = \{a, b, c\}$

(c) Is $\text{length} : \Sigma^* \rightarrow \mathbb{N}$ onto? Yes: $\text{length}^{\leftarrow}(\{n\}) = \Sigma^n \neq \emptyset$

(d) $\text{length}^{\leftarrow}(2) = \{aa, ab, ac, bb, \dots, cc\}$

Exercise

1.7.12 Verify that $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ defined by $f(x, y) = (x + y, x - y)$ is invertible.

The inverse is $f^{-1}(a, b) = (\frac{a+b}{2}, \frac{a-b}{2})$; substituting shows that $f \circ f^{-1} = \text{Id}_{\mathbb{R} \times \mathbb{R}}$

Exercise

1.7.12 Verify that $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ defined by $f(x, y) = (x + y, x - y)$ is invertible.

The inverse is $f^{-1}(a, b) = (\frac{a+b}{2}, \frac{a-b}{2})$; substituting shows that $f \circ f^{-1} = \text{Id}_{\mathbb{R} \times \mathbb{R}}$

19



20



Supplementary Exercises

Exercise

1.8.16 $\Sigma = \{a, b\}$; relate it to Σ^*

(a) Is there an onto $\Sigma \rightarrow \Sigma^*$?

(b) Is there an onto $\Sigma^* \rightarrow \Sigma$?

Supplementary Exercises

Exercise

1.8.16 $\Sigma = \{a, b\}$; relate it to Σ^* :

(a) Is there an onto $\Sigma \rightarrow \Sigma^*$? No: $|\Sigma| = 2, |\Sigma^*| = \infty$.

(b) Is there an onto $\Sigma^* \rightarrow \Sigma$? Yes, eg $f(\omega) = a$ when $\text{length}(\omega)$ is odd, $f(\omega) = b$ when $\text{length}(\omega)$ is even.

The following is **not** completely correct $f: \omega \mapsto \langle \text{first letter of } \omega \rangle$
Reason: $f(\lambda)$ is not defined.

21



Matrices

An $m \times n$ **matrix** is a rectangular array with m horizontal rows and n vertical columns.

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

NB

Matrices are important objects in Computer Science, e.g. for

- optimisation
- graphics and computer vision
- cryptography
- information retrieval and web search
- machine learning

22



Basic Matrix Operations

The **transpose** \mathbf{A}^T of an $m \times n$ matrix $\mathbf{A} = [a_{ij}]$ is the $n \times m$ matrix whose entry in the i th row and j th column is a_{ji} .

Example

$$\mathbf{A} = \begin{bmatrix} 2 & -1 & 0 & 4 \\ 3 & 2 & -1 & 2 \\ 4 & 0 & 1 & 3 \end{bmatrix} \quad \mathbf{A}^T = \begin{bmatrix} 2 & 3 & 4 \\ -1 & 2 & 0 \\ 0 & -1 & 1 \\ 4 & 2 & 3 \end{bmatrix}$$

NB

A matrix \mathbf{M} is called symmetric if $\mathbf{M}^T = \mathbf{M}$

23



24



The **sum** of two $m \times n$ matrices $\mathbf{A} = [a_{ij}]$ and $\mathbf{B} = [b_{ij}]$ is the $m \times n$ matrix whose entry in the i th row and j th column is $a_{ij} + b_{ij}$.

Example

$$\mathbf{A} = \begin{bmatrix} 2 & -1 & 0 & 4 \\ 3 & 2 & -1 & 2 \\ 4 & 0 & 1 & 3 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 1 & 0 & 5 & 3 \\ 2 & 3 & -2 & 1 \\ 4 & -2 & 0 & 2 \end{bmatrix}$$

$$\mathbf{A} + \mathbf{B} = \begin{bmatrix} 3 & -1 & 5 & 7 \\ 5 & 5 & -3 & 3 \\ 8 & -2 & 1 & 5 \end{bmatrix}$$

Fact

$$\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A} \text{ and } (\mathbf{A} + \mathbf{B}) + \mathbf{C} = \mathbf{A} + (\mathbf{B} + \mathbf{C})$$

Given $m \times n$ matrix $\mathbf{A} = [a_{ij}]$ and $c \in \mathbb{R}$, the **scalar product** $c\mathbf{A}$ is the $m \times n$ matrix whose entry in the i th row and j th column is $c \cdot a_{ij}$.

Example

$$\mathbf{A} = \begin{bmatrix} 2 & -1 & 0 & 4 \\ 3 & 2 & -1 & 2 \\ 4 & 0 & 1 & 3 \end{bmatrix} \quad 2\mathbf{A} = \begin{bmatrix} 4 & -2 & 0 & 8 \\ 6 & 4 & -2 & 4 \\ 8 & 0 & 2 & 6 \end{bmatrix}$$

25

The **product** of an $m \times n$ matrix $\mathbf{A} = [a_{ij}]$ and an $n \times p$ matrix $\mathbf{B} = [b_{jk}]$ is the $m \times p$ matrix $\mathbf{C} = [c_{ik}]$ defined by

$$c_{ik} = \sum_{j=1}^n a_{ij} b_{jk} \quad \text{for } 1 \leq i \leq m \text{ and } 1 \leq k \leq p$$

Example

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \cdot \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{bmatrix}$$

NB

The **rows** of \mathbf{A} must have the same number of entries as the **columns** of \mathbf{B} .

The product of a $1 \times n$ matrix and an $n \times 1$ matrix is usually called the **inner product** of two **n -dimensional vectors**.

26

Exercise

Consider

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 2 & -1 \\ -6 & 3 \end{bmatrix}$$

Calculate \mathbf{AB} , \mathbf{BA}

$$\mathbf{AB} = \begin{bmatrix} -10 & 5 \\ -20 & 10 \end{bmatrix} \quad \mathbf{BA} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

NB

In general, $\mathbf{A} \cdot \mathbf{B} \neq \mathbf{B} \cdot \mathbf{A}$


27

28

Example: Computer Graphics

Rotating an object w.r.t. the x axis by degree α :

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \alpha & -\sin \alpha \\ 0 & \sin \alpha & \cos \alpha \end{bmatrix} \cdot \begin{bmatrix} 5 & 5 & 7 & 7 & 5 & 7 & 5 & 7 \\ 1 & 1 & 1 & 1 & 3 & 3 & 3 & 3 \\ 9 & 7 & 7 & 9 & 7 & 7 & 9 & 9 \end{bmatrix}$$



Exercise

Consider

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 2 & -1 \\ -6 & 3 \end{bmatrix}$$

Calculate \mathbf{AB} , \mathbf{BA}

$$\mathbf{AB} = \begin{bmatrix} -10 & 5 \\ -20 & 10 \end{bmatrix} \quad \mathbf{BA} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

NB

In general, $\mathbf{A} \cdot \mathbf{B} \neq \mathbf{B} \cdot \mathbf{A}$

29

30

Relations and their Representation

Relations are an abstraction used to capture the idea that the objects from certain domains (often the same domain for several objects) are *related*. These objects may

- influence one another (each other for binary relations; self(?) for unary)
- share some common properties
- correspond to each other precisely when some constraints are satisfied

In general, relations formalise the concept of interaction among objects from various domains; however, there must be a specified domain for each type of objects.

An **n-ary relation** is a subset of the cartesian product of n sets.

$$\mathcal{R} \subseteq S_1 \times S_2 \times \dots \times S_n$$

$$x \in \mathcal{R} \Rightarrow x = (x_1, x_2, \dots, x_n) \text{ where each } x_i \in S_i$$

If $n = 2$ we have a **binary** relation $\mathcal{R} \subseteq S \times T$.

(mostly we consider binary relations)

equivalent notations: $(x_1, x_2, \dots, x_n) \in \mathcal{R} \iff \mathcal{R}(x_1, x_2, \dots, x_n)$

for binary relations: $(x, y) \in \mathcal{R} \iff \mathcal{R}(x, y) \iff x\mathcal{R}y$.

31

32

Database Examples

Example (course enrolments)

S = set of CSE students
(S can be a subset of the set of all students)
 C = set of CSE courses
(likewise)
 E = enrolments = $\{ (s, c) : s \text{ takes } c \}$

$$E \subseteq S \times C$$

In practice, almost always there are various 'onto' (nonemptiness) and 1-1 (uniqueness) constraints on database relations.

Example (class schedule)

C = CSE courses
 T = starting time (hour & day)
 R = lecture rooms
 S = schedule =

$$\{ (c, t, r) : c \text{ is at } t \text{ in } r \} \subseteq C \times T \times R$$

Example (sport stats)

$$\mathcal{R} \subseteq \text{competitions} \times \text{results} \times \text{years} \times \text{athletes}$$

33

Navigation icons

34

Applications

Relations are ubiquitous in Computer Science

- Databases are collections of relations
- Common data structures (e.g. graphs) are relations
- Any ordering is a relation
- Functions/procedures/programs compute relations between their input and output

Relations are therefore used in most problem specifications and to describe formal properties of programs.

For this reason, studying relations and their properties helps with formalisation, implementation and verification of programs.

35

Navigation icons

n -ary Relations

Relations can be defined linking $k \geq 1$ domains D_1, \dots, D_k simultaneously.

In database situations one also allows for *unary* ($n = 1$) relations.

Most common are **binary** relations

$$\mathcal{R} \subseteq S \times T; \quad \mathcal{R} = \{ (s, t) : \text{"some property that links } s, t" \}$$

For related s, t we can write $(s, t) \in \mathcal{R}$ or $s\mathcal{R}t$; for unrelated items either $(s, t) \notin \mathcal{R}$ or $s \not\mathcal{R}t$.

\mathcal{R} can be defined by

- explicit enumeration of interrelated k -tuples (ordered pairs in case of binary relations);
- properties that identify relevant tuples within the entire $D_1 \times D_2 \times \dots \times D_k$;
- construction from other relations.

Navigation icons

36

Functions as Relations

Any function $f : S \rightarrow T$ can be viewed as a binary relation

$$\{ (s, f(s)) : s \in S \} \subseteq S \times T$$

If a subset of $S \times T$ corresponds to a function, it must satisfy certain conditions w.r.t. S and T (which?)

Binary Relations

A binary relation, say $\mathcal{R} \subseteq S \times T$, can be presented as a matrix with rows enumerated by (the elements of) S and the columns by T ; eg. for $S = \{s_1, s_2, s_3\}$ and $T = \{t_1, t_2, t_3, t_4\}$ we may have

$$\begin{bmatrix} \bullet & \circ & \bullet & \bullet \\ \circ & \bullet & \bullet & \bullet \\ \bullet & \bullet & \circ & \circ \end{bmatrix}$$

37

Navigation icons

38

Example

Exercise

3.1.2(e) Write the following relation on $A = \{0, 1, 2\}$ as a matrix.

$(m, n) \in \mathcal{R}$ if $m \cdot n = m$

$$\begin{array}{c} 0 \\ 1 \\ 2 \end{array} \begin{bmatrix} & 0 & 1 & 2 \\ \bullet & \bullet & \bullet \\ \circ & \bullet & \circ \\ \circ & \bullet & \circ \end{bmatrix}$$

39

Navigation icons

Example

Exercise

3.1.2(e) Write the following relation on $A = \{0, 1, 2\}$ as a matrix.

$(m, n) \in \mathcal{R}$ if $m \cdot n = m$

$$\begin{array}{c} 0 \\ 1 \\ 2 \end{array} \begin{bmatrix} & 0 & 1 & 2 \\ \bullet & \bullet & \bullet \\ \circ & \bullet & \circ \\ \circ & \bullet & \circ \end{bmatrix}$$

40

Navigation icons

Relations on a Single Domain

Special (Trivial) Relations

Particularly important are binary relationships between the elements of the same set. We say that ' \mathcal{R} is a binary relation on S ' if

$$\mathcal{R} \subseteq S \times S$$

(all w.r.t. set S)

Identity (diagonal, equality)

$$E = \{ (x, x) : x \in S \}$$

Empty \emptyset

Universal $U = S \times S$

41

Navigation icons

42

Navigation icons

Important Properties of Binary Relations $\mathcal{R} \subseteq S \times S$

Examples

- (R) reflexive $(x, x) \in \mathcal{R} \quad \forall x \in S$
- (AR) antireflexive $(x, x) \notin \mathcal{R} \quad \forall x \in S$
- (S) symmetric $(x, y) \in \mathcal{R} \Rightarrow (y, x) \in \mathcal{R} \quad \forall x, y \in S$
- (AS) antisymmetric $(x, y), (y, x) \in \mathcal{R} \Rightarrow x = y \quad \forall x, y \in S$
- (T) transitive $(x, y), (y, z) \in \mathcal{R} \Rightarrow (x, z) \in \mathcal{R} \quad \forall x, y, z \in S$

NB

An object, notion etc. is considered to satisfy a property if none of its instances violates any defining statement of that property.

(R) reflexive $(x, x) \in \mathcal{R}$ for all $x \in S$ $\begin{bmatrix} \circ & \circ & \circ \\ \circ & \circ & \circ \\ \circ & \circ & \circ \end{bmatrix}$

(AR) antireflexive $(x, x) \notin \mathcal{R}$ $\begin{bmatrix} \circ & \circ & \circ \\ \circ & \circ & \circ \\ \circ & \circ & \circ \end{bmatrix}$

(S) symmetric $(x, y) \in \mathcal{R} \Rightarrow (y, x) \in \mathcal{R}$ $\begin{bmatrix} \circ & \circ & \circ \\ \circ & \circ & \circ \\ \circ & \circ & \circ \end{bmatrix}$

(AS) antisymmetric $(x, y), (y, x) \in \mathcal{R} \Rightarrow x = y$ $\begin{bmatrix} \circ & \circ & \circ \\ \circ & \circ & \circ \\ \circ & \circ & \circ \end{bmatrix}$

(T) transitive $(x, y), (y, z) \in \mathcal{R} \Rightarrow (x, z) \in \mathcal{R}$ $\begin{bmatrix} \circ & \circ & \circ \\ \circ & \circ & \circ \\ \circ & \circ & \circ \end{bmatrix}$

43

Navigation icons

44

Navigation icons

Example

Exercise

3.1.1 The following relations are on $S = \{1, 2, 3\}$. Which of the properties (R), (AR), (S), (AS), (T) does each satisfy?

(a) $(m, n) \in \mathcal{R}$ if $m + n = 3$
(AR) and (S)

(e) $(m, n) \in \mathcal{R}$ if $\max\{m, n\} = 3$
(S)

3.1.2(b) $(m, n) \in \mathcal{R}$ if $m < n$
(AR), (AS), (T)

Example

Exercise

3.1.1 The following relations are on $S = \{1, 2, 3\}$. Which of the properties (R), (AR), (S), (AS), (T) does each satisfy?

(a) $(m, n) \in \mathcal{R}$ if $m + n = 3$
(AR) and (S)

(e) $(m, n) \in \mathcal{R}$ if $\max\{m, n\} = 3$
(S)

3.1.2(b) $(m, n) \in \mathcal{R}$ if $m < n$
(AR), (AS), (T)

45

46

Interaction of Properties

A relation *can* be both symmetric and antisymmetric. Namely, when \mathcal{R} consists only of some pairs (x, x) , $x \in S$.

A relation *cannot* be simultaneously reflexive and antireflexive (unless $S = \emptyset$).

NB

nonreflexive
nonsymmetric } is not the same as { antireflexive/irreflexive
antisymmetric

Most important kinds of relations on S

- total order $\begin{bmatrix} \bullet & \bullet & \bullet \\ \circ & \circ & \circ \\ \circ & \circ & \bullet \end{bmatrix}$
- partial order $\begin{bmatrix} \bullet & \bullet & \bullet \\ \circ & \circ & \circ \\ \circ & \circ & \bullet \end{bmatrix}, \begin{bmatrix} \bullet & \bullet & \circ \\ \circ & \bullet & \bullet \\ \circ & \circ & \bullet \end{bmatrix}$
- equivalence $\begin{bmatrix} \bullet & \bullet & \circ \\ \bullet & \bullet & \circ \\ \circ & \circ & \bullet \end{bmatrix}$
- identity $\begin{bmatrix} \bullet & \circ & \circ \\ \circ & \bullet & \circ \\ \circ & \circ & \bullet \end{bmatrix}$

NB

Some of those are special cases of the others, eg. 'total order' of a 'partial order', 'identity' of an 'equivalence'.

47

48

Relation \mathcal{R} as Correspondence From S to T

$\mathcal{R}(A) \stackrel{\text{def}}{=} \{t \in T : (s, t) \in \mathcal{R} \text{ for some } s \in A \subseteq S\}$
 $\mathcal{R}^{\leftarrow}(B) \stackrel{\text{def}}{=} \{s \in S : (s, t) \in \mathcal{R} \text{ for some } t \in B \subseteq T\}$
 Converse relation \mathcal{R}^{\leftarrow}

$$\mathcal{R}^{\leftarrow} = \{(t, s) \in T \times S : (s, t) \in \mathcal{R}\}$$

Note that $\mathcal{R}^{\leftarrow} \subseteq T \times S$.
 Observe that $(\mathcal{R}^{\leftarrow})^{\leftarrow} = \mathcal{R}$.

NB

Viewed this way \mathcal{R} becomes a function from $\text{Pow}(S)$ to $\text{Pow}(T)$. However, *not* every $g : \text{Pow}(S) \rightarrow \text{Pow}(T)$ can be matched to a relation.

(Why? Using a small domain like $S = \{a, b\}$, provide an example of a function $g : \text{Pow}(S) \rightarrow \text{Pow}(S)$ which does not correspond to any relation on S ! Can you even do it with $S' = \{a\}$?)

NB

The order of axes — S and T — is important. For $\mathcal{R} \subseteq S \times S$, its converse \mathcal{R}^{\leftarrow} is usually quite different from \mathcal{R} .

Example: divisibility relation on \mathbb{P}

$$\begin{aligned}
 D &\stackrel{\text{def}}{=} \{(p, q) : p \mid q\} = \{(1, 1), (1, 2), \dots, (2, 2), (2, 4), \dots\} \\
 D^{\leftarrow} &= \{(p, q) : p \in q\mathbb{P}\} \\
 &= \{(1, 1), (2, 1), (2, 2), (3, 1), (3, 3), (4, 1), (4, 2), \dots\}
 \end{aligned}$$

For every $n \in \mathbb{P}$, $D(\{n\})$ is infinite, $D^{\leftarrow}(\{n\})$ is finite.

49

50

Exercise

f^{\leftarrow} is a relation; when is it a function?

Exercise

3.1.9 Find the properties of the *empty relation* $\emptyset \subset S \times S$ and the *universal relation* $U = S \times S$. Assume that S is a nonempty domain.

Exercise

f^{\leftarrow} is a relation; when is it a function?

When f is 1-1 and onto.

Exercise

3.1.9 Find the properties of the *empty relation* $\emptyset \subset S \times S$ and the *universal relation* $U = S \times S$. Assume that S is a nonempty domain.

- (a) \emptyset is (AR), (S), (AS), (T); if $S = \emptyset$ itself then \emptyset is also (R).
- (b) U is (R), (S), (T); if $|S| \leq 1$ then also (AS)

51

52

Examples

Exercise

3.1.10(a) Give examples of relations with specified properties.
(AS), (T), $\neg(R)$.

Examples

Exercise

3.1.10(a) Give examples of relations with specified properties.
(AS), (T), $\neg(R)$.

Examples over \mathbb{N} , $\text{Pow}(\mathbb{N})$:

- strict order of numbers $x < y$
- simple (weak) order, but with some pairs (x, x) removed from \mathcal{R}
- being a prime divisor
 $(p, n) \in \mathcal{R}$ iff p is prime and $p|n$
 - not reflexive: $(1, 1) \notin \mathcal{R}, (4, 4) \notin \mathcal{R}, (6, 6) \notin \mathcal{R}$
 - transitivity is meaningful only for the pairs $(p, p), (p, n), p|n$ for p prime

More Examples

Exercise

3.1.10(b) Give examples of relations with specified properties.
(S), $\neg(R)$, $\neg(T)$.

More Examples

Exercise

3.1.10(b) Give examples of relations with specified properties.
(S), $\neg(R)$, $\neg(T)$.

Easiest examples: inequality

- $\mathcal{R} = \{(x, y) | x \neq y, x, y \in \mathbb{N}\}$
- $\mathcal{R} = \{(A, B) | A \neq B, A, B \subseteq S\}$

3.1.14 Which properties carry from individual relations to their union?

(a) $\mathcal{R}_1, \mathcal{R}_2 \in (R) \Rightarrow \mathcal{R}_1 \cup \mathcal{R}_2 \in (R)$

(b) $\mathcal{R}_1, \mathcal{R}_2 \in (S) \Rightarrow \mathcal{R}_1 \cup \mathcal{R}_2 \in (S)$

(c) $\mathcal{R}_1, \mathcal{R}_2 \in (T) \not\Rightarrow \mathcal{R}_1 \cup \mathcal{R}_2 \in (T)$

Eg. $S = \{a, b, c\}, a\mathcal{R}_1b, b\mathcal{R}_2c$

and no other relationships

- Functions
(co-)domain, image, composition $f \circ g, f^{-1}, f^{\leftarrow}$
- Properties of functions: onto, 1-1
- Matrix operations: transposition, sum, scalar product, product
- Properties of binary relations: (R), (AR); (S), (AS); (T)