

Stability of general class of hairy black holes in $D = 2 + 1$

Weyner Ccuiro,^a David Choque,^b Raúl Rojas^b

^a *Universidad Nacional de San Antonio Abad del Cusco, Av. La Cultura 733, Cusco, Perú.*

^b *Pontificia Universidad Católica de Valparaíso, Instituto de Física, Av. Brasil 2950, Valparaíso, Chile*

E-mail: xxxxx@pucv.cl, david.choque@pucv.cl, raulox.2012@gmail.com

ABSTRACT: We present an exact.

Contents

1	Theory	1
2	Solution	1
3	Thermodynamic stability	2
4	Mechanics stability	2
	References	3

1 Theory

We are interested in asymptotically AdS hairy black hole solutions with a spherical horizon. The action is [1]

$$I[g_{\mu\nu}, \phi] = \frac{1}{2\kappa} \int_{\mathcal{M}} d^3x \sqrt{-g} \left[R - \frac{(\partial\phi)^2}{2} - V(\phi) \right] + \frac{1}{\kappa} \int_{\partial\mathcal{M}} d^2x K \sqrt{-h} \quad (1.1)$$

where $V(\phi)$ is the scalar potential, $\kappa = 8\pi G_N$, and the last term is the Gibbons-Hawking boundary term. Here, h_{ab} is the boundary metric and K is the trace of the extrinsic curvature.

$$I[g_{\mu\nu}, \phi] = \frac{1}{2\kappa} \int_{\mathcal{M}} d^3x \sqrt{-g} [R + L_M] \quad (1.2)$$

$$G_{\alpha\beta} = \frac{1}{2} T_{\alpha\beta} = g_{\alpha\beta} L_M - 2 \frac{\partial L_M}{\partial g^{\alpha\beta}}, \quad L_M = -\frac{(\partial\phi)^2}{2} - V(\phi) \quad (1.3)$$

Then the equation of motion for the metric are

$$G_{\alpha\beta} = \frac{1}{2} T_{\alpha\beta} = \partial_\alpha \phi \partial_\beta \phi - g_{\alpha\beta} \left[\frac{(\partial\phi)^2}{2} + V(\phi) \right] \quad (1.4)$$

And the variation of the action respect to scalar field we have

$$\delta I = \frac{1}{2\kappa} \int d^3x \sqrt{-g} \left[\frac{1}{\sqrt{-g}} \partial_\alpha (\sqrt{-g} g^{\alpha\beta} \partial_\beta \phi) - \frac{\partial V}{\partial \phi} \right] \delta \phi - \frac{1}{2\kappa} \int d^2x \sqrt{-h} n^\alpha \partial_\alpha \phi \delta \phi \quad (1.5)$$

$$\frac{1}{\sqrt{-g}} \partial_\alpha (\sqrt{-g} g^{\alpha\beta} \partial_\beta \phi) - \frac{\partial V}{\partial \phi} = 0 \quad (1.6)$$

2 Solution

$$ds^2 = \Omega(x) \left[-f(x) dt^2 + \frac{\eta^2 dx^2}{f(x)} + d\varphi^2 \right] \quad (2.1)$$

$$E_t{}^t - E_x{}^x = 0 \Rightarrow (\phi')^2 = \frac{3\Omega'^2 - 2\Omega\Omega''}{2\Omega^2} \quad (2.2)$$

$$E_t{}^t - E_\phi{}^\phi = 0 \Rightarrow (\Omega^{1/2} f')' = 0$$

$$E_t{}^t + E_\phi{}^\phi = 0 \Rightarrow V = -\frac{1}{4\eta^2\Omega(x)^3} [2\Omega'' f\Omega + 2f''\Omega^2 - f\Omega'^2 + 3\Omega\Omega' f']$$

$$\Omega(x) = \frac{\nu^2 x^{\nu-1}}{\eta^2 (x^\nu - 1)^2} \quad (2.3)$$

$$f(x) = \frac{1}{L^2} + \frac{\alpha}{2\nu} \left[2\nu + (3-\nu)x^{\frac{3+\nu}{2}} - (3+\nu)x^{\frac{3-\nu}{2}} \right] \quad (2.4)$$

$$\phi(x) = \ell^{-1} \ln x, \quad \ell^{-1} = \frac{\sqrt{2}}{2} \sqrt{\nu^2 - 1} \quad (2.5)$$

And the on-shell potential is

$$V(\phi) = -\frac{3(\alpha L^2 + 1) \exp(-\phi\ell)}{4\nu^2 L^2} \left[(1+\nu) \left(1 + \frac{\nu}{3} \right) \exp(\phi\ell\nu) + (1-\nu) \left(1 - \frac{\nu}{3} \right) \exp(-\phi\ell\nu) \right. \quad (2.6)$$

$$\left. - 2(1-\nu^2) \right] - \frac{\alpha \exp(\phi\ell/2)}{\nu} \left[(1-\nu) \exp\left(\frac{\phi\ell\nu}{2}\right) - (1+\nu) \exp\left(-\frac{\phi\ell\nu}{2}\right) \right] \quad (2.7)$$

$$V(\nu) = V(-\nu), \quad V(\phi) = V(-\phi), \quad V(\phi, \nu) = V(-\phi, -\nu) \quad (2.8)$$

The non-hair limit is well defined in

$$\nu = 1 \Rightarrow V = -\frac{2}{L^2}, \quad \Omega(x) = \frac{1}{\eta^2 (x-1)^2}, \quad f(x) = \frac{1}{L^2} + \alpha(x-1)^2 \quad (2.9)$$

$$V(\phi) = -\frac{2}{L^2} - \frac{\ell^2(\nu^2 - 1)}{4L^2} \phi^2 - \frac{\ell^3(\nu^2 - 1)}{12L^2} \phi^3 - \frac{\phi^4 \ell^4}{192} (\nu^2 - 1) \left[3\alpha(\nu^2 - 9) + \frac{4(\nu^2 - 6)}{L^2} \right] + \mathcal{O}(\phi^4) \quad (2.10)$$

$$x = 1 \pm \frac{1}{\eta r} - \frac{\nu^2 - 1}{24\eta^3 r^3} + \frac{\nu^2 - 1}{24\eta^4 r^4} + \mathcal{O}(r^{-5}) \quad (2.11)$$

For the negative and positive branch respectively

$$\phi(r) = \pm \frac{1}{\eta \ell r} - \frac{1}{2\eta^2 \ell r^2} + \mathcal{O}(r^{-3}) \quad (2.12)$$

3 Thermodynamic stability

4 Mechanics stability

Acknowledgments

Research of AA is supported in part by Fondecyt Grants 1141073 and 1170279. The work of DA is supported by the Fondecyt Grants 1161418 and 1171466. DC is supported by Fondecyt Postdoc Grant 3180185. The work of J.D.E. is supported by the Ministry of Science grant FPA2017-84436-P, Xunta de Galicia ED431C 2017/07, FEDER, and the María de Maeztu Unit of Excellence MDM-2016-0692. He wishes to thank Pontificia Universidad Católica de Valparaíso and Universidad Adolfo Ibáñez for hospitality, during the visit funded by CONICYT MEC 80150093, at the initial stages of the article.

References

- [1] D. Astefanesei, J.L. Blázquez-Salcedo, F. Gómez and R. Rojas, *Thermodynamically stable asymptotically flat hairy black holes with a dilaton potential: the general case*, [2009.01854](#).