

Numerical Methods in Engineering and Applied Science.

Assignment 2.

Student: Vyacheslav Kovalev

$$u'(x_i) \approx \frac{1}{h} \left(\frac{1}{6}u_{i-2} - \frac{4}{3}u_{i-1} + \frac{11}{6}u_i - \frac{5}{3}u_{i+1} + \frac{4}{3}u_{i+2} - \frac{1}{3}u_{i+3} \right) \quad (1)$$

(1). Is (1) a consistent approximation to the first derivative of $u(x)$?

Yes. Answer why in the next point.

(2). Use the Taylor series expansion to find the order of this approximation.

$$u(x_i + sh) = u(x_i) + sh u'(x_i) + \frac{(sh)^2}{2} u''(x_i) + \frac{(sh)^3}{6} u'''(x_i) + O(h^4)$$

Let's consider coefficients at $u(x_i)$.

$$\begin{aligned} u'(x_i) &\approx \frac{1}{h} \left(\frac{1}{6}u_{i-2} - \frac{4}{3}u_{i-1} + \frac{11}{6}u_i - \frac{5}{3}u_{i+1} + \frac{4}{3}u_{i+2} - \frac{1}{3}u_{i+3} \right) = \\ &= \frac{1}{h} \left(\frac{1}{6}u(x_i) - \frac{4}{3}u(x_i) + \frac{11}{6}u(x_i) - \frac{5}{3}u(x_i) + \frac{4}{3}u(x_i) - \frac{1}{3}u(x_i) + O(h) \right) = \frac{1}{h} (O(h)) \end{aligned}$$

Coefficient at $u(x_i)$ is zero.

Performing this procedure for other members in the Taylor series:

$$\begin{aligned} u'(x_i) &\approx \frac{1}{h} \left(\frac{1}{6}u_{i-2} - \frac{4}{3}u_{i-1} + \frac{11}{6}u_i - \frac{5}{3}u_{i+1} + \frac{4}{3}u_{i+2} - \frac{1}{3}u_{i+3} \right) \\ &= u'(x_i) + \frac{h^3 u''''(x_i)}{4} + O(h^4) \end{aligned}$$

Therefore:

$$\begin{aligned} &\frac{1}{h} \left(\frac{1}{6}u_{i-2} - \frac{4}{3}u_{i-1} + \frac{11}{6}u_i - \frac{5}{3}u_{i+1} + \frac{4}{3}u_{i+2} - \frac{1}{3}u_{i+3} \right) - u'(x_i) \\ &= \frac{h^3 u''''(x_i)}{4} + O(h^4) \end{aligned}$$

3rd order of approximation.

While h is small then it is a consistent approximation.

(3).

$$u' - \frac{\sin(x)}{3 + \cos(x)} u = \frac{\cos(2x) + 3 \cos(x)}{3 + \cos(x)}$$

Homogenous:

$$\begin{aligned} u' - \frac{\sin(x)}{3 + \cos(x)} u &= 0 \\ \int \frac{du}{u} &= \int \frac{\sin(x)}{3 + \cos(x)} dx \end{aligned}$$

$$\begin{aligned}\ln(u) &= \left\{ \frac{t = 3 + \cos(x)}{dt = -\sin(x) dx} \right\} = \int \frac{\sin(x)}{t} \frac{dt}{-\sin(x)} = - \int \frac{1}{t} dt \\ &= -\ln(t) + \ln(C) = \ln\left(\frac{C}{t}\right) = \ln\left(\frac{C}{3 + \cos(x)}\right) \\ u_h &= \frac{C}{3 + \cos(x)}\end{aligned}$$

Particular solution of the equation.

$$u' - \frac{\sin(x)}{3 + \cos(x)} u = \frac{\cos(2x) + 3 \cos(x)}{3 + \cos(x)}$$

Let $u = A \sin(x) + B \cos(x)$

$$\begin{aligned}u' - \frac{\sin(x)}{3 + \cos(x)} u &= A \cos(x) - B \sin(x) - \frac{\sin(x) (A \sin(x) + B \cos(x))}{3 + \cos(x)} \\ &= \frac{3A \cos(x) - 3B \sin(x) + A \cos^2(x) - B \sin(x) \cos(x) - \sin^2(x) A - \sin(x) B \cos(x)}{3 + \cos(x)} \\ &= \frac{3A \cos(x) - 3B \sin(x) + A \cos(2x) - 2B \sin(2x)}{3 + \cos(x)} = \frac{\cos(2x) + 3 \cos(x)}{3 + \cos(x)}\end{aligned}$$

get: $B = 0, A = 1$

$$u_p = \sin(x)$$

Summing up:

$$u = \frac{C}{3 + \cos(x)} + \sin(x)$$

Periodic boundary conditions: $x \in [0, 2\pi], u(0) = 0$

$$u(0) = \frac{C}{3 + \cos(0)} + \sin(0) = \frac{C}{3 + 1} = 0 \Rightarrow C = 0$$

Summing up:

$$u = \sin(x), \quad x \in [0, 2\pi]$$

- (4). Use the finite-difference approximation (1) to solve this problem numerically. Plot the exact and the approximate $u(x)$ calculated on a uniform grid with step $h = \pi/10$ over the interval from 0 to 2π .

$$u' - \frac{\sin(x)}{3 + \cos(x)} u = \frac{\cos(2x) + 3 \cos(x)}{3 + \cos(x)}$$

Represent as:

$$\begin{aligned}Au + Bu &= F \\ u^T &= [u_1, u_2, \dots, u_N], \quad u_0 = 0\end{aligned}$$

From the task:

$$h = \frac{2\pi}{N} \Rightarrow N = \frac{2\pi}{h} = 20$$

Keep in mind that $u_0 = 0$ and neglect every u_0 .

$$A = \frac{1}{6h} \begin{bmatrix} 11 & -10 & 8 & -2 & 0 & \dots & 0 & 1 \\ -8 & 11 & -10 & 8 & -2 & 0 & \dots & 0 \\ 1 & -8 & 11 & -10 & 8 & -2 & 0 & \dots & 0 \\ 0 & 1 & -8 & 11 & -10 & 8 & -2 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & 1 & -8 & 11 & -10 & 8 & -2 \\ 0 & \dots & 0 & 1 & -8 & 11 & -10 & 8 & \\ -2 & 0 & \dots & 0 & 1 & -8 & 11 & -10 & \\ 8 & -2 & 0 & \dots & 0 & 1 & -8 & 11 & \end{bmatrix}$$


$$x_i = ih \Rightarrow B = - \begin{bmatrix} \frac{\sin(h)}{3 + \cos(h)} & 0 & \dots & 0 \\ 0 & \frac{\sin(2h)}{3 + \cos(2h)} & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & \frac{\sin(Nh)}{3 + \cos(Nh)} \end{bmatrix}$$

$$F = \begin{bmatrix} \frac{\cos(2h) + 3 \cos(h)}{3 + \cos(h)} \\ \frac{\cos(2 * 2h) + 3 \cos(2h)}{3 + \cos(2h)} \\ \dots \\ \frac{\cos(2 * Nh) + 3 \cos(Nh)}{3 + \cos(Nh)} \end{bmatrix}$$


Solve $(A + B)U = F$:

$$U_{approx} = (A + B)^{-1}F$$

Plot the exact and the approximate $u(x)$ calculated on a uniform grid with step $h = \pi/10$ over the interval from 0 to 2π .



- (5). Vary h from a suitably small to a suitably large value to plot the convergence error norms (pointwise maximum and r.m.s.) as functions of h . Comment on the rate of convergence: compare your numerical observation with your theoretical result.



It is clear that $\max|e_k| \approx O(h)$, $(\log(L_\infty) \approx \log(C) + p \log(h))$, where $p = 1$) So if we divide h by 2, $\max|e_k|$ becomes twice smaller. The numerical result gives the same as theoretical.