Air quality impacts: Impact Pathway Analysis


Milan Ščasný

Charles University

Environment Center

Institute of Economic Studies FSV

Top-down vs. Bottom-up



Phases of pathway	Example
Utilization of sources	Production of electricity
Emissions	SO_2 emissions
Pollution transport	Addition to the atmospheric concentration
Course	Acidification of forest ecosystems
Impact	Disruption of ecosystem stability
Damage	Damages on forest production and non-production functions
Monetary valuation	Market and non - market valuation of forest functions

ExternE project series

Project ExternE = Externalities of Energy launched in 1991, financed by DG Research within the Joule programme


- **airborne pollutants** from power plants, later extended to energy, transport, industry, chemicals, …
- the most developed **bottom-up approach** to calculate the external costs originally for airborne pollutants from power plants
- DPSIR framework: **D**riving force → **P**ressure → **S**tate → **Impact** → **R**esponse
- development of **the Impact Pathway Approach**

  --- I = not physical impact, but **monetised impacts**

Empact Pathway Analysis

- **assessment of impacts** (e.g. on human health), not **pressures** (e.g. emissions, concentrations)

- damage associated with **certain process** depends on
  - ✓ technology
  - ✓ character of operation
  - ✓ site (location)
  - ✓ time
  - ✓ scope of fuel cycle



bottom-up approach
for complex pathways>
‘impact pathway approach’

- **preference structure** of the population

Impact pathway approach

POLLUTANT & NOISE EMISSIONS

TRANSPORT & CHEMICAL TRANSFORMATION


DIFFERENCES OF PHYSICAL IMPATS


MONETARY VALUATION


















PRESSURE


Change in sulfates due to emissions of SO2 emitted from power plant in Lauffen (Stuttgart), Germany



Source: IER

RECEPTORS


Population potentially affected by pollution



Source: IER

IMPACT

Change in premature mortality (LYL, life-year-lost) due to increases in pollution




Source: IER

From Exposure to Response (IMPACT)

Exposure Response Function (ERF) / Dose-Response (DRF) / Concentration-Response (CRF):

For instance, ***Additional Years of Life Lost*** $= 3.9 \cdot 10^{-5} \cdot \Delta Sulfate \cdot Population$



Concentration-response functions

Table 1: Evaluated impacts from airborne pollution and noise exposure and their valuation.

Pollutant	Endpoint (unit)	CRF (per 10 µg/m³)	EU-wide monetary value (€$_{2010}$)
Air pollution: long-term exposure			
$PM_{2.5}$	Adult mortality (loss of life expectancy, per YOLL)	1.062	65,066
PM_{10}	Post-neonatal infant mortality (per case)	1.04	2,762,767
PM_{10}	Incidence of chronic bronchitis in adults (per case)	1.117	60,443
PM_{10}	Prevalence of bronchitis in children (per case)	1.08	663
Air pollution: short-term exposure			
$PM_{2.5}$	Restricted activity days (per day)	1.047	104
$PM_{2.5}$	Work days lost (per day)	1.046	147
$PM_{2.5}$	Hospital admissions, respiratory diseases (per case)	1.019	2503
$PM_{2.5}$	Hospital admissions, cardiovascular diseases (per case)	1.0091	2503
PM_{10}	Incidence of asthma symptoms in asthmatic children (per case)	1.028	47
O_3	Adult mortality (per YOLL)	1.0029	65,066
O_3	Hospital admissions (65+years), respiratory diseases (per case)	1.0044	2503
O_3	Hospital admissions (65+years), cardiovascular diseases (per case)	1.0089	2503
O_3	Minor restricted activity days (per day)	1.0154	47
Noise exposure		ERF (per dB L_{den})	
	Slightly annoyed (per year)	$0.02815 * L_{den}^2 - 1.130 * L_{den} + 11.477$	56
	Annoyed (per year)	$0.03270 * L_{den}^2 - 2.121 * L_{den} + 36.854$	112
	Highly annoyed (per year)	$0.02523 * L_{den}^2 - 1.886 * L_{den} + 36.307$	187


CRF, Concentration-response function; ERF, exposure-response function; YOLL, year of life lost; L_{den}, weighted day-evening-night noise level.
Source: adapted from (5, 6).

Máca, Melichar, Ščasný, Braun Kohlová *Rev Enviro Health* 2017

Impacts included in the current ExternE projects

Pollutant/burden

- Particulate matters
- SO_2, NOx, NH3, VOC
- CO_2
- Heavy metals
- Noise
- Odour



Impact category

- Human health
  - morbidity
  - mortality
  - development imp IQ
- Building materials
- Crops
- Climate change

- Landscape Amenities
- Visibility
- Cultural heritage

Put Monetary Values on Impacts

elicit preference structure of the population

How much are you willing-to-pay to avoid adverse impact (or receive desired impact) ?

- **market price**

  -- building materials, crops, medical treatment,…

- **non-market valuation**

  -- suffer due to illness, premature death, recreation, aesthetic values, landscape amenities,….

JEM218 "Environmental Economics and Behavior"

Put Monetary Values on Health Impacts

I. Premature mortality

- WTP for reducing the risk of dying
- WTP for changing life expectancy is very difficult to derive
- WTP per Quality-Adjusted-Life-Yer (or Disability-….) is not conclusive, it depends on the context and cause

II. Morbidity, illness

- Medical treatment costs (direct & indirect) + Loss of earning = Cost-of-illness
- WTP for suffer, pain, inconveniences


III. Development impairment, (in-)fertility

- WTP for reducing the risk of VLBW, birth defects, chance to conceive, IQ
- Cost of IVF, cost of training, loss of earnings (IQ)

IV. Mental health


- WTP studies

External costs of electricity generation, CZE




Melichar, Máca, Ščasný, 2012

External costs of electricity generation in the Czech republic, Kč per kWh (2008 prices)




Sources to premature deaths in DK in 2021




Approx. 3,000 premature deaths

Approx. 1,000 premature deaths





Approx. 220 premature deaths



Approx. 70 premature deaths

Approx. 70 premature deaths



DK's contribution to premature deaths abroad approx 2,300

Approx. 330 premature deaths (SNAP2_households) Approx. 240 from wood stoves)

Ellermann et al, 2023. Luftkvalitet 2021

Contribution from pollutants

EVAv7.1 – Average for periode of 2019-2021 for Denmark

Health effect	SO_2	O_3	NO_2	$PM_{2.5}$	Total
Premature death (short-term exposure)	~25	~85	~14	~302	~427
Premature deaths (**long-term exposure**)			~15	~3,440	~3,460
Total	**~25**	**~85**	**~30**	**~3,740**	**~3,890**

- $PM_{2.5}$ is the dominant pollutant to premature death
- $PM_{2.5}$ includes primary emitted particles and secondary particles
- Secondary particles like ammonium nitrate and ammonium sulfate are formed in the atmosphere from emissions of gasses like NO_x, SO_2 and NH_4
- Secondary particles dominate $PM_{2.5}$

Ellermann et al, 2023. Luftkvalitet 2021

Local versus long-range contributions


- What is a city's contribution to itself? And what are the largest local sources?
- Premature death due to local sources: Copenhagen (12%, 5%, 9%), Frederiksberg (3%), Aarhus (8%), Odense (11%), Region Hovedstaden (11%)
- Two largest sources are road traffic and wood stoves.


Table 1.2. Percentage of premature deaths in Nordic capital cities in 2030 distributed on emission contributions from city to city, from country and from abroad

	From city	From country	From abroad	Total
København	15%	23%	63%	100%
Stockholm	35%	18%	47%	100%
Helsinki	25%	21%	53%	100%
Reykavík	43%	27%	30%	100%

Jensen et al., 2023. Air Quality, Health Effects and External Costs in Selected Cities in Nordic Countries. TemaNord 2023:548

Economic standing of envi/health externalities





Which impacts should BCA include?

- on country's residents
- in the EU
- worldwide

Which values to attach to the impact (*i.e. social planner perspective*)?

- CZ values
- EU values
- Country-specific WTPs

Figure 2: Total external costs by the variant of mining limit revision and policy-making perspective.

Máca, Melichar, Ščasný, Braun Kohlová *Rev Enviro Health* 2017

Environmental Economics

Economics of Climate Change: Carbon & Greenhouse effects

Milan Ščasný


Charles University

Environment Center

Institute of Economic Studies FSV


WHY GREENHOUSE?

Greenhouse Effect



Obrázek 5: Zjednodušený model skleníkového efektu.
Zdroj: Le Treut et al. (2007).

The greenhouse effect



Why carbon dioxide matters?

- Carbon dioxide is Earth's most important greenhouse gas: **a gas that absorbs and radiates heat.** Unlike oxygen or nitrogen (which make up most of our atmosphere), **greenhouse gases absorb heat radiating from the Earth's surface and re-release it in all directions**—including back toward Earth's surface.
  - Without carbon dioxide, Earth's natural greenhouse effect would be too weak to keep the average global surface temperature above **freezing**.
  - By adding more carbon dioxide to the atmosphere, people are supercharging the natural greenhouse effect, causing **global temperature to rise**.
  - According to observations by the NOAA Global Monitoring Lab, in 2021 **carbon dioxide (CO_2)** alone was responsible for *about two-thirds* of the total heating influence of all human-produced greenhouse gases.

Greenhouse effect

Primary human-produced **greenhouse gases**

- **carbon dioxide (CO_2)**
- **methane (CH4)**
- **nitrous oxide (N_2O)**
- chlorofluorocarbons (CFCs), (SFs, black carbon, HFCs)
- all usually expressed in CO_2 (or C) equivalent

Greenhouse gasses are gases in an atmosphere that **absorb and emit radiation** within the thermal infrared range. In the 1860s Irish physicist John Tyndall proved that the Earth's atmosphere has a **greenhouse effect**.

Greenhouse effect causes **climate change** (sometimes *"global warming"*).

Climate change is more about **an excess of energy** (the cause), rather than about "warming" (a consequence)

GWP - Global warming potential

is a measure of **how much a given mass of greenhouse gas is estimated to contribute to global warming**.

It is a relative scale which compares the gas in question to that of **the same mass of CO_2 [carbon dioxide]** whose GWP is by definition **1**.

A GWP is calculated **over a specific time interval** and the value of this must be stated whenever a GWP is quoted or else the value is meaningless.


For example, the GWP for methane over 100 years is 25 and for N_2O 298. This means that emissions of 1 million metric tones of methane and nitrous oxide respectively is equivalent to emissions of 25 and 298 million metric tones of carbon dioxide

	Lifetime (years)	GWP time horizon		
		20 years	100 years	500 years
Methane (CH4)	12	72	25	7.6
Nitrous oxide (N2O)	114	289	298	153
HFC-23 (hydrofluorocarbon)	270	12000	14800	12200
HFC-134a (hydrofluorocarbon)	14	3830	1430	425
sulfur hexafluoride	3200	16300	22800	32600

GWP values and lifetimes from 2007 IPCC AR4

IS ATMOSHERIC CO2 (CONCENTRATION) RISING?

ATMOSPHERIC CARBON DIOXIDE




Based on the annual report from NOAA's Global Monitoring Lab, **global average atmospheric carbon dioxide was 419.3 parts per million in 2023** (417.06 ppm in 2022), setting a new record high.

The increase between 2021 and 2022 was 2.13 ppm—**the 12th year in a row** where the amount of carbon dioxide in the atmosphere **increased by more than 2 ppm**.

Update: **November 1, 2025**, the daily average at the Mauna Loa Observatory was **425.68 ppm** (notes the NOAA Global Monitoring Laboratory)

This graph shows the station's monthly average carbon dioxide measurements since 1958 in parts per million (ppm). **The seasonal cycle of highs and lows** (small peaks and valleys) is driven by Northern Hemisphere summer vegetation growth, which reduces atmospheric carbon dioxide, and winter decay, which increases it. **The long-term trend of rising** carbon dioxide levels is driven by human activities. At Mauna Loa, the highest monthly value each year occurs in May. In May 2023, carbon dioxide hit **424 ppm**—**a new record**.

Source: https://www.climate.gov/news-features/understanding-climate/climate-change-atmospheric-carbon-dioxide (Lindsey April 2024)




Based on the annual report from NOAA's Global Monitoring Lab, **global average atmospheric carbon dioxide was 422.7 parts per million in 2024** (417.06 ppm in 2022).

The increase over 2023 amounts was **3.75 ppm**—the largest one-year increase on record

Update: **November 1, 2025**, the daily average at the Mauna Loa Observatory was **425.68 ppm** (notes the NOAA Global Monitoring Laboratory)

This graph shows the station's monthly average carbon dioxide measurements since 1958 in parts per million (ppm). **The seasonal cycle of highs and lows** (small peaks and valleys) is driven by Northern Hemisphere summer vegetation growth, which reduces atmospheric carbon dioxide, and winter decay, which increases it. **The long-term trend of rising** carbon dioxide levels is driven by human activities. At Mauna Loa, the highest monthly value each year occurs in May. In May 2024, carbon dioxide hit **427 ppm**—**a new record**.


Source: https://www.climate.gov/news-features/understanding-climate/climate-change-atmospheric-carbon-dioxide (Lindsey April 2024)



Atmospheric carbon dioxide (ppm) for the past 800,000 years based on ice-core data (light purple line) compared to 2022 concentration (bright purple dot). The peaks and valleys in the line show ice ages (low CO_2) and warmer interglacials (higher CO_2). Throughout that time, CO_2 **was never higher than 300 ppm** (light purple dot, between 300,000 and 400,000 years ago). The increase over the last 60 years is **100 times faster than previous natural increases**. In fact, on the geologic time scale, the increase from the end of the last ice age to the present (dashed purple line) looks virtually instantaneous. Graph by NOAA Climate.gov based on data from Lüthi, et al., 2008, via NOAA NCEI.


IS GLOBAL TEMPERATURE CHANGING ?


Earth's Temperature



Obrázek 3: Rekonstrukce průběhu teploty v geologické minulosti Země, nehomogenní časová osa.
Poznámka: Km = Kambrium, O = Ordovik, S = Silur, D = Devon, K = Karbon, P = Perm, Tr = Trias, J = Jura, Kř = Křída, Pal = Paleocén, Eo = Eocén, Ol = Oligocén, Mio = Miocén.
Zdroj: Zachos et al. (2001).


Hockey stick dispute (zoom in)




GFS 1-day Avg
2m Temperature (°C)
Sunday, Feb 25, 2018
50
40
30
20
10
0
-10
-20
-30
-40
-50
-60
ClimateReanalyzer.org
University of Maine | Climate Change Institute

Average temperature anomaly, Global

Global average land-sea temperature anomaly relative to the 1961-1990 average temperature.



Data source: Met Office Hadley Centre (2023)

OurWorldInData.org/co2-and-greenhouse-gas-emissions | CC BY

Note: The gray lines represent the upper and lower bounds of the 95% confidence intervals.

GREENHOUSE EFFECT RESULTS IN EARTH ENERGY IMBALANCE




Figure 8 Schematic presentation on the Earth heat inventory for the current anthropogenically driven positive Earth energy imbalance at the top of the atmosphere. **The relative partition (in %) of the Earth heat inventory** presented in Fig. 6 for the different components is given for the ocean (upper: 0–700 m, intermediate: 700–2000 m, deep: >2000 m), land, cryosphere (grounded and floating ice) and atmosphere, for the periods **1971–2018** and **2010–2018** (for the latter period values are provided in parentheses), as well as for the EEI. The total heat gain (in red) over the period 1971–2018 is obtained from the Earth heat inventory as presented in Fig. 6. **To reduce the 2010–2018 EEI of 0.87±0.12 W m^{-2} towards zero, current atmospheric CO_2 would need to be reduced by −57±8 ppm**.

Source: von Schuckmann, K., Cheng, L., Palmer, M. D., Hansen, J., Tassone, C., Aich, V., Adusumilli, S., Beltrami, H., Boyer, T., Cuesta-Valero, F. J., Desbruyères, D., Domingues, C., García-García, A., Gentine, P., Gilson, J., Gorfer, M., Haimberger, L., Ishii, M., Johnson, G. C., Killick, R., King, B. A., Kirchengast, G., Kolodziejczyk, N., Lyman, J., Marzeion, B., Mayer, M., Monier, M., Monselesan, D. P., Purkey, S., Roemmich, D., Schweiger, A., Seneviratne, S. I., Shepherd, A., Slater, D. A., Steiner, A. K., Straneo, F., Timmermans, M.-L., and Wijffels, S. E.: Heat stored in the Earth system: where does the energy go?, Earth Syst. Sci. Data, 12, 2013–2041, https://doi.org/10.5194/essd-12-2013-2020, 2020.





Figure 1 Ensemble mean time series and ensemble standard deviation (2σ, shaded) of **global ocean heat content (OHC) anomalies** relative to the 2005–2017 climatology for the 0–300 m (gray), 0–700 m (blue), 0–2000 m (yellow) and 700–2000 m depth layer (green). The ensemble mean is an outcome of an international assessment initiative, and all products used are referenced in the legend of Fig. 2. The trends derived from the time series are given in Table 1. Note that values are given for the ocean surface area between 60∘ S and 60∘ N and are limited to the 300 m bathymetry of each product.

Source: von Schuckmann, K., Cheng, L., Palmer, M. D., Hansen, J., Tassone, C., Aich, V., Adusumilli, S., Beltrami, H., Boyer, T., Cuesta-Valero, F. J., Desbruyères, D., Domingues, C., García-García, A., Gentine, P., Gilson, J., Gorfer, M., Haimberger, L., Ishii, M., Johnson, G. C., Killick, R., King, B. A., Kirchengast, G., Kolodziejczyk, N., Lyman, J., Marzeion, B., Mayer, M., Monier, M., Monselesan, D. P., Purkey, S., Roemmich, D., Schweiger, A., Seneviratne, S. I., Shepherd, A., Slater, D. A., Steiner, A. K., Straneo, F., Timmermans, M.-L., and Wijffels, S. E.: Heat stored in the Earth system: where does the energy go?, Earth Syst. Sci. Data, 12, 2013–2041

Another reason carbon dioxide is important in the Earth system is that it **dissolves into the ocean like the fizz in a can of soda**. It reacts with water molecules, producing carbonic acid and lowering the ocean's pH (raising its acidity). Since the start of the Industrial Revolution, the pH of the ocean's surface waters has dropped from 8.21 to 8.10. This drop in pH is called ***ocean acidification***.



(**left**) A healthy ocean snail has a transparent shell with smoothly contoured ridges.

(**right**) A shell exposed to more acidic, corrosive waters is cloudy, ragged, and pockmarked with 'kinks' and weak spots. Source: https://www.climate.gov/news-features/understanding-climate/climate-change-atmospheric-carbon-dioxide

HOW CLIMATE CHANGE SCENARIOS ARE DEFINED THEN ?

How scenarios are defined? What SSP & RCPs Are?

- **Baseline scenario** ⇒ *"A Middle-of-the-road Scenario" (**SSP2**)*
  - **Shared Socio-economic Pathway** built on the assumption of continuation of all major trends that we observe today (Ebi et al. 2014; Van Vuuren et al. 2014)
  - projects current trends into the future, without major changes in economic growth, use and availability of resources, technological trends, population growth, economic and envi policies


- **Climate change mitigation policy scenarios** ⇒ represent the challenges of reaching two long term radiative forcing targets corresponding to **Representative Concentration Pathways (RCPs)**, as in van Vuuren et al. 2011:

  - **RCP2.6** – radiative forcing is declining to **2.6 W/m^2** by 2100, ≈ ***490 ppm CO_2-eq***
  - **RCP4.5** - radiative forcing is declining to **4.5 W/m^2** by 2100, ≈ ***650 ppm CO_2-eq***
  - **RCP6.0** - radiative forcing is **6.0 W/m^2** in 2100 ≈ ***850 ppm CO_2-eq***; close to SSP2
  - RCP8.5 …

RCPs & Global Temperature Increase

AR5 global warming increase (°C) projections *(IPCC 2014)*

	2046-2065	2081-2100
Scenario	Mean and *likely* range	Mean and *likely* range
RCP2.6	1.0 (0.4 to 1.6)	1.0 (0.3 to 1.7)
RCP4.5	1.4 (0.9 to 2.0)	1.8 (1.1 to 2.6)
RCP6.0	1.3 (0.8 to 1.8)	2.2 (1.4 to 3.1)
RCP8.5 (close to SSP2)	**2.0** (1.4 to 2.6)	**3.7** (2.6 to 4.8)




NOAA Climate.gov, adapted from IPCC AR6 Technical Summary, Figure TS.4

Plausible future socioeconomic pathways for **annual carbon dioxide emissions** (left) and **the resulting atmospheric carbon dioxide concentrations** (right) through the end of the century. A *shared socioeconomic pathway* is an internally consistent set of assumptions about future population growth, global and regional economic activity, and technological advances. Models use these pathways to project a range of possible future carbon dioxide emissions; for simplicity, the image only shows the only the mean value. NOAA Climate.gov graphic adapted from figure TS.4 in the IPCC Sixth Assessment Report Technical Summary.

Climate Change Policy

- UNFCCC → 1996 Kyoto Protocol
  - Annex I, Annex II countries, DCM, JI
- UN Annual Climatic Conferences
  - **COP 21, Paris, Nov-Dec 2015**
    - a long-term goal of keeping the increase in global average temperature to **well below 2°C** above pre-industrial levels;
    - to aim to limit the increase to **1.5°C**, since this would significantly reduce risks and the impacts of climate change;
    - on the need for **global emissions to peak as soon as possible**, recognising that this will take longer for developing countries;
    - to undertake **rapid reductions thereafter** in accordance with the best available science.
    - Before and during the Paris conference, countries submitted comprehensive **national climate action plans** (INDCs). These are not yet enough to keep global warming below 2°C, but the agreement traces the way to achieving this target.
- **The Paris Agreement** has been ratified Nov 4th, 2016
- COP 22 (Marrakesh, 2016), COP 23 (Bonn, 2017), COP 24 (Katowice, 2018), COP 25 (Madrid, 2019), *"COVID-2020"*, COP 26 (Glasgow, 2021), COP 27 (Cairo, 2022), COP 28 (Dubai, 2023), COP 29 (Baku, 2024), COP 30 (Brasil, 2025)

A well known fact





Source: IPCC AR5 WG II (2014)


The recent Paris NDCs, or even a fully successful post-Paris mitigation process will still leave unavoidable climate change impacts → our societies will have to **«adapt»**


Bosello (2017)

Voluntary readings


Carbon Dioxide
(CO_2)
Parts per million (ppm)
410
400
390
380
370
360
350
340
330
1975 1980 1985 1990 1995 2000 2005 2010 2015 2020


Nitrous Oxide
(N_2O)
Parts per billion (ppb)
335
330
325
320
315
310
305
300
295
1975 1980 1985 1990 1995 2000 2005 2010 2015 2020


Methane
(CH_4)
Parts per billion (ppb)
1900
1850
1800
1750
1700
1650
1600
1550
1975 1980 1985 1990 1995 2000 2005 2010 2015 2020


CFC-12
CFC-11
HCFC-22
HFC-134a
Parts per trillion (ppt)
600
500
400
300
200
100
0
1975 1980 1985 1990 1995 2000 2005 2010 2015 2020

GHG emissions (and a bit of chemistry)

Greenhouse gases

- carbon dioxide (CO_2)
- methane (CH4)
- nitrous oxide (N_2O)
- chlorofluorocarbons (CFCs), (SFs, black carbon, HFCs)

Molecular weight

CO_2 = 1*__Carbon__ + 2*__0xygen__ = 1*12 + 2*16 = **44**


C share: 12/44 *(=0.2727)* t **C** in 1 t **CO_2**

1 t **C** ≈ 44/12 *(3.667)* t of **CO_2**

100$ per t **C** = 27.27$ per t **CO_2**


100$ per t **CO_2** = 366.67 $ per t **C**

Pre-1978 changes in the CO_2-equivalent abundance and AGGI



Pre-1978 changes in the CO_2-equivalent abundance and AGGI based on the ongoing measurements of all greenhouse gases reported here, measurements of CO_2 going back to the 1950s from C.D. Keeling *[Keeling et al., 1958]*, and atmospheric changes derived from air trapped in ice and snow above glaciers *[Machida et al., 1995, Battle et al., 1996, Etheridge, et al., 1996; Butler, et al., 1999]*. Equivalent CO_2 atmospheric amounts (in ppm) are derived with the relationship (Table 1) between CO_2 concentrations and radiative forcing from all long-lived greenhouse gases. **AGGI = Annual GHG Gas Index** tracks the increasing amount of heat being added to the atmosphere by human-related GHG emissions.


Source: https://www.esrl.noaa.gov/gmd/aggi/aggi.html




The amount of carbon dioxide in the atmosphere (blue line) has increased along with human emissions (gray line) since the start of the Industrial Revolution in 1750. Emissions rose slowly to about **5 gigatons**—one gigaton is a billion metric tons—per year in the mid-20th century before skyrocketing to more than **35 billion tons per year** by the end of the century.

NOAA Climate.gov graph, adapted from original by Dr. Howard Diamond (NOAA ARL). Atmospheric CO_2 data from NOAA and ETHZ. CO_2 emissions data from Our World in Data and the Global Carbon Project.

Why athmospheric carbon (concentration) keep growing even when carbon emissions are (hope) stabilizing?




Per capita CO_2 emissions by sector, World, 2020



Data source: Climate Watch (2023); Population based on various sources (2023)
OurWorldInData.org/co2-and-greenhouse-gas-emissions | CC BY

Methane emissions by sector, World

Methane (CH_4) emissions are measured in tonnes of carbon dioxide-equivalents[1].




Data source: Climate Watch (2023)

OurWorldInData.org/co2-and-greenhouse-gas-emissions | (

1. Carbon dioxide equivalents (CO_2eq): Carbon dioxide is the most important greenhouse gas, but not the only one. To capture all greenhouse gas emissions, researchers express them in "carbon dioxide equivalents" (CO_2eq). This takes all greenhouse gases into account, not just CO_2. To express all greenhouse gases in carbon dioxide equivalents (CO_2eq), each one is weighted by its global warming potential (GWP) value. GWP measures the amount of warming a gas creates compared to CO_2. CO_2 is given a GWP value of one. If a gas had a GWP of 10 then one kilogram that gas would generate ten times the warming effect as one kilogram of CO_2. Carbon dioxide equivalents are calculated for each gas by multiply the mass of emissions of a specific greenhouse gas by its GWP factor. This warming can be stated over different timescales. To calculate CO_2eq over 100 years, we'd multiply each gas by its GWP over a 100-year timescale (GWP100). Total greenhouse gas emissions – measured in CO_2eq – are then calculated by summing each gas' CO_2eq value.

Nitrous oxide emissions by sector, World

Nitrous oxide (N_2O) emissions are measured in tonnes of carbon dioxide equivalents[1].



Data source: Climate Watch (2023)


OurWorldInData.org/co2-and-greenhouse-gas-emissions | CC BY

1. Carbon dioxide equivalents (CO_2eq): Carbon dioxide is the most important greenhouse gas, but not the only one. To capture all greenhouse gas emissions, researchers express them in "carbon dioxide equivalents" (CO_2eq). This takes all greenhouse gases into account, not just CO_2. To express all greenhouse gases in carbon dioxide equivalents (CO_2eq), each one is weighted by its global warming potential (GWP) value. GWP measures the amount of warming a gas creates compared to CO_2. CO_2 is given a GWP value of one. If a gas had a GWP of 10 then one kilogram of that gas would generate ten times the warming effect as one kilogram of CO_2. Carbon dioxide equivalents are calculated for each gas by multiplying the mass of emissions of a specific greenhouse gas by its GWP factor. This warming can be stated over different timescales. To calculate CO_2eq over 100 years, we'd multiply each gas by its GWP over a 100-year timescale (GWP100). Total greenhouse gas emissions – measured in CO_2eq – are then calculated by summing each gas' CO_2eq value.

Greenhouse gas emissions by sector, World

Our World in Data


Greenhouse gas emissions[1] are measured in tonnes of carbon dioxide-equivalents[2] over a 100-year timescale.



Data source: Climate Watch (2023)

OurWorldInData.org/co2-and-greenhouse-gas-emissions | CC BY


1. Greenhouse gas emissions: A greenhouse gas (GHG) is a gas that causes the atmosphere to warm by absorbing and emitting radiant energy. Greenhouse gases absorb radiation that is radiated by Earth, preventing this heat from escaping to space. Carbon dioxide (CO_2) is the most


Carbon dioxide emissions per person
Per person emissions (tCO2e)
20
15
10
5
0
Countries with largest total emissions
Areas of rectangles
show total emissions
Saudi Arabia
Australia
USA
Canada
S. Korea
Russia
Iran
Japan
Germany
China
S. Africa
Turkey
Brazil
Indonesia
India
. . .
1B
2B
3B
4B
Population (shown by widths of rectangles)

Total greenhouse gas emissions per capita, 2020

Our World in Data

Greenhouse gas emissions[1] are measured in tonnes of carbon dioxide-equivalents[2] per person. Contributions from land-use change and forestry are included.




Data source: Climate Watch (2023); Population based on various sources (2023)
OurWorldInData.org/co2-and-greenhouse-gas-emissions | CC BY

1. Greenhouse gas emissions: A greenhouse gas (GHG) is a gas that causes the atmosphere to warm by absorbing and emitting radiant energy.

Per capita greenhouse gas emissions, 2021

Our World in Data

Greenhouse gas emissions[1] include carbon dioxide, methane and nitrous oxide from all sources, including land-use change. They are measured in tonnes of carbon dioxide-equivalents[2] over a 100-year timescale.




Data source: Jones et al. (2023); Population based on various sources (2023)
Note: Land-use change emissions can be negative.
OurWorldInData.org/co2-and-greenhouse-gas-emissions | CC BY

1. Greenhouse gas emissions: A greenhouse gas (GHG) is a gas that causes the atmosphere to warm by absorbing and emitting radiant energy. Greenhouse gases absorb radiation that is radiated by Earth, preventing this heat from escaping to space. Carbon dioxide (CO_2) is the most

CO_2 emissions per capita vs GDP per capita, 2018

This measures CO_2 emissions from fossil fuels and cement production only – land use change is not included. Gross domestic product (GDP) per capita is measured in international-$ in 2011 prices to adjust for price differences between countries and adjust for inflation.

Our World in Data



Source: Global Carbon Project; Maddison Project Database 2020 (Bolt and van Zanden (2020))
OurWorldInData.org/co2-and-other-greenhouse-gas-emissions/ • CC BY