

Part 1: Groups

0 Preliminaries

- 0.1 Properties of Integers
- 0.2 Modular Arithmetic
- 0.3 Complex Numbers
- 0.4 Mathematical Induction
- 0.5 Equivalence Relations
- 0.6 Functions (Mappings)
- 0.7 Exercise

1 Introduction to Groups

- 1.1 Symmetries of a Square
- 1.2 The Dihedral Groups
- 1.3 Bibliography of Niels Abel

2 Groups

- 2.1 Definition and Examples of Groups
- 2.2 Elementary Properties of Groups
- 2.3 Historical Note
- 2.4 Exercises

3 Finite Groups; Subgroups

- 3.1 Terminology and Notation
- 3.2 Subgroup Tests
- 3.3 Examples of Subgroups
- 3.4 Exercises

4 Cyclic Groups

- 4.1 Properties of Cyclic Groups
- 4.2 Classification of Subgroups of Cyclic Groups
- 4.3 Exercise
- 4.3 Bibliography of James Joseph Sylvester

5 Permutation Groups

- 5.1 Definitions and Notation
- 5.2 Cycle Notation
- 5.3 Properties of Permutations
- 5.4 A Check-Digit Scheme Based on D_5
- 5.5 Exercise
- 5.5 Bibliography of Augustin Cauchy
- 5.6 Bibliography of Alan Turing

6 Isomorphisms

- 6.1 Motivation
- 6.2 Definition and Examples
- 6.3 Properties of Isomorphisms
- 6.4 Automorphisms
- 6.5 Cayley's Theorem
- 6.6 Exercise
- 6.7 Bibliography of Arthur Cayley

7 Cosets and Lagrange's Theorem

- 7.1 Properties of Cosets
- 7.2 Lagrange's Theorem and Consequences
- 7.3 An Application of Cosets to Permutation Groups
- 7.4 The Rotation Group of a Cube and a Soccer Ball

7.5 An Application of Cosets to the Rubik's Cube

7.6 Exercises

7.7 Bibliography of Joseph Lagrange

8 External Direct Products

8.1 Definition and Examples

8.2 Properties of External Direct Products

8.3 The Group of Units Modulo n as an External Direct Product

8.4 RSA Public Key Encryption Scheme

8.5 Exercises

8.6 Bibliography of Leonard Adleman

9 Normal Subgroups and Factor Groups

9.1 Normal Subgroups

9.2 Factor Groups

9.3 Applications of Factor Groups

9.4 Internal Direct Products

9.5 Exercises

9.6 Bibliography of Evariste Galois

10 Group Homomorphisms

10.1 Definition and Examples

10.2 Properties of Homomorphisms

10.3 The First Isomorphism Theorem

10.4 Exercises

10.5 Bibliography of Camille Jordan

11 Fundamental Theorem of Finite Abelian Groups

11.1 The Fundamental Theorem

11.2 The Isomorphism Classes of Abelian Groups

11.3 Proof of the Fundamental Theorem

11.4 Exercises

0 Preliminaries

0.1 Properties of Integers

Universal Product Code (UPC)

$$(a_1, a_2, \dots, a_{12}) \cdot (3, 1, 3, 1, \dots, 3, 1) \pmod{10} = 0.$$

The 10-digit International Standard Book Number (ISBN-10) has the property

$$(a_1, a_2, \dots, a_{10}) \cdot (10, 9, 8, 7, 6, 5, 4, 3, 2, 1) \pmod{11} = 0. \text{ As for } a_{10}, X \text{ stands for } 10.$$

0.2 Modular Arithmetic

Logic Gates & modulo 2 arithmetic

0.3 Complex Numbers

norm of $a+bi$

Closure under division

conjugation

0.4 Mathematical Induction

e.g. Prove that $2^n 3^{2n} - 1$ is always divisible by 17.

0.5 Equivalence Relations

1. reflexive property: $a \sim a$.
2. symmetric property: $a \sim b \Rightarrow b \sim a$.
3. transitive property: $a \sim b, b \sim c \Rightarrow a \sim c$.

e.g.

- $(a, b) \cong (c, d)$ if $ad = bc, b, d \neq 0$.

Partition

0.6 Functions (Mappings)

To verify that a correspondence is a function:

$$x_1 = x_2 \Rightarrow \phi(x_1) = \phi(x_2).$$

One-to-one function:

$$\phi(x_1) = \phi(x_2) \Rightarrow a_1 = a_2.$$

Function from A onto B

Properties

1. associativity: $\gamma(\beta\alpha) = (\gamma\beta)\alpha$.
2. If α and β is one-to-one, then $\beta\alpha$ is one-to-one.
3. If α and β is onto, then $\beta\alpha$ is onto.
4. If α is one-to-one and onto, then there is a function α^{-1} from B onto A such that $(\alpha^{-1}\alpha)(a) = a$ for all a in A and $(\alpha\alpha^{-1})(b) = b$ for all b in B .

0.7 Exercise

1. If $a \bmod st = b \bmod st$, show that $a \bmod s = b \bmod s$ and $a \bmod t = b \bmod t$. The converse is true if s and t are relatively prime.
2. If n is an integer greater than 1 and $(n - 1)! = 1 \bmod n$, prove that n is prime.
3. Prove that 3, 5, and 7 are the only three consecutive odd integers that are prime.

1 Introduction to Groups

1.1 Symmetries of a Square

Cayley table

- closure
- identity
- inverse
- associativity

commutative (Abelian)

1.2 The Dihedral Groups

~~cross-cancellation~~

1.3 Bibliography of Niels Abel

2 Groups

2.1 Definition and Examples of Groups

Group	Operation	Identity	Form of Element	Inverse	Abelian
$GL(n, F)$	Matrix multiplication	E	$ A \neq 0$		No
$SL(n, F)$	Matrix multiplication	E	$ A = 1$		No
$U(n)$	Multiplication mod n	1	$\gcd(k, n) = 1$		Yes
\mathbb{R}^n	Componentwise addition	$(0, 0, \dots, 0)$	(a_1, a_2, \dots, a_n)		Yes

2.2 Elementary Properties of Groups

- Uniqueness of the Identity
- Cancellation
- Uniqueness of Inverses
- Socks-Shoes Property: $(ab)^{-1} = b^{-1}a^{-1}$.

2.3 Historical Note

2.4 Exercises

1. Left-right cancellation implies commutativity, and [cross cancellation](#) implies [Abelian](#) property.
2. Law of Exponents for Abelian Groups: $(ab)^n = a^n b^n$.
3. $ab = ba \iff (ab)^2 = a^2 b^2 \iff (ab)^{-2} = b^{-2} a^{-2}$.
4. Suppose F_1 and F_2 are distinct reflections in a dihedral group D_n . Prove that $F_1 F_2 \neq R_0$. If $F_1 F_2 = F_2 F_1$, then $F_1 F_2 = R_{180}$.

3 Finite Groups; Subgroups

3.1 Terminology and Notation

Order of a group

Order of an element

Subgroup

Proper subgroup: $H \subset G$.

3.2 Subgroup Tests

- To prove that a subset is a subgroup
 - **One-Step Test:** $ab^{-1} \in H$.
 - **Two-Step Test:** $ab, a^{-1} \in H$.
 - **Finite Subgroup Test:** $ab \in H$.
- To prove that a subset is not a subgroup
 - Show that the **identity** is not in the set.
 - Exhibit an element of the set whose **inverse** is not in the set.
 - Exhibit two elements of the set whose **product** is not in the set.

3.3 Examples of Subgroups

- $\langle a \rangle$ is an Abelian subgroup, where a is called a *generator* of G .
- $\langle S \rangle$ is the smallest subgroup of G containing S .
- Gaussian Integers: $\langle 1, i \rangle = \{a + bi \mid a, b \in \mathbb{Z}\}$.
- Center is a subgroup. $Z(G) = \{a \in G \mid ax = xa \text{ for all } x \text{ in } G\}$.
- For $n \geq 3$,

$$Z(D_n) = \begin{cases} \{R_0, R_{180}\}, & n \text{ is even,} \\ \{R_0\}, & n \text{ is odd.} \end{cases}$$

- Centralizer of a in G is a subgroup: $C(a) = \{g \in G \mid ga = ag\}$.
- Centralizer of H in G is a subgroup: $C(H) = \{g \in G \mid xh = hx \text{ for all } h \in H\}$.
- $Z(G) \in C(a)$, $Z(G) = \bigcap_{a \in G} C(a)$.
- G is Abelian if and only if $C(a) = G$ for all a in G .

3.4 Exercises

1. For elements a, b in group \mathbb{Z}_n , $|a + b| = (|a| + |b|) \mod n$.
2. Prove that if a is the only element of order 2 in a group, then a lies in the center of the group.
Proof. $(x^{-1}ax)^2 = x^{-1}ax = a \Rightarrow ax = xa$.
3. No group is the union of two proper subgroups, but some groups are the union of three proper subgroups.
4. Let G be a group and let H be a subgroup of G . For any fixed x in G , define the **conjugate** of H : $xHx^{-1} = \{xhx^{-1} \mid h \in H\}$, which preserves structure.
5. Compute the probability that two randomly chosen elements (they can be the same) from D_4 commute:

$$P = \begin{cases} \frac{n+3}{4n}, & n \text{ is odd,} \\ \frac{n+6}{4n}, & n \text{ is even.} \end{cases}$$

4 Cyclic Groups

4.1 Properties of Cyclic Groups

If a and b belong to a finite group and $ab = ba$, then $|ab|$ divides $|a| |b|$.

- $|ab| = |a| |b|$ if and only if $(|a|, |b|) = 1$.

Theorem 4.2 ★

$$|a| = n, d = \gcd(n, k) \Rightarrow \langle a^k \rangle = \langle a^d \rangle, |a^k| = \frac{n}{d}.$$

- In a finite cyclic group, the order of an element divides the order of the group.

$$\gcd(n, i) = \gcd(n, j) \Leftrightarrow \langle a^i \rangle = \langle a^j \rangle \Leftrightarrow |a^i| = |a^j|.$$

- $\gcd(n, j) = 1 \Leftrightarrow \langle a \rangle = \langle a^j \rangle \Leftrightarrow |a| = |\langle a^j \rangle|$.

4.2 Classification of Subgroups of Cyclic Groups

Theorem 4.3 ★ Fundamental Theorem of Cyclic Groups

Every subgroup of a cyclic group is cyclic. Moreover, if $|\langle a \rangle| = n$, then the order of any subgroup of $\langle a \rangle$ is a divisor of n ; and, for each positive divisor k of n , the group $\langle a \rangle$ has exactly one subgroup of order k , namely, $\langle a^{n/k} \rangle$.

Theorem 4.4 Number of Elements of Each Order in a Cyclic Group

If d is a positive divisor of n , the number of elements of order d in a cyclic group of order n is $\phi(d)$.

- Notice that for a finite cyclic group of order n , the number of elements of order d for any divisor d of n depends only on d .

Corollary 4.1 Number of Elements of Order d in a Finite Group.

In a finite group, the number of elements of order d is a multiple of $\phi(d)$.

$$\begin{aligned} \phi(p^n) &= p^n - p^{n-1} \\ \phi(p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}) &= \phi(p_1^{k_1}) \phi(p_2^{k_2}) \cdots \phi(p_m^{k_m}) \end{aligned}$$

subgroup lattice

4.3 Exercise

1. If a is a group element of infinite order, then

$$\begin{aligned} \langle a^i \rangle \cap \langle a^j \rangle &= \langle a^{[i,j]} \rangle \\ \langle a^i \rangle \cup \langle a^j \rangle &= \langle a^{(i,j)} \rangle \end{aligned}$$

2. Prove that a finite group is the union of proper subgroups if and only if the group is not cyclic.

4.3 Bibliography of James Joseph Sylvester

5 Permutation Groups

5.1 Definitions and Notation

5.2 Cycle Notation

5.3 Properties of Permutations

Theorem 5.1 Products of Disjoint Cycles

Every permutation of a finite set can be written as a cycle or as a product of disjoint cycles.

Theorem 5.2 Disjoint Cycles Commute

If the pair of cycles $\alpha = (a_1, a_2, \dots, a_m)$ and $\beta = (b_1, b_2, \dots, b_n)$ have no entries in common, then $\alpha\beta = \beta\alpha$.

Theorem 5.3 Order of a Permutation

The order of a permutation of a finite set written in disjoint cycle form is the least common multiple of the lengths of the cycles.

Theorem 5.4 Product of 2-Cycles

Every permutation in $S_n, n > 1$, is a product of 2-cycles.

Lemma If $\varepsilon = \beta_1\beta_2 \cdots \beta_r$, where β_i are 2-cycles, then r is even.

Theorem 5.5 Always Even or Always Odd

Theorem 5.6 Even Permutations Form a Group

The set of even permutations in S_n forms a subgroup of S_n , which is called the alternating group of degree n , A_n .

Theorem 5.7 For $n \geq 1$, A_n has order $n!/2$.

5.4 A Check-Digit Scheme Based on D_5

5.5 Exercise

1. Stabilizer of a in G is a subgroup: $\text{stab}(a) = \{\alpha \in G \mid \alpha(a) = a\}$.
2. Let α belong to S_n , Prove that $|\alpha|$ divides $n!$.

5.5 Bibliography of Augustin Cauchy

5.6 Bibilography of Alan Turing

6 Isomorphisms

6.1 Motivation

6.2 Definition and Examples

Definition Group Isomorphism

An isomorphism ϕ from a group G_1 to a group G_2 is a one-to-one onto mapping (or function) from G_1 to G_2 that preserves the group operation. That is,

$$\forall a, b \in G_1, \phi(ab) = \phi(a)\phi(b).$$

If there is an isomorphism from G_1 onto G_2 , we say that G_1 and G_2 are isomorphic and write $G_1 \approx G_2$.

To prove a group G_1 is isomorphic to a group G_2 :

1. "Mapping": Define a function ϕ from G_1 to G_2 ;
2. "1-1": Assume that $\phi(a) = \phi(b)$, prove that $a = b$;
3. "Onto": For any element \bar{g} in G_2 , find an element g in G_1 such that $\phi(g) = \bar{g}$;
4. "O.P.": Prove that ϕ is operation-preserving; that is, show that $\phi(ab) = \phi(a)\phi(b)$.

Example

- Conjugation by M: $\phi_M = MAM^{-1}$.

6.3 Properties of Isomorphisms

Theorem 6.1 Properties of Isomorphisms Acting on Elements

Suppose that ϕ is an isomorphism from a group G_1 onto a group G_2 . Then

1. ϕ carries the identity of G_1 to the identity of G_2 .
2. For every integer n and for every group element a in G_1 , $\phi(a^n) = [\phi(a)]^n$. (Additive form: $\phi(na) = n\phi(a)$.)
3. For any elements a and b in G_1 , a and b commute if and only if $\phi(a)$ and $\phi(b)$ commute.
4. $G_1 = \langle a \rangle$ if and only if $G_2 = \langle \phi(a) \rangle$.
5. $|a| = |\phi(a)|$ for all a in G_1 (isomorphisms preserve orders).
6. For a fixed integer k and a fixed group element b in G_1 , the equation $x^k = b$ has the same number of solutions in G_1 as does the equation $x^k = \phi(b)$ in G_2 .
7. If G_1 is finite, then G_1 and G_2 have exactly the same number of elements of every order.

Theorem 6.2 Properties of Isomorphisms Acting on Groups

Suppose that ϕ is an isomorphism from a group G_1 onto a group G_2 . Then

1. ϕ^{-1} is an isomorphism from G_2 onto G_1 .
2. G_1 is Abelian if and only if G_2 is Abelian.
3. G_1 is cyclic if and only if G_2 is cyclic.
4. If K is a subgroup of G_1 , then $\phi(K) = \{\phi(k) \mid k \in K\}$ is a subgroup of G_2 .
5. If K is a subgroup of G_2 , then $\phi^{-1}(K) = \{g \in G_1 \mid \phi(g) \in K\}$ is a subgroup of G_1 .
6. $\phi(Z(G_1)) = Z(G_2)$.

To prove groups G_1 and G_2 are not isomorphic:

- Observe that $|G_1| \neq |G_2|$.
- Observe that G_1 is cyclic but G_2 is not.
- Observe that G_1 is Abelian but G_2 is not.
- show that the largest order of any element in G_1 is not the same as that in G_2 .
- Show that the number of elements of some specific order in G_1 is not the same as G_2 .

6.4 Automorphisms

Definition Inner Automorphism Induced by a

Let G be a group, and let $a \in G$. The function ϕ_a defined by $\phi_a(x) = axa^{-1}$ for all x in G is called the **inner** automorphism of G induced by a .

$\text{Aut}(G)$: the set of all automorphisms of G .

$\text{Inn}(G)$: the set of all inner automorphisms of G .

Theorem 6.3 $\text{Aut}(G)$ and $\text{Inn}(G)$ are groups.

Theorem 6.4 $\text{Aut}(Z_n) \approx U(n)$.

6.5 Cayley's Theorem

Theorem 6.5 Cayley's Theorem

Every group is isomorphic to a group of permutations.

The *left regular representation* of G : $\{T_g \mid T_g(x) = gx, g \in G\}$.

6.6 Exercise

1. $U(8) \approx U(12)$.
2. For all finite groups, the order of a subgroup divides the order of the group.
3. $|\text{Aut}(D_n)| = n |U(n)|$.
4. Prove that

$$|\text{Inn}(D_n)| = \begin{cases} 2n, & n \text{ is odd,} \\ n, & n \text{ is even.} \end{cases}$$

6.7 Bibliography of Arthur Cayley

7 Cosets and Lagrange's Theorem

7.1 Properties of Cosets

Definition Coset of H in G

Let G be a group and let H be a nonempty subset of G . For any $a \in G$, $aH = \{ah \mid h \in H\}$, which is called the **left coset** of H in G containing a . The element a is called the **coset representative** of aH .

Lemma 7.1 Properties of Cosets

Let H be a subgroup of G , and $a, b \in G$, then

1. $a \in aH$.
2. $aH = H \iff a \in H$.
3. $(ab)H = a(bH)$, $H(ab) = (Ha)b$.

4. $aH = bH \Leftrightarrow a \in bH$.
5. $aH = bH$ or $aH \cap bH = \emptyset$.
6. $aH = bH \Leftrightarrow a^{-1}b \in H$.
7. $|aH| = |bH| = |H|$.
8. $aH = Ha \Leftrightarrow H = aHa^{-1} \Leftrightarrow H = a^{-1}Ha$.
9. $aH \subset G \Leftrightarrow a \in H$.

7.2 Lagrange's Theorem and Consequences

Theorem 7.1 Lagrange's Theorem: $|H|$ Divides $|G|$

If G is a finite group and H is a subgroup of G , then $|H|$ divides $|G|$. Moreover, the number of distinct left cosets of H in G is $|G|/|H|$.

- The **index** of a subgroup H in G is the number of distinct left cosets of H in G , denoted by $|G : H|$.

Corollary 1 $|G : H| = |G|/|H|$.

Corollary 2 $|a|$ Divides $|G|$.

Corollary 3 Groups of Prime Order Are Cyclic.

Corollary 4 $a^{|G|} = e$.

Corollary 5 Fermat's Little Theorem: $a^p \equiv a \pmod{p}$.

- $a^{p^n} \equiv a \pmod{p}$.

Theorem 7.2 $|HK| = |H||K|/|H \cap K|$.

For two finite subgroups H and K of a group, define the set $HK = \{hk \mid h \in H, k \in K\}$. Then $|HK| = |H||K|/|H \cap K|$.

- HK and hK may not be a subgroup.
- ★ HK may not be a subgroup of G , but $HK \in G$, so $|HK| < |G|$, but need not divide $|G|$.

Theorem 7.3 Classification of Groups of Order $2p$.

Let G be a group of order $2p$, where p is a prime greater than 2. Then G is isomorphic to Z_{2p} or D_p .

- $D_3 \approx S_3 \approx \text{GL}(2, Z_2)$.

7.3 An Application of Cosets to Permutation Groups

Definition Stabilizer of a Point

Let G be a group of permutations of a set S . For each i in S , let $\text{stab}_G(i) = \{\phi \in G \mid \phi(i) = i\}$. We call $\text{stab}_G(i)$ the **stabilizer** of i in G .

Definition Orbit of a Point

Let G be a group of permutations of a set S . For each i in S , let $\text{orb}_G(i) = \{\phi(i) \mid \phi \in G\}$. The set $\text{orb}_G(i)$ is a subset of S called the **orbit** of i under G .

Theorem 7.4 Orbit-Stabilizer Theorem

Let G be a finite group of permutations of a set S . Then, for any i from S ,
 $|G| = |\text{orb}_G(i)| \cdot |\text{stab}_G(i)|$.

7.4 The Rotation Group of a Cube and a Soccer Ball

Theorem 7.5 The Rotation Group of a Cube

The group of rotations of a cube is isomorphic to S_4 .

- The rotation group of a pyramid is isomorphic to A_4 .
- The rotation group of a soccer ball (truncated octahedron) is isomorphic to A_5 .

7.5 An Application of Cosets to the Rubik's Cube

7.6 Exercises

1. Let a and b be elements of a group G , and H and K be subgroups of G . If $aH = bK$, prove that $H = K$.
2. Let H and K be subgroups of G and g belongs to G , show that $g(H \cap K) = gH \cap gK$.
3. If G is a finite group of order n with the property that G has exactly one subgroup of order d for each positive divisor d of n , then G is cyclic.
4. Let H and K be subgroups of a finite group G with $H \subseteq K \subseteq G$. Prove that $|G : H| = |G : K| |K : H|$.
5. If a finite group G has subgroups H and K such that $K \subseteq H \subseteq G$ with $[G : K] = p$ where p is prime, prove that $H = G$ or $H = K$.
6. Prove that if G is a finite group, the index of $Z(G)$ cannot be prime.
7. Prove that A_5 has no subgroup of order 15, 20 or 30, and S_5 has no subgroup of order 30.

7.7 Bibliography of Joseph Lagrange

8 External Direct Products

8.1 Definition and Examples

Definition External Direct Product

Let G_1, G_2, \dots, G_n be a finite collection of groups. The external direct product of them, written as $G_1 \oplus G_2 \oplus \dots \oplus G_n$, is the set of all n -tuples for which the i^{th} component is an element of G_i and the operation is componentwise.

- $|G_1 \oplus G_2 \oplus \dots \oplus G_n| = |G_1| |G_2| \dots |G_n|$.
- $Z_m \oplus Z_n \approx Z_{mn}$ if and only if $\gcd(m, n) = 1$.

8.2 Properties of External Direct Products

Theorem 8.1 Order of an Element in a Direct Product

The order of an element in a direct product of a finite number of finite groups is the least common multiple of the orders of the components of the element. In symbols,

$$|(g_1, g_2, \dots, g_n)| = \text{lcm}(|g_1|, |g_2|, \dots, |g_n|).$$

- If m and n be positive integers that are divisible by a prime p , then the number of elements of order p in $Z_m \oplus Z_n$ is $p^2 - 1$.

Theorem 8.2 Criterion for $G \oplus H$ to be Cyclic.

Let G and H be finite cyclic groups. Then $G \oplus H$ is cyclic if and only if $|G|$ and $|H|$ are relatively prime.

Corollary 1 Criterion for $G_1 \oplus G_2 \oplus \cdots \oplus G_n$ to be Cyclic.

Corollary 2 Criterion for $Z_{n_1 n_2 \cdots n_k} \approx Z_{n_1} \oplus Z_{n_2} \oplus \cdots \oplus Z_{n_k}$.

$Z_{n_1 n_2 \cdots n_k} \approx Z_{n_1} \oplus Z_{n_2} \oplus \cdots \oplus Z_{n_k}$ if and only if n_i and n_j are relatively prime when $i \neq j$.

$$\begin{aligned} Z_2 \oplus Z_{30} &\approx Z_2 \oplus Z_6 \oplus Z_5 \\ &\approx Z_2 \oplus Z_2 \oplus Z_3 \oplus Z_5 \\ &\approx Z_2 \oplus Z_6 \oplus Z_5 \\ &\approx Z_2 \oplus Z_3 \oplus Z_2 \oplus Z_5 \\ &\approx Z_6 \oplus Z_{10}. \end{aligned}$$

8.3 The Group of Units Modulo n as an External Direct Product

$U_k(n) \equiv \{x \in U(n) \mid x \equiv 1 \pmod{k}\}$ is a subgroup of $U(n)$.

Theorem 8.3 $U(n)$ as an External Direct Product

Suppose s and t are relatively prime. Then $U(st)$ is isomorphic to the external direct product of $U(s)$ and $U(t)$. In short,

$$U(st) \approx U(s) \oplus U(t).$$

Moreover, $U_s(st)$ is isomorphic to $U(t)$, and $U_t(st)$ is isomorphic to $U(s)$.

$$\left| \begin{array}{l} U(st) \rightarrow U(s) \oplus U(t) \\ x \mapsto (x \pmod{s}, x \pmod{t}) \end{array} \right| \quad \left| \begin{array}{l} U_s(st) \rightarrow U(t) \\ x \mapsto x \pmod{t} \end{array} \right|$$

Corollary

Let $m = n_1 n_2 \cdots n_k$, where $\gcd(n_i, n_j) = 1$ for $i \neq j$. Then

$$U(m) \approx U(n_1) \oplus U(n_2) \oplus \cdots \oplus U(n_k).$$

$$\begin{cases} U(2) \approx \{0\}, \\ U(4) \approx Z_2, \\ U(2^n) \approx Z_{2^{n-2}} \oplus Z_2, & \text{for } n \geq 3, \\ U(p^n) \approx Z_{p^{n-1}} \oplus Z_{p^{n-2}} \oplus \cdots \oplus Z_p, & \text{for } p \text{ an odd prime.} \end{cases}$$

e.g. $|\text{Aut}^4(Z_{27})| = 1$.

8.4 RSA Public Key Encryption Scheme

Receiver

1. Pick very large primes p and q and compute $n = pq$.
2. Compute the least common multiple of $p - 1$ and $q - 1$; let us call it m .
3. Pick e relatively prime to m .

- Find d such that $ed \bmod m = 1$.
- Publicly announce n and e .

Sender

- Convert the message to a string of digits.
- Break up the message into uniform blocks of digits; call them M_1, M_2, \dots, M_k . (The integer value of each M_i must be less than n . In practice, n is so large that this is not a concern.)
- Check to see that the greatest common divisor of each M_i and n is 1. If not, n can be factored and our code is broken. (In practice, the primes p and q are so large that they exceed all M_i , so this step may be omitted.)
- Calculate and send $R_i = M_i^e \bmod n$.

Receiver

- For each received message R_i , calculate $R_i^d \bmod n$.
- Convert the string of digits back to a string of characters.

Principles

$$U(n) \approx U(p) \oplus U(q) \approx Z_{p-1} \oplus Z_{q-1}.$$

$$R_i^d = (M_i^e)^d = M_i^{ed} = M_i^{1+km} = M_i.$$

8.5 Exercises

- $G \oplus H$ is Abelian if and only if G and H are Abelian.
- $G_1 \approx G_2, H_1 \approx H_2 \Rightarrow G_1 \oplus H_1 \approx G_2 \oplus H_2$.
- $A \oplus B \approx A \oplus C \Leftrightarrow B \approx C$.
- $U(8) \approx U(12), U(55) \approx U(75), U(144) \approx U(140), U_{50}(200) \approx U(4)$.
- $U_p(p^n) \approx Z_{p^{n-1}}$.
- For relatively prime positive integers $s \leq n$ and $t \leq n$, show that $U_{st}(n) = U_s(n) \cap U_t(n)$.

8.6 Bibliography of Leonard Adleman

9 Normal Subgroups and Factor Groups

9.1 Normal Subgroups

Definition Normal Subgroup

A subgroup H of a group G is called a normal subgroup of G if $aH = Ha$ for all a in G . We denote this by $H \triangleleft G$.

Theorem 9.1 Normal Subgroup Test

A subgroup H of G is normal in G if and only if $xHx^{-1} \subseteq H$ for all x in G .

e.g.

- Every subgroup of an Abelian group is normal.
- The center $Z(G)$ of a group is normal.

- A_n is a normal subgroup of S_n .
- Every subgroup of D_n consisting solely of rotations is normal.
- $\text{SL}(2, \mathbb{R})$ is a normal subgroup of $\text{GL}(2, \mathbb{R})$.

Properties:

- If H and K are subgroups of G and H is normal, then HK is a subgroup of G .
- If a group G has a unique subgroup H of some finite order, then H is normal in G .
- Normality is not transitive: $K \triangleleft L \triangleleft G \nRightarrow K \triangleleft G$.
- If N and M are normal, then $N \cap M$ and NM are normal.
- $K/N \triangleleft G/N \Rightarrow K \triangleleft G$.

9.2 Factor Groups

Theorem 9.2 Factor (Quotient) Groups

Let G be a group and let H be a **normal** subgroup of G . The set $G/H = \{aH \mid a \in G\}$ is a group under the operation $(aH)(bH) = abH$.

- The converse is also true: if $aHbH = abH$ defines a group operation on the set of left cosets of H in G , then H is normal in G .

9.3 Applications of Factor Groups

Theorem 9.3 G/Z Theorem

Let G be a group and let $Z(G)$ be the center of G . If $G/Z(G)$ is cyclic, then G is **Abelian**, thus $G/Z(G)$ is trivial.

- If G/H is cyclic, where H is a subgroup of $Z(G)$, then G is Abelian.
- If G is non-Abelian, then $G/Z(G)$ is not cyclic.
 - A non-Abelian group of order pq , where p and q are primes, must have a trivial center.
- If $K = \{H, a_1H, a_2H, a_3H\}$ is a subgroup of the factor group G/H , then the set $K = H \cup a_1H \cup a_2H \cup a_3H$ is a subgroup of G of order $4|H|$, called the **pull back** of K to G .
- Suppose that G is a finite group and a factor group G/H has an element aH of order n , then G has an element of order n .

Theorem 9.4 $G, G/Z(G)$

For any group G , $G/Z(G)$ is isomorphic to $\text{Inn}(G)$.

It can be proved by the First Isomorphism Theorem in chapter 10 easily.

- $|Z(D_6)| = 2 \Rightarrow |D_6/Z(D_6)| = 6 \Rightarrow D_6/Z(D_6) \approx D_3$ or Z_6 . By Theorem 9.3 and 9.4, we know that $\text{Inn}(D_6) \approx D_3$.
- $\text{Inn}(D_{2n}) \approx D_n$, $\text{Inn}(D_{2n+1}) \approx D_{2n+1}$.

Theorem 9.5 Cauchy's Theorem for Abelian Groups

Let G be a finite Abelian group and let p be a prime that divides the order of G , then G has an element of order p .

9.4 Internal Direct Products

Definition Internal Direct Product of H and K

We say that G is the internal direct product of H and K and write $G = H \times K$ if H and K are normal subgroups of G and

$$G = HK \text{ and } H \cap K = \{e\}.$$

- If s and t are relatively prime positive integers then $U(st) = U_s(st) \times U_t(st)$.
- $D_6 = \{R_0, R_{120}, R_{240}, F, R_{120}F, R_{240}F\} \times \{R_0, R_{180}\} \approx D_3 \oplus Z_2$.

Definition Internal Direct Product $H_1 \times H_2 \times \cdots \times H_n$

Let H_1, H_2, \dots, H_n be a finite collection of normal subgroups of G . We say that G is the internal direct product of H_1, H_2, \dots, H_n and write $G = H_1 \times H_2 \times \cdots \times H_n$ if

1. $G = H_1 H_2 \cdots H_n = \{h_1 h_2 \cdots h_n \mid h_i \in H_i\}$,
2. $(H_1 H_2 \cdots H_i) \cap H_{i+1} = \{e\}$ for $i = 1, 2, \dots, n-1$.

Theorem 9.6 $H_1 \times H_2 \times \cdots \times H_n \approx H_1 \oplus H_2 \oplus \cdots \oplus H_n$

If a group G is the internal direct product of a finite number of subgroups H_1, H_2, \dots, H_n , then G is isomorphic to the external direct product of H_1, H_2, \dots, H_n .

- To prove this
 - $\forall h_i \in H_i, h_j \in H_j, h_i h_j = h_j h_i$.
 - Each member of G can be expressed uniquely in the form $h_1 h_2 \cdots h_n$.
 - Mapping: $\phi(h_1 h_2 \cdots h_n) = (h_1, h_2, \dots, h_n)$.
- If $m = n_1 n_2 \cdots n_k$, $(n_i, n_j) = 1$ for $i \neq j$, then

$$\begin{aligned} U(m) &= U_{m/n_1}(m) \times U_{m/n_2}(m) \times \cdots \times U_{m/n_k}(m) \\ &\approx U(n_1) \oplus U(n_2) \oplus \cdots \oplus U(n_k) \end{aligned}$$

Classification Theorems

- Classification of subgroups of finite cyclic groups: There is exactly one subgroup for each divisor of the order of the group and no others.
- Classification of groups of prime order: Every group of prime order p is isomorphic to Z_p .
- Classification of groups of $2p$ where p is an odd prime: Every group of $2p$ is isomorphic to Z_{2p} or D_p .
- Classification of groups of 4: Every group of order 4 is isomorphic to Z_4 or $Z_2 \oplus Z_2$.

Theorem 9.7 Classification of finite Abelian groups of squarefree order

Every Abelian group of order $p_1 p_2 \cdots p_k$ where p_i are distinct primes is cyclic.

- $G = H_1 \times H_2 \times \cdots \times H_k$.

Theorem 9.8 Classification of Groups of Order p^2

Every group of order p^2 , where p is a prime, is isomorphic to Z_{p^2} or $Z_p \oplus Z_p$.

- Let G be a group of order p^2 , then every subgroup of the form $\langle a \rangle$ is normal in G .

Corollary

If G is a group of order p^2 , where p is a prime, then G is Abelian.

9.5 Exercises

1. Prove that if H has index 2 in G , then H is normal in G .

2. Prove that a factor group of a cyclic group is cyclic, a factor group of an Abelian group is Abelian.
3. H is normal in G , a is an element of G . Then the order of the element aH in the factor group G/H is the smallest positive integer n such that a^n is in H . Moreover, $|gH|$ divides $|g|$.
4. $H \approx K \not\Rightarrow G/H \approx G/K$.
5. Groups of order 2 or 4 are all Abelian.
6. Let G be a group and let $S = \{x^{-1}y^{-1}xy \mid x, y \in G\}$, $G' = [G, G] = \langle S \rangle$. Then
 1. G' is normal in G .
 2. G/G' is Abelian.
 3. If G/N is Abelian, then $G' \subseteq N$.
 4. If H is a subgroup of G and $G' \subseteq H$, then H is normal.
7. $\text{Inn}(G)$ is normal in $\text{Aut}(G)$.

Question: 66.

9.6 Bibliography of Evariste Galois

10 Group Homomorphisms

10.1 Definition and Examples

Definition Group Homomorphism

A homomorphism ϕ from a group G_1 to a group G_2 is a mapping from G_1 into G_2 that preserves the group operation; that is, $\phi(ab) = \phi(a)\phi(b)$ for all a, b in G .

Definition Kernel of a Homomorphism

The **kernel** of a homomorphism ϕ from a group G to a group with identity e is the set $\text{Ker } \phi = \{x \in G \mid \phi(x) = e\}$.

- Any isomorphism is a homomorphism that is also [onto](#) and [one-to-one](#), the kernel of which is a trivial subgroup.
- Let $\phi : \text{GL}(n, \mathbb{R}) \rightarrow \mathbb{R}^*$, $A \mapsto \det A$, then $\text{Ker } \phi = \text{SL}(n, \mathbb{R})$.
- $U(st) = U_s(st)U_t(st)$, $\phi(ab) = a$, then $\text{Ker } \phi = U_t(st)$.
- Every linear transformation is a group homomorphism and the null-space is the same as the kernel. An invertible linear transformation is a group isomorphism.

10.2 Properties of Homomorphisms

Theorem 10.1 Properties of Elements Under Homomorphisms

Let ϕ be a homomorphism from a group G_1 to a group G_2 and let g be an element of G_1 . Then

1. ϕ carries the identity of G_1 to the identity of G_2 .
2. $\phi(g^n) = \phi(g)^n$ for all n in \mathbb{Z} .

3. If $|g|$ is finite, then $|\phi(g)|$ divides $|g|$ and if $|G_1|$ is finite, then $|\phi(g)|$ divides $|g|$ and $|\phi(G_1)|$.
4. $\text{Ker } \phi$ is a subgroup of G_1 .
5. $\phi(a) = \phi(b)$ if and only if $a \text{Ker } \phi = b \text{Ker } \phi$.
6. If $\phi(g) = g'$, then $\phi^{-1}(g') = \{x \in G_1 \mid \phi(x) = g'\} = g \text{Ker } \phi$.


- The particular solution to $Ax = b$ is x_0 , the entire solution to $Ax = 0$ is S , then the entire solution to $Ax = b$ is $x_0 + S$. It's a special case of property 6.

Theorem 10.2 Properties of Subgroups Under Homomorphisms

Let ϕ be a homomorphism from a group G_1 to a group G_2 and let H be a subgroup of G_1 . Then

1. $\phi(H) = \{\phi(h) \mid h \in H\}$ is a subgroup of G_2 .
2. If H is cyclic, then $\phi(H)$ is **cyclic**.
3. If H is Abelian, then $\phi(H)$ is **Abelian**.
4. If H is normal in G_1 , then $\phi(H)$ is **normal** in $\phi(G_1)$.
5. If $|\text{Ker } \phi| = n$, then ϕ is an n -to-1 mapping from G_1 onto $\phi(G_1)$.
6. If H is finite, then $|\phi(H)|$ divides $|H|$.
7. $\phi(Z(G_1))$ is a subgroup of $Z(\phi(G_1))$.
8. If K is a subgroup of G_2 , then $\phi^{-1}(K) = \{k \in G_1 \mid \phi(k) \in K\}$ is a **subgroup** of G_1 .
9. If K is a normal subgroup of G_2 , then $\phi^{-1}(K) = \{k \in G_1 \mid \phi(k) \in K\}$ is a **normal subgroup** of G_1 .
10. If ϕ is onto and $\text{Ker } \phi = \{e\}$, then ϕ is an isomorphism from G_1 to G_2 .

- $|\phi^{-1}(H)| = |H| |\text{Ker } \phi|$.
- The inverse image of an element is a coset of the kernel and that every element in that coset has the same image.



Corollary Kernels are Normal

Let ϕ be a group homomorphism from G_1 to G_2 , then $\text{Ker } \phi$ is a normal subgroup of G_1 .

- The number of homomorphisms from \mathbb{Z}_m to \mathbb{Z}_n is $d = \gcd(m, n)$, since such a homomorphism is completely specified by the image a of 1, and $|a|$ divides both m and n , and $d = \sum \phi(a)$ for all divisor a of d .

10.3 The First Isomorphism Theorem

Theorem 10.3 First Isomorphism Theorem

Let ϕ be a group homomorphism from G_1 to G_2 , then the mapping from $G_1/\text{Ker } \phi$ to $\phi(G_1)$, given by $g\text{Ker } \phi \rightarrow \phi(g)$, is an isomorphism. In symbols, $G_1/\text{Ker } \phi \approx \phi(G_1)$.


Corollary 1

If ϕ is a homomorphism from a finite group G_1 to G_2 , then $|G_1|/|\text{Ker } \phi| = |\phi(G_1)|$.

Corollary 2

If ϕ is a homomorphism from a finite group G_1 to G_2 , then $|\phi(G_1)|$ divides $|G_1|$ and $|G_2|$.

The commutative diagram of Theorem 10.3 is:



$\gamma : G \rightarrow G/\text{Ker } \phi$, $g \mapsto g\text{Ker } \phi$ is called the **natural mapping** from G to $G/\text{Ker } \phi$. The diagram is commutative since $\psi\gamma = \phi$.

- $\mathbb{Z}/\langle n \rangle \approx \mathbb{Z}_n$ since $\phi(m) = m \pmod n$ is a homomorphism with $\text{Ker } \phi = \langle n \rangle$. Likewise, $\mathbb{Z}[i]/\{sn + tni \mid s, t \in \mathbb{Z}\} \approx \mathbb{Z}_n[i]$.
- $\text{GL}(2, \mathbb{R})/\text{SL}(2, \mathbb{R}) \approx \mathbb{R}^*$ since $\phi(A) = \det A$ from $\text{GL}(2, \mathbb{R})$ onto \mathbb{R}^* is a homomorphism with $\text{Ker } \phi = \text{SL}(2, \mathbb{R})$. Likewise, $\text{SL}^\pm(2, \mathbb{R}) = \{A \in \text{GL}(2, \mathbb{R}) \mid \det A = \pm 1\} \approx \mathbb{R}^+$ since we have $\phi(A) = (\det A)^2$.
- For an Abelian group G and a positive integer k , let G^k denote the subgroup $\{x^k \mid x \in G\}$ and $G^{(k)}$ the subgroup $\{x \in G \mid x^k = e\}$. Then $G/G^{(k)} \approx G^k$ since we have $\phi(x) = x^k$, but $G/G^k \not\approx G^{(k)}$ since $\phi(x^k) = x$ may be not well-defined.

Theorem N/C Theorem

Let H be a subgroup of a group G . Noting that the normalizer of H in G , $N(H) = \{x \in G \mid xHx^{-1} = H\}$, and the centralizer of H in G , $C(H) = \{x \in G \mid \forall h \in H, xhx^{-1} = h\}$, are subgroups of G , consider the mapping from $N(H)$ to $\text{Aut}(H)$ given by $g \mapsto \phi_g$, where $\phi_g(h) = ghg^{-1}$. This mapping is a homomorphism with $\text{Ker } \phi_g = C(H)$. So, $N(H)/C(H)$ is isomorphic to a subgroup of $\text{Aut}(H)$, in fact, $N(H)/C(H) \approx \text{Inn}(H)$.

Theorem 10.4 Normal Subgroups Are Kernels

Every normal subgroup N of a group G is the kernel of a **natural homomorphism** of G defined by $\phi : G \rightarrow G/N$, $g \mapsto gN$.

10.4 Exercises

1. $G \xrightarrow{\phi} H \xrightarrow{\sigma} K$, then $\text{Ker } \phi$ is a normal subgroup of $\text{Ker } \sigma\phi$, and $[\text{Ker } \sigma\phi : \text{Ker } \phi] = |H|/|K|$.
2. $U(st)/U_s(st) \approx U(s)$.
3. If $G = \langle S \rangle$ and ϕ is a homomorphism from G to some group, prove that $\phi(G) = \langle \phi(S) \rangle$.
4. Let N be a normal subgroup of a group G . Prove that every subgroup of G/N has the form H/N , where H is a subgroup of G .
5. For any two primes p and q with $p < q$ where $p \nmid q - 1$, a group of order pq is cyclic.

Theorem First Isomorphism Theorem

Let ϕ be a group homomorphism from G_1 onto G_2 , then the mapping ψ from $G_1/\text{Ker } \phi$ to G_2 , given by $g\text{Ker } \phi \rightarrow \phi(g)$, is an isomorphism. In symbols, $G_1/\text{Ker } \phi \approx G_2$.

Proof $\psi(x\text{Ker } \phi y\text{Ker } \phi) = \psi(xy\text{Ker } \phi) = \phi(xy) = \phi(x)\phi(y) = \psi(x\text{Ker } \phi)\psi(y\text{Ker } \phi)$.
□

Theorem Second Isomorphism Theorem

If K is a subgroup of G and N is a normal subgroup of G , then $K/(K \cap N) \approx KN/N$.

Proof Let $\phi : K \rightarrow KN/N$, $k \mapsto kN$, then $\text{Ker } \phi = K \cap N$. □

Theorem Third Isomorphism Theorem

If M and N are normal subgroups of G and $N \subseteq M$, then $(G/N)/(M/N) \approx G/M$.

Proof Let $\phi : G/N \rightarrow G/M$, $gN \mapsto gM$, then $\text{Ker } \phi = M/N$. □

10.5 Bibliography of Camile Jordan

11 Fundamental Theorem of Finite Abelian Groups

11.1 The Fundamental Theorem

Theorem 11.1 Fundamental Theorem of Finite Abelian Groups

Every finite Abelian group is a direct product of cyclic groups of prime-power order. Moreover, the number of terms in the product and the orders of the cyclic groups are uniquely determined by the group.

Writing an Abelian group G in the form $\mathbb{Z}_{p_1^{n_1}} \oplus \mathbb{Z}_{p_2^{n_2}} \oplus \cdots \oplus \mathbb{Z}_{p_k^{n_k}}$, is called determining the isomorphism class of G .

- If $k = n_1 + n_2 + \cdots + n_t$, then $\mathbb{Z}_{p^{n_1}} \oplus \mathbb{Z}_{p^{n_2}} \oplus \cdots \oplus \mathbb{Z}_{p^{n_t}}$ is an Abelian group of order p^k .

11.2 The Isomorphism Classes of Abelian Groups

Corollary Existence of Subgroups of Abelian Groups

If m divides the order of a finite Abelian group G , then G has a subgroup of order m .

11.3 Proof of the Fundamental Theorem

Lemma 1

Let G be a finite Abelian group of order $p^n m$, where p is a prime that does not divide m . Then $G = H \times K$, where $H = \{x \in G \mid x^{p^n} = e\}$ and $K = \{x \in G \mid x^m = e\}$. Moreover, $|H| = p^n$.

- Given an Abelian group G with $|G| = p_1^{n_1} p_2^{n_2} \cdots p_k^{n_k}$, where p 's are distinct primes, let $G(p_i) = \{x \in G \mid x^{p_i^{n_i}} = e\}$, then $G = G(p_1) \times G(p_2) \times \cdots \times G(p_k)$ and $|G(p_i)| = p_i^{n_i}$.

Lemma 2

Let G be an Abelian group of prime-power order and let a be an element of maximum order in G , then G can be written in the form $\langle a \rangle \times K$.

Lemma 3

A finite Abelian group of prime-order is an internal direct product of cyclic groups.

Lemma 4

Suppose that G is a finite Abelian group of prime-power order. If $G = H_1 \times H_2 \times \cdots \times H_m$ and $G = K_1 \times K_2 \times \cdots \times K_n$, where the H 's and K 's are nontrivial cyclic subgroups with $|H_1| \geq |H_2| \geq \cdots \geq |H_m|$ and $|K_1| \geq |K_2| \geq \cdots \geq |K_n|$, then $m = n$ and $|H_i| = |K_i|$ for all i .

11.4 Exercises

1. The number of elements in $\mathbb{Z}_{p^{n_1}} \oplus \mathbb{Z}_{p^{n_2}} \oplus \cdots \oplus \mathbb{Z}_{p^{n_k}}$ of order p is $p^{n-1} + p^{n-2} + \cdots + p + 1 = \frac{p^n - 1}{p - 1}$.
2. Dirichlet's Theorem says that, for every pair of relatively prime integers a and b , there are infinitely many primes of the form $at + b$. Use **Dirichlet's Theorem** to prove that every finite Abelian group is isomorphic to a subgroup of a U -group. (Hint: $U(p_i^{n_i}t + 1) \approx \mathbb{Z}_{p_i^{n_i}t}$)