

Contents

2	Quantum Fields	2
Exercise 2.1		2
Exercise 2.2		2
Exercise 2.3		3
Exercise 2.4		4
Exercise 2.5		7
Exercise 2.6		10
Exercise 2.7		12
Exercise 2.8		13
3	Conserved Particles	15
Exercise 3.1		15
Exercise 3.2		15
Exercise 3.3		17
Exercise 3.4		18
Exercise 3.5		19
Exercise 3.6		20

2 Quantum Fields

Exercise 2.1

The partition function is

$$\begin{aligned} Z &= \sum_n e^{-\beta\hbar\omega(n+\frac{1}{2})} \\ &= e^{-\beta\hbar\omega/2} \frac{1}{1 - e^{-\beta\hbar\omega}}. \end{aligned} \quad (2.1.1)$$

Let $x = \beta\hbar\omega$,

$$Z = e^{-x/2} (1 - e^{-x})^{-1}. \quad (2.1.2)$$

The internal energy is

$$\begin{aligned} E &= N_A \left(-\frac{\partial \ln Z}{\partial \beta} \right) \\ &= -N_A \hbar\omega \frac{1}{Z} \frac{\partial Z}{\partial x} \\ &= \frac{1}{2} N_A \hbar\omega \left(1 + \frac{e^{-x/2}}{\sinh \frac{x}{2}} \right). \end{aligned} \quad (2.1.3)$$


The heat capacity is

$$\begin{aligned} C_V &= \frac{\partial E}{\partial T} = \frac{dx}{dT} \frac{\partial E}{\partial x} \\ &= \frac{1}{4} R x^2 \frac{1}{\sinh^2(\frac{x}{2})} \\ &= R F(x) \end{aligned} \quad (2.1.4)$$

where

$$F(x) = \left(\frac{x/2}{\sinh(x/2)} \right)^2 \rightarrow 1 \quad (2.1.5)$$

in the $x \rightarrow 0$ limit ($k_B T \gg \hbar\omega$), where it recovers the Dulong–Petit law.



Exercise 2.2

$$\begin{aligned} \sum_j \langle q_m | j \rangle \langle j | q_n \rangle &= \frac{1}{N_S} \sum_{j=0}^{N_S-1} e^{i(q_n - q_m)ja} \\ &= \frac{1}{N_S} \sum_{j=0}^{N_S-1} e^{i(n-m)\frac{2\pi}{N_S}j}. \end{aligned} \quad (2.2.1)$$

If $n \neq m$, then¹

$$\begin{aligned}
 \text{LHS} &= \frac{1}{N_S} \frac{e^{i2\pi(n-m)} - 1}{e^{i2\pi(n-m)/N_S} - 1} \\
 &= \frac{1}{N_S} \frac{\sin(\pi(n-m))}{\sin\left(\frac{\pi}{N_S}(n-m)\right)} \frac{e^{-i\pi(n-m)/N_S}}{e^{-i\pi(n-m)}} \\
 &= \frac{1}{N_S} \frac{\sin(\pi(n-m))}{\sin\left(\frac{\pi}{N_S}(n-m)\right)} (-1)^{\frac{N_S-1}{N_S}(n-m)} \\
 &= 0.
 \end{aligned} \tag{2.2.2}$$

If $n = m$, then

$$\begin{aligned}
 \text{LHS} &= \frac{1}{N_S} \sum_{j=0}^{N_S-1} 1 \\
 &= 1.
 \end{aligned} \tag{2.2.3}$$

Therefore,

$$\langle q_m | q_n \rangle = \delta_{mn}. \tag{2.2.4}$$

Exercise 2.3

Have

$$\hat{x}(0) = \sqrt{\frac{\hbar}{2m\omega}} (\hat{a} + \hat{a}^\dagger) \tag{2.3.1}$$

$$\hat{x}(t) = \sqrt{\frac{\hbar}{2m\omega}} (\hat{a}e^{i\omega t} + \hat{a}^\dagger e^{-i\omega t}). \tag{2.3.2}$$

The symmetrised correlation function can be expanded as

$$\begin{aligned}
 S_\beta(t) &= \frac{1}{2} \langle \hat{x}(t)\hat{x}(0) + \hat{x}(0)\hat{x}(t) \rangle \\
 &= \frac{\hbar}{4m\omega} [2e^{i\omega t} \langle \hat{a}\hat{a} \rangle + 2e^{-i\omega t} \langle \hat{a}^\dagger \hat{a}^\dagger \rangle + 2 \cos \omega t (\langle \hat{a}\hat{a}^\dagger \rangle + \langle \hat{a}^\dagger \hat{a} \rangle)].
 \end{aligned} \tag{2.3.3}$$

At (inverse) temperature β ,

$$\begin{cases} \langle \hat{a}^\dagger \hat{a} \rangle = \langle \hat{n} \rangle \\ \langle \hat{a}\hat{a}^\dagger \rangle = \langle \hat{a}^\dagger \hat{a} + 1 \rangle = \langle \hat{a} \rangle + 1 \\ \langle \hat{a}\hat{a} \rangle = \langle \hat{a}^\dagger \hat{a}^\dagger \rangle = 0, \end{cases} \tag{2.3.4}$$

where

$$\langle \hat{n} \rangle = n(\beta) = \frac{1}{e^{\beta\hbar\omega} - 1}. \tag{2.3.5}$$

Substituting in (2.3.3) gives

$$S_\beta(t) = \frac{\hbar}{2m\omega} \cos(\omega t) \left[\underbrace{2n(\beta)}_{\text{finite-}T \text{ excitation}} + \underbrace{1}_{\text{zero-point fluctuation}} \right]. \tag{2.3.6}$$

¹I believe the stated result is missing the red N_S^{-1} .

Exercise 2.4

(a)

$$\begin{aligned} [\hat{b}, \hat{b}] &= u^2[\hat{a}, \hat{a}] + uv[\hat{a}, \hat{a}^\dagger] + uv[\hat{a}^\dagger, \hat{a}] + v^2[\hat{a}^\dagger, \hat{a}^\dagger] \\ &= 0 \end{aligned} \quad (2.4.1)$$

$$\begin{aligned} [\hat{b}^\dagger, \hat{b}^\dagger] &= u^2[\hat{a}^\dagger, \hat{a}^\dagger] + uv[\hat{a}^\dagger, \hat{a}] + uv[\hat{a}, \hat{a}^\dagger] + v^2[\hat{a}, \hat{a}] \\ &= 0 \end{aligned} \quad (2.4.2)$$

$$\begin{aligned} [\hat{b}, \hat{b}^\dagger] &= u^2[\hat{a}, \hat{a}^\dagger] + uv[\hat{a}, \hat{a}] + uv[\hat{a}^\dagger, \hat{a}^\dagger] + v^2[\hat{a}^\dagger, \hat{a}] \\ &= u^2 - v^2. \end{aligned} \quad (2.4.3)$$

The first two canonical commutation relations of $\hat{b}$ are guaranteed by the canonical commutation relations of $\hat{a}$, while the last one satisfies $[\hat{b}, \hat{b}^\dagger] = 1$ if and only if $u^2 - v^2 = 1$.

This means that it is convenient to parameterise u, v as $u = \cosh \theta$, $v = \sinh \theta$.

(b) Given $u^2 - v^2 = 1$, the inverse transform is

$$\hat{a} = u\hat{b} - v\hat{b}^\dagger\hat{a}^\dagger = u\hat{b}^\dagger - v\hat{b}. \quad (2.4.4)$$

Then we can expand

$$\begin{aligned} \hat{a}^\dagger\hat{a} + \frac{1}{2} &= u^2\hat{b}^\dagger\hat{b} - uv\hat{b}^\dagger\hat{b}^\dagger - uv\hat{b}\hat{b} + v^2\hat{b}^\dagger\hat{b}^\dagger + \frac{1}{2} \\ &= (u^2 + v^2) \left(\hat{b}^\dagger\hat{b} + \frac{1}{2} \right) - uv(\hat{b}^\dagger\hat{b}^\dagger + \hat{b}\hat{b}) \end{aligned} \quad (2.4.5)$$

and similarly

$$\hat{a}\hat{a} + \hat{a}^\dagger\hat{a}^\dagger = (u^2 + v^2)(\hat{b}\hat{b} + \hat{b}^\dagger\hat{b}^\dagger) - 4uv \left(\hat{b}^\dagger\hat{b} + \frac{1}{2} \right) \quad (2.4.6)$$

using $u^2 - v^2 = 1$. Therefore,

$$\hat{H} = [\omega(u^2 + v^2) - 2\Delta uv] \left(\hat{b}^\dagger\hat{b} + \frac{1}{2} \right) + \left[\frac{\Delta}{2}(u^2 + v^2) - \omega uv \right] (\hat{b}^\dagger\hat{b}^\dagger + \hat{b}\hat{b}). \quad (2.4.7)$$

For $\hat{H}$ to be diagonalisable, there must be some θ such that for $u = \cosh \theta$, $v = \sinh \theta$, the latter term vanish,

$$\Delta(u^2 + v^2) = 2\omega uv. \quad (2.4.8)$$

In θ , this is

$$\Delta \cosh 2\theta = \omega \sinh 2\theta, \quad (2.4.9)$$

so

$$\tanh 2\theta = \frac{\Delta}{\omega}. \quad (2.4.10)$$

Therefore, we must have

$$|\Delta| < \omega \quad (2.4.11)$$

for $\hat{H}$ to be diagonalisable.

The new frequency is

$$\begin{aligned} \tilde{\omega} &= \omega(u^2 + v^2) - 2\Delta uv \\ &= \omega \cosh 2\theta - \Delta \sinh 2\theta. \end{aligned} \quad (2.4.12)$$

Given $\tanh 2\theta = \frac{\Delta}{\omega}$, we have

$$\begin{cases} \sinh 2\theta = \frac{\Delta}{\sqrt{\omega^2 - \Delta^2}} \\ \cosh 2\theta = \frac{\omega}{\sqrt{\omega^2 - \Delta^2}}, \end{cases} \quad (2.4.13)$$

so

$$\tilde{\omega} = \sqrt{\omega^2 - \Delta^2}. \quad (2.4.14)$$

Since

$$\begin{cases} u^2 + v^2 = \cosh 2\theta = \frac{\omega}{\tilde{\omega}} \\ u^2 - v^2 = 1, \end{cases} \quad (2.4.15)$$

we have

$$\begin{cases} u^2 = \frac{1}{2} \left(\frac{\omega}{\tilde{\omega}} + 1 \right) \\ v^2 = \frac{1}{2} \left(\frac{\omega}{\tilde{\omega}} - 1 \right). \end{cases} \quad (2.4.16)$$

We need to take care of the signs when taking the square roots.

$$\begin{cases} u = \sqrt{\frac{1}{2} \left(\frac{\omega}{\sqrt{\omega^2 - \Delta^2}} + 1 \right)} \\ v = \operatorname{sgn}(\Delta) \sqrt{\frac{1}{2} \left(\frac{\omega}{\sqrt{\omega^2 - \Delta^2}} - 1 \right)}. \end{cases} \quad (2.4.17)$$

When $\Delta = \omega$, the transition is singular. As $\theta \rightarrow \infty$, $u, v \rightarrow \infty$ but $\tilde{\omega} \rightarrow 0$. However, this does not mean that $\hat{H}$ is 0 (as it does not, apparently). In fact,

$$\begin{aligned} \hat{H} &= \frac{1}{2} \omega (\hat{a} + \hat{a}^\dagger)^2 \\ &= \omega \hat{x}^2. \end{aligned} \quad (2.4.18)$$

The momentum term vanishes.

(c) We have

$$e^{-\alpha \hat{a}^\dagger \hat{a}^\dagger} = \sum_{k=0}^{\infty} \frac{1}{k!} (-\alpha \hat{a}^\dagger \hat{a}^\dagger)^k. \quad (2.4.19)$$

The commutation relations are

$$[\hat{a}^\dagger, e^{-\alpha \hat{a}^\dagger \hat{a}^\dagger}] = 0 \quad (2.4.20)$$

and

$$\begin{aligned} [\hat{a}, e^{-\alpha \hat{a}^\dagger \hat{a}^\dagger}] &= \sum_{k=0}^{\infty} \frac{1}{k!} (-\alpha)^k [\hat{a}, (\hat{a}^\dagger)^{2k}] \\ &= \sum_{k=1}^{\infty} \frac{1}{k!} (-\alpha)^k \cdot 2k (\hat{a}^\dagger)^{2k-1}, \end{aligned} \quad (2.4.21)$$

where in evaluating $[\hat{a}, (\hat{a}^\dagger)^n]$, notice that when you move $\hat{a}$ to the right of a string of $\hat{a}^\dagger$'s, each time you move through an $\hat{a}^\dagger$, you get a $[\hat{a}, \hat{a}^\dagger] = 1$ multiplied with the other $(n-1)$ $\hat{a}^\dagger$'s.

Therefore,

$$\begin{aligned}
\hat{b}|\tilde{0}\rangle &= (u\hat{a} + v\hat{a}^\dagger)e^{-\alpha\hat{a}^\dagger\hat{a}}|0\rangle \\
&= \left[u \left(e^{-\alpha\hat{a}^\dagger\hat{a}}\hat{a} + [\hat{a}, e^{-\alpha\hat{a}^\dagger\hat{a}}] \right) + v e^{-\alpha\hat{a}^\dagger\hat{a}}\hat{a}^\dagger \right] |0\rangle \\
&= \left\{ u \sum_{k=1}^{\infty} \frac{1}{k!} (-\alpha)^k 2k (\hat{a}^\dagger)^{2k-1} + v \sum_{k=0}^{\infty} \frac{1}{k!} (-\alpha)^k (\hat{a}^\dagger)^{2k+1} \right\} |0\rangle \\
&= \sum_{k=0}^{\infty} \frac{1}{k!} (-\alpha)^k \left[-\frac{\alpha}{k+1} \cdot 2(k+1)u + v \right] (\hat{a}^\dagger)^{2k+1} |0\rangle.
\end{aligned} \tag{2.4.22}$$

For this to be 0, each term in the sum must be zero, and hence

$$\alpha = \frac{v}{2u}. \tag{2.4.23}$$

Alternatively: suppose

$$|\tilde{0}\rangle = \sum_{n=0}^{\infty} c_n |n\rangle. \tag{2.4.24}$$

Then

$$\begin{aligned}
\hat{b}|\tilde{0}\rangle &= (u\hat{a} + v\hat{a}^\dagger) \sum_{n=0}^{\infty} c_n |n\rangle \\
&= u \sum_{n=1}^{\infty} c_n \sqrt{n} |n-1\rangle + v \sum_{n=0}^{\infty} c_n \sqrt{n+1} |n+1\rangle \\
&= uc_1 |0\rangle + \sum_{n=1}^{\infty} (uc_{n+1}\sqrt{n+1} + vc_{n-1}\sqrt{n}) |n\rangle.
\end{aligned} \tag{2.4.25}$$

Each term must vanish, so

$$\begin{cases} uc_1 = 0 \\ uc_{n+1}\sqrt{n+1} + vc_{n-1}\sqrt{n} = 0 \quad \text{for } n \geq 1. \end{cases} \tag{2.4.26}$$

The first condition implies $c_1 = 0$, while the second condition gives the iterative formula for $n \geq 2$:

$$c_n = -\frac{v}{u} \sqrt{\frac{n-1}{n}} c_{n-2}. \tag{2.4.27}$$

This means that all the odd terms must vanish:

$$c_{2k+1} = 0 \tag{2.4.28}$$

for $k \in \mathbb{N}$, while the even coefficients are

$$c_{2k} = \left(-\frac{v}{u}\right)^k \sqrt{\frac{(2k-1)!!}{(2k)!!}} c_0. \tag{2.4.29}$$

Therefore,

$$\begin{aligned}
|\tilde{0}\rangle &= \sum_{k=0}^{\infty} c_{2k} |2k\rangle \\
&= \sum_{k=0}^{2k} \left(-\frac{v}{u}\right)^k \sqrt{\frac{(2k-1)!!}{(2k)!!}} \frac{(\hat{a}^\dagger)^{2k}}{(2k)!} |0\rangle \\
&= \sum_{k=0}^{\infty} \left(-\frac{v}{u}\right)^k \frac{1}{(2k)!!} (\hat{a}^\dagger)^{2k} |0\rangle \\
&= \left\{ \sum_{k=0}^{\infty} \frac{1}{k!} \left[-\frac{v}{2u} (\hat{a}^\dagger)^2\right]^k \right\} |0\rangle \\
&= \exp\left(-\frac{v}{2u} \hat{a}^\dagger \hat{a}^\dagger\right) |0\rangle.
\end{aligned} \tag{2.4.30}$$

Note that the new vacuum is not the old vacuum. It contains indefinite, even number of the original bosons.

Exercise 2.5

(a) Let $u_\ell = \phi_{2k}$, $v_\ell = \phi_{2k+1}$. The equation of motion is

$$\begin{cases} m\ddot{u}_\ell = -k(2u_\ell - v_{\ell-1} - v_\ell) \\ M\ddot{v}_\ell = -k(2v_\ell - u_\ell - u_{\ell+1}). \end{cases} \tag{2.5.1}$$

Transforming into the Fourier basis

$$\begin{cases} U_q = \frac{1}{\sqrt{N}} \sum_\ell e^{-2i\ell a q} u_\ell \\ V_q = \frac{1}{\sqrt{N}} \sum_\ell e^{-2i\ell a q} v_\ell, \end{cases} \tag{2.5.2}$$

where q lives in the first Brillouin zone $q \in (-\frac{\pi}{2a}, \frac{\pi}{2a}]$, the equation of motion is

$$\begin{cases} m\ddot{U}_q = -2kU_q + 2k \cos(aq)V_q \\ M\ddot{V}_q = -2kV_q + 2k \cos(aq)U_q. \end{cases} \tag{2.5.3}$$

In matrix notation,

$$\begin{pmatrix} m & 0 \\ 0 & M \end{pmatrix} \begin{pmatrix} \ddot{U}_q \\ \ddot{V}_q \end{pmatrix} = - \begin{pmatrix} 2k & -2k \cos(qa) \\ -2k \cos(qa) & 2k \end{pmatrix} \begin{pmatrix} U_q \\ V_q \end{pmatrix}, \tag{2.5.4}$$

or

$$\mathbf{M}\ddot{\mathbf{U}}_q = -\mathbf{K}_q \mathbf{U}_q. \tag{2.5.5}$$

Using the ansatz $\mathbf{U}_q(t) = \mathbf{A}_q e^{i\omega_q t}$, this is

$$\omega_q^2 \mathbf{M} \mathbf{A}_q = \mathbf{K}_q \mathbf{A}_q. \tag{2.5.6}$$

This is a generalised eigenvalue equation, which is not the easiest to solve. This inspires us to introduce the mass-scaled coordinates

$$\mathbf{X}_q = \begin{pmatrix} X_q \\ Y_q \end{pmatrix} = \begin{pmatrix} \sqrt{m} U_q \\ \sqrt{M} V_q \end{pmatrix} \tag{2.5.7}$$

such that

$$\mathbf{U}_q = \mathbf{M}^{-1/2} \mathbf{X}_q. \tag{2.5.8}$$

Then we have

$$\ddot{\mathbf{X}} = -\mathbf{D}_q \mathbf{X}_q \quad (2.5.9)$$

where

$$\mathbf{D} = \mathbf{M}^{-1/2} \mathbf{K} \mathbf{M}^{-1/2} = \begin{pmatrix} \frac{2k}{m} & -\frac{2k \cos(qa)}{\sqrt{mM}} \\ -\frac{2k \cos(qa)}{\sqrt{mM}} & \frac{2k}{M} \end{pmatrix}. \quad (2.5.10)$$

Now suppose $\mathbf{X}_q(t) = \mathbf{A}_q e^{i\omega_q t}$, we have the eigenvalue equation

$$\mathbf{D}_q \mathbf{A}_q = \omega_q^2 \mathbf{A}_q. \quad (2.5.11)$$

Solving for the eigenvalue equation

$$\det(\mathbf{D}_q - \omega_q^2 \mathbf{I}) = 0 \quad (2.5.12)$$

gives the two eigenvalues

$$\omega_{\pm}^2(q) = k \left(\frac{1}{m} + \frac{1}{M} \right) \pm k \sqrt{\left(\frac{1}{m} + \frac{1}{M} \right)^2 - \frac{4}{mM} \sin^2(qa)}. \quad (2.5.13)$$

We denote the two normalised eigenvectors

$$\mathbf{e}_{\pm}(q) = \begin{pmatrix} e_{\pm,1} \\ e_{\pm,2} \end{pmatrix}. \quad (2.5.14)$$

They satisfy

$$\left(\frac{2k}{m} - \omega_{\pm}^2 \right) e_{\pm,1} - \frac{2k \cos(qa)}{\sqrt{mM}} e_{\pm,2} = 0, \quad (2.5.15)$$

which gives

$$\frac{e_{\pm,2}}{e_{\pm,1}} = \frac{\sqrt{mM} \left(\frac{2k}{m} - \omega_{\pm}^2 \right)}{2k \cos(qa)}. \quad (2.5.16)$$

This should be solved along with the normalisation condition

$$e_{\pm,1}^2 + e_{\pm,2}^2 = 1, \quad (2.5.17)$$

but I do not bother to.

Let

$$\mathbf{E}_q = \begin{pmatrix} | & | \\ \mathbf{e}_+(q) & \mathbf{e}_-(q) \\ | & | \end{pmatrix}, \quad (2.5.18)$$

then

$$\mathbf{E}_q^T \mathbf{D}_q \mathbf{E}_q = \begin{pmatrix} \omega_+^2(q) & 0 \\ 0 & \omega_-^2(q) \end{pmatrix} =: \Omega_q, \quad (2.5.19)$$

and so

$$\ddot{\mathbf{X}}_q = -\mathbf{E}_q \Omega_q \mathbf{E}_q^T \mathbf{X}_q. \quad (2.5.20)$$

Therefore

$$\frac{d^2}{dt^2} (\mathbf{E}_q^T \mathbf{X}_q) = -\Omega_q (\mathbf{E}_q^T \mathbf{X}_q). \quad (2.5.21)$$

This is a diagonal equation of motion, and

$$\mathbf{Q}_q = \mathbf{E}_q^T \mathbf{X}_q \quad (2.5.22)$$


are the normal coordinates. The two normal coordinates are

$$Q_{q,\pm} = \mathbf{e}_{q,\pm}^T \mathbf{X}_q. \quad (2.5.23)$$

They satisfy

$$\ddot{Q}_{q,\pm} = -\omega_{q,\pm}^2 Q_{q,\pm}, \quad (2.5.24)$$

so that each is oscillating independently with frequency $\omega_{q,\pm}$.



(b) At the band edge $q = \frac{\pi}{2a}$,

$$\omega_{\pm}^2 = \frac{k}{mM} [(m+M) \pm (m-M)], \quad (2.5.25)$$

so

$$\begin{cases} \omega_+ = \sqrt{\frac{2k}{M}} \\ \omega_- = \sqrt{\frac{2k}{m}} \end{cases} . \quad (2.5.26)$$

The band gap is

$$\Delta\omega = \sqrt{2k} \left| \frac{1}{\sqrt{M}} - \frac{1}{\sqrt{m}} \right|. \quad (2.5.27)$$

(c) We will denote the momentum conjugate to $\mathbf{X}_q$ as $\mathbf{P}$, and that to $\mathbf{Q}_q$ as $\mathbf{\Pi}_q$. The kinetic energy is

$$\begin{aligned} T &= \frac{1}{2} \sum_q m \left| \dot{\mathbf{U}}_q \right|^2 + M \left| \dot{\mathbf{V}}_q \right|^2 \\ &= \frac{1}{2} \sum_q \dot{\mathbf{X}}_q^\dagger \dot{\mathbf{X}}_q \\ &= \frac{1}{2} \sum_q \mathbf{P}_q^\dagger \mathbf{P}_q \\ &= \frac{1}{2} \sum_q \mathbf{\Pi}_q^\dagger \mathbf{\Pi}_q, \end{aligned} \quad (2.5.28)$$

and the potential energy is

$$\begin{aligned} V &= \frac{1}{2} \sum_q \mathbf{X}_q^\dagger \mathbf{D}_q \mathbf{X}_q \\ &= \frac{1}{2} \sum_q \mathbf{Q}_q^\dagger \Omega_q \mathbf{Q}_q. \end{aligned} \quad (2.5.29)$$

The Hamiltonian is therefore

$$H = \sum_q \sum_{\sigma=\pm} \left[\frac{1}{2} |\Pi_{q,\sigma}|^2 + \frac{1}{2} \omega_{q,\sigma}^2 |Q_{q,\sigma}|^2 \right]. \quad (2.5.30)$$

Quantising:

$$\hat{a}_{q,\sigma} = \sqrt{\frac{\omega_{q,\sigma}}{2\hbar}} Q_{q,\sigma} + \frac{i}{\sqrt{2\hbar\omega_{q,\sigma}}} \Pi_{q,\sigma} \quad (2.5.31)$$

such that

$$[\hat{a}_{q,\sigma}, \hat{a}_{q',\sigma'}^\dagger] = \delta_{qq'} \delta_{\sigma\sigma'}. \quad (2.5.32)$$

The Hamiltonian is

$$\hat{H} = \sum_q \sum_{\sigma=\pm} \left[\hbar\omega_{q,\sigma} \left(\hat{a}_{q,\sigma}^\dagger \hat{a}_{q,\sigma} + \frac{1}{2} \right) \right]. \quad (2.5.33)$$

Exercise 2.6

(a) The RMS displacement

$$\Delta r_{\text{RMS}} = \sqrt{\Delta x^2 + \Delta y^2 + \Delta z^2}, \quad (2.6.1)$$

where in each direction,

$$\Delta x \sim \sqrt{\frac{\hbar}{m\omega}} \quad (2.6.2)$$

by the uncertainty principle. Therefore, by Linderman criterion, the crystal is unstable if

$$\Delta r_{\text{RMS}} \sim \sqrt{\frac{\hbar}{m\omega}} \gtrsim \frac{1}{3}a \implies \frac{\hbar}{m\omega a^2} > \zeta_c \quad (2.6.3)$$

for some dimensionless ζ_c .

(b) Have

$$\begin{aligned} \hat{\Phi}_j &= \frac{1}{\sqrt{N_S}} \sum_{\mathbf{q}} e^{i\mathbf{q} \cdot \mathbf{R}_j} \hat{\Phi}_{\mathbf{q}} \\ &= \sqrt{\frac{\hbar}{2mN_S}} \sum_{\mathbf{q}} e^{i\mathbf{q} \cdot \mathbf{R}_j} \frac{1}{\sqrt{\omega_{\mathbf{q}}}} (\hat{a}_{\mathbf{q}} + \hat{a}_{-\mathbf{q}}^\dagger). \end{aligned} \quad (2.6.4)$$

Take $j = 0$,

$$\hat{\Phi} = \sqrt{\frac{\hbar}{2mN_S}} \sum_{\mathbf{q}} \frac{1}{\sqrt{\omega_{\mathbf{q}}}} (\hat{a}_{\mathbf{q}} + \hat{a}_{-\mathbf{q}}^\dagger). \quad (2.6.5)$$

To work out $\omega_{\mathbf{q}}$, consider

$$\Phi_{\mathbf{R}+\mathbf{a}} - \Phi_{\mathbf{R}} = \frac{1}{\sqrt{N_S}} \sum_{\mathbf{q}} \Phi_{\mathbf{q}} e^{i\mathbf{q} \cdot \mathbf{R}} (e^{i\mathbf{q} \cdot \mathbf{a}} + 1). \quad (2.6.6)$$

By Fourier orthogonality

$$\sum_{\mathbf{R}} e^{i(\mathbf{q}+\mathbf{q}') \cdot \mathbf{R}} = N \delta_{\mathbf{q}', -\mathbf{q}}, \quad (2.6.7)$$

we have

$$\begin{aligned} \sum_{\mathbf{R}} (\Phi_{\mathbf{R}+\mathbf{a}} - \Phi_{\mathbf{R}})^2 &= \sum_{\mathbf{q}} \Phi_{\mathbf{q}} \Phi_{-\mathbf{q}} (e^{i\mathbf{q} \cdot \mathbf{a}} - 1)(e^{-i\mathbf{q} \cdot \mathbf{a}} - 1) \\ &= \sum_{\mathbf{q}} 4 \sin^2 \left(\frac{\mathbf{q} \cdot \mathbf{a}}{2} \right) \Phi_{\mathbf{q}}^\dagger \Phi_{\mathbf{q}}. \end{aligned} \quad (2.6.8)$$

Summing over the three directions $\mathbf{a} \in \{a\hat{\mathbf{x}}, a\hat{\mathbf{y}}, a\hat{\mathbf{z}}\}$,

$$V = \frac{1}{2} m \omega^2 \sum_{\mathbf{q}} 4 \sin \left[\sin^2 \frac{q_x a}{2} + \sin^2 \frac{q_y a}{2} + \sin^2 \frac{q_z a}{2} \right] \Phi_{\mathbf{q}}^\dagger \Phi_{\mathbf{q}}, \quad (2.6.9)$$

and hence

$$\omega_{\mathbf{q}} = 2\omega \sqrt{\sin^2 \frac{q_x a}{2} + \sin^2 \frac{q_y a}{2} + \sin^2 \frac{q_z a}{2}}. \quad (2.6.10)$$

Therefore,

$$\begin{aligned}
\langle 0 | \hat{\Phi}^2 | 0 \rangle &= \frac{\hbar}{2mN_S} \sum_{\mathbf{q}, \mathbf{q}'} \frac{1}{\sqrt{\omega_{\mathbf{q}} \omega_{\mathbf{q}'}}} \langle 0 | (\hat{a}_{\mathbf{q}} + \hat{a}_{-\mathbf{q}}^\dagger)(\hat{a}_{\mathbf{q}'} + \hat{a}_{-\mathbf{q}'}^\dagger) | 0 \rangle \\
&= \frac{\hbar}{2mN_S} \sum_{\mathbf{q}, \mathbf{q}'} \frac{1}{\sqrt{\omega_{\mathbf{q}} \omega_{\mathbf{q}'}}} \underbrace{\langle 0 | \hat{a}_{\mathbf{q}} \hat{a}_{-\mathbf{q}'}^\dagger | 0 \rangle}_{\delta_{\mathbf{q}, -\mathbf{q}'}} \\
&= \frac{\hbar}{2mN_S} \sum_{\mathbf{q}} \frac{1}{\omega_{\mathbf{q}}} .
\end{aligned} \tag{2.6.11}$$

We will approximate the sum by an integral

$$\frac{1}{N_S} \sum_{\mathbf{q}} \longrightarrow \left(\frac{a}{2\pi} \right)^3 \int_{-\pi/a}^{\pi/a} d^3 \mathbf{q} . \tag{2.6.12}$$

Then

$$\begin{aligned}
\langle 0 | \hat{\Phi}^2 | 0 \rangle &= \frac{\hbar}{4\pi\omega} \int_{-\pi}^{\pi} \frac{-\pi}{\pi} \frac{d^3 \mathbf{x}}{(2\pi)^3} \frac{1}{\sqrt{\sin^2(\frac{x}{2}) + \sin^2(\frac{y}{2}) + \sin^2(\frac{z}{2})}} \\
&= \frac{\hbar}{4m\omega} I ,
\end{aligned} \tag{2.6.13}$$

where

$$I = 0.910688 \dots \tag{2.6.14}$$

For the crystal to be unstable,

$$\langle 0 | \hat{\Phi}^2 | 0 \rangle > \frac{\hbar}{4m\omega} I > \frac{1}{9} a^2 , \tag{2.6.15}$$

and so

$$\frac{\hbar}{m\omega a^2} > \frac{4}{9I} =: \zeta_c \approx 0.4880 \dots \tag{2.6.16}$$

(c) Under the unrealistic assumption that ω is independent of a , Helium is *stable* if

$$\frac{\hbar}{m\omega a^2} < \zeta_c , \tag{2.6.17}$$

giving

$$a > a_c = \sqrt{\frac{\hbar}{m\omega \zeta_c}} \approx 2.878 \times 10^{-11} \text{ m} . \tag{2.6.18}$$

This is apparently unrealistic since crystalline Helium should be *unstable* at low density (for a greater than some a_c).

Using the realistic assumption that $\omega \sim a^{-\alpha}$ for some $\alpha > 2$, we now have that Helium is *stable* if

$$\frac{\hbar}{m\omega a^2} = \frac{\hbar}{m\omega_0} a^\gamma < \zeta_c \tag{2.6.19}$$

for some $\gamma = \alpha - 2 > 0$. This is

$$a < \left(\frac{\zeta_c m \omega_0}{\hbar} \right)^{1/\gamma} =: a_c . \tag{2.6.20}$$

Exercise 2.7

(a) Transforming into the Fourier basis,

$$\begin{aligned}
 \sum_j \hat{a}_{j+1}^\dagger \hat{a}_j &= \sum_j \sum_q \sum_p e^{-i(q-p)ja} e^{-iqa} \hat{a}_q^\dagger \hat{a}_p \\
 &= \sum_q \sum_p \delta_{pq} e^{-iqa} \hat{a}_q^\dagger \hat{a}_p \\
 &= \sum_q e^{-iqa} \hat{a}_q^\dagger \hat{a}_q,
 \end{aligned} \tag{2.7.1}$$

and its Hermitian conjugate is

$$\sum_q e^{+iqa} \hat{a}_q^\dagger \hat{a}_q. \tag{2.7.2}$$

Therefore, the first term in the Hamiltonian is

$$\hat{H}_q = J_1 \sum_q 2 \cos(qa) \hat{a}_q^\dagger \hat{a}_q. \tag{2.7.3}$$

Next,

$$\begin{aligned}
 \sum_j \hat{a}_{j+1}^\dagger \hat{a}_j^\dagger &= \sum_j \sum_q \sum_p e^{-i(q+p)ja} e^{-iqa} \hat{a}_q^\dagger \hat{a}_p^\dagger \\
 &= \sum_q \sum_p \delta_{p(-q)} e^{-iqa} \hat{a}_q^\dagger \hat{a}_p^\dagger \\
 &= \sum_q e^{-iqa} \hat{a}_q^\dagger \hat{a}_{-q}^\dagger,
 \end{aligned} \tag{2.7.4}$$

Since $\hat{a}_q^\dagger$ commutes with $\hat{a}_{-q}^\dagger$, we can set $q \rightarrow -q$ and write

$$\sum_j \hat{a}_{j+1}^\dagger \hat{a}_j^\dagger = \sum_q e^{+iqa} \hat{a}_q^\dagger \hat{a}_{-q}^\dagger. \tag{2.7.5}$$

Taking the average of the two expressions,

$$\sum_j \hat{a}_{j+1}^\dagger \hat{a}_j^\dagger = \sum_q \cos(qa) \hat{a}_q^\dagger \hat{a}_{-q}^\dagger, \tag{2.7.6}$$

and its Hermitian conjugate is

$$\sum_q \cos(qa) \hat{a}_q \hat{a}_{-q}, \tag{2.7.7}$$

so the second part of the Hamiltonian is

$$\hat{H}_2 = J_2 \sum_j \cos(qa) (\hat{a}_q^\dagger \hat{a}_{-q}^\dagger + \hat{a}_q \hat{a}_{-q}). \tag{2.7.8}$$

The full Hamiltonian is

$$\hat{H} = \sum_q \left[\underbrace{2J_1 \cos(qa)}_{\omega_q} \hat{a}_q^\dagger \hat{a}_q + \underbrace{J_2 \cos(qa)}_{\frac{1}{2}\Delta_q} (\hat{a}_q^\dagger \hat{a}_{-q}^\dagger + \hat{a}_q \hat{a}_{-q}) \right] \tag{2.7.9}$$

Now we carry out canonical transformation as we did in Exercise 2.4:

$$\hat{b}_q = u_q \hat{a}_q + v_q \hat{a}_{-q}^\dagger. \tag{2.7.10}$$

This is slightly different but you can check that it does transform the Hamiltonian in the same way. Note that the q -dependent factor in ω_q and $\frac{1}{2}\Delta_q$ is identical and hence can be factor out, so if the canonical transformation were to diagonalise $\hat{H}$, the u_q and v_q should be independent of q . We will denote them as $u = \cosh \theta$ and $v = \sinh \theta$.

Now the condition for $\hat{H}$ to be diagonalisable is

$$\tanh 2\theta = \frac{\omega_q}{\Delta_q} = \frac{J_2}{J_1} \in (-1, 1). \quad (2.7.11)$$

This gives

$$\hat{H} = \sum_q \tilde{\omega}_q \left(\hat{b}_q \hat{b}_{-q} + \frac{1}{2} \right), \quad (2.7.12)$$

where

$$\tilde{\omega}_q = \sqrt{\omega_q^2 - \Delta_q^2} = 2 \cos(qa) \sqrt{J_2^2 - J_1^2}. \quad (2.7.13)$$

As discussed in Exercise 2.4, when $J_1 = J_2$, the transition is singular and

$$\begin{aligned} \hat{H} &= J \sum_j (\hat{a}_{j+1} + \hat{a}_{j+1}^\dagger)(\hat{a}_j + \hat{a}_j^\dagger) \\ &= 2J \sum_j \hat{x}_{j+1} \hat{x}_j. \end{aligned} \quad (2.7.14)$$

Exercise 2.8

(a) From Schrödinger equation,

$$i\hbar \partial_t e^{-i\hat{H}_0 t/\hbar} |\psi_I(t)\rangle = (\hat{H}_0 + \hat{V}(t)) e^{-i\hat{H}_0 t/\hbar} |\psi_I(t)\rangle. \quad (2.8.1)$$

Expanding the derivative,

$$i\hbar \left[-\frac{i\hat{H}_0}{\hbar} e^{-i\hat{H}_0 t/\hbar} + e^{-i\hat{H}_0 t/\hbar} \partial_t \right] |\psi_I(t)\rangle = (\cancel{\hat{H}_0} + \hat{V}(t)) e^{-i\hat{H}_0 t/\hbar} |\psi_I(t)\rangle, \quad (2.8.2)$$

giving

$$i\hbar e^{-i\hat{H}_0 t/\hbar} \partial_t |\psi_I(t)\rangle = \hat{V}(t) e^{-i\hat{H}_0 t/\hbar} |\psi_I(t)\rangle, \quad (2.8.3)$$

and so

$$i\hbar \partial_t |\psi_I(t)\rangle = \hat{V}_I(t) |\psi_I(t)\rangle, \quad (2.8.4)$$

where

$$\begin{aligned} \hat{V}_I(t) &= e^{+i\hat{H}_0 t/\hbar} (-f(t)\hat{x}) e^{-i\hat{H}_0 t/\hbar} \\ &= -f(t)\hat{x}(t). \end{aligned} \quad (2.8.5)$$

(b) Solving

$$i\hbar \partial_t |\psi_I(t)\rangle = \hat{V}_I(t) |\psi_I(t)\rangle \quad (2.8.6)$$

iteratively gives

$$\begin{aligned} |\psi_I(t)\rangle &= |\psi_I(-\infty)\rangle + \int_{-\infty}^t dt' \partial_{t'} |\psi_I(t')\rangle \\ &= |0\rangle - \frac{i}{\hbar} \int_{-\infty}^t dt' \hat{V}_I(t') \underbrace{|\psi_I(t')\rangle}_{|0\rangle - \frac{i}{\hbar} \int_{-\infty}^{t'} dt'' \hat{V}_I(t'') |\psi_I(t'')\rangle}. \end{aligned} \quad (2.8.7)$$

Truncating this series to the $O(\hat{V}_I)$ gives

$$\begin{aligned} |\psi_I(t)\rangle &= |0\rangle - \frac{i}{\hbar} \int_{-\infty}^t dt' \hat{V}_I(t') |0\rangle + O(\hat{V}_I^2) \\ &= |0\rangle + \frac{i}{\hbar} \int_{-\infty}^t dt' f(t') \hat{x}(t') |0\rangle + O(f^2), \end{aligned} \quad (2.8.8)$$

and

$$\langle \psi_I(t) | = \langle 0 | - \frac{i}{\hbar} \int_{-\infty}^t dt' f(t') \langle 0 | \hat{x}(t') + O(f^2). \quad (2.8.9)$$

(c)

$$\begin{aligned} \langle \hat{x}(t) \rangle &= \langle \psi_I(t) | \hat{x}(t) | \psi_I(t) \rangle \\ &= \langle 0 | \hat{x}(t) | 0 \rangle - \frac{i}{\hbar} \int_{-\infty}^t dt' f(t') \langle 0 | \hat{x}(t') \hat{x}(t) | 0 \rangle + \frac{i}{\hbar} \int_{-\infty}^t dt'' f(t'') \langle 0 | \hat{x}(t) \hat{x}(t'') | 0 \rangle + O(f^2) \\ &= \underbrace{\langle 0 | \hat{x}(t) | 0 \rangle}_0 + \frac{i}{\hbar} \int_{-\infty}^t dt' f(t') \underbrace{\langle 0 | [\hat{x}(t) \hat{x}(t')] | 0 \rangle}_{R(t-t')} + O(f^2) \\ &= \int_{-\infty}^t dt' R(t-t') f(t') + O(f^2). \end{aligned} \quad (2.8.10)$$

If we define $f(t)$ such that it is switched on only after $t = 0$, and define $R(\tau) = 0$ for $\tau < 0$, then we can rewrite

$$\langle \hat{x}(t) \rangle = \int_0^\infty dt' R(t-t') f(t') + O(f^2). \quad (2.8.11)$$

3 Conserved Particles

Exercise 3.1

(a) For $\hat{c}_3^\dagger \hat{c}_6 \hat{c}_4 \hat{c}_6^\dagger \hat{c}_3 |11111000 \dots\rangle$, the first $\hat{c}_3$ removes the particle at state 3, but then $\hat{c}_6^\dagger$ creates a particle at state 6, which is already occupied. This is not allowed for fermions, so

$$\hat{c}_3^\dagger \hat{c}_6 \hat{c}_4 \hat{c}_6^\dagger \hat{c}_3 |11111000 \dots\rangle = 0. \quad (3.1.1)$$

(b) To get from $|1111100000 \dots\rangle$ to $|1101100100 \dots\rangle$, we would like to remove a particle at state 3, then create a particle at state 8, so

$$|1101100100 \dots\rangle = (-1)^P \hat{c}_8^\dagger \hat{c}_3 |1111100000 \dots\rangle, \quad (3.1.2)$$

with the sign yet to be determined. While acting $\hat{c}_3$ on the Fermi sea state,

$$\hat{c}_3 |1111100000 \dots\rangle \equiv \hat{c}_3 \hat{c}_5^\dagger \hat{c}_4^\dagger \hat{c}_3^\dagger \hat{c}_2^\dagger \hat{c}_1^\dagger |0\rangle, \quad (3.1.3)$$

we need to move $\hat{c}_3$ through two creation operators of higher state, each time we get a minus sign. Therefore, the overall parity to destroy the particle at state 3 is positive. Then we can create the particle at state 8, which is the highest state so we don't need to move anything. Therefore,

$$|1101100100 \dots\rangle = +\hat{c}_8^\dagger \hat{c}_3 |1111100000 \dots\rangle. \quad (3.1.4)$$

(c)

$$\begin{aligned} \langle \Psi | \hat{N} | \Psi \rangle &= |A|^2 \langle 100 | \hat{N} | 100 \rangle + |B|^2 \langle 111000 | \hat{N} | 111000 \rangle \\ &\quad + A^* B \langle 100 | \hat{N} | 111000 \rangle + B^* A \langle 111000 | \hat{N} | 100 \rangle \\ &= |A|^2 + 3 |B|^2. \end{aligned} \quad (3.1.5)$$

If the state is normalised (as it should to make this problem meaningful), $|A|^2 + |B|^2 = 1$, so

$$\langle \Psi | \hat{N} | \Psi \rangle = 1 + 2 |B|^2. \quad (3.1.6)$$

Exercise 3.2

This is similar to Exercise 2.4, just now we have fermions instead of bosons.

(a) We have a lot of canonical anticommutation relations to check between 4 operators! But luckily, most of them vanish (as they should) by the anticommutation relations of a . The only possibly non-vanishing ones are those pairing $\hat{a}_i$ with $\hat{a}_i^\dagger$: they are

$$\{\hat{c}_1, \hat{c}_2\}, \{\hat{c}_1, \hat{c}_1^\dagger\}, \{\hat{c}_2, \hat{c}_2^\dagger\}, \{\hat{c}_1^\dagger, \hat{c}_2^\dagger\}, \quad (3.2.1)$$

in which we want $\{\hat{c}_1, \hat{c}_1^\dagger\} = \{\hat{c}_2, \hat{c}_2^\dagger\} = 1$ and the other two to be zero.

We have

$$\begin{aligned} \{\hat{c}_1, \hat{c}_2\} &= -uv \underbrace{\{\hat{a}_1, \hat{a}_1^\dagger\}}_1 + uv \underbrace{\{\hat{a}_2, \hat{a}_2^\dagger\}}_2 \\ &= 0 \end{aligned} \quad (3.2.2)$$

and

$$\{\hat{c}_1^\dagger, \hat{c}_2^\dagger\} = (\{\hat{c}_1, \hat{c}_2\})^\dagger = 0 \quad (3.2.3)$$

as expected.

$$\begin{aligned} \{\hat{c}_1, \hat{c}_1^\dagger\} &= u^2 \{\hat{a}_1, \hat{a}_1^\dagger\} + v^2 \{\hat{a}_2^\dagger, \hat{a}_2\} \\ &= u^2 + v^2, \end{aligned} \quad (3.2.4)$$

$$\begin{aligned} \{\hat{c}_2, \hat{c}_2^\dagger\} &= v^2 \{\hat{a}_1^\dagger, \hat{a}_1\} + u^2 \{\hat{a}_2, \hat{a}_2^\dagger\} \\ &= u^2 + v^2. \end{aligned} \quad (3.2.5)$$

Therefore the anticommutator relations of c is satisfied if and only if $u^2 + v^2 = 1$.

(b) The Hamiltonian is

$$\hat{H} = \epsilon(\hat{a}_1^\dagger \hat{a}_1 - \hat{a}_2 \hat{a}_2^\dagger) + \Delta(\hat{a}_1^\dagger \hat{a}_2^\dagger + \text{H.C.}). \quad (3.2.6)$$

The first term is

$$\begin{aligned} \hat{a}_1^\dagger \hat{a}_1 &= (u\hat{c}_1^\dagger - v\hat{c}_2)(u\hat{c}_1 - v\hat{c}_2^\dagger) \\ &= u^2 \hat{c}_1^\dagger \hat{c}_1 - uv \hat{c}_1^\dagger \hat{c}_2^\dagger - vu \hat{c}_2 \hat{c}_1 + v^2 \underbrace{\hat{c}_2 \hat{c}_2^\dagger}_{1 - \hat{c}_2^\dagger \hat{c}_2}, \end{aligned} \quad (3.2.7)$$

and

$$\begin{aligned} \hat{a}_2 \hat{a}_2^\dagger &= (v\hat{c}_1^\dagger + u\hat{c}_2)(v\hat{c}_1 + u\hat{c}_2^\dagger) \\ &= v^2 \hat{c}_1^\dagger \hat{c}_1 + uv \hat{c}_1^\dagger \hat{c}_2^\dagger + uv \hat{c}_2 \hat{c}_1 + u^2 \hat{c}_2 \hat{c}_2^\dagger, \end{aligned} \quad (3.2.8)$$

and so

$$\hat{a}_1^\dagger \hat{a}_1 - \hat{a}_2 \hat{a}_2^\dagger = (u^2 - v^2)(\hat{c}_1^\dagger \hat{c}_1 + \hat{c}_2^\dagger \hat{c}_2 - 1) - 2uv(\hat{c}_1^\dagger \hat{c}_2^\dagger + \hat{c}_2 \hat{c}_1). \quad (3.2.9)$$

The second term is

$$\begin{aligned} \hat{a}_1^\dagger \hat{a}_2^\dagger &= (u\hat{c}_1^\dagger - v\hat{c}_2)(v\hat{c}_1 + u\hat{c}_2^\dagger) \\ &= uv \hat{c}_1^\dagger \hat{c}_1 + u^2 \hat{c}_1^\dagger \hat{c}_2^\dagger - v^2 \hat{c}_2 \hat{c}_1 - \underbrace{uv \hat{c}_2 \hat{c}_2^\dagger}_{+uv \hat{c}_2^\dagger \hat{c}_2 - uv}, \end{aligned} \quad (3.2.10)$$

and its Hermitian conjugate is

$$\hat{a}_2 \hat{a}_1 = uv \hat{c}_1^\dagger \hat{c}_1 + u^2 \hat{c}_2 \hat{c}_1 - v^2 \hat{c}_1^\dagger \hat{c}_2^\dagger - vu \hat{c}_2 \hat{c}_2^\dagger, \quad (3.2.11)$$

so

$$\hat{a}_1^\dagger \hat{a}_2^\dagger + \text{H.C.} = 2uv(\hat{c}_1^\dagger \hat{c}_1 + \hat{c}_2^\dagger \hat{c}_2 - 1) + (u^2 - v^2)(\hat{c}_1^\dagger \hat{c}_2^\dagger + \hat{c}_2 \hat{c}_1). \quad (3.2.12)$$

Therefore, the Hamiltonian is

$$\hat{H} = [\epsilon(u^2 - v^2) + 2\Delta uv](\hat{c}_1^\dagger \hat{c}_1 + \hat{c}_2^\dagger \hat{c}_2 - 1) + [\Delta(u^2 - v^2) - 2\epsilon uv](\hat{c}_1^\dagger \hat{c}_2^\dagger + \hat{c}_2 \hat{c}_1). \quad (3.2.13)$$

The factor for the latter term should vanish for $\hat{H}$ to be diagonalisable. Since $u^2 + v^2 = 1$, we set $u = \cos \theta$ and $v = \sin \theta$, so

$$\Delta \cos 2\theta = \epsilon \sin 2\theta, \quad (3.2.14)$$

and so

$$\frac{\Delta}{\epsilon} = \tan 2\theta \in \mathbb{R}. \quad (3.2.15)$$

Therefore, the Hamiltonian for $\epsilon \neq 0$.

We have

$$\sin 2\theta = \frac{\Delta}{\sqrt{\Delta^2 + \epsilon^2}} \quad \cos 2\theta = \frac{\epsilon}{\sqrt{\Delta^2 + \epsilon^2}} \quad (3.2.16)$$

and so

$$\epsilon(u^2 - v^2) + 2\Delta uv = \sqrt{\Delta^2 + \epsilon^2}. \quad (3.2.17)$$

The Hamiltonian is

$$\hat{H} = \sqrt{\Delta^2 + \epsilon^2}(\hat{c}_1^\dagger \hat{c}_1 + \hat{c}_2^\dagger \hat{c}_2 - 1). \quad (3.2.18)$$

(c)

$$\hat{H} |\tilde{0}\tilde{0}\rangle = -\sqrt{\Delta^2 + \epsilon^2} |\tilde{0}\tilde{0}\rangle. \quad (3.2.19)$$

(d) I believe that there is a typo in this question. It should be expanding the c -vacuum state $|\tilde{0}\tilde{0}\rangle$ in terms of the a -fermion states $|n_1 n_2\rangle$:

$$|\tilde{0}\tilde{0}\rangle = b_1 |00\rangle + b_2 |10\rangle + b_3 |01\rangle + b_4 |11\rangle. \quad (3.2.20)$$

We require that $\hat{c}_i |\tilde{0}\tilde{0}\rangle = 0$ for $i = 1, 2$.

$$\begin{aligned} \hat{c}_1 |\tilde{0}\tilde{0}\rangle &= (u\hat{a}_1 + v\hat{a}_2) |\tilde{0}\tilde{0}\rangle \\ &= ub_2 |00\rangle + (vb_1 - ub_4) |01\rangle + vb_2 |11\rangle, \end{aligned} \quad (3.2.21)$$

which requires

$$\begin{cases} b_2 = 0 \\ ub_4 - vb_1 = 0. \end{cases} \quad (3.2.22)$$

$$\begin{aligned} \hat{c}_2 |\tilde{0}\tilde{0}\rangle &= (-v\hat{a}_1 + u\hat{a}_2) |\tilde{0}\tilde{0}\rangle \\ &= ub_3 |00\rangle - (vb_1 - ub_4) |01\rangle + vb_3 |11\rangle, \end{aligned} \quad (3.2.23)$$

which requires

$$\begin{cases} b_3 = 0 \\ ub_4 - vb_1 = 0. \end{cases} \quad (3.2.24)$$

This is solved by $b_1 = u$ and $b_4 = v$. Therefore

$$\begin{aligned} |\tilde{0}\tilde{0}\rangle &= u |00\rangle + v |11\rangle \\ &= (u + v\hat{a}_2^\dagger \hat{a}_1^\dagger) |00\rangle. \end{aligned} \quad (3.2.25)$$

Exercise 3.3

(a) Substituting

$$\rho(\mathbf{x}) = \sum_{j=1}^N \delta(\mathbf{x} - \mathbf{x}_j), \quad (3.3.1)$$

the right hand side can be expanded as

$$\begin{aligned}
\text{RHS} &= \frac{1}{2} \int d^3\mathbf{x} d^3\mathbf{y} V(\mathbf{x} - \mathbf{y}) \left[\sum_i \delta(\mathbf{x} - \mathbf{x}_i) \right] \left[\sum_j \delta(\mathbf{y} - \mathbf{x}_j) \right] - \frac{1}{2} V(0)N \\
&= \sum_{j < i} \int d^3\mathbf{x} d^3\mathbf{y} V(\mathbf{x} - \mathbf{y}) \delta(\mathbf{x} - \mathbf{x}_i) \delta(\mathbf{y} - \mathbf{x}_j) \\
&\quad + \frac{1}{2} \sum_{i=1}^N \int d^3\mathbf{x} d^3\mathbf{y} V(\mathbf{x} - \mathbf{y}) \delta(\mathbf{x} - \mathbf{x}_i) \delta(\mathbf{y} - \mathbf{x}_i) - \frac{1}{2} V(0)N \\
&= \sum_{i < j} V(\mathbf{x}_i - \mathbf{x}_j) + \frac{1}{2} V(0)N - \frac{1}{2} V_0 N \\
&= \sum_{i < j} V(\mathbf{x}_i - \mathbf{x}_j) = V.
\end{aligned} \tag{3.3.2}$$

We see that $\frac{1}{2} V_0 N$ is exactly the self-interaction energy.

(b) Direct second quantisation gives

$$\begin{aligned}
\hat{V} &= \frac{1}{2} \int d^3\mathbf{x} d^3\mathbf{y} V(\mathbf{x} - \mathbf{y}) \hat{\rho}(\mathbf{x}) \hat{\rho}(\mathbf{y}) - \frac{1}{2} \int d^3\mathbf{x} V(0) \hat{\rho}(\mathbf{x}) \\
&= \frac{1}{2} \int d^3\mathbf{x} d^3\mathbf{y} V(\mathbf{x} - \mathbf{y}) \hat{\phi}^\dagger(\mathbf{x}) \hat{\phi}(\mathbf{x}) \hat{\phi}^\dagger(\mathbf{y}) \hat{\phi}(\mathbf{y}) - \frac{1}{2} \int d^3\mathbf{x} V(0) \hat{\phi}^\dagger(\mathbf{x}) \hat{\phi}(\mathbf{x}).
\end{aligned} \tag{3.3.3}$$

We have

$$\begin{aligned}
\hat{\phi}^\dagger(\mathbf{x}) \hat{\phi}(\mathbf{x}) \hat{\phi}^\dagger(\mathbf{y}) \hat{\phi}(\mathbf{y}) &= \hat{\phi}^\dagger(\mathbf{x}) \hat{\phi}(\mathbf{y}) \delta(\mathbf{x} - \mathbf{y}) \pm \hat{\phi}^\dagger(\mathbf{x}) \hat{\phi}^\dagger(\mathbf{y}) \hat{\phi}(\mathbf{x}) \hat{\phi}(\mathbf{y}) \\
&= \hat{\phi}^\dagger(\mathbf{x}) \hat{\phi}(\mathbf{y}) \delta(\mathbf{x} - \mathbf{y}) + \underbrace{\hat{\phi}^\dagger(\mathbf{y}) \hat{\phi}^\dagger(\mathbf{x}) \hat{\phi}(\mathbf{x}) \hat{\phi}(\mathbf{y})}_{:\hat{\rho}(\mathbf{x})\hat{\rho}(\mathbf{y}):}.
\end{aligned} \tag{3.3.4}$$

Therefore,

$$\begin{aligned}
\hat{V} &= \frac{1}{2} \int d^3\mathbf{x} d^3\mathbf{y} V(\mathbf{x} - \mathbf{y}) : \hat{\rho}(\mathbf{x}) \hat{\rho}(\mathbf{y}) : \\
&\quad + \frac{1}{2} \int d^3\mathbf{x} d^3\mathbf{y} V(\mathbf{x} - \mathbf{y}) \hat{\phi}^\dagger(\mathbf{x}) \hat{\phi}(\mathbf{y}) \delta(\mathbf{x} - \mathbf{y}) - \frac{1}{2} \int d^3\mathbf{x} d^3\mathbf{y} V(0) \hat{\rho}(\mathbf{x}) \\
&= \frac{1}{2} \int d^3\mathbf{x} d^3\mathbf{y} V(\mathbf{x} - \mathbf{y}) : \hat{\rho}(\mathbf{x}) \hat{\rho}(\mathbf{y}) :,
\end{aligned} \tag{3.3.5}$$

where the last two terms cancel.

Exercise 3.4

(a)

$$\hat{V} = \frac{1}{2} \int d^3\mathbf{x} d^3\mathbf{y} V(\|\mathbf{x} - \mathbf{y}\|) : \hat{\rho}(\mathbf{x}) \hat{\rho}(\mathbf{y}) :. \tag{3.4.1}$$

(b) Commutation relation in momentum basis:

$$\begin{aligned}
[\hat{c}_{\mathbf{k}}, \hat{c}_{\mathbf{k}'}^\dagger] &= \int d^3\mathbf{r} d^3\mathbf{r}' \underbrace{[\hat{\phi}(\mathbf{r}), \hat{\phi}^\dagger(\mathbf{r}')]_{\pm}}_{\delta(\mathbf{r} - \mathbf{r}')} e^{-i\mathbf{k} \cdot \mathbf{r}} e^{i\mathbf{k}' \cdot \mathbf{r}'} \\
&= \int d^3\mathbf{r} e^{-i\mathbf{k} \cdot \mathbf{r}} e^{i\mathbf{k}' \cdot \mathbf{r}} \\
&= (2\pi)^3 \delta(\mathbf{k} - \mathbf{k}').
\end{aligned} \tag{3.4.2}$$

Fourier transform of potential:

$$\begin{aligned}
 V(\mathbf{q}) &= \int d^3\mathbf{r} V(\mathbf{r}) e^{-i\mathbf{q}\cdot\mathbf{r}} \\
 &= \int r^2 \sin\theta dr d\theta d\phi V(r) e^{-iqr \cos\theta} \\
 &= 2\pi \int_0^\infty dr r^2 V(r) \int_{-1}^1 d(\cos\theta) e^{-iqr \cos\theta} \\
 &= \frac{4\pi U}{q} \int_0^R dr r \sin qr \\
 &= \frac{4\pi U}{q^3} [\sin qR - qR \cos qR] .
 \end{aligned} \tag{3.4.3}$$

Exercise 3.5

(a) Expanding the trace in the N -particle energy eigenstate,

$$\begin{aligned}
 F &= -k_B T \ln \text{tr}[e^{-\beta(\hat{H}-\mu\hat{N})}] \\
 &= -k_B T \ln \left(\sum_n e^{-\beta(E_n - \mu N)} \right) .
 \end{aligned} \tag{3.5.1}$$

The partial derivative with respect to μ is

$$\begin{aligned}
 -\frac{\partial F}{\partial \mu} &= k_B T \cdot \frac{\sum_n \frac{\partial}{\partial \mu} e^{-\beta(E_n - \mu N)}}{\sum_n e^{-\beta(E_n - \mu N)}} \\
 &= k_B T \cdot \frac{\sum_n \beta N e^{-\beta(E_n - \mu N)}}{\sum_n e^{-\beta(E_n - \mu N)}} \\
 &= N .
 \end{aligned} \tag{3.5.2}$$

(b) Considering the single level as a grand canonical ensemble, the grand partition function is

$$\Xi = \sum_{n=0}^{\infty} e^{-\beta(\epsilon - \mu)n} = \sum_{n=0}^{\infty} x^n = \frac{1}{1-x} , \tag{3.5.3}$$

where we denoted $x = e^{-\beta(\epsilon - \mu)}$. We must have $x < 1$, or $\mu < \epsilon$, for convergence.

The average particle number in this grand ensemble is given by

$$\langle \hat{n} \rangle = -\frac{\partial \Omega}{\partial \mu} = \frac{1}{\beta} \frac{\partial \ln \Xi}{\partial \mu} , \tag{3.5.4}$$

where $\Omega = -k_B T \ln \Xi$ is the grand (Landau) potential.² Therefore,

$$\begin{aligned}
 \langle \hat{n} \rangle &= \frac{1}{\beta \Xi} \frac{\partial \Xi}{\partial \mu} \\
 &= \frac{1}{e^{\beta(\epsilon - \mu)} - 1} .
 \end{aligned} \tag{3.5.5}$$

²The proof is similar to the above one in canonical ensemble with conserved particle, just now we also need to sum over the particle numbers in expanding the trace.

Exercise 3.6

(a) Writing the delta functions in Laplace transforms, the microcanonical density of state is

$$\begin{aligned} W(E, N) &= \text{tr}[\delta(E - \hat{H})\delta(N - \hat{N})] \\ &= \int_{\beta_0 - i\infty}^{\beta_0 + i\infty} \frac{d\beta}{2\pi i} \int_{\lambda_0 - i\infty}^{\lambda_0 + i\infty} \frac{d\lambda}{2\pi i} e^{+\beta E} e^{-\lambda N} \text{tr}[e^{-\beta \hat{H}} e^{+\lambda \hat{N}}]. \end{aligned} \quad (3.6.1)$$

If we identify $\lambda = \beta\mu$, then

$$\text{tr}[e^{-\beta(\hat{H} - \mu \hat{N})}] = \Xi(\beta, \mu) \quad (3.6.2)$$

is the grand partition function, and

$$\begin{aligned} W(E, N) &= \int \frac{d\beta}{2\pi i} \int \frac{d\lambda}{2\pi i} e^{\beta(E - \mu N)} \Xi \\ &= \int \frac{d\beta}{2\pi i} \int \frac{d\lambda}{2\pi i} e^{\beta(E - \mu N) + \ln \Xi} \\ &=: \int \frac{d\beta}{2\pi i} \int \frac{d\lambda}{2\pi i} e^{\Phi(\beta, \mu)}. \end{aligned} \quad (3.6.3)$$

All terms in $\Phi = \beta(E - \mu N) + \ln \Xi$ are extensive, so in a macroscopic system, this integral is dominated by its very sharp peak at $\Phi(\beta^*, \mu^*)$, where

$$\left. \frac{\partial \Phi}{\partial \beta} \right|_{(\beta^*, \mu^*)} = \left. \frac{\partial \Phi}{\partial \lambda} \right|_{(\beta^*, \mu^*)} = 0, \quad (3.6.4)$$

i.e. when

$$E = -\frac{\partial \ln \Xi}{\partial \beta} = \langle \hat{H} \rangle, \quad (3.6.5)$$

$$N = -\frac{\partial \ln \Xi}{\partial (\beta\mu)} = \langle \hat{N} \rangle. \quad (3.6.6)$$

This is to say that in the thermodynamic limit, a microcanonical and a grand canonical ensemble is equivalent, with (E, N) in the microcanonical ensemble the same as $(\langle \hat{H} \rangle, \langle \hat{N} \rangle)$ in the grand canonical ensemble.

Therefore,

$$\begin{aligned} \ln W &= \Phi(\beta^*, \lambda^*) + O(\ln N) \\ &= \ln \Xi - \beta^*(E - \mu^* N) + O(\ln N). \end{aligned} \quad (3.6.7)$$

For thermodynamics, we have

$$\Omega = -k_B T \ln \Xi = -k_B T \ln \Xi = E - TS - \mu N. \quad (3.6.8)$$

Therefore,

$$\begin{aligned} S &= k_B (\ln \Xi - \beta E - \beta \mu N) \\ &= k_B \ln W. \end{aligned} \quad (3.6.9)$$

(b) The microcanonical average is

$$\begin{aligned}
\langle A \rangle_M &= \frac{\text{tr}[\delta(E - \hat{H})\delta(N - \hat{N})\hat{A}]}{W(E, N)} \\
&= \frac{\int \frac{d\beta d\lambda}{(2\pi i)} e^{\beta E - \lambda N} \text{tr}[e^{-\beta \hat{H} + \lambda \hat{N}} \hat{A}]}{\int \frac{d\beta d\lambda}{(2\pi i)} e^{\beta E - \lambda N} \text{tr}[e^{-\beta \hat{H} + \lambda \hat{N}}]} \\
&= \frac{\int \frac{d\beta d\lambda}{(2\pi i)} e^{\beta E - \lambda N} \Xi(\beta, \mu) \langle \hat{A} \rangle_{\beta, \lambda}}{\int \frac{d\beta d\lambda}{(2\pi i)} e^{\beta E - \lambda N} \Xi(\beta, \mu)} \\
&= \frac{\Xi(\beta^*, \mu^*) \langle \hat{A} \rangle_{\beta^*, \lambda^*}}{\Xi(\beta^*, \mu^*)} \\
&= \langle \hat{A} \rangle_{\beta^*, \mu^*}, \tag{3.6.10}
\end{aligned}$$

which is the grand canonical average at equilibrium, where we again assumed $e^{\Phi(\beta, \lambda)}$ to be sharply peaked in the thermodynamic limit.

Also,

$$\begin{aligned}
\langle A \rangle_M &= \frac{\Xi(\beta^*, \mu^*) \langle \hat{A} \rangle_{\beta^*, \lambda^*}}{\Xi(\beta^*, \mu^*)} \\
&= \frac{1}{\Xi(\beta^*, \mu^*)} \text{tr}[e^{-\beta^* (\hat{H} - \mu^* \hat{N})} \hat{A}] \\
&= \text{tr}[\hat{\rho}_B \hat{A}]. \tag{3.6.11}
\end{aligned}$$