

Problem 1a, p20:

Let $f : A \rightarrow B$. Let $A_0 \subset A$, $B_0 \subset B$.

Let $a \in A_0$. Then $f(a) \in f(A_0)$. Now, $a \in f^{-1}(f(A_0))$ if there is $a' \in f(A_0)$ with $f(a) = a'$. Because $f(a) \in f(A_0)$, there is $a' \in f(A_0)$ with $f(a) = a'$. So $a \in f^{-1}(f(A_0))$.

So, $A_0 \subset f^{-1}(f(A_0))$.

Now, let f be injective and $a \in f^{-1}(f(A_0))$. Then there is $b \in f(A_0)$ with $f(a) = b$. Now, because $b \in f(A_0)$, we have that $b = f(a')$ for some $a' \in A_0$. Because f is injective, $f(a) = b = f(a')$ implies that $a = a'$. So, because $a' \in A_0$, this means that $a \in A_0$.

So, $A_0 \supset f^{-1}(f(A_0))$ if f is injective. So, $A_0 = f^{-1}(f(A_0))$ if f is injective.

Problem 1b, p20:

Let $f : A \rightarrow B$. Let $A_0 \subset A$, $B_0 \subset B$.

Let $b \in f(f^{-1}(B_0))$. Then there is an $a \in f^{-1}(B_0)$ with $f(a) = b$. Now, because $a \in f^{-1}(B_0)$, there is $b' \in B_0$ with $f(a) = b'$. Now, f is a function. So $f(a) = b' = b$. So $b \in B_0$.

So, $f(f^{-1}(B_0)) \subset B_0$.

Now, let f be surjective and $b \in B_0$. Then there is $a \in A$ with $f(a) = b$, as f is surjective. That is, $f^{-1}(B_0)$ is nonempty, and there is $a \in f^{-1}(B_0)$ with $f(a) = b$. Now, $f(a) = b$, so $b \in f(f^{-1}(B_0))$.

So, $f(f^{-1}(B_0)) \supset B_0$ if f is surjective. So, $f(f^{-1}(B_0)) = B_0$ if f is surjective.

Problem 2g, p20:

Let $f : A \rightarrow B$ and let $A_i \subset A$ and $B_i \subset B$ for $i = 0$ and $i = 1$.

Let $b \in f(A_0 \cap A_1)$. Then there is $a \in A_0 \cap A_1$ with $f(a) = b$. So there is $a \in A_0$ with $f(a) = b$; that is, $b \in f(A_0)$. Also there is $a \in A_1$ with $f(a) = b$; that is, $b \in f(A_1)$. So $b \in f(A_0) \cap f(A_1)$, because $b \in f(A_0)$ and $b \in f(A_1)$.

So, $f(A_0 \cap A_1) \subset f(A_0) \cap f(A_1)$.

Now, let f be injective and $b \in f(A_0) \cap f(A_1)$. Then there is $a_0 \in A_0$ with $f(a_0) = b$. Also, there is $a_1 \in A_1$ with $f(a_1) = b$. Because f is injective, $f(a_0) = b = f(a_1)$ implies that $a_0 = a_1$. So $a_0 \in A_1$. So $a_0 \in A_0 \cap A_1$. So because $f(a_0) = b$, we have that $b = f(a)$ for some $a \in A_0 \cap A_1$. So $b \in f(A_0 \cap A_1)$.

So, $f(A_0 \cap A_1) \supset f(A_0) \cap f(A_1)$ if f is injective. So if f is injective, $f(A_0 \cap A_1) = f(A_0) \cap f(A_1)$.

Problem 3, p83:

Let X be a set. Consider the collection \mathcal{T}_C , the collection of all subsets U of X such that $X \setminus U$ is countable or all of X .

Then $\emptyset \in \mathcal{T}_C$, as $X \setminus \emptyset = X$, which is all of X . Also, $X \in \mathcal{T}_C$, as $X \setminus X = \emptyset$, which is countable.

That is, $\emptyset \in \mathcal{T}_C$ and $X \in \mathcal{T}_C$.

Next, let $\mathcal{C} \subset \mathcal{T}_C$ be nonempty. If the only element of \mathcal{C} is \emptyset , then $X \setminus \bigcup_{U \in \mathcal{C}} U = X$, so that $\bigcup_{U \in \mathcal{C}} U \in \mathcal{T}_C$. Else, pick a nonempty element of \mathcal{C} ; call it U_0 . Then $\bigcup_{U \in \mathcal{C}} U \supset U_0$, so $X \setminus U_0 \supset X \setminus \bigcup_{U \in \mathcal{C}} U$. Because $X \setminus U_0$ is finite (as $U_0 \in \mathcal{T}_C$ so that $X \setminus U_0$ is either finite or all of X , and $X \setminus U_0$ is not all of X , as U_0 is nonempty), this means that $X \setminus \bigcup_{U \in \mathcal{C}} U$ is finite. That is, $X \setminus \bigcup_{U \in \mathcal{C}} U \in \mathcal{T}_C$.

That is, arbitrary unions of elements in \mathcal{T}_C are contained in \mathcal{T}_C .

Last, let $\{U_0, U_1, \dots, U_n\} \subset \mathcal{T}_C$. If $\bigcap_{i=0}^n U_i = \emptyset$, then $X \setminus \bigcap_{i=0}^n U_i = X \setminus \emptyset = X$. Else, recall that $X \setminus \bigcap_{i=0}^n U_i = \bigcup_{i=0}^n X \setminus U_i$, which is a finite union of finite sets (None of $X \setminus U_i$ are all of X , else $\bigcap_{i=0}^n U_i = \emptyset$). So, $X \setminus \bigcap_{i=0}^n U_i$ is finite, so that $\bigcap_{i=0}^n U_i \in \mathcal{T}_C$.

That is, finite intersections of elements in \mathcal{T}_C are contained in \mathcal{T}_C .

So \mathcal{T}_C is a topology.

Problem 5, p83:

Let \mathcal{A} be a basis for a topology, call it \mathcal{T} , on X .

Consider $\mathcal{T}' = \bigcap_{\mathcal{T}'': \mathcal{T}'' \supset \mathcal{A}} \mathcal{T}''$; the intersection of all topologies containing \mathcal{A} .

First, $\mathcal{T}' \subset \mathcal{T}$: Let $U \in \mathcal{T}'$. Then U is in the intersection of all topologies

containing \mathcal{A} . So U is in any topology containing \mathcal{A} . Now, \mathcal{T} is a topology containing \mathcal{A} . So $\mathcal{A} \in \mathcal{T}$.

Next, $\mathcal{T} \subset \mathcal{T}'$: Let $U \in \mathcal{T}$, let \mathcal{T}'' be a topology containing \mathcal{A} . Then U is a union of elements in \mathcal{A} . So U is a union of elements in \mathcal{T}'' . So $U \in \mathcal{T}''$. So U is in any topology containing \mathcal{A} . So U is in the intersection of all topologies containing \mathcal{A} . So $U \in \mathcal{T}'$.

So, $\mathcal{T} = \mathcal{T}'$. So, the topology generated by \mathcal{A} is the intersection of all topologies containing \mathcal{A} .

Next, let \mathcal{A} be a subbasis for a topology, call it \mathcal{T} , on X .

Consider $\mathcal{T}' = \bigcap_{\mathcal{T}'' : \mathcal{T}'' \supset \mathcal{A}} \mathcal{T}''$; the intersection of all topologies containing \mathcal{A} .

First, $\mathcal{T}' \subset \mathcal{T}$: Let $U \in \mathcal{T}'$. Then U is in the intersection of all topologies containing \mathcal{A} . So U is in any topology containing \mathcal{A} . Now, \mathcal{T} is a topology containing \mathcal{A} . So $\mathcal{A} \in \mathcal{T}$.

Next, $\mathcal{T} \subset \mathcal{T}'$: Let $U \in \mathcal{T}$, let \mathcal{T}'' be a topology containing \mathcal{A} . Then U is a union of finite intersections of elements of \mathcal{A} . So U is a union of finite intersections of elements of \mathcal{T}'' . So $U \in \mathcal{T}''$. So U is in any topology containing \mathcal{A} . So U is in the intersection of all topologies containing \mathcal{A} . So $U \in \mathcal{T}'$.

So, $\mathcal{T} = \mathcal{T}'$. So, the topology generated by \mathcal{A} is the intersection of all topologies containing \mathcal{A} .

Problem 8b, p83:

Consider $\mathcal{C} = \{[a, b) : a < b, a, b \in \mathbb{Q}\}$.

For each $x \in \mathbb{R}$, $[\lfloor x \rfloor, \lfloor x \rfloor + 1) \in \mathcal{C}$ contains x .

That is, for each $x \in \mathbb{R}$, there is a $B \in \mathcal{C}$ containing x .

Now, let $B_1, B_2 \in \mathcal{C}$; say that $B_1 = [a, b)$ and $B_2 = [c, d)$ with $a, b, c, d \in \mathbb{Q}$. Let $x \in B_1 \cap B_2$. Define $e = \max(a, c)$ and $f = \min(b, d)$. Then it is clear that $[e, f) = B_1 \cap B_2$ (so that $[e, f) \subset B_1 \cap B_2$), and also $[e, f)$ contains x .

Problem 1, p91:

Let Y be a subspace of X , and $A \subset Y$. Let \mathcal{T} be the topology A inherits as a subspace of Y , and \mathcal{T}' be the topology A inherits as a subspace of X .

First, $\mathcal{T} \subset \mathcal{T}'$: Let $U \in \mathcal{T}$. Then $U = A \cap U'$ for some open U' in X . Now, $A = A \cap Y$, so $U = A \cap Y \cap U' = A \cap (Y \cap U')$. That is, $U = A \cap U''$ for some U'' open in Y ; $U \in \mathcal{T}'$.

Next, $\mathcal{T}' \subset \mathcal{T}$: Let $U \in \mathcal{T}'$. Then $U = A \cap U'$ for some open U' in Y . Now, $U' = Y \cap U''$ for some open U'' in X , so $U = A \cap Y \cap U'' = A \cap U''$. That is, $U = A \cap U''$ for some U'' open in X ; $U \in \mathcal{T}$.

So, $\mathcal{T} = \mathcal{T}'$.

Problem 4, p91:

Problem 6, p91:

Consider $\mathcal{B} = \{(a, b) \times (c, d) : a < b, c < d, \{a, b, c, d\} \subset \mathbb{Q}\}$.

For each $(x, y) \in \mathbb{R}^2$, the element $(\lfloor x - 1 \rfloor, \lceil x + 1 \rceil) \times (\lfloor y - 1 \rfloor, \lceil y + 1 \rceil) \in \mathcal{B}$ contains (x, y) . (Where $\lfloor x \rfloor$ and $\lceil x \rceil$ are the standard floor and ceiling functions.)

That is, for each $(x, y) \in \mathbb{R}^2$, at least one element in \mathcal{B} contains (x, y) .

Next, let $B_1, B_2 \in \mathcal{B}$, with $B_1 = (a, b) \times (c, d)$ and $B_2 = (e, f) \times (g, h)$. Define $i = \max(a, e)$, $j = \min(b, f)$, $k = \max(c, g)$, and $l = \min(d, h)$. Then it is clear that $B_1 \cap B_2 = (i, j) \times (k, l) \in \mathcal{B}$. Let $x \in B_1 \cap B_2$. Then $B_3 = (i, j) \times (k, l)$ contains x and $B_3 \subset B_1 \cap B_2$.

That is, if $B_1, B_2 \in \mathcal{B}$ and $x \in B_1 \cap B_2$, there is a $B_3 \in \mathcal{B}$ with $x \in B_3$ and $B_3 \subset B_1 \cap B_2$.

Thus, \mathcal{B} is a basis.

Problem 9, p91: