

MATH440 PA Thursday Week 2

Josh Derrow

August 28, 2025

- 1 **Determine the equilibrium temperature distribution for a one-dimensional rod with constant thermal properties with the following sources and boundary conditions:**

1.1 (1.4.2b) $Q = 0, u(0) = T, u(L) = 0$

Heat Equation: $\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2} + Q$

Constant thermal properties ($\frac{\partial u}{\partial t} = 0$) and no sources/sinks ($Q = 0$)

$$\Rightarrow k \frac{\partial^2 u}{\partial x^2} = 0$$

$$\Rightarrow \frac{\partial^2 u}{\partial x^2} = 0$$

$$\Rightarrow \frac{\partial u}{\partial x} = c_1$$

$$\Rightarrow u(x) = c_1 x + c_2$$

$$u(0) = c_1(0) + c_2 = T$$

$$\Rightarrow c_2 = T$$

$$u(L) = c_1(L) + c_2 = 0$$

$$\Rightarrow c_1(L) + T = 0$$


$$\Rightarrow c_1 L = -T$$

$$\Rightarrow c_1 = -\frac{T}{L}$$

$$u(x) = c_1 x + c_2 = -\frac{T}{L}x + T$$

Heat distribution: $u(x) = -\frac{T}{L}x + T$

Below is a diagram of this distribution:



1.2 (1.4.2c) $Q = 0, \frac{\partial u}{\partial x}(0) = 0, u(L) = T$

Heat Equation: $\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2} + Q$

Constant thermal properties ($\frac{\partial u}{\partial t} = 0$) and no sources/sinks ($Q = 0$)

$$\Rightarrow k \frac{\partial^2 u}{\partial x^2} = 0$$

$$\Rightarrow \frac{\partial^2 u}{\partial x^2} = 0$$

$$\frac{\partial u}{\partial x}(0) = c_1 = 0$$

$$\Rightarrow c_1 = 0$$

$$u(x) = c_1 x + c_2$$

$$\Rightarrow u(x) = c_2$$

$$u(L) = T = c_2$$

$$\Rightarrow c_2 = T$$

$$u(x) = c_2 = T$$

Heat distribution: $u(x) = T$

Below is a diagram of this distribution:

