AGRISENSE AI


Intelligent Agriculture Decision Support Platform

DIAGNOSTIC REPORT
Report ID: 53760AE0
Date: 19 Aug 2026, 03:10 PM


Farmer: Ramesh Gowda | **Location:** Mysuru, Karnataka | **Crop Analyzed:** Rice

DIAGNOSED CONDITION:	**Rice Blast *(Magnaporthe oryzae)***
Confidence Level:	**94.5%** (EfficientNetB3 Saliency)
Severity Rating:	**MODERATE** (Estimated affected canopy: **22.0%**)
Pathogen Class:	Fungal

Original Leaf Specimen



Explainable Grad-CAM Heatmap Attention



Clinical Symptoms & Biological Cause

Symptoms: Spindle-shaped or diamond-shaped lesions with grey or whitish centers and brown/reddish margins on leaves, nodes, and panicles.

Etiology/Cause: Airborne spores exacerbated by high nitrogen fertilizer application and cloudy, humid weather.

Recommended Treatment Protocol

Treatment Type	Prescription & Application Details
Organic / Bio-Control	Foliar spray of Pseudomonas fluorescens @ 5g/L or Cow urine concoction (10%) with neem oil.
Chemical / Fungicide	Tricyclazole 75% WP @ 0.6g/L (Beam) or Isoprothiolane 40% EC @ 1.5ml/L or Kasugamycin 3% SL @ 2ml/L. **Dosage:** Tricyclazole 75% WP @ 0.6g per liter water (120g/acre). **Frequency:** Spray at tillering and panicle emergence stages.
Preventive Measures	Avoid excessive split applications of nitrogen; maintain proper water depth in paddy fields; use blast-resistant cultivars.

Note: This clinical report was generated by AgriSense AI Diagnostic Engine using deep learning visual analysis. For epidemic outbreaks or severe crop stress, please consult your nearest Krishi Vigyan Kendra (KVK) or Regional Agriculture Extension Officer.