# AGRISENSE AI


**Intelligent Agriculture Decision Support Platform**

**DIAGNOSTIC REPORT**
**Report ID:** 5A3571F7
**Date:** 18 Aug 2026, 04:05 PM


**Farmer:** Ramesh Gowda | **Location:** Mysuru, Karnataka | **Crop Analyzed:** Potato

| | |
|---|---|
| **DIAGNOSED CONDITION:** | **Late Blight *(Phytophthora infestans)*** |
| **Confidence Level:** | **93.5%** (EfficientNetB3 Saliency) |
| **Severity Rating:** | **SEVERE** (Estimated affected canopy: **28.7%**) |
| **Pathogen Class:** | Oomycete / Water Mold |

**Original Leaf Specimen**



**Explainable Grad-CAM Heatmap Attention**



## Clinical Symptoms & Biological Cause

**Symptoms:** Dark brown water-soaked blotches starting from leaf edges and tips, rapid blight of canopy with white sporulation on undersides.

**Etiology/Cause:** Rapidly virulent water mold thriving during humid, foggy, and cool weather.

## Recommended Treatment Protocol

| Treatment Type | Prescription & Application Details |
|---|---|
| **Organic / Bio-Control** | Bordeaux mixture 1% foliar spray, immediate removal of blighted foliage before rain. |
| **Chemical / Fungicide** | Dimethomorph 50% WP @ 1g/L or Cymoxanil 8% + Mancozeb 64% WP @ 2g/L or Fenamidone + Mancozeb @ 2.5g/L.<br>**Dosage:** Dimethomorph 1g + Mancozeb 2g per liter water.<br>**Frequency:** Every 5 days during critical danger window. |
| **Preventive Measures** | Ensure deep hilling to protect tubers, monitor local late-blight weather warnings, harvest in dry conditions. |

*Note: This clinical report was generated by AgriSense AI Diagnostic Engine using deep learning visual analysis. For epidemic outbreaks or severe crop stress, please consult your nearest Krishi Vigyan Kendra (KVK) or Regional Agriculture Extension Officer.*