

Attach this page to the front of your submission. Remember to sign the declaration.

I hereby state that the work contained in this assignment has not previously been submitted for assessment, either in whole or in part, by either myself or any other student at either The University of Queensland or at any other tertiary institution except where explicitly acknowledged. To the best of my knowledge and belief, the assignment contains no material that has been previously published or written by another person except where due reference is made. I make this Statement in full knowledge of an understanding that, should it be found to be false, I will be subject to disciplinary action under Student Integrity and Misconduct Policy 3.60.04 of the University of Queensland. The University of Queensland 's policy on plagiarism can be found at

<http://ppl.app.uq.edu.au/content/3.60.04-student-integrity-and-misconduct>  
(Reference 3.60.04).

Name ..

Samuel Allpass

.....Student ID .

48030504

Signed .

Samuel Allpass

.....Date

14/5/23

..

Question	Mark
1	
2	
3	
4	
Total	

$$Q1) \sum_{n=1}^{\infty} \frac{(n!)^2}{(2n)!} (x-1)^n$$

a)

Radius of convergence,  $L = \lim_{n \rightarrow \infty} \left( \frac{n+1}{n} \right)$

$$\lim_{n \rightarrow \infty} \left| \frac{\frac{(n+1)!^2}{(2n+2)!} (x-1)^{n+1}}{\frac{(n!)^2}{(2n)!} (x-1)^n} \right| = \lim_{n \rightarrow \infty} \left| \frac{\frac{(n+1)!^2}{(2n+2)!} (x-1)(x-1)^n}{\frac{(n!)^2}{(2n)!} (x-1)^n} \right|$$

$$= |x-1| \lim_{n \rightarrow \infty} \left| \frac{(n+1)^2 (2n)!}{(n!)^2 (2n+2)!} \right| = |x-1| \lim_{n \rightarrow \infty} \left| \frac{(n+1)!^2 (2n)!}{(n!)^2 (2n+2)!} \right|$$

$$= |x-1| \lim_{n \rightarrow \infty} \left| \frac{\left( \frac{n! \times (n+1)}{n!} \right)^2 (2n)!}{(2n)! \times (2n+1) \times (2n+2)} \right|$$

$$= |x-1| \lim_{n \rightarrow \infty} \left| \frac{(n+1)^2 \times 1}{(2n+1) \times (2n+2)} \right| = |x-1| \lim_{n \rightarrow \infty} \left| \frac{n^2 + 2n + 1}{4n^2 + 6n + 2} \right|$$

L'Hopital's

$$= |x-1| \lim_{n \rightarrow \infty} \left| \frac{2n+2}{8n+6} \right|$$

$$= |x-1| \lim_{n \rightarrow \infty} \left| \frac{2}{8} \right|$$

$$= |x-1| \times \frac{1}{4}$$

$$\therefore |x-1| \frac{1}{4} < 1$$

$$\therefore -1 < |x-1| \frac{1}{4} < 1$$

$$-4 < |x-1| < 4$$

$\therefore$  The radius of convergence is 4

b)

$$\therefore \text{Centre} = +1 \text{ as } 0 = x-1$$

$$x=1$$

Interval: as the centre is at  $x=1$  and the radius is 4,  $(-3, 5)$  is the interval of convergence



Q2)

$$a_n = \int_0^{2\pi n} f(x) \sin x \, dx$$

$$b_n = \int_0^{n+2\pi n} f(x) \sin(x) \, dx$$

a)  $f(x) = 1$

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \int_0^{2\pi n} \sin(x) \, dx$$

$$= \lim_{n \rightarrow \infty} [-\cos x]_0^{2\pi n}$$

$$= \lim_{n \rightarrow \infty} (-\cos(2\pi n) + 1)$$

$\Rightarrow$  undefined

as  $\lim_{n \rightarrow \infty} \cos(n)$   
is undefined

$$\lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} \int_0^{n+2\pi n} \sin(x) \, dx$$

$$= \lim_{n \rightarrow \infty} [-\cos(x)]_0^{n+2\pi n}$$

$$= \lim_{n \rightarrow \infty} (-\cos(n+2\pi n) + 1)$$

$\Rightarrow$  undefined

b)  $\int_0^{\infty} \sin x \, dx$  will converge if it has a limit:

$$\lim_{x \rightarrow \infty} \int_0^{\infty} \sin x \, dx = \lim_{x \rightarrow \infty} [-\cos x]_0^{\infty}$$

$$= \lim_{x \rightarrow \infty} (-\cos \infty - (-1))$$

this is undefined as  $\cos \infty$  is  
undefined

$\therefore$  This integral does not converge

c)  $f(x) = e^{-x}$

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \int_0^{2\pi n} e^{-x} \sin x \, dx$$

LIATE

let  $u = \sin x \quad v' = e^{-x}$   
 $u' = \cos x \quad v = -e^{-x}$

$$\int e^{-x} \sin x \, dx = -e^{-x} \sin x + \int +e^{-x} \cos x \, dx$$

let  $u = \cos x \quad v' = +e^{-x}$   
 $u' = -\sin x \quad v = -e^{-x}$

$$= -e^{-x} \sin x - e^{-x} \cos x - \int e^{-x} \sin x \, dx$$

$$2 \int e^{-x} \sin x \, dx = -e^{-x} (\sin x + \cos x)$$

$$\int e^{-x} \sin x \, dx = \frac{-e^{-x} (\sin x + \cos x)}{2}$$

$$\begin{aligned} \therefore \lim_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} \left[ \frac{-e^{-x} (\sin(x) + \cos(x))}{2} \right]_0^{2\pi n} \\ &= \lim_{n \rightarrow \infty} \left( \frac{-e^{-2\pi n} (\sin(2\pi n) + \cos(2\pi n))}{2} - - \frac{1(0+1)}{2} \right) \\ &= \frac{1}{2} \end{aligned}$$

$$\lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} \int_0^{2\pi n + \pi} e^{-x} \sin x \, dx$$

$$\begin{aligned} &= \lim_{n \rightarrow \infty} \left[ \frac{-e^{-x} (\sin x + \cos x)}{2} \right]_0^{\pi + 2\pi n} \\ &= \lim_{n \rightarrow \infty} \left( \frac{-e^{-(\pi + 2\pi n)} (\sin(\pi + 2\pi n) + \cos(\pi + 2\pi n))}{2} - - \frac{1(0+1)}{2} \right) \\ &= \frac{1}{2} \end{aligned}$$



$$d) f(x) = e^{-x} \sin^3 x$$

$\therefore$  for  $\int_0^{\infty} f(x) \sin x \, dx$  to be convergent

$\lim_{x \rightarrow \infty} \int_0^{\infty} e^{-x} \sin^4 x \, dx$  must have a solution.

Using Identities

$$\lim_{x \rightarrow \infty} \int_0^{\infty} e^{-x} \sin^4 x \, dx = \lim_{x \rightarrow \infty} \int_0^{\infty} e^{-x} \left( \frac{1}{2}(1 - \cos(2x)) \right)^2 dx$$

$$= \lim_{x \rightarrow \infty} \int_0^{\infty} e^{-x} \left( \frac{1}{2} - \frac{1}{2} \cos(2x) \right)^2 dx$$

$$= \lim_{x \rightarrow \infty} \int_0^{\infty} e^{-x} \left( \frac{1}{4} - \frac{1}{2} \cos(2x) + \frac{1}{4} \cos^2(2x) \right) dx$$

$$= \lim_{x \rightarrow \infty} \int_0^{\infty} \frac{e^{-x}}{4} - \frac{e^{-x}}{2} \cos(2x) + \frac{e^{-x}}{4} \left( \frac{1}{2} (\cos(4x) + 1) \right) dx$$

$$= \lim_{x \rightarrow \infty} \int_0^{\infty} \frac{e^{-x}}{4} - \frac{e^{-x}}{2} \cos(2x) + \frac{e^{-x}}{8} \cos(4x) + \frac{e^{-x}}{8} dx$$

$$= \lim_{x \rightarrow \infty} \int_0^{\infty} \frac{e^{-x} \cos(4x)}{8} - \frac{e^{-x} \cos(2x)}{2} + \frac{3e^{-x}}{8} dx$$

$$\text{let } e^{-x} = v' \quad u = \cos(4x) \\ v = -e^{-x} \quad u' = -4\sin(4x)$$

$$\int \frac{e^{-x} \cos(4x)}{8} dx = \frac{-e^{-x} \cos(4x)}{8} - \int -e^{-x} \times -4\sin(4x)$$

$$\text{let } e^{-x} = v' \quad u = \cos(2x) \\ v = -e^{-x} \quad u' = -2\sin(2x)$$

$$\int \frac{e^{-x} \cos(2x)}{2} dx = \frac{-e^{-x} \cos(2x)}{2} - \int -e^{-x} \times -2\sin(2x)$$

$$\text{let } u = -4\sin(4x) \quad v' = e^{-x} \\ u' = -16\cos(4x) \quad v = -e^{-x}$$

$$\int e^{-x} \cos(4x) dx = \frac{-e^{-x} \cos(4x)}{16} + \frac{e^{-x} \sin(4x)}{4} - \int -16e^{-x} \cos(4x)$$

$$\text{let } u = -2\sin(2x) \quad v' = e^{-x} \\ u' = -4\cos(2x) \quad v = -e^{-x}$$

$$\int e^{-x} \cos(2x) dx = \frac{-e^{-x} \cos(2x)}{4} + \frac{2\sin(2x)e^{-x}}{2} - \int e^{-x} \times 4\cos(2x) dx$$

$$\int e^{-x} \cos(4x) dx = \frac{e^{-x}(4\sin(4x) + \cos(4x))}{17} + C$$

$$= \frac{e^{-x}(2\sin(2x) - \cos(2x))}{5} + C$$



$$\therefore \lim_{x \rightarrow \infty} \int_0^{\infty} e^{-x} \sin^4(x) dx$$

$$= \lim_{x \rightarrow \infty} \left[ \frac{1}{8} \left( \frac{e^{-x}(4\sin(4x) - \cos(4x))}{17} \right) - \frac{1}{2} \left( \frac{e^{-x}(2\sin(2x) - \cos(2x))}{5} \right) - \frac{3e^{-x}}{8} \right]_0^{\infty}$$

$$= \lim_{x \rightarrow \infty} \left( \frac{e^{-\infty}(4\sin(\infty) - \cos(\infty))}{136} - \frac{e^{-\infty}(2\sin(\infty) - \cos(\infty))}{10} - \frac{3e^{-\infty}}{8} \right)$$

$$- \left( \frac{e^{-0}(4\sin(0) - \cos(0))}{136} - \frac{e^{-0}(2\sin(0) - \cos(0))}{10} - \frac{3e^0}{8} \right)$$

$$= \frac{24}{85} \approx 0.28 = \lim_{x \rightarrow \infty} \int_0^{\infty} e^{-x} \sin^4(x) dx$$

$\therefore$  as the limit has a solution, the improper integral  $\int_0^{\infty} f(x) \sin^4(x) dx$  converges.



e) Example of nonconstant  $f(x)$  for which  $\int f(x) \sin(x) dx$  diverges implies that:

$$\lim_{x \rightarrow \infty} \int_0^x f(x) \sin(x) dx \text{ has no solution}$$

Example.  $f(x) = x$

$$\lim_{x \rightarrow \infty} \int_0^x x \sin(x) dx$$

$$\text{let } x = u \quad u' = \sin x$$


$$u' = 1 \quad u = -\cos x$$

$$= \lim_{x \rightarrow \infty} \left[ -x \cos x - \int -\cos x dx \right]_0^{\infty}$$

$$= \lim_{x \rightarrow \infty} \left[ -x \cos x + \sin x \right]$$

as  $\cos(\infty)$  has no solution, it cannot have a limit and must therefore be divergent.

Q3)



As parallelogram, B must be at (4, 1)

$$\vec{OB} = \vec{OA} + \vec{AB} = (4, 1) = \begin{pmatrix} 4 \\ 1 \end{pmatrix}$$

$y = -\frac{1}{2}x$  For the sake of this question can be vector  $f$  where:  $\vec{f} = \begin{pmatrix} 4 \\ -4 \end{pmatrix}$

$$\text{Proj}_{\vec{f}} \vec{OB} = \frac{\vec{OB} \cdot \vec{f}}{|\vec{f}|} \times \frac{\vec{f}}{|\vec{f}|} = \frac{(4 \times 4) + (1 \times -4)}{\sqrt{4^2 + 4^2}} \times \frac{\begin{pmatrix} 4 \\ -4 \end{pmatrix}}{\sqrt{4^2 + 4^2}}$$

$$\text{Proj}_{\vec{f}} \vec{OB} = \begin{pmatrix} 4 \\ -4 \end{pmatrix} \times \frac{3}{8} = \begin{pmatrix} \frac{3}{2} \\ -\frac{3}{2} \end{pmatrix}$$


$\therefore$  The projection is line from the origin to  $(\frac{3}{2}, -\frac{3}{2})$  following  $y = -x$ .



Q4)

$$A = \begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix}$$

a) Considering the columns, the two vectors are using standard basis of  $\mathbb{R}^2$ .  
 $\begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix} \times \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$  and  $\begin{pmatrix} 3 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix} \times \begin{pmatrix} 0 \\ 1 \end{pmatrix}$   
 these are displayed as:




b) Rotating by some angle anticlockwise. The formula is:

$$B = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

as  $\theta = 45^\circ$  or  $\frac{\pi}{4}$  radians.

$$B = \begin{pmatrix} \cos \frac{\pi}{4} & -\sin \frac{\pi}{4} \\ \sin \frac{\pi}{4} & \cos \frac{\pi}{4} \end{pmatrix}$$



$$B = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} = \begin{pmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \end{pmatrix}$$

$$c) AB = \begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix} \begin{pmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \end{pmatrix} = \begin{pmatrix} \frac{\sqrt{2}}{2} + 3\frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} + 3\frac{\sqrt{2}}{2} \\ 3\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2} & -3\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2} \end{pmatrix} = \begin{pmatrix} 2\sqrt{2} & \sqrt{2} \\ 2\sqrt{2} & -\sqrt{2} \end{pmatrix}$$

$$BA = \begin{pmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \end{pmatrix} \begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix} = \begin{pmatrix} \frac{\sqrt{2}}{2} - 3\frac{\sqrt{2}}{2} & 3\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} + 3\frac{\sqrt{2}}{2} & 3\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2} \end{pmatrix} = \begin{pmatrix} -\sqrt{2} & \sqrt{2} \\ 2\sqrt{2} & 2\sqrt{2} \end{pmatrix}$$

No they are not the same.

d)  $\begin{pmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \end{pmatrix} C = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \rightarrow$  This is the case when  $B$  is transformed to  $\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$   
 $\therefore$  Clockwise by  $45^\circ$  ( $-\frac{\pi}{4}$ )  $\leftarrow$  when  $\theta = 0$ .

$B \times C$

$$\therefore C = \begin{pmatrix} \cos(-\frac{\pi}{4}) & -\sin(-\frac{\pi}{4}) \\ \sin(-\frac{\pi}{4}) & \cos(-\frac{\pi}{4}) \end{pmatrix} \text{ Proof: } \begin{pmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \end{pmatrix} \begin{pmatrix} \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \\ -\frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$