

# BESOV-ISH SPACES THROUGH ATOMIC DECOMPOSITION: THE LEAN 4 FORMALIZATION

DANIEL SMANIA

ABSTRACT. This document describes, without proofs, exactly those results of the paper *Besov-ish spaces through atomic decomposition* (Analysis & PDE 15 (2022), no. 1) that have been formally verified in the Lean 4 repository `BesovSpacesGoodGrid`, together with a few results obtained in the formalization that go beyond the paper. It was produced from a prerelease version of the published paper (`preprint-source/besovish-2020-07-18.tex` in the repository), but all references to numbered statements of the paper follow the *published* version. Unlike the paper, the document is organized to mirror the repository: Part I covers the weak-grid library (`WeakGrid/`) and Part II the good-grid library (`GoodGrid/`), with sections ordered by file dependency; each section opens with a box naming the Lean file it documents and the project files that file imports. Statements follow the notation of the paper; each result is followed, in the running text, by a *Formalization* paragraph naming the corresponding Lean declarations, describing what each of them proves, and recording every deviation from the published statements, together with a snapshot of the Lean code. The document also explains the purpose and architecture of the formalization and ends with a guide to the repository files. The repository compiles with no `sorry`, and the main theorems depend only on the standard axioms `propext`, `Classical.choice`, `Quot.sound`.

## CONTENTS

|                                                             |    |
|-------------------------------------------------------------|----|
| 1. Introduction                                             | 2  |
| 1.1. Purpose of the formalization                           | 2  |
| 1.2. How this document was produced                         | 4  |
| 2. Notation                                                 | 5  |
| 3. Measure spaces and weak grids                            | 6  |
| 4. Atoms                                                    | 8  |
| 5. Besov-ish spaces                                         | 10 |
| 6. Limits of representations, compactness, and completeness | 13 |

---

2020 *Mathematics Subject Classification.* 43A99, 43A85, 30H25, 42B15, 42B35, 42C15, 42C40, 28C99.

*Key words and phrases.* atomic decomposition, Besov space, harmonic analysis, wavelets, multipliers, Haar wavelets, atoms, formal verification, Lean.

|                                                                                 |    |
|---------------------------------------------------------------------------------|----|
| 7. Scales of spaces                                                             | 16 |
| 8. Induced spaces                                                               | 17 |
| 9. Transmutation of atoms                                                       | 18 |
| 10. Further weak-grid infrastructure                                            | 22 |
| 11. Good grids                                                                  | 23 |
| 12. Souza’s atoms                                                               | 25 |
| 13. Besov spaces in a measure space with a good grid                            | 26 |
| 14. Besov’s atoms and the Souza–Besov comparison                                | 27 |
| 15. Unbalanced Haar wavelets                                                    | 29 |
| 16. Alternative norms: standard, Haar, and mean oscillation                     | 30 |
| 17. Positive cone                                                               | 34 |
| 18. Dirac’s approximations                                                      | 38 |
| 19. Pointwise multipliers acting on $\mathcal{B}_{p,q}^s$                       | 39 |
| 19.1. The selfs classes and their seminorms                                     | 40 |
| 19.2. Restriction to a cell                                                     | 41 |
| 19.3. The multipliers of $\mathcal{B}_{1,1}^s$                                  | 42 |
| 19.4. <i>selfs</i> multipliers are bounded                                      | 43 |
| 19.5. Non-Archimedean behaviour in $\mathcal{B}_{p,\tilde{q},selfs}^\beta$      | 44 |
| 19.6. The tail selfs classes consist of multipliers                             | 51 |
| 19.7. Strongly regular domains and Pointwise Multipliers I                      | 53 |
| 19.8. $\mathcal{B}_{p,\infty}^{1/p} \cap L^\infty$ and Pointwise Multipliers II | 56 |
| 20. Regular domains                                                             | 59 |
| 21. The bounded Souza–Besov space as a quasi-algebra                            | 62 |
| 22. A remarkable description of $\mathcal{B}_{1,1}^s$                           | 63 |
| 23. Left compositions                                                           | 65 |
| 24. Guide to the repository files                                               | 66 |
| 24.1. Weak-grid layer (Part I)                                                  | 66 |
| 24.2. Good-grid layer (Part II)                                                 | 67 |
| 24.3. Applications and multiplier layer (Part II)                               | 68 |
| 24.4. External dependencies                                                     | 69 |
| 24.5. Repository status                                                         | 69 |
| References                                                                      | 69 |

## 1. INTRODUCTION

**1.1. Purpose of the formalization.** The repository `BesovSpacesGoodGrid` formalizes, in Lean 4 over `mathlib`, a substantial part of the atomic decomposition theory of [1]. The goals are twofold. First, to obtain machine-checked proofs of the main results of the paper — the Banach structure, compactness and completeness of Besov-ish spaces, the transmutation-of-atoms machinery, the equivalence of the atomic, Haar,

and mean-oscillation norms, and the theory of pointwise multipliers up to the non-Archimedean property and its positive-cone refinements. Second, and just as important, to produce a *reusable* Lean library: the basic theory is stated and proved for abstract *weak grids* and abstract atom families, and the good-grid / Souza-atom spaces of the paper are then obtained as a specialization. The weak-grid layer (grids, atoms, Besov-ish spaces, scales, completeness, transmutation) is independent of the good-grid geometry and is intended to be usable in later developments.

This document is organized to mirror the repository rather than the paper. Part I documents the weak-grid library (`BesovSpacesGoodGrid/WeakGrid/`) and Part II the good-grid library (`BesovSpacesGoodGrid/GoodGrid/`). The early sections follow the import order of the Lean files; the later application sections follow the mathematical narrative of the paper more closely, with each section's file box recording the exact project imports. Each section opens with a box naming the file it documents and the project files it imports. The mathematical pipeline, in this order, is:

- (1) define quantitative good grids and general weak grids;
- (2) package weak-grid cells and local spaces, and define atom families with support, convexity, phase invariance, and atom-size estimates;
- (3) build levelwise atomic blocks and infinite  $L^p$  representations, define the Besov-ish space and the coefficient-cost gauge `Norm_Costpq`;
- (4) prove the  $L^t$  embedding, scale inclusions, representation-limit theorems, compactness statements, and completeness for the cost norm;
- (5) prove the transmutation claims moving atomic representations from one weak grid to another, and specialize them to compare Souza's atoms, Besov's atoms, and sandwiched atom families on good grids;
- (6) introduce the positive cone, the Haar and standard representation gauges, and the mean-oscillation norm, with finite-constant comparison theorems;
- (7) prove the Dirac-approximation identities: the partial sums of the standard representation evaluate as averages of the function against the grid Dirac kernels, with the resulting  $L^\infty$  bounds on ancestor coefficient sums (paper Proposition 17.1);
- (8) develop the pointwise-multiplier layer: the `selfs` classes, boundedness of `selfs` multipliers, the  $\mathcal{B}_{1,1}^s$  multiplier characterization, the non-Archimedean multiplier estimates, including their positive-cone versions, and their consequences — the inclusion of the tail `selfs` classes in the multiplier space, strongly regular domains and Pointwise Multipliers I, and Pointwise Multipliers II with its positive-cone version;

- (9) prove the quasi-algebra property of the bounded Souza–Besov space:  $\mathcal{B}_{p,q}^s \cap L^\infty$  is closed under pointwise multiplication, with a bilinear bound (Pointwise Multipliers III).
- (10) formalize the regular-domain applications: regular families, the indicator bound `estG`, localized restriction representations `pdd/hiip1`, and the alternative description of  $\mathcal{B}_{1,1}^s$  by normalized indicators of regular domains.

A guide to the repository files is given in Section 24.

**1.2. How this document was produced.** This is a mathematical description of the current state (June 17, 2026) of the Lean 4 formalization of [1]. It was produced from a *prerelease version of the published paper* (the LaTeX source `preprint-source/besovish-2020-07-18.tex`, included in the repository) by the following rules.

- Results that are not present in the repository were deleted. These are: the elementary estimates of paper Section 4 (Hölder-like and convolution tricks, inlined in the Lean proofs without standalone statements); the Hölder and bounded variation atoms (paper Sections 11B–11C) and the corresponding characterizations (paper Propositions 16.2 and 16.3); and the full paper Section 21 left-composition theorem (only the mean-oscillation gauge estimate is currently incorporated). Dirac’s approximations (paper Section 17), the whole multiplier Section 18 — Corollary 18.6, strongly regular domains and Pointwise Multipliers I (18.7–18.9), Pointwise Multipliers II (18.10) together with its positive-cone version, and the regular-domains subsection `cf` (regular families, the indicator bound `estG`, and the localized restriction representations `pdd/hiip1`) — the quasi-algebra result of Section 19 (Proposition 19.1, Pointwise Multipliers III), and the regular-domain description of  $\mathcal{B}_{1,1}^s$  from Section 20 are formalized and documented in Sections 18, 19, 21, 20, and 22.
- Statements were updated to match the precision of the formalization; every change with mathematical content is recorded in the *Formalization* paragraph that follows the statement.
- All references of the form “paper Proposition  $x.y$ ” use the numbering of the *published* version, Analysis & PDE **15** (2022), no. 1 (which the reader should consult alongside this document); the prerelease source in the repository has the same section structure but its statement numbers may differ.
- The material was reordered to follow the repository and the application pipeline (see above), so the section numbers of this document do *not* match the section numbers of the paper.

- No proofs are included. Each statement is followed by a *Formalization* paragraph giving the corresponding Lean declarations and files, with a short description of what each declaration proves, and then by a snapshot of the Lean code that formalizes it (the formal statement as it appears in the repository, with the proof omitted; an ellipsis  $\dots$  marks the omitted proof or, in a few long definitions, omitted structural fields).

A global change must be singled out: *throughout this document*  $p \in [1, \infty)$ ,  $q \in [1, \infty]$  and  $u \in [1, \infty]$ . The paper develops Part I for the quasi-Banach range  $p, q \in (0, \infty]$ ; the formalization restricts to the Banach range, so  $\rho = \min\{1, p, q\} = 1$  everywhere and all the  $\rho$ -norms of the paper are genuine norms. In Lean the standing hypotheses are  $1 \leq p, p \neq \infty$ ,  $\text{Fact } (1 \leq q), 1 \leq u, 0 < s$ .

## 2. NOTATION

TABLE 1. Most important notation/symbols/constants

| Symbol                                                                 | Description                                                           |
|------------------------------------------------------------------------|-----------------------------------------------------------------------|
| $I$                                                                    | phase space                                                           |
| $m$                                                                    | finite measure in the phase space $I$                                 |
| $a_P, b_P$                                                             | an atom supported on $P$                                              |
| $\mathcal{A}$                                                          | a class of atoms                                                      |
| $\mathcal{A}(Q)$                                                       | set of atoms of class $\mathcal{A}$ supported on $Q$                  |
| $\mathcal{A}_{s,p}^{sz}$                                               | class of $(s, p)$ -Souza's atoms                                      |
| $\mathcal{A}_{s,\beta,p,q}^{bs}$                                       | class of $(s, \beta, p, q)$ -Besov's atoms                            |
| $\mathcal{P}$                                                          | grid of $I$                                                           |
| $\mathcal{P}^k$                                                        | family of subsets of $I$ at the $k$ -th level of $\mathcal{P}$        |
| $\lambda_1 \leq \lambda_2$                                             | describes geometry of the good grid $\mathcal{P}$                     |
| $\mathcal{B}_{p,q}^s(\mathcal{A})$                                     | $(s, p, q)$ -Banach space defined by the class of atoms $\mathcal{A}$ |
| $\mathcal{B}_{p,q}^s$ or $\mathcal{B}_{p,q}^s(\mathcal{A}_{s,p}^{sz})$ | $(s, p, q)$ -Banach space defined by Souza's atoms                    |
| $P, Q, W$                                                              | elements of the grid $\mathcal{P}$                                    |
| $L^p$ or $L^p(m)$                                                      | Lebesgue spaces of $(I, \mathbb{A}, m)$                               |
| $ \cdot _p$                                                            | norm in $L^p$ , $p \in (0, \infty]$                                   |
| $p'$                                                                   | $1/p + 1/p' = 1$ , with $p \in [1, \infty]$                           |

We removed from the paper's table the rows for Hölder atoms, bounded variation atoms, the constants  $\rho$  and  $\hat{t}$  (no longer needed in the Banach range) since the corresponding material is not formalized. Constants denoted  $C$ ,  $C(\cdot)$  below depend only on the displayed arguments; the Lean statements carry explicit values for most of them.

# I. THE WEAK-GRID LIBRARY

BesovSpacesGoodGrid/WeakGrid/

In Part I we assume  $s > 0$ ,  $p \in [1, \infty)$ ,  $u \in [1, \infty]$  and  $q \in [1, \infty]$ .<sup>a</sup>

<sup>a</sup>Paper:  $s > 0$ ,  $p \in (0, \infty)$ ,  $q \in (0, \infty]$ . See the Introduction for the restriction to the Banach range.

## 3. MEASURE SPACES AND WEAK GRIDS

**File:** WeakGrid/Definition.lean  
**Imports (within the project):** none (external:  
 UnbalancedHaarWavelet, LaminarFamiliesMaximalBinaryTrees)

The starting point of the whole library is deliberately poor in structure: no metric, no topology, no nesting between scales — only a finite measure space together with a sequence of finite families of measurable sets (“cells”), one family per scale  $k$ , satisfying a single quantitative hypothesis: a uniform bound  $C_1$  on how many cells of the *same* level can meet a given cell. This bounded-overlap condition  $G_1$  is exactly what is needed to make levelwise  $\ell^p$  bookkeeping work: when a level- $k$  sum of locally supported functions is integrated, each point is covered by at most  $C_1$  summands, so  $L^t$  norms of levelwise sums can be compared with  $\ell^p$  norms of their coefficients at the cost of powers of  $C_1$  (this is the computation behind inequality (6-5) of the paper). Everything in Part I — the Besov-ish spaces, their completeness, compactness, and the transmutation machinery — is built on this hypothesis alone.

Let  $I$  be a measure space with a  $\sigma$ -algebra  $\mathbb{A}$  and  $m$  be a measure on  $(I, \mathbb{A})$ ,  $m(I) < \infty$ . Given a measurable set  $J \subset I$  denote  $|J| = m(J)$ . We denote the Lebesgue spaces of  $(I, \mathbb{A}, m)$  by  $L^p$ . A **weak grid** is a sequence of finite families of measurable sets with positive measure  $\mathcal{P} = (\mathcal{P}^k)_{k \in \mathbb{N}}$ , so that at least one of these families is non empty and

$G_1$ . Given  $Q \in \mathcal{P}^k$ , let

$$\Omega_Q^k = \{P \in \mathcal{P}^k : P \cap Q \neq \emptyset\}.$$

Then

$$C_1 = \sup_k \sup_{Q \in \mathcal{P}^k} \#\Omega_Q^k < \infty.$$

Define  $|\mathcal{P}^k| = \sup\{|Q| : Q \in \mathcal{P}^k\}$ . We often abuse notation using  $\mathcal{P}$  for both  $(\mathcal{P}^k)_{k \in \mathbb{N}}$  and  $\cup_k \mathcal{P}^k$ .

**Formalization.** A grid is the structure `WeakGrid` in `WeakGrid/Definition.lean` (fields `partitions`, `measurable`, `positive_measure`, `exists_nonempty`, `Cmult1`, `overlap_card_le`), bundled with its ambient measurable space as `WeakGridSpace`. The paper calls this simply a “grid”; in the repository it is called a *weak grid*, to distinguish it from the good grids of Section 11. The paper’s convention that  $P \neq Q$  for cells at different levels is replaced by indexing cells by (level, cell) pairs (the structure `WeakGridCell`). Implementation notes: each level is a `Finset` (`Set  $\alpha$` ), so all levelwise sums are finite sums and no convergence issues arise inside a level; the exponents  $p, q, u$  live in  $\mathbb{R}_{\geq 0} \cup \{\infty\}$  (the type `ENNReal`), which lets the endpoint  $q = \infty$  or  $u = \infty$  be handled by the same statements instead of by parallel “usual modification” versions; and the overlap neighborhood  $\Omega_Q^k$  is the definition `overlapFinset`, on which  $G_1$  is imposed as a cardinality bound.

```

structure WeakGrid where
  /-- The measure on ' $\alpha$ '. -/
   $\mu$  : MeasureTheory.Measure  $\alpha$ 
  /-- The measure is finite on the whole space. -/
  isFinite : MeasureTheory.IsFiniteMeasure  $\mu$ 
  /-- The finite family ' $P^k$ ' at each level ' $k$ '. -/
  partitions :  $\mathbb{N} \rightarrow \text{Finset } (\text{Set } \alpha)$ 
  /-- Every cell is measurable. -/
  measurable :  $\forall k Q, Q \in \text{partitions } k \rightarrow \text{MeasurableSet } Q$ 
  /-- Every cell has positive measure. -/
  positive_measure :  $\forall k Q, Q \in \text{partitions } k \rightarrow 0 < \mu Q$ 
  /-- At least one level is nonempty. -/
  exists_nonempty :  $\exists k, (\text{partitions } k). \text{Nonempty}$ 
  /-- Uniform same-level overlap multiplicity bound. -/
  Cmult1 :  $\mathbb{N}$ 
  /-- ' $\# \Omega_Q^k \leq Cmult1$ ' for every ' $Q \in P^k$ '. -/
  overlap_card_le :
     $\forall k Q, Q \in \text{partitions } k \rightarrow (\text{overlapFinset } (\text{partitions } k) Q). \text{card} \leq Cmult1$ 

structure WeakGridSpace where
  grid : WeakGrid ( $\alpha := \alpha$ )

```

**Remark 3.1.** There are plenty of examples of spaces with grids; the simplest one is  $[0, 1)$  with the Lebesgue measure and the dyadic grid  $\mathcal{D}^k = \{[i/2^k, (i+1)/2^k), 0 \leq i < 2^k\}$ , and similarly the dyadic and  $d$ -adic grids of  $[0, 1)^n$ . Any measure space with a finite non-atomic measure can be endowed with a grid made of a nested sequence of partitions whose elements at the same level have precisely the same measure.

## 4. ATOMS

**File:** WeakGrid/Atoms.lean

**Imports (within the project):** WeakGrid/Definition.lean

An atom family assigns to every cell  $Q$  a set  $\mathcal{A}(Q)$  of model functions supported on  $Q$ , normalized so that an atom of a cell of measure  $|Q|$  has size at most  $|Q|^{s-1/(u'p)}$  in  $L^{pu}$ . The normalization is chosen so that the smoothness parameter  $s$  plays, on an abstract measure space, the role that derivative order plays on  $\mathbb{R}^n$ : smaller cells must carry atoms with larger norm before they contribute unit cost, exactly as in the classical atomic decompositions of Besov spaces. Two structural axioms deserve comment. Convexity and phase invariance of  $\mathcal{A}(Q)$  ( $A_3$ ) are what make the infimum norm of Section 5 a genuine norm: given representations of  $f$  and of  $g$ , the triangle inequality is proved by merging them cellwise, and the merged “atom” is a convex combination of phase-rotated atoms, hence again an atom (this is the proof of paper Proposition 6.3). The optional compactness axiom  $A_5$  is what later turns boundedness in cost into compactness in  $L^p$ . The flexibility of the abstraction — a possibly different local model space  $\mathcal{B}(Q)$  for every cell — is used in earnest in Part II, where the same interface is instantiated by one-dimensional spaces of constants (Souza’s atoms) and by infinite-dimensional induced Besov spaces (Besov’s atoms).

Let  $\mathcal{P}$  be a grid. Let  $p \in [1, \infty)$ ,  $u \in [1, \infty]$ , and  $s > 0$ . A family of **atoms** associated with  $\mathcal{P}$  of type  $(s, p, u)$  is an indexed family  $\mathcal{A}$  of pairs  $(\mathcal{B}(Q), \mathcal{A}(Q))_{Q \in \cup_k \mathcal{P}^k}$  where

- $A_1$ .  $\mathcal{B}(Q)$  is a complex linear space of functions contained in  $L^{pu}$ .
- $A_2$ . If  $\phi \in \mathcal{B}(Q)$  then  $\phi(x) = 0$  for every  $x \notin Q$ .
- $A_3$ .  $\mathcal{A}(Q)$  is a nonempty convex subset of  $\mathcal{B}(Q)$  such that  $\phi \in \mathcal{A}(Q)$  implies  $\sigma\phi \in \mathcal{A}(Q)$  for every  $\sigma \in \mathbb{C}$  satisfying  $|\sigma| = 1$ .
- $A_4$ . We have

$$|\phi|_{pu} \leq |Q|^{s - \frac{1}{u'p}}$$

for every  $\phi \in \mathcal{A}(Q)$ , where  $1/u + 1/u' = 1$ .

We will say that  $\phi \in \mathcal{A}(Q)$  is an  $\mathcal{A}$ -atom of type  $(s, p, u)$  supported on  $Q$ . Sometimes we also assume

- $A_5$ . For every  $Q \in \mathcal{P}$  we have that  $\mathcal{A}(Q)$  is a compact subset in the strong topology of  $L^p$ ,

or

- $A_6$ . every  $\mathcal{A}(Q)$  is a sequentially compact subset in the weak topology of  $L^p$ .

**Formalization.** Atom families are the structure `AtomFamily` in `WeakGrid/Atoms.lean`, whose fields formalize  $A_1$ – $A_4$ . The local space  $\mathcal{B}(Q)$  is the structure

**LocalVectorSpace** (field **localSpace**): in the paper  $A_1$  asks for a complex *Banach* space contained in  $L^{p_u}$ , while Lean only requires a complex vector space together with an injective linear map into the functions  $\alpha \rightarrow \mathbb{C}$ , all whose members belong to  $L^{p_u}$  (field **local\_memLp**); no completeness of the local space is assumed, and it is never used.  $A_2$  is the field **local\_support**;  $A_3$  corresponds to **atoms\_nonempty**, **atoms\_convex** and **atoms\_phase\_invariant** (the paper states the phase invariance as an “if and only if”; the two formulations are equivalent);  $A_4$  is the field **atom\_bound**. The compactness assumption  $A_5$  appears as the predicate **AssumptionA5**. The paper’s alternative assumption  $A_7$  ( $B(Q)$  finite dimensional and  $\mathcal{A}(Q)$  closed) is used in the paper only to derive  $A_5$ ; in the formalization the finite-dimensional case is instantiated directly for Souza’s atoms. Implementation notes: the Hölder conjugate  $u'$  is a structure field (**uConj**) constrained by the mathlib-style predicate **ENNReal.HolderConjugate**, rather than a computed value, which avoids case splits at  $u \in \{1, \infty\}$ ; the local space construction **LocalVectorSpace** is universe-polymorphic, so the same **AtomFamily** interface accepts both small concrete models (the constants  $\mathbb{C}$ , for Souza’s atoms) and large function-space models (induced  $L^p$  spaces, for Besov’s atoms); and the atom-size bound is stated with **MeasureTheory.eLpNorm**, the  $\mathbb{R}_{\geq 0} \cup \{\infty\}$ -valued  $L^p$  seminorm of mathlib, so no integrability side conditions are needed to state  $A_4$ .

```

structure AtomFamily
  (G : WeakGridSpace (α := α)) (s : ℝ) (p u : ℝ ≥ 0 ∞) where
    /-- The Hölder conjugate exponent 'u'. -/
    uConj : ℝ ≥ 0 ∞
    /-- The smoothness parameter is positive. -/
    s_pos : 0 < s
    /-- 'p ∈ [1, ∞)'. -/
    one_le_p : 1 ≤ p
    /-- 'p < ∞', expressed in 'ℝ ≥ 0 ∞' as 'p ≠ ∞'. -/
    p_ne_top : p ≠ ∞
    /-- 'u ∈ [1, ∞]'. -/
    one_le_u : 1 ≤ u
    /-- 'uConj' is the Hölder conjugate of 'u'. -/
    holder_conjugate : ENNReal.HolderConjugate u uConj
    /-- The local vector space 'B(Q)'. -/
    localSpace : WeakGridCell G → LocalVectorSpace.{u, v} α
    /-- The chosen atoms 'A(Q)', as elements of 'B(Q)'. -/
    atoms : ∀ Q, Set ((localSpace Q).carrier)
    /-- Every cell has at least one atom. -/
    atoms_nonempty : ∀ Q, (atoms Q).Nonempty
    /-- 'B(Q)' is contained in 'L^{p_u}'. -/
    local_memLp : ∀ Q φ, MeasureTheory.MemLp ((localSpace Q).toFun φ) (p * u)
      G.measure
    /-- Local functions are supported on 'Q'. -/
    local_support : ∀ Q φ, ∀ x, x ∉ Q.cell → (localSpace Q).toFun φ x = 0
    /-- 'A(Q)' is convex in the local vector space. -/
    atoms_convex : ∀ Q, Convex ℝ (atoms Q)
    /-- 'A(Q)' is invariant under multiplication by complex scalars of modulus one. -/
    atoms_phase_invariant :
      ∀ Q φ (σ : ℂ), φ ∈ atoms Q → ||σ|| = (1 : ℝ) → σ • φ ∈ atoms Q

```

```

/-- The atom size estimate in ' $L^{\{pu\}}$ '. -/
atom_bound :
  ∀ Q φ, φ ∈ atoms Q →
    MeasureTheory.eLpNorm ((localSpace Q).toFun φ) (p * u) G.measure ≤
      atomMeasureScale G s p uConj Q

```

## 5. BESOV-ISH SPACES

**File:** WeakGrid/BesovishSpaces.lean  
**Imports (within the project):** WeakGrid/Atoms.lean

This is the core file of the library. The Besov-ish space is defined by *declaring* what an atomic decomposition is and taking the functions that admit one: a representation is a choice, for every level  $k$  and cell  $Q$ , of a coefficient  $s_Q$  and an atom  $a_Q$ , such that the levelwise sums converge to  $g$  in  $L^p$ ; its *cost* is the  $\ell^q$  norm over levels of the  $\ell^p$  norms over the cells of each level — precisely the sequence-space norm of the classical Frazier–Jawerth atomic decomposition, transplanted to an abstract grid. Since representations are far from unique, the norm is the infimum of the cost over all of them. Two analytic inputs drive the section. The bounded overlap  $G_1$  gives the levelwise estimate (6-5) of the paper, and the summability hypothesis  $G_2$  (a weighted geometric-type series over the levels) lets the levelwise estimates be summed; together they yield the embedding  $\mathcal{B}_{p,q}^s(\mathcal{A}) \subset L^t$  of Proposition 5.1, which in particular ( $t = p$ ) shows that a finite-cost series really does define an element of  $L^p$  and that the infimum norm separates points. Convexity and phase invariance of the atom sets then give the triangle inequality, so the cost gauge is a norm.

Let  $p \in [1, \infty)$ ,  $u \in [1, \infty]$ ,  $q \in [1, \infty]$ ,  $s > 0$ ,  $\mathcal{P} = (\mathcal{P}^k)_{k \geq 0}$  be a grid and let  $\mathcal{A}$  be a family of atoms of type  $(s, p, u)$ . We will also assume

$G_2$ . We have

$$\left( \sum_k |\mathcal{P}^k|^{sq^*} \right)^{1/q^*} < \infty \quad (\text{usual modification if } q^* = \infty), \quad \lim_k |\mathcal{P}^k| = 0,$$

where  $q^*$  is the exponent produced by the Hölder-like estimate of the paper (for  $q > 1$ ,  $q^* = q'$ ).

**Formalization.**  $G_2$  is the predicate `AssumptionG2` in `WeakGrid/BesovishSpaces.lean`, combining the finiteness of the coefficient-weight series needed for the  $t = p$  embedding with  $\lim_k |\mathcal{P}^k| = 0$ . The paper writes this condition as  $C(p, q, (|\mathcal{P}^k|^s)_k) < \infty$  using the constant of its Proposition 4.1; since paper Section 4 has no standalone Lean counterpart, the formalization states the required summability of the weighted level measures directly.

The **Besov-ish space**  $\mathcal{B}_{p,q}^s(I, \mathcal{P}, \mathcal{A})$  is the set of all complex valued functions  $g \in L^p$  that can be represented by an absolutely convergent series in

$L^p$

$$(1) \quad g = \sum_{k=0}^{\infty} \sum_{Q \in \mathcal{P}^k} s_Q a_Q$$

where  $a_Q$  is in  $\mathcal{A}(Q)$ ,  $s_Q \in \mathbb{C}$  and with finite **cost**

$$(2) \quad \left( \sum_{k=0}^{\infty} \left( \sum_{Q \in \mathcal{P}^k} |s_Q|^p \right)^{q/p} \right)^{1/q} < \infty$$

(usual modification when  $q = \infty$ ). The series in (1) is called a  $\mathcal{B}_{p,q}^s(I, \mathcal{P}, \mathcal{A})$ -**representation** of the function  $g$ . Define

$$|g|_{\mathcal{B}_{p,q}^s(I, \mathcal{P}, \mathcal{A})} = \inf \left( \sum_{k=0}^{\infty} \left( \sum_{Q \in \mathcal{P}^k} |s_Q|^p \right)^{q/p} \right)^{1/q},$$

where the infimum runs over all possible representations of  $g$  as in (1).

**Formalization.** All in `WeakGrid/BesovishSpaces.lean`. For each level  $k$ , the structure `LevelBlock` chooses one coefficient and one atom per cell of  $\mathcal{P}^k$ ; its value in  $L^p$  (the finite sum  $\sum_{Q \in \mathcal{P}^k} s_Q a_Q$ ) is `LevelBlock.toLp`. A representation (1) is the structure `LpGridRepresentation`: a sequence of level blocks together with a `HasSum` witness in  $L^p$ . The cost (2) is `pqCost` (with the  $\mathbb{R}_{\geq 0} \cup \{\infty\}$ -valued variant `pqCostENNReal`), and its finiteness is the predicate `FinitePQCost`, with the endpoint  $q = \infty$  represented by a supremum bound. Membership with some finite-cost representation is `MemBesovishCoeffCost`; `BesovishSpace` is the corresponding `Submodule` of  $L^p$ ; and the norm  $|\cdot|_{\mathcal{B}_{p,q}^s(I, \mathcal{P}, \mathcal{A})}$  is the cost gauge `Norm_Costpq`, an infimum over representation costs. Implementation notes: convergence of the series (1) is expressed by `mathlib`’s `HasSum` over the level index, inside the Banach space `Lp` (over  $\mathbb{C}$ , exponent  $p$ ), which encapsulates the paper’s “absolutely convergent in  $L^p$ ” once the levelwise sums are grouped into blocks; the hypotheses  $1 \leq p, 1 \leq q$  are carried as `Fact` instances so that `mathlib`’s  $L^p$  machinery picks them up by instance resolution; and the normed structure on the space is provided as a *local* instance (`costNormedAddCommGroup`) rather than a global one, because the same underlying  $L^p$  element may belong to several Besov-ish spaces with different gauges.

When the choice of  $I$  and/or  $\mathcal{P}$  is obvious we write  $\mathcal{B}_{p,q}^s(\mathcal{P}, \mathcal{A})$  or  $\mathcal{B}_{p,q}^s(\mathcal{A})$ ; whenever we write just  $\mathcal{B}_{p,q}^s$  we choose the Souza’s atoms  $\mathcal{A}_{s,p}^{sz}$  of Section 12.

```

structure LpGridRepresentation
  (A : AtomFamily G s p u) (g : Lp ℂ p G.measure) where
  block : (k : ℕ) → LevelBlock A k
  hasSum : HasSum (fun k => (block k).toLp A) g

def BesovishSpace (A : AtomFamily G s p u) (q : ℝ ≥ 0 ∞)
  [Fact (1 ≤ q)]
  : Submodule ℂ (Lp ℂ p G.measure) where
  -- Carrier: all 'L^p' elements admitting a Besov-ish atomic
  -- representation with finite p q cost.

```

```

carrier := { g | MemBesovishCoeffCost A q g }
...

noncomputable def Norm_Costpq
  (A : AtomFamily G s p u) (q : ℝ≥0∞) [Fact (1 ≤ q)]
  (g : BesovishSpace A q) : ℝ :=
  pqPseudoNorm A q g

```

**Proposition 5.1** (Embedding into  $L^t$ ). *Assume  $G_1$ – $G_2$  and  $A_1$ – $A_4$ . Let  $t \in [p, \infty)$  be such that*

$$(3) \quad s - \frac{1}{p} + \frac{1}{t} \geq 0, \quad p \leq t \leq pu,$$

*and suppose*

$$(4) \quad C_2(t) = C_1^{1+1/t} C(t, q, (|\mathcal{P}^k|^{t(s-\frac{1}{p}+\frac{1}{t})})_k) < \infty.$$

*Then for every  $g$  with a  $\mathcal{B}_{p,q}^s(\mathcal{A})$ -representation (1) of cost  $K$  the series converges in  $L^t$  and*

$$(5) \quad |g|_t \leq C_2(t) K.$$

*In particular*

$$|g|_t \leq C_2(t) |g|_{\mathcal{B}_{p,q}^s(\mathcal{A})},$$

*and  $\mathcal{B}_{p,q}^s(\mathcal{A}) \subset L^t$  continuously.*

**Formalization.** The theorem `lp_embedding_adapted_statement` in `WeakGrid/BesovishSpaces.lean` proves the per-representation inequality (5): for any finite-cost representation  $R$  of  $g$ , the  $L^t$  norm of  $g$  is bounded by the explicit constant  $C_1^{1+1/t} \cdot \text{cCoefficient}$  times the cost of  $R$ . The norm inequality follows by taking the infimum over representations. Compared with paper Proposition 6.1 the hypotheses are identical except for  $q \geq 1$  and  $t \geq p$  being imposed globally, and the levelwise estimate (6.2) of the paper is internal to the proof rather than part of the statement.

```

theorem lp_embedding_adapted_statement
  {A : AtomFamily G s p u} {t : ℝ≥0∞}
  [Fact (1 ≤ t)]
  (hp_ne_top : p ≠ ∞) (ht_ne_top : t ≠ ∞)
  (hq_one : 1 ≤ q)
  (hp_le_t : p ≤ t) (ht_le_pu : t ≤ p * u)
  (hs_nonneg : 0 ≤ s - 1 / p.toReal + 1 / t.toReal)
  {g : Lp ℂ p G.measure} (R : LpGridRepresentation A g)
  (hRfin : LpGridRepresentation.FinitePQCost (q := q) R)
  (hCco_fin : cCoefficientFinite t q (fun k =>
    (levelMeasureWeight G s p t k) ^ t.toReal)) :
    (MeasureTheory.eLpNorm (g : α → ℂ) t G.measure).toReal ≤
    ((G.grid.Cmult1 : ℝ) ^ (1 + 1 / t.toReal)) *

```

```

cCoefficient t q (fun k => (levelMeasureWeight G s p t k) ^
t.toReal) *
LpGridRepresentation.pqCost (q := q) R := ...

```

**Proposition 5.2** (Linear structure). *Assume  $G_1$ – $G_2$  and  $A_1$ – $A_4$ . Then  $\mathcal{B}_{p,q}^s(\mathcal{A})$  is a complex linear subspace of  $L^p$  and  $|\cdot|_{\mathcal{B}_{p,q}^s(\mathcal{A})}$  is a norm on it. Moreover the inclusion*

$$\iota : (\mathcal{B}_{p,q}^s(\mathcal{A}), |\cdot|_{\mathcal{B}_{p,q}^s(\mathcal{A})}) \rightarrow (L^p, |\cdot|_p)$$

*is continuous.*

**Formalization.** `BesovishSpace` is by construction a `Submodule` of  $L^p$ , which gives the linear structure. The normed structure is the definition

`BesovishSpace.costNormedAddCommGroup` (requiring  $p \neq \infty$  and  $G_2$ ), which installs `Norm.Costpq` as a norm — nonnegativity, triangle inequality and homogeneity are packaged, together with the  $L^t$  embedding estimate, in the summary theorem `BesovishSpace.normedSpace_and_lp_embedding`. Continuity of  $\iota$  is the case  $t = p$  of Proposition 5.1. Compared with paper Proposition 6.3,  $\rho = \min\{1, p, q\} = 1$  since  $p, q \geq 1$ , so the  $\rho$ -norm of the paper is a norm.

```

noncomputable def costNormedAddCommGroup
  (hp_top : p ≠ ∞)
  (hCco_fin : LpGridRepresentation.cCoefficientFinite p q (fun k =>
    (LpGridRepresentation.levelMeasureWeight G s p p k) ^ p.toReal)) :
  NormedAddCommGroup (BesovishSpace A q) where
norm := Norm_Costpq A q
dist x y := Norm_Costpq A q (-x + y)
...

```

## 6. LIMITS OF REPRESENTATIONS, COMPACTNESS, AND COMPLETENESS

**File:** `WeakGrid/Completeness.lean`

**Imports (within the project):** `WeakGrid/Atoms.lean`, `WeakGrid/BesovishSpaces.lean`

The compactness and completeness theory rests on a single mechanism, the *limit of representations*: if functions  $g_n$  have representations with uniformly bounded cost, one extracts — by a Cantor diagonal argument over the countably many cells — a subsequence along which every coefficient converges and, using the compactness of the atom sets ( $A_5$  or  $A_6$ ), every atom converges as well; the limiting coefficients and atoms then form a representation of the limit function, and lower semicontinuity of the cost under cellwise convergence gives the same cost bound  $C$ . This one proposition yields, in cascade: sequential compactness of cost balls in the  $L^p$  topology, completeness of

the Besov-ish space (a Cauchy sequence converges in  $L^p$  by the embedding, and the limit stays in the space with the right norm), and later all the good-grid compactness statements of Part II. A technical point that the formalization makes explicit: to even *state* the limit representation one must know that a block sequence with finite abstract cost is summable in  $L^p$ ; this is `formalBlockSeq_summable`, proved from  $G_2$ , and it is the reason  $G_2$  appears in all the statements of this section.

**Proposition 6.1** (Limits of representations). *Assume  $G_1$ – $G_2$  and  $A_1$ – $A_4$ . Suppose that  $g_n$  are functions in  $\mathcal{B}_{p,q}^s(\mathcal{A})$  with  $\mathcal{B}_{p,q}^s(\mathcal{A})$ -representations*

$$g_n = \sum_{k=0}^{\infty} \sum_{Q \in \mathcal{P}^k} s_Q^n a_Q^n$$

satisfying

- i. *There is  $C$  such that for every  $n$*

$$\left( \sum_{k=0}^{\infty} \left( \sum_{Q \in \mathcal{P}^k} |s_Q^n|^p \right)^{q/p} \right)^{1/q} \leq C.$$

- ii. *For every  $Q \in \mathcal{P}$  the limit  $s_Q = \lim_n s_Q^n$  exists.*
- iii. *For every  $Q \in \mathcal{P}$  there is  $a_Q \in \mathcal{A}(Q)$  such that either  $a_Q^n \rightarrow a_Q$  in the strong topology of  $L^p$ , or  $a_Q^n \rightarrow a_Q$  weakly in  $L^p$ .*

*Then  $g_n$  converges (strongly, respectively weakly) in  $L^p$  to a function  $g \in \mathcal{B}_{p,q}^s(\mathcal{A})$  which has the  $\mathcal{B}_{p,q}^s(\mathcal{A})$ -representation*

$$g = \sum_{k=0}^{\infty} \sum_{Q \in \mathcal{P}^k} s_Q a_Q$$

*of cost at most  $C$ .*

**Formalization.** In `WeakGrid/Completeness.lean`, matching paper Proposition 6.4. The theorem `representation_limit` proves the weak version: under uniform cost bounds (i), cellwise convergence of the coefficients (ii) and weak convergence of the atoms (iii), the limiting blocks define a Besov-ish function of cost at most  $C$  and the  $g_n$  converge weakly. The theorem `representation_limit_strong` is the strong  $L^p$  version: strong convergence of the atoms gives strong convergence of the represented functions.

```

theorem representation_limit
  (A : AtomFamily G s p u) (hG2 : AssumptionG2 G s p u)
  {gseq : ℕ → Lp C p G.measure}
  {gLim : Lp C p G.measure} {C : ℝ}
  (H : RepresentationLimitHypotheses A q gseq gLim C)
  (hp_ne_top : p ≠ ∞) (hs_pos : 0 < s) (hu_one : 1 ≤ u)
  [Fact (1 ≤ u)] (hC : 0 ≤ C) :
  MemBesovishCoeffCost A q gLim ∧

```

```

LpGridRepresentation.FinitePQCost (q := q) H.Rlim ∧
LpGridRepresentation.pqCost (q := q) H.Rlim ≤ C ∧
Tendsto (fun n => toWeakSpace C (Lp C p G.measure) (gseq n)) atTop
(N (toWeakSpace C (Lp C p G.measure) gLim)) := ...

```

**Corollary 6.2** (Compactness and completeness). *Assume  $G_1$ – $G_2$ ,  $A_1$ – $A_4$ , and either  $A_5$  or  $A_6$ . Then*

- i. *If  $g_n \in \mathcal{B}_{p,q}^s(\mathcal{A})$  satisfy  $|g_n|_{\mathcal{B}_{p,q}^s(\mathcal{A})} \leq C$  for every  $n$ , then there is a subsequence converging (strongly under  $A_5$ , weakly under  $A_6$ ) in  $L^p$  to some  $g \in \mathcal{B}_{p,q}^s(\mathcal{A})$  with  $|g|_{\mathcal{B}_{p,q}^s(\mathcal{A})} \leq C$ .*
- ii. *Under  $A_5$ , every closed ball of  $(\mathcal{B}_{p,q}^s(\mathcal{A}), |\cdot|_{\mathcal{B}_{p,q}^s(\mathcal{A})})$  is sequentially compact in the strong topology of  $L^p$ .*
- iii. *Under  $A_5$ ,  $(\mathcal{B}_{p,q}^s(\mathcal{A}), |\cdot|_{\mathcal{B}_{p,q}^s(\mathcal{A})})$  is a complex Banach space.*

**Formalization.** All in `WeakGrid/Completeness.lean`. Item i. is

`representation_limit_strong_existence` and

`representation_limit_weak_existence` (paper Corollary 6.5(i)): they start from a bare sequence of limit blocks, prove it is summable, and package the resulting limit as a genuine representation of cost at most  $C$ . Item ii. is

`exists_strongly_convergent_subseq_of_uniform_pqCost` (extraction of a strongly convergent subsequence from any uniformly cost-bounded sequence; the extended version also returns the convergence of the coefficients and of the atoms) and

`closed_Norm_Costpq_ball_strongly_seqCompact` (sequential compactness of closed cost balls in the strong  $L^p$  topology); paper Corollary 6.5(iii) states the compactness of the inclusion  $\iota$  under  $A_5$ , and the formalization states the sequential compactness of closed cost balls, which is the equivalent form used downstream. Item iii. is `besovishSpace_costNorm_completeSpace`, completeness of the Besov-ish space for the cost norm (paper Corollary 6.5(ii); for  $p, q \geq 1$  the  $\rho$ -Banach space of the paper is a Banach space, and paper Corollary 6.7 is the finite-dimensional criterion through which the Souza case of Part II is obtained). The paper’s Corollary 6.6 (existence of a representation attaining the infimum cost) is not formalized and is omitted.

```

theorem closed_Norm_Costpq_ball_strongly_seqCompact
  (hp_ne_top : p ≠ ∞) (hs_pos : 0 < s) (hu_one : 1 ≤ u)
  (A : AtomFamily G s p u) (hG2 : AssumptionG2 G s p u q)
  (hA5 : AssumptionA5 A)
  {C : ℝ}
  [Fact (1 ≤ u)]
  (hC : 0 ≤ C)
  (gseq : ℕ → BesovishSpace A q)
  (hball : ∀ n, BesovishSpace.Norm_Costpq A q (gseq n) ≤ C) :
  ∃ (φ : ℕ → ℕ) (_hφ : StrictMono φ) (gLim : BesovishSpace A q),
    BesovishSpace.Norm_Costpq A q gLim ≤ C ∧

```

```

Tendsto
  (fun n => ((gseq (φ n) : BesovishSpace A q) : Lp ℂ p G.measure))
  atTop
  (λ (gLim : BesovishSpace A q) : Lp ℂ p G.measure)) := ...

theorem besovishSpace_costNorm_completeSpace
  (hp_ne_top : p ≠ ∞) (hs_pos : 0 < s) (hu_one : 1 ≤ u)
  (A : AtomFamily G s p u) (hG2 : AssumptionG2 G s p u q)
  (hA5 : AssumptionA5 A)
  [Fact (1 ≤ u)] :
  @CompleteSpace (BesovishSpace A q)
    (BesovishSpace.costNormedAddCommGroup
      (A := A) (q := q) hp_ne_top
      hG2.1).toMetricSpace.toPseudoMetricSpace.toUniformSpace := ...

```

## 7. SCALES OF SPACES

**File:** WeakGrid/Scales.lean

**Imports (within the project):** WeakGrid/BesovishSpaces.lean

Changing the smoothness parameter from  $s$  to  $\tilde{s}$  amounts to rescaling every atom of a cell  $Q$  by the deterministic factor  $|Q|^{\tilde{s}-s}$  — there is no analysis in the rescaling itself, only in comparing the resulting spaces. When  $s \leq \tilde{s}$  and the level measures decay fast enough (a weighted summability over levels, the same shape as  $G_2$ ), a representation in the finer scale converts levelwise into a representation in the coarser scale with a summable loss, giving the continuous inclusion  $\mathcal{B}_{p,\tilde{q}}^{\tilde{s}} \subset \mathcal{B}_{p,q}^s$  — the abstract analogue of the classical monotonicity of Besov scales in the smoothness parameter.

A family of atoms  $\mathcal{A}$  of type  $(s, p, u)$  induces a one-parameter scale  $\tilde{s} \mapsto \mathcal{A}_{\tilde{s},p}$ , where  $\mathcal{A}_{\tilde{s},p}$  is the family of atoms of type  $(\tilde{s}, p, u)$  defined by

$$\mathcal{A}_{\tilde{s},p}(Q) = \{|Q|^{\tilde{s}-s} a_Q : a_Q \in \mathcal{A}(Q)\}.$$

**Proposition 7.1** (Scale inclusion). *Assume  $G_1$ – $G_2$  and that  $\mathcal{A}$  satisfies  $A_1$ – $A_4$ . Let  $0 < s \leq \tilde{s}$  and  $q, \tilde{q} \in [1, \infty]$ . Suppose*

$$\left( \sum_k |\mathcal{P}^k|^{q^*(\tilde{s}-s)} \right)^{1/q^*} < \infty.$$

*Then  $\mathcal{B}_{p,\tilde{q}}^{\tilde{s}}(\mathcal{A}_{\tilde{s},p}) \subset \mathcal{B}_{p,q}^s(\mathcal{A}_{s,p})$ , the inclusion is a continuous linear map, and*

$$|g|_{\mathcal{B}_{p,q}^s(\mathcal{A}_{s,p})} \leq C(q, (|\mathcal{P}^k|^{\tilde{s}-s})_k) |g|_{\mathcal{B}_{p,\tilde{q}}^{\tilde{s}}(\mathcal{A}_{\tilde{s},p})}.$$

**Formalization.** In `WeakGrid/Scales.lean`. The cellwise rescaling by the deterministic factor  $\mu(Q)^{\tilde{s}-s}$  is implemented by

`smoothnessScaleFactor`, `smoothnessScaleAtoms` and `smoothnessScaleAtomFamily`. The theorem `smoothnessScaleBesovishSpace_subset` proves the set inclusion under the

deterministic summability hypothesis `scaleCoefficientFinite` and  $s \leq \tilde{s}$ ; the definition `smoothnessScaleBesovishSpaceInclusion` packages it as a linear map, and the theorem `smoothnessScaleBesovishSpaceInclusion_Norm_Costpq_le` gives the norm bound with the explicit constant `scaleCoefficient`. This is item A of paper Proposition 7.1; items B (compactness of bounded sets of the finer space in the coarser norm) and C (compactness of the inclusion under  $A_7$ ) are not formalized and are omitted. For  $u = \infty$  the two-parameter scaling map `smoothnessIntegrabilityScaleAtomFamily` (scale factor  $\mu(Q)^{\tilde{s}-s+1/p-1/\tilde{p}}$ ) is also defined, but the paper’s two-parameter scale inclusion is not formalized.

```

theorem smoothnessScaleBesovishSpace_subset
  [Fact (1 ≤ p)]
  [Fact (1 ≤ u)] [Fact (1 ≤ q)] [Fact (1 ≤ qTilde)]
  (A : AtomFamily G s p u) (sTilde : ℝ) (hsTilde_pos : 0 < sTilde)
  (hs_le : s ≤ sTilde)
  (hqfin : scaleCoefficientFinite q (smoothnessScaleLevelWeight G s
    sTilde p))
  {g : Lp ℂ p G.measure}
  (hg : g ∈ SmoothnessScaleBesovishSpace
    (A.smoothnessScaleAtomFamily sTilde hsTilde_pos) qTilde) :
  g ∈ BesovishSpace A q := ...

```

## 8. INDUCED SPACES

**File:** `WeakGrid/InducedGrid.lean`

**Imports (within the project):** `WeakGrid/BesovishSpaces.lean`

Induced spaces formalize the idea of *zooming into a cell*: fixing a cell  $Q$  at level  $k_0$ , the cells of deeper levels contained in  $Q$  form a grid of the measure space  $(Q, m|_Q)$ , with levels re-indexed to start at 0, and the atoms supported in  $Q$  form an atom family on this induced grid. Reading a representation on the induced grid as a representation on the ambient grid costs nothing (the coefficients are the same, placed at levels  $k_0 + i$ ), which is why the inclusion below is a weak contraction. This construction is the basic localization tool of the library: Besov’s atoms (Section 14) are *defined* through induced norms, and the multiplier theory (Section 19) constantly moves between a cell and the ambient space through it. The paper introduces induced spaces for good grids (Section 10 of the published version); the formalization develops them for arbitrary weak grids, since no good-grid geometry is needed.

Consider a Besov-ish space  $\mathcal{B}_{p,q}^s(I, \mathcal{P}, \mathcal{A})$ , where  $\mathcal{P}$  is a grid. Given  $Q \in \mathcal{P}^{k_0}$ , consider the sequence of finite families of subsets  $\mathcal{P}_Q = (\mathcal{P}_Q^i)_{i \geq 0}$  of  $Q$  given by

$$\mathcal{P}_Q^i = \{P \in \mathcal{P}^{k_0+i}, P \subset Q\},$$

and let  $\mathcal{A}_Q$  be the restriction of  $\mathcal{A}$  to indices in  $\mathcal{P}_Q$ . Then we can consider the **induced** Besov-ish space  $\mathcal{B}_{p,q}^s(Q, \mathcal{P}_Q, \mathcal{A}_Q)$ , and the inclusion

$$i: \mathcal{B}_{p,q}^s(Q, \mathcal{P}_Q, \mathcal{A}_Q) \rightarrow \mathcal{B}_{p,q}^s(I, \mathcal{P}, \mathcal{A})$$

is well defined and a weak contraction:

$$|f|_{\mathcal{B}_{p,q}^s(I, \mathcal{P}, \mathcal{A})} \leq |f|_{\mathcal{B}_{p,q}^s(Q, \mathcal{P}_Q, \mathcal{A}_Q)}.$$

**Formalization.** In `WeakGrid/InducedGrid.lean`. The induced grid and its ambient space are the definitions `inducedWeakGrid` and `inducedWeakGridSpace`; the restricted atom family is `inducedAtomFamily`; and the transfer of a representation from the induced grid to the ambient one, with control of cost (which gives the weak contraction above), is `inducedRepresentationToAmbient`. For Souza's atoms the formalization also transfers positivity along this map (`inducedRepresentationToAmbient.atom_toFunction_pos`); this is the form needed by the multiplier theory of Section 19. The reverse restriction map is treated, for cells of the grid, in Lemma 19.1 below.

```
def inducedRepresentationToAmbient
  (G : WeakGridSpace (α := α)) {k₀ : ℕ} (Q : LevelCell G k₀)
  {s : ℝ} {p u : ℝ≥0∞} [Fact (1 ≤ p)] (A : AtomFamily G s p u)
  {f : Lp ℂ p (inducedWeakGridSpace G Q).measure}
  (R : LpGridRepresentation (inducedAtomFamily G Q A) f) :
  LpGridRepresentation A (inducedLpToAmbient G Q f) := ...
```

## 9. TRANSMUTATION OF ATOMS

```
File: WeakGrid/Transmutation.lean
Imports (within the project): WeakGrid/Atoms.lean, WeakGrid/
BesovishSpaces.lean, WeakGrid/Completeness.lean
```

The key result in Part I (paper Proposition 8.1) states that if we replace each atom of an atomic decomposition by a combination of atoms of a possibly different class, in such way that we do not change the location of the supports and the mass of the new representation concentrates pretty much on the original scale, then we obtain a new atomic decomposition whose cost is controlled by the cost of the original one.

The mechanism is worth describing, because the formalization follows it closely. Each source atom  $a_Q$ ,  $Q \in \mathcal{G}^i$ , is expanded in target atoms  $b_{P,Q}$  with coefficients  $s_{P,Q}$  whose levelwise  $\ell^p$  mass decays geometrically away from the level  $k_i$  attached to  $i$  (hypothesis III); the level selector  $i \mapsto k_i$  is only assumed *almost linear* (hypothesis II), which is the right invariance for comparing two grids whose scales are exponentially related. Summing over  $Q$  with coefficients  $c_Q$  and collecting, for each target cell  $P$ , all contributions, one obtains target coefficients  $m_P = \sum_Q |c_Q s_{P,Q}|$  and normalized target

atoms  $d_P$  (a convex combination, by  $A_3$  again). Estimating the cost of  $(m_P)_P$  is then a discrete convolution inequality across levels: the geometric decay  $\lambda^{k-k_i}$  acts as a convolution kernel in the level variable, and the paper's convolution trick (Proposition 4.2) bounds the  $\ell^{q/p}$  norm of the convolution by the kernel mass times the source cost. This is where the explicit constant  $C_1 C_6^{1/p} \lambda^{-B/p} C_3(p, q, b) C_7^{1/q}$  of (8-17) comes from. The truncated sums (over source levels  $i \leq N$ ) are exact identities of finite character (Claim I); each truncation is a genuine representation with the right cost (Claim II); and the passage  $N \rightarrow \infty$  uses the limit-of-representations theorem of Section 6 (Claim III) — this is why the file imports **Completeness**.

**Proposition 9.1** (Transmutation of atoms). *Assume*

- I.  $\mathcal{A}_2$  is a class of  $(s, p, u_2)$ -atoms for a grid  $\mathcal{W}$ , satisfying  $A_1$ – $A_4$  and  $G_1$ – $G_2$ . Let  $\mathcal{G}$  be also a grid satisfying  $G_1$ – $G_2$ .
- II.  $k_i \in \mathbb{N}$ ,  $i \geq 0$ , is a sequence such that there are  $\alpha > 0$  and  $A, B \in \mathbb{R}$  with

$$\alpha i + A \leq k_i \leq \alpha i + B \quad \text{for every } i.$$

- III. There is  $\lambda \in (0, 1)$  such that the following holds. For every  $Q \in \mathcal{G}^i$  there are atoms  $b_{P,Q} \in \mathcal{A}_2(P)$  and coefficients  $s_{P,Q} \in \mathbb{C}$ , indexed by the cells  $P \in \mathcal{W}$  with  $P \subset Q$ , such that

$$h_Q = \sum_k \sum_{P \in \mathcal{W}^k, P \subset Q} s_{P,Q} b_{P,Q}$$

is a  $\mathcal{B}_{p,q}^s(\mathcal{A}_2)$ -representation of a function  $h_Q$ , with  $s_{P,Q} = 0$  whenever  $P \in \mathcal{W}^k$  with  $k < k_i$ , and moreover

$$(6) \quad \sum_{P \in \mathcal{W}^k, P \subset Q} |s_{P,Q}|^p \leq C_3 \lambda^{k-k_i} \quad \text{for every } k \geq k_i.$$

Then

- A. For all coefficients  $(c_Q)_{Q \in \mathcal{G}}$  with

$$\left( \sum_i \left( \sum_{Q \in \mathcal{G}^i} |c_Q|^p \right)^{q/p} \right)^{1/q} < \infty,$$

the sequence  $N \mapsto \sum_{i \leq N} \sum_{Q \in \mathcal{G}^i} c_Q h_Q$  converges in  $L^p$  to a function in  $\mathcal{B}_{p,q}^s(\mathcal{A}_2)$  that has a  $\mathcal{B}_{p,q}^s(\mathcal{A}_2)$ -representation  $\sum_k \sum_P m_P d_P$  with  $m_P \geq 0$  and

$$\left( \sum_k \left( \sum_{P \in \mathcal{W}^k} |m_P|^p \right)^{q/p} \right)^{1/q} \leq C_4 \left( \sum_i \left( \sum_{Q \in \mathcal{G}^i} |c_Q|^p \right)^{q/p} \right)^{1/q},$$

where  $C_4 = C_4(C_1, C_3, \lambda, \alpha, A, B, p, q)$  is explicit.

- B. Suppose, in addition to the assumptions of A., that the  $s_{P,Q}$  are nonnegative real numbers and  $b_{P,Q} > 0$  on  $P$  for all  $P, Q$ . If  $m_P \neq 0$  and  $d_P \neq 0$  somewhere on  $P$ , then  $P \subset Q$  for some cell  $Q \in \mathcal{G}$  with

$c_Q \neq 0$  and  $s_{P,Q} > 0$ , and the corresponding  $h_Q$  is a positive multiple of a positive function a.e. on  $P$ . If moreover  $c_Q \geq 0$  for every  $Q$ , then  $m_P \neq 0$  implies  $d_P > 0$  a.e. on  $P$ .

- C. Let  $\mathcal{A}_1$  be a class of  $(s, p, u_1)$ -atoms for the grid  $\mathcal{G}$  satisfying  $A_1$ – $A_4$ . Suppose that for every atom  $a_Q \in \mathcal{A}_1(Q)$  we can find  $s_{P,Q}$  and  $b_{P,Q}$  as in III. such that  $h_Q = a_Q$ . Then

$$\mathcal{B}_{p,q}^s(\mathcal{A}_1) \subset \mathcal{B}_{p,q}^s(\mathcal{A}_2)$$

and this inclusion is continuous, with  $|\phi|_{\mathcal{B}_{p,q}^s(\mathcal{A}_2)} \leq C_4 |\phi|_{\mathcal{B}_{p,q}^s(\mathcal{A}_1)}$  for every  $\phi \in \mathcal{B}_{p,q}^s(\mathcal{A}_1)$ .

**Formalization.** In `WeakGrid/Transmutation.lean`. The transmuted objects (target coefficients  $m_{P,N}$ , normalized target atoms  $d_{P,N}$ , target level blocks, and their stable limits) are the definitions `TransmutationCoeff`, `TransmutationAtom`, `TransmutationBlock` and `TransmutationCoeffLimit`, `TransmutationAtomLimit`, `TransmutationBlockLimit`; the hypothesis III is the predicate `RepresentationWsubGandALS`, with the almost-linear level control II packaged as `AlmostLinearSequence`. The internal pipeline is formalized by the theorems `ClaimI` (exact bookkeeping identity: the transmuted representation at truncation  $N$  sums to the source partial expansion), `ClaimII` (each finite transmutation block sequence is a genuine representation of the source partial expansion, with cost controlled by the source cost) and `ClaimIII` (passage to the stable limiting data, producing the  $L^p$  limit and its representation), together with their endpoint versions `ClaimI_top`, `ClaimII_top`, `ClaimIII_top` for  $q = \infty$ . Item A is the theorem `Transmutation_of_Atoms_Claim_A`, with the explicit constant of the paper (product of  $C_1$ ,  $C_6^{1/p}$ ,  $\lambda^{-B/p}$ , the coefficient of the convolution trick — paper Proposition 4.2, which in Lean is the constant `cCoefficientInt` of the kernel `transmutationKernelZ` — and the integer-part factor  $\max\{\ell \in \mathbb{N} : \ell < \alpha\} + 1$ ). Item B is `Transmutation_of_Atoms_Claim_B` and `Transmutation_of_Atoms_Claim_B_sharp` (support and positivity of the transmuted data); compared with paper Proposition 8.1.B, the pointwise positivity conclusions are stated almost everywhere on  $P$  (with positivity witnessed by positive real multiples), which is the natural formulation in  $L^p$ . Item C is `Transmutation_of_Atoms_Claim_C_explicit` and the continuous-embedding form `Transmutation_of_Atoms_continuous_embedding_explicit`. For the identity level selector, the theorem `LpGridRepresentation.cCoefficientInt_transmutationKernelZ_zero_one` computes the integer transmutation coefficient of the Claim C kernel as the geometric sum  $(1 - \rho)^{-1}$ ,  $\lambda = \rho^p$ ; this is the constant simplification used in the Souza/Besov atom comparison of Section 14.

```

theorem Transmutation_of_Atoms_Claim_A
  (G W : WeakGridSpace (α := α))
  (AW : AtomFamily W s p u)
  (k : ℕ → ℕ)
  (A_als B_als r_als : ℝ)
  (hr_als : 0 < r_als)

```

```

(hk_bound : ∀ i : ℕ,
  (k i : NNReal) ≤ r_als * (i : NNReal) + B_als ∧
  r_als * (i : NNReal) + A_als ≤ (k i : NNReal))
(lam : ℝ) (hlam_pos : 0 < lam) (hlam_lt : lam < 1)
(C : ℝ) (hC : 0 ≤ C)
(h : (i : ℕ) → LevelCell G i → Lp ℂ p W.measure)
(R : (i : ℕ) → (Q : LevelCell G i) → LpGridRepresentation AW (h i Q))
(hR : RepresentationWsubGandALS (p := p) (q := q) G W AW k
  ⟨A_als, B_als, r_als, hr_als, hk_bound⟩ lam hlam_pos hlam_lt C hC h R)
(c : (i : ℕ) → LevelCell G i → ℂ)
(hc : CoeffFinitePQCost (p := p) (q := q) G c)
(hq_ne_top : q ≠ ∞)
(hG2_W : AssumptionG2 W s p u q)
(hp_ne_top : p ≠ ∞)
(hs_pos : 0 < s) :
∃ gLim : Lp ℂ p W.measure,
  ∃ hsum :
    HasSum (fun j => (TransmutationBlockLimit G W AW h R c A_als r_als j).toLp
AW) gLim,
    MemBesovishCoeffCost AW q gLim ∧
    LpGridRepresentation.FinitePQCost (q := q)
    ({ block := TransmutationBlockLimit G W AW h R c A_als r_als
      hasSum := hsum } : LpGridRepresentation AW gLim) ∧
    Tendsto (fun N => PartialSumLevels G W h c N) atTop (ℵ gLim) ∧
    LpGridRepresentation.pqCost (q := q)
    ({ block := TransmutationBlockLimit G W AW h R c A_als r_als
      hasSum := hsum } : LpGridRepresentation AW gLim) ≤
    (G.grid.Cmult1 : ℝ) *
    C ^ (1 / p.toReal) *
    lam ^ (-(B_als : ℝ) / p.toReal) *
    LpGridRepresentation.cCoefficientInt p ∞
    (transmutationKernelZ lam A_als r_als) *
    (Nat.ceil (r_als : ℝ) : ℝ) ^ (1 / q.toReal) *
    CoeffPQCost (p := p) (q := q) G c := ...

theorem Transmutation_of_Atoms_Claim_B (G W : WeakGridSpace (α := α))
  (AW : AtomFamily W s p u)
  (k : ℕ → ℕ)
  (A_als B_als r_als : ℝ)
  (hr_als : 0 < r_als)
  (hk_bound : ∀ i : ℕ,
    (k i : NNReal) ≤ r_als * (i : NNReal) + B_als ∧
    r_als * (i : NNReal) + A_als ≤ (k i : NNReal))
  (lam : ℝ) (hlam_pos : 0 < lam) (hlam_lt : lam < 1)
  (C : ℝ) (hC : 0 ≤ C)
  (h : (i : ℕ) → LevelCell G i → Lp ℂ p W.measure)
  (R : (i : ℕ) → (Q : LevelCell G i) → LpGridRepresentation AW (h i Q))
  (hR : RepresentationWsubGandALS_pos (p := p) (q := q) G W AW k
    ⟨A_als, B_als, r_als, hr_als, hk_bound⟩ lam hlam_pos hlam_lt C hC h R)
  (c : (i : ℕ) → LevelCell G i → ℂ)
  (hc : CoeffFinitePQCost (p := p) (q := q) G c)
  (hG2_W : AssumptionG2 W s p u q)
  (hp_ne_top : p ≠ ∞)
  (hs_pos : 0 < s) :
  (∃ gLim : Lp ℂ p W.measure,

```

```

HasSum (fun j => (TransmutationBlockLimit G W AW h R c A_als r_als j).toLp
AW) gLim ^
MemBesovishCoeffCost AW q gLim ^
Tendsto (fun N => PartialSumLevels G W h c N) atTop (λ gLim) ^
(∀ j : ℕ, ∀ P : LevelCell W j,
TransmutationCoeffLimit G W AW h R c A_als r_als P ≠ 0 →
(∃ x, x ∈ P.1 →
AW.toFunction (levelCellToWeakGridCell W j P)
(TransmutationAtomLocalLimit G W AW h R c A_als r_als P) x ≠ 0) →
∃ i ∈ Finset.range (transmutationStabilizationIndex A_als r_als j),
∃ Q ∈ (G.grid.partitions i).attach,
P.1 ⊆ Q.1 ∧ c i Q ≠ 0 ∧
∃ r : NNReal, 0 < r ∧ ((R i Q).block j).coeff P = (r : ℂ) ∧
∀m x ∂ W.measure.restrict P.1,
∃ r : NNReal, 0 < r ∧ h i Q x = (r : ℂ)) := ...

theorem Transmutation_of_Atoms_continuous_embedding_explicit
(G : WeakGridSpace (α := α))
(u1 u2 : ℝ≥0∞)
[Fact (1 ≤ u2)]
(AG1 : AtomFamily G s p u1)
(AG2 : AtomFamily G s p u2)
(lam : ℝ) (hlam_pos : 0 < lam) (hlam_lt : lam < 1)
(C : ℝ) (hC : 0 ≤ C)
(hAG1_to_AG2 : ∀ i : ℕ, ∀ Q : LevelCell G i,
∀ g : Lp ℂ p G.measure,
(∃ φ : (AG1.localSpace (levelCellToWeakGridCell G i Q)).carrier,
AG1.IsAtom (levelCellToWeakGridCell G i Q) φ ∧
g = atomLp AG1 (levelCellToWeakGridCell G i Q) φ) →
∃ Rg : LpGridRepresentation AG2 g,
CoeffFinitePQCost (p := p) (q := q) G
(fun j S => (Rg.block j).coeff S) ∧
(∀ j : ℕ, ∀ S : LevelCell G j,
(¬ S.1 ⊆ Q.1 → (Rg.block j).coeff S = 0) ∧
(j < i → (Rg.block j).coeff S = 0)) ∧
∀ j : ℕ, i ≤ j → Rg.levelCoeffPower j ≤ C * lam ^ (j - i))
(hG2_G : AssumptionG2 G s p u2 q)
(hp_ne_top : p ≠ ∞)
(hs_pos : 0 < s) :
0 ≤ transmutationClaimCEmbeddingConstant G p q lam C ^
∀ g : BesovishSpace AG1 q,
∃ hg2 : MemBesovishCoeffCost AG2 q (g : Lp ℂ p G.measure),
BesovishSpace.Norm_Costpq AG2 q
((g : Lp ℂ p G.measure), hg2) : BesovishSpace AG2 q ≤
transmutationClaimCEmbeddingConstant G p q lam C *
BesovishSpace.Norm_Costpq AG1 q g := ...

```

## 10. FURTHER WEAK-GRID INFRASTRUCTURE

Two more files complete the weak-grid layer; they contain no statement of the paper but are imported by Part II.

**File:** WeakGrid/Multipliers.lean  
**Imports (within the project):** WeakGrid/Completeness.lean, WeakGrid/InducedGrid.lean

This file defines, at the weak-grid level of generality, the predicates on which the multiplier theory of Section 19 is built: a measurable  $g$  is a *pointwise multiplier* of a Besov-ish space (`IsPointwiseMultiplier`: multiplication by  $g$  maps the space to itself boundedly for the cost gauge) and  $g$  is *in the selfs class* (`PointwiseSelfsClass`: the products of  $g$  with all atoms have uniformly bounded Besov-ish norm). Keeping these definitions abstract means that the implication “multiplier  $\Rightarrow$  selfs class”, which holds for any atom family, lives here, while everything genuinely good-grid-ish is deferred to `GoodGrid/Multipliers/`.

**File:** Sums.lean  
**Imports (within the project):** none

Reusable indexing types and notation for sums over grid blocks: the global pair type `BlockIndex` for  $(k, Q)$ , the types `ChildIndex` and `ParentIndex` for strict descendants and ancestors of a cell, and the corresponding `tsum` notations (`sumb`, `sum+`, `sum-` in the source file).

## II. THE GOOD-GRID LIBRARY

BesovSpacesGoodGrid/GoodGrid/

In Part II we suppose that  $s > 0$ ,  $p \in [1, \infty)$  and  $q \in [1, \infty]$ ; from Section 19 on (the multiplier theory) we further assume

$$0 < s < 1/p.$$

### 11. GOOD GRIDS

**File:** GoodGrid/Definition.lean  
**Imports (within the project):** none (external: `UnbalancedHaarWavelet`)

A good grid adds to a weak grid the geometry of a nested dyadic-like structure: levels partition the space  $(G_4 - G_5)$ , each cell sits inside a parent at the previous level  $(G_6)$ , and — the quantitative heart — the measure of a child is comparable to the measure of its parent, with ratio pinched between  $\lambda_1$  and  $\lambda_2$  ( $G_7$ ). Two consequences are used constantly. First, cell measures decay geometrically in the level,  $|Q| \leq \lambda_2^k |I|$  for  $Q \in \mathcal{P}^k$ , so the hypothesis  $G_2$  of Part I holds *automatically* for every  $s > 0$ : the entire weak-grid theory

applies to a good grid with no extra summability assumptions. Second, a cell has at most  $1/\lambda_1$  children, which bounds all the branching constants in the Haar construction of Section 15. The simplest example is the dyadic grid of  $[0, 1)$  with  $\lambda_1 = \lambda_2 = 1/2$ ; a good grid, however, need not be self-similar, and the two-sided ratio bound is the only regularity retained.

A  $(\lambda_1, \lambda_2)$ -**good grid**, with  $0 < \lambda_1 \leq \lambda_2 < 1$ , is a grid  $\mathcal{P} = (\mathcal{P}^k)_{k \in \mathbb{N}}$  with the following properties:

$G_3$ .  $\mathcal{P}^0 = \{I\}$ .

$G_4$ .  $I = \cup_{Q \in \mathcal{P}^k} Q$  up to a set of zero  $m$ -measure.

$G_5$ . The elements of  $\mathcal{P}^k$  are pairwise disjoint.

$G_6$ . For every  $Q \in \mathcal{P}^k$ ,  $k > 0$ , there exists  $P \in \mathcal{P}^{k-1}$  such that  $Q \subset P$ .

$G_7$ . We have

$$\lambda_1 \leq \frac{|Q|}{|P|} \leq \lambda_2$$

for every  $Q \subset P$  with  $Q \in \mathcal{P}^{k+1}$  and  $P \in \mathcal{P}^k$ ,  $k \geq 0$ .

$G_8$ . The family  $\cup_k \mathcal{P}^k$  generates the  $\sigma$ -algebra  $\mathbb{A}$ .

**Formalization.** Good grids are the structure `GoodGrid` in `GoodGrid/Definition.lean`, which extends the structure `UnbalancedHaarWavelet.Grid` of the external repository (where the nested-partition properties  $G_3$ – $G_6$ ,  $G_8$  live) with the ratio bounds  $G_7$  (fields `ratio_lower`, `ratio_upper`). The ambient object used by the good-grid files is `GoodGridSpace`, a measurable space with a fixed good grid; the definition `GoodGridSpace.toWeakGrid` turns a good grid into a weak grid (since good-grid partitions are disjoint, the same-level overlap multiplicity is 1), so that the whole of Part I applies. The paper requires  $\lambda_1 < \lambda_2$ ; the formalization allows  $\lambda_1 \leq \lambda_2$ .

```

structure GoodGrid extends UnbalancedHaarWavelet.Grid (α := α) where
  lambda1 : ℝ
  /-- Upper ratio bound (strictly less than 1). -/
  lambda2 : ℝ
  /-- 'λ₁' is strictly positive. -/
  hlambda1_pos : 0 < lambda1
  /-- 'λ₂' is strictly less than '1'. -/
  hlambda2_lt_one : lambda2 < 1
  /-- 'λ₁ ≤ λ₂'. -/
  hlambda1_le_lambda2 : lambda1 ≤ lambda2
  /-- Each child cell's measure is at least 'λ₁' times its parent's
  measure. -/
  ratio_lower : ∀ (n : ℕ) (s t : Set α),
    s ∈ grid.partitions (n + 1) → t ∈
grid.partitions n → s ⊆ t →
    ENNReal.ofReal lambda1 * μ t ≤ μ s
  /-- Each child cell's measure is at most 'λ₂' times its parent's
  measure. -/
  ratio_upper : ∀ (n : ℕ) (s t : Set α),

```

```

s ∈ grid.partitions (n + 1) → t ∈
grid.partitions n → s ⊆ t →
μ s ≤ ENNReal.ofReal lambda2 * μ t

```

## 12. SOUZA'S ATOMS

**File:** GoodGrid/BesovSpace.lean  
**Imports (within the project):** WeakGrid/Atoms.lean, WeakGrid/  
BesovishSpaces.lean, WeakGrid/Completeness.lean, WeakGrid/  
InducedGrid.lean, GoodGrid/Definition.lean

Souza's atoms (paper Section 11A) are the simplest conceivable atoms: a multiple of the indicator of a cell, normalized in  $L^\infty$ . Their local model space is one-dimensional ( $\mathcal{B}(Q) \cong \mathbb{C}$ ), so the atom set of every cell is a closed bounded subset of a one-dimensional space, hence compact: assumption  $A_5$  comes for free (this is the finite-dimensional criterion of paper Corollary 6.7), and the whole compactness/completeness theory of Part I applies. Their simplicity is the point of the construction: products, restrictions, and positivity of Souza's atoms are trivial to control, which is what makes the multiplier theory of Section 19 tractable.

Let  $Q \in \mathcal{P}$ . A  $(s, p)$ -**Souza's atom** supported on  $Q$  is a function  $a: I \rightarrow \mathbb{C}$  such that  $a(x) = 0$  for every  $x \notin Q$  and  $a$  is constant on  $Q$ , with

$$|a|_\infty \leq |Q|^{s-1/p}.$$

The set of Souza's atoms supported on  $Q$  will be denoted by  $\mathcal{A}_{s,p}^{sz}(Q)$ . The **canonical Souza's atom** on  $Q$  is the Souza's atom with  $a(x) = |Q|^{s-1/p}$  for every  $x \in Q$ . Souza's atoms are  $(s, p, \infty)$ -type atoms and satisfy  $A_1$ – $A_5$ .

**Formalization.** In GoodGrid/BesovSpace.lean. The predicate `IsSouzaAtom` says that a function is supported on a good-grid cell, constant on it, and of size at most  $|Q|^{s-1/p}$ ; the canonical atom is `canonicalSouzaAtom`, with `canonicalSouzaAtom_isSouzaAtom` proving that it is indeed a Souza's atom. The definitions `souzaLocalVectorSpace`, `souzaAtomsSet` and `souzaAtomFamily` package Souza's atoms as an `AtomFamily` with  $u = \infty$  satisfying  $A_1$ – $A_5$ . The paper's further examples — Hölder atoms (Section 11B) and bounded variation atoms (Section 11C) — are not formalized and are omitted.

```

def IsSouzaAtom (G : GoodGridSpace (α := α)) (s : ℝ) (p : ℝ ≥ 0 ∞)
  (Q : GoodGridCell G) (a : α → ℂ) : Prop :=
  (∀ x ∉ Q.cell, a x = 0) ∧
  ∃ c : ℂ, (∀ x ∈ Q.cell, a x = c) ∧
  ||c|| ≤ (G.grid.μ Q.cell).toReal ^ (s - (p.toReal)⁻¹)

```

## 13. BESOV SPACES IN A MEASURE SPACE WITH A GOOD GRID

**File:** GoodGrid/BesovSpace.lean

**Imports (within the project):** (same file as the previous section)

The paper’s Section 12 introduces the spaces  $\mathcal{B}_{p,q}^s = \mathcal{B}_{p,q}^s(\mathcal{P}, \mathcal{A}_{s,p}^{sz})$  and observes that the general theory applies; it states no numbered theorem there. The theorem below is the formalization’s packaging of the resulting Souza-space facts, obtained by instantiating Proposition 5.1 and Corollary 6.2 (paper Proposition 6.1 and Corollaries 6.5 and 6.7). The density statements in item C deserve a remark: a function constant on the cells of level  $k$  is, by definition, a finite sum of level- $k$  Souza’s atoms, and as  $k \rightarrow \infty$  such “grid-simple” functions approximate any  $L^p$  function (this is where  $G_8$ , the generation of the  $\sigma$ -algebra by the cells, enters); for  $s < 1/p$  the cost of these approximations stays finite, giving density — so, in contrast with classical Besov spaces of positive smoothness on  $\mathbb{R}^n$ , here the space is a *dense* compactly embedded subspace of  $L^p$ .

We study the Besov-ish spaces  $\mathcal{B}_{p,q}^s = \mathcal{B}_{p,q}^s(\mathcal{P}, \mathcal{A}_{s,p}^{sz})$  associated with a measure space with a good grid  $(I, \mathcal{P}, m)$ . If  $p \in [1, \infty)$ ,  $q \in [1, \infty]$  and  $0 < s < \frac{1}{p}$  we say that  $\mathcal{B}_{p,q}^s$  is a **Besov space**. The general theory of Part I specializes as follows.

**Theorem 13.1.** *Let  $(I, \mathcal{P}, m)$  be a measure space with a good grid,  $s > 0$ ,  $p \in [1, \infty)$ ,  $q \in [1, \infty]$ . Then*

- A.  $(\mathcal{B}_{p,q}^s, |\cdot|_{\mathcal{B}_{p,q}^s})$  is a complex Banach space.
- B. Closed balls of  $\mathcal{B}_{p,q}^s$  are compact subsets of  $L^p$ , and also of  $L^1$ .
- C. If  $0 < s < 1/p$  then  $\mathcal{B}_{p,q}^s$  is dense in  $L^p$ , and dense in  $L^1$ .

**Formalization.** All in GoodGrid/BesovSpace.lean. Item A is `souzaBesovSpace_costNorm_completeSpace`, the specialization of the weak-grid completeness theorem to Souza’s atoms on a good grid. Item B is `souza_closedCostBallInLp_isSeqCompact`, `souza_closedCostBallInLp_isCompact` and `souza_closedCostBallInL1_isCompact`: compactness of closed cost balls in the ambient  $L^p$  topology and, after the finite-measure inclusion, in  $L^1$  — the Souza-atom instance of Corollary 6.2 together with the embedding of Proposition 5.1. Item C is `souzaBesovSpace_dense` and `souzaBesovSpace_dense_inL1`, density of the Souza Besov space in  $L^p$  and of its image in  $L^1$ ; the density statements do not appear explicitly as theorems in the paper, and are recorded here because they are formalized.

```
theorem souzaBesovSpace_costNorm_completeSpace
  (G : GoodGridSpace (α := α))
  (s : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)] :
```

```

@CompleteSpace
  (SouzaBesovSpace G s p q hs hp hp_top)
  (WeakGridSpace.BesovishSpace.costNormedAddCommGroup
    (A := souzaAtomFamily G s p hs hp hp_top) (q := q)
    hp_top (souza_hCco G s p q hs hp
      hp_top)).toMetricSpace.toPseudoMetricSpace.toUniformSpace := ...

theorem souza_closedCostBallInLp_isCompact
  (G : GoodGridSpace (α := α))
  (s : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)]
  {C : ℝ} (hC : 0 ≤ C) :
  IsCompact
    {f : MeasureTheory.Lp C p G.toWeakGridSpace.measure |
      ∃ hf : f ∈ SouzaBesovSpace G s p q hs hp hp_top,
        WeakGridSpace.BesovishSpace.Norm_Costpq
          (souzaAtomFamily G s p hs hp hp_top) q (f, hf) ≤ C} := ...

theorem souzaBesovSpace_dense
  (G : GoodGridSpace (α := α))
  (s : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)] :
  Dense (SouzaBesovSpace G s p q hs hp hp_top :
    Set (MeasureTheory.Lp C p G.toWeakGridSpace.measure)) := ...

```

#### 14. BESOV’S ATOMS AND THE SOUZA–BESOV COMPARISON

**File:** GoodGrid/BesovAtoms.lean

**Imports (within the project):** GoodGrid/BesovSpace.lean, WeakGrid/  
InducedGrid.lean, WeakGrid/Transmutation.lean

This is the paper’s “Alternative characterizations, II: messing with atoms” (Section 16 of the published version), and the first serious application of the transmutation machinery. A Besov atom on  $Q$  is not required to be constant — it is any function supported on  $Q$  whose *induced* Besov norm of a higher smoothness  $\beta > s$  is controlled by the scale  $|Q|^{s-\beta}$ ; the induced spaces of Section 8 are exactly what makes this definition possible. The comparison theorem (paper Proposition 16.1) says that any atom family sandwiched between Souza’s atoms and Besov’s atoms generates the *same* space with equivalent norms. The nontrivial inclusion is proved by transmutation: each Besov atom is expanded, inside its own cell, into Souza’s atoms with geometrically decaying levelwise mass (rate  $\lambda_2^{(k-j)(\beta-s)}$  — this is where  $\beta > s$  and the good-grid ratio bound enter), which is precisely hypothesis III of paper Proposition 8.1 with the identity level selector; Claim C of that proposition then converts every representation by Besov atoms into one by Souza’s atoms with controlled cost. In practice this proposition is the

standard tool for checking that a concrete function belongs to  $\mathcal{B}_{p,q}^s$ : it suffices to exhibit atoms in any sandwiched class.

Let  $s < \beta$  and  $\tilde{q} \in [1, \infty]$ . A  $(s, \beta, p, \tilde{q})$ -**Besov atom** on  $Q$  is a function  $a \in \mathcal{B}_{p,\tilde{q}}^\beta(Q, \mathcal{P}_Q, \mathcal{A}_{\beta,p}^{sz})$  such that  $a(x) = 0$  for  $x \notin Q$  and

$$|a|_{\mathcal{B}_{p,\tilde{q}}^\beta(Q, \mathcal{P}_Q, \mathcal{A}_{\beta,p}^{sz})} \leq \frac{1}{C_7} |Q|^{s-\beta},$$

where  $C_7 = C_7(C_1, \lambda_2, \beta, \tilde{q}, p) \geq 1$  is the normalizing constant of the paper. The family of  $(s, \beta, p, \tilde{q})$ -Besov atoms supported on  $Q$  is denoted  $\mathcal{A}_{s,\beta,p,\tilde{q}}^{bs}(Q)$ , and  $\mathcal{A}_{s,p}^{sz}(Q) \subset \mathcal{A}_{s,\beta,p,\tilde{q}}^{bs}(Q)$ ; Besov atoms are atoms of type  $(s, p, 1)$ .

**Proposition 14.1** (Souza's atoms and Besov's atoms). *Let  $\mathcal{P}$  be a good grid and  $0 < s < 1/p$ . Let  $\mathcal{A}$  be a class of  $(s, p, u)$ -atoms, with  $u \geq 1$ , such that for some  $s < \beta$ ,  $\tilde{q} \in [1, \infty]$ , and  $C_8, C_9 \geq 0$  we have, for every  $Q \in \mathcal{P}$ ,*

$$\frac{1}{C_8} \mathcal{A}_{s,p}^{sz}(Q) \subset \mathcal{A}(Q) \subset C_9 \mathcal{A}_{s,\beta,p,\tilde{q}}^{bs}(Q).$$

Then

$$\mathcal{B}_{p,q}^s(\mathcal{P}, \mathcal{A}_{s,p}^{sz}) = \mathcal{B}_{p,q}^s(\mathcal{P}, \mathcal{A}) = \mathcal{B}_{p,q}^s(\mathcal{P}, \mathcal{A}_{s,\beta,p,\tilde{q}}^{bs}),$$

with

$$|f|_{\mathcal{B}_{p,q}^s(\mathcal{A})} \leq C_8 |f|_{\mathcal{B}_{p,q}^s(\mathcal{A}_{s,p}^{sz})} \quad \text{and} \quad |f|_{\mathcal{B}_{p,q}^s(\mathcal{A}_{s,\beta,p,\tilde{q}}^{bs})} \leq C |f|_{\mathcal{B}_{p,q}^s(\mathcal{A})}.$$

**Formalization.** This is paper Proposition 16.1. The sandwich proposition is the theorem `atoms_between_souza_atoms_and_besov_atoms` in `GoodGrid/BesovAtoms.lean`: it proves the equality of the three Besov-ish spaces and both continuous norm bounds, the second with the explicit constant  $C_2/(1 - \lambda_2^{\beta-s})$ . It is supported by the decay claims `besovAtom_to_souza_representation_decay` and `besovAtom_to_induced_souzaS_representation_decay_claimC` (the Lean counterparts of the Claim in the paper's proof: a single Besov atom, read on the weak grid induced on its supporting cell, admits a Souza-atom representation with geometric coefficient decay rate  $\lambda_2^{(k-j)(\beta-s)}$ ), by the normalizing constant `besovAtomConstant` (with `one_le_besovAtomConstant` giving the lower bound  $1 \leq C_7$  used to simplify the final constant), and by the identity-level geometric-sum computation `LpGridRepresentation.cCoefficientInt_transmutationKernelZ_zero_one` of Section 9. Paper Propositions 16.2 (Hölder atoms) and 16.3 (bounded variation atoms) are not formalized and are omitted.

```

theorem atoms_between_souza_atoms_and_besov_atoms
  (G : GoodGridSpace (α := α)) (s β : ℝ) (p u q qtilde : ℝ ≥ 0 ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ u)] [Fact (1 ≤ q)] [Fact (1 ≤ qtilde)]
  (hs : 0 < s) (hβ : 0 < β) (hβs : s < β) (hp_top : p ≠ ∞)
  (A : WeakGridSpace.AtomFamily G.toWeakGridSpace s p u)
  (C1 C2 : ℝ)
  (hSandwich :
    SouzaBesovSandwich G s β p u qtilde hs hβ (inferInstance : Fact (1 ≤ p))
    hp_top A C1 C2) :

```

```

(WeakGridSpace.BesovishSpace (souzaAtomFamily G s p hs (Fact.out : 1 ≤ p)
hp_top) q =
  WeakGridSpace.BesovishSpace A q) ∧
(WeakGridSpace.BesovishSpace A q =
  WeakGridSpace.BesovishSpace
    (besovAtomFamily G s β p qtilde hs hβ inferInstance hp_top) q) ∧
(∀ f : WeakGridSpace.BesovishSpace
  (souzaAtomFamily G s p hs (Fact.out : 1 ≤ p) hp_top) q,
  ∃ hfA : WeakGridSpace.MemBesovishCoeffCost A q
    (f : Lp ℂ p G.toWeakGridSpace.measure),
    WeakGridSpace.BesovishSpace.Norm_Costpq A q
      ((f : Lp ℂ p G.toWeakGridSpace.measure), hfA) :
      WeakGridSpace.BesovishSpace A q)
    ≤ C1 *
      WeakGridSpace.BesovishSpace.Norm_Costpq
        (souzaAtomFamily G s p hs (Fact.out : 1 ≤ p) hp_top) q f) ∧
(∀ f : WeakGridSpace.BesovishSpace A q,
  ∃ hfS : WeakGridSpace.MemBesovishCoeffCost
    (souzaAtomFamily G s p hs (Fact.out : 1 ≤ p) hp_top) q
    (f : Lp ℂ p G.toWeakGridSpace.measure),
    WeakGridSpace.BesovishSpace.Norm_Costpq
      (souzaAtomFamily G s p hs (Fact.out : 1 ≤ p) hp_top) q
      ((f : Lp ℂ p G.toWeakGridSpace.measure), hfS) :
      WeakGridSpace.BesovishSpace
        (souzaAtomFamily G s p hs (Fact.out : 1 ≤ p) hp_top) q)
    ≤ (C2 / (1 - G.grid.lambda2 ^ (β - s))) *
      WeakGridSpace.BesovishSpace.Norm_Costpq A q f) := ...

```

## 15. UNBALANCED HAAR WAVELETS

**File:** GoodGrid/AlternativeRepresentationsAndNorms/  
 HaarRepresentationNorm.lean  
**Imports (within the project):** GoodGrid/BesovSpace.lean (external:  
 UnbalancedHaarWavelet, UnconditionalSchauderBasis, Burkholder)

On a good grid there is a genuine wavelet system: since each cell has between 2 and  $1/\lambda_1$  children, one can follow Girardi and Sweldens and organize the children of each cell into a balanced binary tree of subsets (the “logarithmic subtrees”), attaching to each internal node a two-valued zero-mean function — an *unbalanced Haar function*. The measure-ratio bounds  $G_7$  guarantee that each  $\phi_S$  supported in  $Q$  has size  $\approx |Q|^{-1/2}$  on its support, so the system behaves like the classical Haar basis even though the grid may be very irregular. The unconditionality of this basis in  $L^\beta$ ,  $1 < \beta < \infty$  — a Burkholder-type martingale inequality — is the one substantial analytic ingredient that the repository does not prove internally: it is supplied by the external formalized libraries listed above.

Let  $\mathcal{P} = (\mathcal{P}^k)_k$  be a good grid. Following Girardi and Sweldens, one constructs, from the children structure of the grid (Type I tree with the

logarithmic subtrees construction), a family of **unbalanced Haar functions**  $\{\phi_S\}_{S \in \hat{\mathcal{H}}}$ , where  $\hat{\mathcal{H}} = \{I\} \cup \mathcal{H}$ ,  $\phi_I = 1_I/|I|^{1/2}$ , each  $\phi_S$  with  $S \in \mathcal{H}_Q$  is supported on  $Q$ , has zero mean,  $L^2$ -norm one, and satisfies

$$\frac{C_5(\lambda_1, \lambda_2)}{|Q|^{1/2}} \leq |\phi_S(x)| \leq \frac{C_6(\lambda_1, \lambda_2)}{|Q|^{1/2}}$$

on its support. The family  $\{\phi_S\}_{S \in \hat{\mathcal{H}}}$  is an unconditional basis of  $L^\beta$  for every  $1 < \beta < \infty$ .

**Formalization.** The construction and the unconditional basis property are provided by the external Lean repositories `UnbalancedHaarWavelet`, `UnconditionalSchauderBasis`, `LaminarFamiliesMaximalBinaryTrees` and `Burkholder` (SmaniaD), on which this repository depends. The local API ( $L^2$ -normalized Haar functions `L2normalizedHaar` and Haar coefficients `Coeff`) is in `GoodGrid/AlternativeRepresentationsAndNorms/HaarRepresentationNorm.lean`.

```
def L2normalizedHaar (G : GoodGridSpace (α := α)) [DecidableEq (Set α)]
  (F : UnbalancedHaarWavelet.FullHaarSystem (G := GridOf G))
  (i : F.Index) (x : α) : ℂ :=
  ((L2NormalizationFactor G F i : ℝ) : ℂ) *
  (UnbalancedHaarWavelet.FullHaarSystem.function (GridOf G) F i x : ℂ)

def Coeff (G : GoodGridSpace (α := α)) [DecidableEq (Set α)]
  (F : UnbalancedHaarWavelet.FullHaarSystem (G := GridOf G))
  (f : α → ℂ) (_hf : Integrable f G.grid.μ) (i : F.Index) : ℂ :=
  ∫ x, f x * L2normalizedHaar G F i x ∂G.grid.μ
```

## 16. ALTERNATIVE NORMS: STANDARD, HAAR, AND MEAN OSCILLATION

**File:** `GoodGrid/AlternativeRepresentationsAndNorms/` (eleven files, see Section 24)

**Imports (within the project):** `GoodGrid/BesovSpace.lean`, `WeakGrid/InducedGrid.lean`, `HaarRepresentationNorm.lean`

This is the paper’s “Alternative characterizations, I: messing with norms” (Section 15 of the published version). The Besov norm is an infimum over all representations, which makes it hard to compute for a concrete  $f$ ; this section provides three norms defined directly from  $f$  and proves them equivalent to it. The *Haar norm*  $N_{haar}$  weighs the Haar coefficients  $d_S^f = \int f \phi_S dm$ ; the *standard norm*  $N_{st}$  weighs the coefficients of one distinguished atomic representation, obtained by resumming the Haar expansion cellwise (each  $k_P^f$  is a finite, geometry-bounded combination of the  $d_S^f$  with  $S$  in the Haar tree of the parent of  $P$ ); and the *mean-oscillation norm*  $N_{osc}$  measures, in the spirit

of Dorronsoro, how well  $f$  is approximated by constants at every cell and scale. The equivalences are proved as a cycle  $|\cdot|_{\mathcal{B}_{p,q}^s} \rightarrow N_{st} \rightarrow N_{haar} \rightarrow N_{osc} \rightarrow |\cdot|_{\mathcal{B}_{p,q}^s}$  (inequalities (15-30)–(15-33) of the paper), each step with its own mechanism: testing  $f - c_Q$  against the zero-mean wavelets bounds Haar coefficients by oscillations; the constancy of low-frequency Souza blocks on finer cells means oscillation only sees the high-frequency tail of a representation, which a discrete convolution estimate sums; and the standard representation, being an actual representation, dominates the infimum norm. In particular the standard representation converges to  $f$  for *every*  $f$  in the space — a canonical, linear-in- $f$  decomposition that substitutes for the nonuniqueness of general representations.

For  $f \in L^\beta$ ,  $\beta > 1$ , write  $d_S^f = \int f \phi_S dm$  for the Haar coefficients of  $f$ , and define

$$N_{haar}(f) = |I|^{1/p-s-1/2} |d_I^f| + \left( \sum_k \left( \sum_{Q \in \mathcal{P}^k} |Q|^{1-sp-\frac{p}{2}} \sum_{S \in \mathcal{H}_Q} |d_S^f|^p \right)^{q/p} \right)^{1/q}.$$

From the Haar coefficients one builds, exactly as in the paper, the **standard atomic representation**

$$f = \sum_i \sum_{Q \in \mathcal{P}^i} k_Q^f a_Q$$

( $a_Q$  canonical Souza's atoms) and its norm

$$N_{st}(f) = |k_I^f| + \left( \sum_{k \geq 1} \left( \sum_{Q \in \mathcal{P}^k} |k_Q^f|^p \right)^{q/p} \right)^{1/q}.$$

Finally, with

$$osc_p(f, Q) = \inf_{c \in \mathbb{C}} \left( \int_Q |f - c|^p dm \right)^{1/p},$$

define

$$N_{osc}(f) = |I|^{-s} |f|_p + \left( \sum_k \left( \sum_{Q \in \mathcal{P}^k} |Q|^{-sp} osc_p(f, Q)^p \right)^{q/p} \right)^{1/q}.$$

**Theorem 16.1** (Equivalence of norms). *Suppose  $0 < s < 1/p$ ,  $p \in [1, \infty)$  and  $q \in [1, \infty]$ . Each one of the quantities  $|f|_{\mathcal{B}_{p,q}^s}$ ,  $N_{st}(f)$ ,  $N_{haar}(f)$ ,  $N_{osc}(f)$  is finite if and only if  $f \in \mathcal{B}_{p,q}^s$ , and these norms are equivalent on  $\mathcal{B}_{p,q}^s$ . More precisely there are constants, depending only on the grid geometry and on  $s, p, q$ , such that*

$$\begin{aligned} N_{st}(f) &\leq C |f|_{\mathcal{B}_{p,q}^s}, & N_{st}(f) &\leq C N_{haar}(f), \\ N_{haar}(f) &\leq C N_{osc}(f), & N_{osc}(f) &\leq C |f|_{\mathcal{B}_{p,q}^s}, \end{aligned}$$

and conversely: if  $N_{st}(f) < \infty$  then  $f \in \mathcal{B}_{p,q}^s$  and the standard atomic representation is a  $\mathcal{B}_{p,q}^s$ -representation of  $f$  (so  $|f|_{\mathcal{B}_{p,q}^s} \leq N_{st}(f)$ ); if  $N_{haar}(f) < \infty$  and  $f \in L^1$  then  $f \in L^p$  and the Haar series of  $f$  converges in  $L^p$ .

**Formalization.** In the files of `GoodGrid/AlternativeRepresentationsAndNorms/`. The equivalences are proved as a cycle of separate inequalities, each with a finite existential constant: `exists_standardRepresentationNorm_le_const_mul_souzaBesovNorm` ( $N_{st} \leq C \cdot |\mathcal{B}_{p,q}^s|$ ), `exists_standardRepresentationNorm_le_const_mul_haarL2RepresentationNorm` (the Haar gauge controls the standard atomic gauge), `exists_haarL2RepresentationNorm_le_const_mul_meanOscillationNorm` (mean oscillation controls the Haar gauge; the cellwise estimate is the Lean analogue of testing  $f - c_Q$  against the zero-mean Haar wavelets over a parent cell) and `exists_meanOscillationNorm_le_const_mul_souzaBesovNorm` (the Besov norm controls mean oscillation; the proof separates the  $L^p$  term from the oscillation seminorm and uses that low-frequency Souza blocks are constant on finer cells, so local oscillation is controlled by the high-frequency tail and a geometric discrete convolution estimate). The converse endpoints are `finite_standardRepresentationNorm_implies_memBesov_and_standardRepresentation` (finite  $N_{st}$  gives  $L^p$  membership, the canonical standard representation as a genuine finite-cost  $\mathcal{B}_{p,q}^s$ -representation, and Souza-Besov membership with the expected cost bounds) and `finite_haarL2RepresentationNorm_implies_memLp_and_hasSum` (finite  $N_{haar}$  gives  $f \in L^p$  and convergence of the normalized Haar expansion to  $f$  in  $L^p$ ); the wrap-up of the Haar-side comparisons is `exists_haarRepresentationNorms_equivalent`. Compared with paper Theorem 15.1: the formalization proves the equivalences as a cycle of separate inequalities with existential constants (the explicit constants  $\theta_1, \theta_2$  of the paper are not tracked), it works with the  $L^2$ -normalized Haar coefficients, and there is no single wrap-up statement packaging all four norms at once.

```

theorem exists_standardRepresentationNorm_le_const_mul_haarL2RepresentationNorm
  (G : GoodGridSpace (α := α)) [DecidableEq (Set α)]
  (F : UnbalancedHaarWavelet.FullHaarSystem (G := HaarRepresentation.GridOf G))
  [DecidableEq F.Index]
  (s : ℝ) (hs : 0 < s)
  (p : ℝ≥0∞) [Fact (1 ≤ p)] (hp_top : p < ∞)
  (q : ℝ≥0∞) [Fact (1 ≤ q)] :
  ∃ C : ℝ≥0∞, C ≠ ∞ ∧
    ∀ (f : α → ℂ) (hfint : Integrable f G.grid.μ),
      HaarRepresentation.haarL2RepresentationNorm G F s p q f hfint ≠ ∞ →
        standardRepresentationNorm G F s hs p hp_top q f hfint ≠ ∞ ∧
          standardRepresentationNorm G F s hs p hp_top q f hfint ≤
            C * HaarRepresentation.haarL2RepresentationNorm G F s p q f hfint :=
    ...

theorem exists_haarL2RepresentationNorm_le_const_mul_meanOscillationNorm
  (G : GoodGridSpace (α := α)) [DecidableEq (Set α)]
  (F : UnbalancedHaarWavelet.FullHaarSystem (G := GridOf G))
  (s : ℝ) (p : ℝ≥0∞) [Fact (1 ≤ p)] (hp_top : p < ∞)
  (q : ℝ≥0∞) [Fact (1 ≤ q)] :
  ∃ C : ℝ≥0∞, C ≠ ∞ ∧
    ∀ (f : α → ℂ) (hf : Integrable f G.grid.μ), MemLp f p G.grid.μ →
      MeanOscillation.meanOscillationNorm G s p q f ≠ ∞ →
        haarL2RepresentationNorm G F s p q f hf ≤

```

```

C * MeanOscillation.meanOscillationNorm G s p q f := ...

theorem exists_meanOscillationNorm_le_const_mul_souzaBesovNorm
  (G : GoodGridSpace (α := α)) [DecidableEq (Set α)]
  (s : ℝ) (hs : 0 < s)
  (p : ℝ≥0∞) [Fact (1 ≤ p)] (hp_top : p < ∞)
  (q : ℝ≥0∞) [Fact (1 ≤ q)] :
  ∃ C : ℝ≥0∞, C ≠ ∞ ∧
    ∀ (g : SouzaBesovSpace G s p q hs Fact.out (ne_of_lt hp_top))
      (f : α → ℂ),
      f ==m [G.grid.μ] ((g : Lp ℂ p G.toWeakGridSpace.measure) : α → ℂ) →
        meanOscillationNorm G s p q f ≤
          C * ENNReal.ofReal
            (WeakGridSpace.BesovishSpace.Norm_Costpq
              (souzaAtomFamily G s p hs Fact.out (ne_of_lt hp_top)) q g) := ...

theorem
  finite_standardRepresentationNorm_implies_memBesov_and_standardRepresentation
  (G : GoodGridSpace (α := α)) [DecidableEq (Set α)]
  (F : UnbalancedHaarWavelet.FullHaarSystem (G := HaarRepresentation.GridOf G))
  [DecidableEq F.Index]
  (s : ℝ) (hs : 0 < s)
  (p : ℝ≥0∞) [Fact (1 ≤ p)] (hp_top : p < ∞)
  (q : ℝ≥0∞) [Fact (1 ≤ q)]
  (f : α → ℂ) (hf : Integrable f G.grid.μ)
  (hN : standardRepresentationNorm G F s hs p hp_top q f hf ≠ ∞) :
  ∃ hfLp : MemLp f p G.grid.μ,
    ∃ g : SouzaBesovSpace G s p q hs Fact.out (ne_of_lt hp_top),
    ∃ R :
      WeakGridSpace.LpGridRepresentation
        (souzaAtomFamily G s p hs Fact.out (ne_of_lt hp_top))
        (g : Lp ℂ p G.toWeakGridSpace.measure),
      (g : Lp ℂ p G.toWeakGridSpace.measure) = hfLp.toLp f ∧
      R.block = canonicalStandardBlockSeq G F s hs p hp_top f hf ∧
      WeakGridSpace.LpGridRepresentation.FinitePQCost (q := q) R ∧
      WeakGridSpace.LpGridRepresentation.pqCostENNReal (q := q) R =
        standardRepresentationNorm G F s hs p hp_top q f hf ∧
      WeakGridSpace.LpGridRepresentation.pqCost (q := q) R ≤
        (standardRepresentationNorm G F s hs p hp_top q f hf).toReal ∧
      WeakGridSpace.BesovishSpace.Norm_Costpq
        (souzaAtomFamily G s p hs Fact.out (ne_of_lt hp_top)) q g
    ≤
      (standardRepresentationNorm G F s hs p hp_top q f hf).toReal
  := ...

```

**Corollary 16.2** (Reading the standard coefficients). *For each  $P \in \mathcal{P}$  the coefficient  $k_P^f$  of the standard atomic representation of  $f$  is a (finite) linear combination of the Haar coefficients  $d_S^f$ ,  $S \in \mathcal{H}_Q$ , of  $f$ , and the standard representation of any  $f \in \mathcal{B}_{p,q}^s$  converges to  $f$  in  $L^p$  with*

$$\left( \sum_k \left( \sum_{P \in \mathcal{P}^k} |k_P^f|^p \right)^{q/p} \right)^{1/q} \leq C \|f\|_{\mathcal{B}_{p,q}^s}.$$

**Formalization.** In `GoodGrid/AlternativeRepresentationsAndNorms/HaarRepresentationNorm.lean`, together with the theorems of the previous *Formalization* paragraph. The Haar coefficients are the definition `Coeff`, and the theorem `hasSum_coeff_smul_l2normalizedHaar_toLp` proves that the normalized Haar expansion of any  $f \in L^\beta$ ,  $1 < \beta < \infty$ , converges to  $f$ . Paper Corollary 15.2 states moreover that  $f \mapsto k_P^f$  extends to a bounded linear functional on  $L^1$ ; this extension is not formalized.

```

theorem hasSum_coeff_smul_l2normalizedHaar_toLp
  (G : GoodGridSpace (α := α)) [DecidableEq (Set α)]
  (F : UnbalancedHaarWavelet.FullHaarSystem (G := GridOf G))
  [DecidableEq F.Index]
  (β : ℝ ≥ 0 ∞) (hβ_one : 1 < β) (hβ_top : β < ∞)
  (f : α → ℂ) (hf : MemLp f β G.grid.μ) :
  letI : Fact (1 ≤ β) := ⟨le_of_lt hβ_one⟩
  HasSum
    (fun i : F.Index =>
      Coeff G F f
      (by
        letI : IsFiniteMeasure G.grid.μ := (GridOf G).isFinite
        exact hf.integrable (le_of_lt hβ_one))
      i •
      (l2normalizedHaar_memLp G F β i).toLp (L2normalizedHaar G F i))
    (hf.toLp f) := ...

```

## 17. POSITIVE CONE

**File:** `GoodGrid/PositiveCone.lean`  
**Imports (within the project):** `GoodGrid/BesovSpace.lean`, `GoodGrid/AlternativeRepresentationsAndNorms/StandarRepresentationNormleqBesovNorm.lean`, `WeakGrid/Multipliers.lean`

Positivity is where Souza’s atoms shine: since canonical atoms are *positive multiples of indicators*, a representation with nonnegative coefficients is a sum of nonnegative functions, and no cancellation can occur. This rigidity has strong structural consequences, proved in this file. The support of a positive function is (mod  $m$ ) a countable union of cells, and — a sharpening obtained in the formalization — in *any* positive representation of a function supported in a cell, all active cells must sit inside that cell: a coefficient cannot be “wasted” outside the support, because nothing can cancel it. This support rigidity is the engine of the positive-cone non-Archimedean theorems of Section 19. The original motivation comes from dynamics: the fixed points of transfer operators of expanding maps — invariant densities — are nonnegative and often discontinuous, and the positive cone is their

natural home inside the Besov space. The file depends on the standard-representation comparison of Section 16 (hence its position in the import order): the decomposition  $f = f_+ - f_-$  is obtained by splitting the coefficients of the *standard* representation, whose linearity and canonicity make the positive parts well defined.

We say that  $f$  is  $\mathcal{B}_{p,q}^s$ -positive if there is a  $\mathcal{B}_{p,q}^s$ -representation

$$f = \sum_k \sum_{P \in \mathcal{P}^k} c_P a_P$$

where  $c_P \geq 0$  and  $a_P$  is the canonical  $(s, p)$ -Souza's atom supported on  $P$ . The set of all  $\mathcal{B}_{p,q}^s$ -positive functions is a convex cone in  $\mathcal{B}_{p,q}^s$ , denoted  $\mathcal{B}_{p,q}^{s+}$ , with the “norm”

$$|f|_{\mathcal{B}_{p,q}^{s+}} = \inf \left( \sum_k \left( \sum_{P \in \mathcal{P}^k} c_P^p \right)^{q/p} \right)^{1/q},$$

the infimum running over all positive representations of  $f$ .

**Formalization.** In `GoodGrid/PositiveCone.lean`. A positive representation (canonical atoms, nonnegative coefficients) is the predicate `SouzaPositiveRepresentation`, and the cone of  $\mathcal{B}_{p,q}^s$ -positive functions is `souzaPositiveCone`, with the positive “norm” `souzaPositiveNorm`. The formalization also uses the wider notion of a *cone-positive* representation (`SouzaConePositiveRepresentation`): real coefficients  $\geq 0$  and Souza's atoms which are a.e. nonnegative real-valued, not necessarily canonical. The file also provides the additive infrastructure used in the non-Archimedean theorems of Section 19: sums and scalar multiples of positive representations (`souzaPositiveRepresentationAdd*`, `exists_souzaPositiveRepresentation_finset_sum`) with levelwise Minkowski control of the cost (`souzaPositiveRepresentationAdd_levelCoeffRoot_le`).

```
def SouzaPositiveRepresentation
  (G : GoodGridSpace (α := α)) (s : ℝ) (p : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)]
  {f : Lp ℂ p G.toWeakGridSpace.measure}
  (R : WeakGridSpace.LpGridRepresentation
    (souzaAtomFamily G s p hs hp hp_top) f) : Prop :=
  ∀ k, SouzaPositiveLevelBlock G s p hs hp hp_top (R.block k)

def SouzaConePositiveRepresentation
  (G : GoodGridSpace (α := α)) (s : ℝ) (p : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)]
  {f : Lp ℂ p G.toWeakGridSpace.measure}
  (R : WeakGridSpace.LpGridRepresentation
    (souzaAtomFamily G s p hs hp hp_top) f) : Prop :=
```

$\forall k, \text{SouzaConePositiveLevelBlock } G \text{ s p hs hp hp\_top } (R.\text{block } k)$

**Proposition 17.1.** *Let  $0 < s < 1/p$ .*

- A. *If  $f \in \mathcal{B}_{p,q}^s$  is (a.e.) real valued then one can find  $f_+, f_- \in \mathcal{B}_{p,q}^{s+}$  such that*

$$f = f_+ - f_-,$$

*with the positive costs of  $f_+$  and  $f_-$  controlled in terms of  $|f|_{\mathcal{B}_{p,q}^s}$ ; an analogous four-term decomposition holds for complex valued  $f$ .*

- B. *The positive cone  $\mathcal{B}_{p,q}^{s+}$  is dense in the cone of nonnegative functions of  $L^p$ .*

**Formalization.** In `GoodGrid/PositiveCone.lean`. Item A is `exists_souzaPositive_decomposition_of_aeRealValued` (the decomposition  $f = f_+ - f_-$  with controlled positive costs, for a.e. real valued  $f$ ) and `exists_souzaPositive_decomposition_of_aeComplexValued` (the four-term decomposition for complex valued  $f$ ); in the paper this appears as an unnumbered observation in Section 13, and the complex case is not stated there. Item B is `souzaPositiveCone_dense_in_LpNonnegativeCone`, density of the positive cone in the nonnegative cone of  $L^p$ ; not stated in the paper.

```

theorem exists_souzaPositive_decomposition_of_aeRealValued
  (G : GoodGridSpace (α := α)) (s : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)] :
  ∃ C : ℝ,
    0 ≤ C ∧
    ∀ f : WeakGridSpace.BesovishSpace
      (souzaAtomFamily G s p hs hp hp_top) q,
      SouzaBesovAEEqRealValued G s p q hs hp hp_top f →
      ∃ u v : WeakGridSpace.BesovishSpace
        (souzaAtomFamily G s p hs hp hp_top) q,
        SouzaPositiveElement G s p q hs hp hp_top u ∧
        SouzaPositiveElement G s p q hs hp hp_top v ∧
        f = u - v ∧
        souzaPositiveNorm G s p q hs hp hp_top u ≤
          ENNReal.ofReal C *
            ENNReal.ofReal (WeakGridSpace.BesovishSpace.Norm_Costpq
              (souzaAtomFamily G s p hs hp hp_top) q f) ∧
        souzaPositiveNorm G s p q hs hp hp_top v ≤
          ENNReal.ofReal C *
            ENNReal.ofReal (WeakGridSpace.BesovishSpace.Norm_Costpq
              (souzaAtomFamily G s p hs hp hp_top) q f) := ...

theorem souzaPositiveCone_dense_in_LpNonnegativeCone
  (G : GoodGridSpace (α := α)) (s : ℝ) (β q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hβ : 1 ≤ β) (hβ_top : β ≠ ∞)
  [Fact (1 ≤ β)] [Fact (1 ≤ q)] :
  LpNonnegativeCone G β ⊆
    closure (SouzaPositiveConeInLbeta G s β q hs hβ hβ_top) := ...

```

**Proposition 17.2** (Support of positive functions). *If  $f \in \mathcal{B}_{p,q}^{s+}$  then its support*

$$\text{supp } f = \{x \in I : f(x) \neq 0\}$$

*is, up to a set of zero measure, a countable union of elements of  $\mathcal{P}$ . Moreover, in any positive representation of  $f$ , the coefficient of a cell  $Q$  vanishes unless  $Q$  is contained (mod  $m$ ) in this union.*

**Formalization.** In `GoodGrid/PositiveCone.lean` (paper Proposition 13.1). The first sentence is `support_ae_countable_iUnion_goodGridCells_of_souzaPositiveFunction`, which exhibits the support of a positive function as an a.e. countable union of good-grid cells. The second sentence is the support theorem `souzaPositiveRepresentation_coeff_eq_zero_of_not_subset_cell`, a sharper statement obtained in the formalization: in any positive representation of a function supported in a cell, every active cell is contained in that cell. A companion “linchpin” lemma, `souzaPositiveRepresentation_ae_ne_zero_on_active_cell`, proves that a positively represented function is a.e. nonzero on every active cell of its representation; both are used repeatedly in the positive-cone non-Archimedean arguments of Section 19.

```

theorem support_ae_countable_iUnion_goodGridCells_of_souzaPositiveFunction
  (G : GoodGridSpace (α := α)) (s : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)]
  {f : α → ℂ}
  (hf : SouzaPositiveFunction G s p q hs hp hp_top f) :
  IsAECountableUnionOfGoodGridCells G {x | f x ≠ 0} := ...

theorem souzaPositiveRepresentation_coeff_eq_zero_of_not_subset_cell
  (G : GoodGridSpace (α := α)) (s : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)]
  (Qc : GoodGridCell G)
  {g : Lp ℂ p G.toWeakGridSpace.measure}
  (R : WeakGridSpace.LpGridRepresentation (souzaAtomFamily G s p hs hp
    hp_top) g)
  (hR : SouzaPositiveRepresentation G s p hs hp hp_top R)
  (hsupp : ∀m x ∂ G.toWeakGridSpace.measure, x ∉ Qc.cell → (g : α → ℂ)
    x = 0)
  {n : ℕ} (P : WeakGridSpace.LevelCell G.toWeakGridSpace n)
  (hP : ¬ P.1 ⊆ Qc.cell) :
  (R.block n).coeff P = 0 := ...

```

## 18. DIRAC'S APPROXIMATIONS

**File:** GoodGrid/DiracApproximations.lean  
**Imports (within the project):** GoodGrid/  
 AlternativeRepresentationsAndNorms/standardRepresentation.lean,  
 GoodGrid/PositiveCone.lean

This section documents the formalization of Section 17 of the published paper. For a cell  $Q \in \mathcal{P}^{k_0}$  the *Dirac kernel*  $\mathbb{1}_Q/m(Q)$  is the approximation of the identity at scale  $k_0$  attached to the grid. The theme of the section is that the partial sums of the standard representation of an integrable function  $f$  (Section 16) evaluate, at any point  $x_0 \in Q$ , to the average of  $f$  against this kernel. Two consequences follow at once: the partial sums are bounded by the local  $L^\infty$  norm of  $f$ , and — since at a point  $x_0$  only the unique ancestor cell at each level contributes — the *ancestor coefficient sums* of the standard representation are controlled by  $|f|_\infty$ . This is exactly the input that the proof of Pointwise Multipliers II (Section 19) needs from this section. The proof goes through the unbalanced Haar system: a telescoping identity expands the Dirac kernel in the Haar functions supported on the ancestors of  $Q$ , the evaluation identity follows by strong induction on the level, and the regrouping of the Haar expansion into the standard representation (Section 16) transports the result to the standard partial sums.

**Proposition 18.1.** *Let  $f \in L^1$ . Fix  $k_0 \in \mathbb{N}$ , a point  $x_0$ , and let  $Q \in \mathcal{P}^{k_0}$  be the cell containing  $x_0$ . Write  $f_{k_0}$  for the partial sum, up to scale  $k_0$ , of the standard representation of  $f$  (the Haar partial sum regrouped on the cells, Section 16). Then:*

- A.  $|f_{k_0}(x_0)| \leq |f|_\infty$ . Moreover, if  $f = \sum_k \sum_{P \in \mathcal{P}^k} c_P \chi_P$  is any positive  $(s, p)$ -Souza representation of  $f$ , then

$$\sum_{J \supseteq Q} c_J m(J)^{s-1/p} \leq |f|_\infty,$$

the sum running over the ancestors  $J$  of  $Q$  of level at most  $k_0$ .

- B.  $f_{k_0}(x_0) = \frac{1}{m(Q)} \int_Q f \, dm$ .

**Formalization.** In GoodGrid/DiracApproximations.lean (paper Proposition 17.1), namespace DiracApproximation. The Dirac kernel is `diracKernel`, and the telescoping expansion of the kernel in the Haar system is `diracKernel.eq_alpha_add_sumDown`, transported from `NormalizedIndicator.in_Haar_Wavelets.span` of the dependency `UnbalancedHaarWavelet`. The partial sums are `partialHaarSum` and `partialStandardSum`, identified pointwise by `partialHaarSum.eq_partialStandardSum`. The evaluation identity  $f_{k_0}(x_0) = \int f \mathbb{1}_Q/m(Q) \, dm$  is `partialHaarSum.eq_integral_mul_diracKernel` (proved by strong induction on the level), and item B is its standard-representation form `claimB`.

The first bound of item A is `claimA_standard`; the positive-representation bound is `claimA_positive`, where the ancestor sum is expressed through `ancestorTerm` (the contribution of the unique level- $k$  ancestor, with the canonical normalization  $m(J)^{s-1/p}$ ), and the proof goes through a.e. convergence of a subsequence of the partial sums together with their monotonicity. The levelwise collapse of the standard blocks at a point to the single ancestor contribution is `standardLevelSum_eq_ancestor_term`. Compared with the paper: the partial sums of the standard representation use the *tilde* atoms of Section 16, whose value at  $x_0$  has modulus at most  $m(J)^{s-1/p}$  but need not equal it; the exact form with  $m(J)^{s-1/p}$ , as in item A, holds for canonical atoms (and is the form used in Section 19). Axioms checked: `propext`, `Classical.choice`, `Quot.sound`.

```

theorem claimA_standard (G : GoodGridSpace (α := α))
  [DecidableEq (Set α)]
  (F : UnbalancedHaarWavelet.FullHaarSystem (G :=
    HaarRepresentation.GridOf G))
  (f : α → ℂ) (hf : Integrable f G.grid.μ)
  (Q : GoodGridCell G) {x₀ : α} (hx₀ : x₀ ∈ Q.cell) :
  ENNReal.ofReal ||partialHaarSum G F f hf Q.level x₀|| ≤
    eLpNorm (Set.indicator Q.cell f) ∞ G.grid.μ := ...

theorem claimA_positive (G : GoodGridSpace (α := α))
  (s : ℝ) (p : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞) [Fact (1 ≤ p)]
  {f : Lp ℂ p G.toWeakGridSpace.measure}
  (R : WeakGridSpace.LpGridRepresentation (souzaAtomFamily G s p hs hp
    hp_top) f)
  (hR : SouzaPositiveRepresentation G s p hs hp hp_top R)
  (Q : GoodGridCell G) :
  ENNReal.ofReal (∑ k ∈ Finset.range (Q.level + 1),
    ancestorTerm G s p hs hp hp_top R Q k) ≤
    eLpNorm (Set.indicator Q.cell (↑f)) ∞ G.toWeakGridSpace.measure :=
    ...

theorem claimB (G : GoodGridSpace (α := α)) [DecidableEq (Set α)]
  (F : UnbalancedHaarWavelet.FullHaarSystem (G :=
    HaarRepresentation.GridOf G))
  (p : ℝ ≥ 0 ∞) (s : ℝ) (f : α → ℂ) (hf : Integrable f G.grid.μ)
  (Q : GoodGridCell G) {x₀ : α} (hx₀ : x₀ ∈ Q.cell) :
  partialStandardSum G F p s f hf Q.level x₀ =
    ∫ x, f x * ((diracKernel G Q x : ℝ) : ℂ) ∂G.grid.μ := ...

```

## 19. POINTWISE MULTIPLIERS ACTING ON $\mathcal{B}_{p,q}^s$

This section documents the directory `GoodGrid/Multipliers/` (eight files plus the standalone interface), which formalizes Section 18 of the published paper. From here on  $0 < s < 1/p$ . The guiding question is: for which  $g$  is

$f \mapsto gf$  bounded on  $\mathcal{B}_{p,q}^s$ ? Because the space is built from Souza's atoms, and the product of  $g$  with a Souza's atom is just  $g$  restricted to a cell and rescaled, the natural test is to multiply  $g$  against every atom — this gives the *selfs* classes, and the theory consists in showing that this obviously necessary condition is, in various quantified senses, also sufficient. The subsections below follow the import order of the files.

### 19.1. The selfs classes and their seminorms.

**File:** GoodGrid/Multipliers/Definition.lean

**Imports (within the project):** WeakGrid/Multipliers.lean, GoodGrid/BesovSpace.lean

Let  $g: I \rightarrow \mathbb{C}$  be a measurable function. We say that  $g$  is a **pointwise multiplier** acting on  $\mathcal{B}_{p,q}^s$  if the transformation  $G(f) = gf$  defines a bounded operator in  $\mathcal{B}_{p,q}^s$ . We denote the set of pointwise multipliers by  $M(\mathcal{B}_{p,q}^s)$ , with the norm

$$|g|_{M(\mathcal{B}_{p,q}^s)} = \sup\{|gf|_{\mathcal{B}_{p,q}^s} : |f|_{\mathcal{B}_{p,q}^s} \leq 1\}.$$

Define

$$\mathcal{B}_{p,q,\text{selfs}}^s = \left\{ g \in \mathcal{B}_{p,q}^s : \sup_{a_Q \in \mathcal{A}_{s,p}^{sz}} |ga_Q|_{\mathcal{B}_{p,q}^s} < \infty \right\},$$

with the norm

$$|g|_{\mathcal{B}_{p,q,\text{selfs}}^s} = \sup_{a_Q \in \mathcal{A}_{s,p}^{sz}} |ga_Q|_{\mathcal{B}_{p,q}^s},$$

and, for  $t \in \mathbb{N}$ , the tail seminorms

$$|g|_{\mathcal{B}_{p,q,\text{selfs}}^{s,t}} = \sup_{\substack{a_Q \in \mathcal{A}_{s,p}^{sz} \\ Q \in \mathcal{P}^j, j \geq t}} |ga_Q|_{\mathcal{B}_{p,q}^s}.$$

**Formalization.** In GoodGrid/Multipliers/Definition.lean. Pointwise multipliers are the predicate `SouzaPointwiseMultiplier`, the class  $\mathcal{B}_{p,q,\text{selfs}}^s$  is `SouzaPointwiseSelfsClass`, and the tail variant is `SouzaPointwiseSelfsTailClass`: the quantitative form `SouzaPointwiseSelfsTailBound`  $g$   $C$  records that for every Souza's atom on a cell of level at least  $t$  the product  $ga_Q$  admits a representation of cost at most  $C$ , and the induced seminorm  $|\cdot|_{\mathcal{B}_{p,q,\text{selfs}}^{s,t}}$  is `souzaPointwiseSelfsTailNorm`.

```
abbrev SouzaPointwiseMultiplier
  (G : GoodGridSpace (α := α)) (s : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)]
  (m : α → ℂ) : Prop :=
  WeakGridSpace.IsPointwiseMultiplier
  (A := souzaAtomFamily G s p hs hp hp_top) q m

abbrev SouzaPointwiseSelfsClass
  (G : GoodGridSpace (α := α)) (s : ℝ) (p q : ℝ ≥ 0 ∞)
```

```

    (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
    [Fact (1 ≤ p)] [Fact (1 ≤ q)]
    (m : α → ℂ) : Prop :=
    WeakGridSpace.PointwiseSelfsClass
    (A := souzaAtomFamily G s p hs hp hp_top) q m

```

## 19.2. Restriction to a cell.

**File:** GoodGrid/Multipliers/Besovspq.lean

**Imports (within the project):** GoodGrid/Multipliers/Definition.lean,  
WeakGrid/Transmutation.lean

The first multiplier theorem is for the simplest candidates: indicators of cells. The proof is another instance of transmutation — restricting a Souza’s atom  $a_Q$  to a cell  $W$  produces either 0, the atom itself, or (when  $W \subset Q$ ) a rescaled atom on  $W$ , with geometric decay in the level gap; Proposition 8.1(A) and (B) of the paper then assemble these local expansions with uniform cost. The lemma is stated on the induced grid of  $W$ , and the uniformity of  $C_{22}$  in  $W$  is what the non-Archimedean arguments below need.

**Lemma 19.1** (Restriction to a cell). *Let  $W \in \mathcal{P}$ . The restriction map*

$$r : \mathcal{B}_{p,q}^s(I, \mathcal{P}, \mathcal{A}_{s,p}^{sz}) \rightarrow \mathcal{B}_{p,q}^s(W, \mathcal{P}_W, \mathcal{A}_{s,p}^{sz}), \quad r(f) = 1_W f,$$

*is continuous: there is  $C_{10} \geq 1$ , independent of  $W$ , such that*

A. *For every  $f \in \mathcal{B}_{p,q}^s$  we have*

$$|1_W f|_{\mathcal{B}_{p,q}^s(W, \mathcal{P}_W, \mathcal{A}_{s,p}^{sz})} \leq C_{10} |f|_{\mathcal{B}_{p,q}^s}.$$

*In particular  $1_W \in M(\mathcal{B}_{p,q}^s)$  with  $|1_W|_{M(\mathcal{B}_{p,q}^s)} \leq C_{10}$ , uniformly in  $W$ .*

B. *For every  $\mathcal{B}_{p,q}^{s+}$ -representation  $f = \sum_k \sum_Q c_Q a_Q$  one can find a positive representation*

$$1_W f = \sum_k \sum_Q d_Q a_Q$$

*on the induced grid with cost at most  $C_{10}$  times the cost of the original one, and such that  $d_Q \neq 0$  implies  $Q \subset \text{supp } f \pmod{m}$ .*

**Formalization.** In GoodGrid/Multipliers/Besovspq.lean (paper Lemma 18.2). The restriction map with its induced-grid representation is souzaCellIndicatorRestrictsToInduced; the cost bounds (uniform in  $W$ ) are the lemmas souzaIndicatorRestrictionBound\_of\_ambientRestrictionTransmutation\_\* and souzaCellIndicatorPointwiseMultiplierBound.uniform, the latter giving the explicit multiplier bound  $2C + 1$  with  $C = \text{souzaAmbientRestrictionMultiplierConstant}$ ; and the theorem souzaCellIndicatorPointwiseMultiplier concludes that  $1_W$  is a point-wise multiplier of  $\mathcal{B}_{p,q}^s$ . The paper’s strongly regular domains (Definition 18.7 and Proposition 18.8), of which the present lemma is the grid-cell case, are not

formalized in general; the multiplier statement for  $1_W$ ,  $W \in \mathcal{P}$ , with a bound uniform in  $W$ , is the formalized special case.

```

theorem souzaCellIndicatorPointwiseMultiplier
  (G : GoodGridSpace (α := α)) (s : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)]
  (hs_lt_inv : s < (p.toReal)-1)
  (W : GoodGridCell G) :
  SouzaPointwiseMultiplier G s p q hs hp hp_top
    (W.cell.indicator fun _ => (1 : ℂ)) := ...

theorem souzaCellIndicatorPointwiseMultiplierBound_uniform
  (G : GoodGridSpace (α := α)) (s : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)]
  (hs_lt_inv : s < (p.toReal)-1)
  (W : GoodGridCell G) :
  SouzaPointwiseMultiplierBound G s p q hs hp hp_top
    (W.cell.indicator fun _ => (1 : ℂ))
    (2 * souzaAmbientRestrictionMultiplierConstant G s p + 1) := ...

```

### 19.3. The multipliers of $\mathcal{B}_{1,1}^s$ .

**File:** GoodGrid/Multipliers/Besovs11.lean

**Imports (within the project):** GoodGrid/Multipliers/Definition.lean

For  $p = q = 1$  the multiplier problem has a complete and clean answer: the necessary condition is sufficient. The reason is that for an  $\ell^1$ -type cost one can substitute the representation of  $f$  into  $gf$  atom by atom,  $gf = \sum c_Q(ga_Q)$ , and the costs of the pieces  $ga_Q$  simply add up — no interaction between cells survives. This is paper Proposition 18.1.

**Proposition 19.2.** *We have that*

$$M(\mathcal{B}_{1,1}^s) = \mathcal{B}_{1,1, selfs}^s.$$

**Formalization.** This is paper Proposition 18.1. The equivalence is the theorem `souzaPointwiseMultiplier_iff_souzaPointwiseSelfsClass_one_one`: the inclusion  $\mathcal{B}_{1,1, selfs}^s \subset M(\mathcal{B}_{1,1}^s)$  is proved in `GoodGrid/Multipliers/Besovs11.lean`, while the converse, which holds for all  $p, q$ , is `souzaPointwiseSelfsClass_of_souzaPointwiseMultiplier` in `GoodGrid/Multipliers/Besovspq.lean`.

```

theorem souzaPointwiseMultiplier_iff_souzaPointwiseSelfsClass_one_one
  (G : GoodGridSpace (α := α)) (s : ℝ)
  (hs : 0 < s) {m : α → ℂ} :
  SouzaPointwiseMultiplier G s (1 : ℝ ≥ 0 ∞) (1 : ℝ ≥ 0 ∞)
    hs le_rfl ENNReal.one_ne_top m ↔

```

```
SouzaPointwiseSelfsClass G s (1 : ℝ≥0∞) (1 : ℝ≥0∞)
  hs le_rfl ENNReal.one_ne_top m := ...
```

#### 19.4. *selfs* multipliers are bounded.

**File:** GoodGrid/Multipliers/MultipliersareBounded.lean  
**Imports (within the project):** GoodGrid/BesovAtoms.lean, GoodGrid/Multipliers/Definition.lean, GoodGrid/Multipliers/Besovsq.lean

A function in a *selfs* class is automatically bounded. The proof is a pretty martingale argument (paper Proposition 18.3): testing  $g$  against the canonical atom of a cell  $Q$  and applying the  $L^p$  embedding of Proposition 5.1 to the product bounds the  $L^p$  average of  $g$  over  $Q$ , uniformly over all cells; since the cells generate the  $\sigma$ -algebra  $(G_8)$ , Lévy's upward theorem lets these averages converge to  $|g(x)|$  at almost every point, giving the sup bound. The formalization sharpens the paper in a direction the non-Archimedean theorems require: the constant is *independent of the tail level  $t$* , so that multipliers known only on deep cells are still uniformly bounded.

**Proposition 19.3** (*selfs* classes are bounded). *There is  $C_{11} = C_{11}(\lambda_1, \lambda_2, C_1, s, p, q)$  such that for every  $t \in \mathbb{N}$  and every  $g$  with  $|g|_{\mathcal{B}_{p,q,\text{selfs}}^{s,t}} < \infty$ ,*

$$|g(x)| \leq C_{11} |g|_{\mathcal{B}_{p,q,\text{selfs}}^{s,t}} \quad \text{for } m\text{-a.e. } x.$$

*In particular  $\mathcal{B}_{p,q,\text{selfs}}^s \subset L^\infty$  and this inclusion is continuous.*

**Formalization.** In GoodGrid/Multipliers/MultipliersareBounded.lean. The pointwise estimate from a single tail bound is `souzaPointwiseSelfsTailBound_norm_ae_le`; the estimate in terms of the tail seminorm, with the explicit constant displayed in the code below, is `souzaPointwiseSelfsTailNorm_norm_ae_le`; and the qualitative  $L^\infty$  membership of the whole class is `souzaPointwiseSelfsTailClass_norm_ae_bounded`. This sharpens paper Proposition 18.3 in two respects: the estimate is proved for the tail seminorms  $|\cdot|_{\mathcal{B}_{p,q,\text{selfs}}^{s,t}}$  with a constant *independent of  $t$*  (this uniformity is what the non-Archimedean arguments below require), and the formalization also provides a *canonical* variant (`SouzaPointwiseCanonicalSelfsTailBound`), in which  $g$  is only tested against products with canonical Souza's atoms, together with the bridges `Bound.toCanonical` and `SouzaPositivePointwiseSelfsTailBound.toCanonical`; the canonical variant is what the infinite positive-cone theorem below consumes.

```
theorem souzaPointwiseSelfsTailNorm_norm_ae_le
  (G : GoodGridSpace (α := α))
  (s : ℝ) (p q : ℝ≥0∞)
  (hs : 0 < s) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)]
  (hs_lt_inv : s < (p.toReal)-1)
  {t : ℕ} {m : α → ℂ}
```

```

(hm : SouzaPointwiseSelfsTailClass G s p q hs hp hp_top t m) :
  ∀m x ∂G.grid.μ,
    ||m x|| ≤
      souzaBesovLpLocalEmbeddingConstant G s p q *
      (2 * souzaAmbientRestrictionMultiplierConstant G s p + 1) *
      souzaPointwiseSelfsTailNorm G s p q hs hp hp_top t m := ...

```

### 19.5. Non-Archimedean behaviour in $\mathcal{B}_{p,\tilde{q},selfs}^\beta$ .

**File:** GoodGrid/Multipliers/NonArchimedeanProperty.lean (statements exposed by NonArchimedeanPropertyPositiveStandalone.lean)  
**Imports (within the project):** GoodGrid/Multipliers/Definition.lean, GoodGrid/BesovAtoms.lean, GoodGrid/Multipliers/Besovspq.lean, GoodGrid/Multipliers/MultipliersareBounded.lean, GoodGrid/PositiveCone.lean, WeakGrid/Scales.lean

These are the deepest theorems of the repository (paper Proposition 18.4 and Remark 18.5). The phenomenon: if the supports of the multipliers  $g_i$  are *separated by the grid* — every cell carrying a nonzero coefficient of  $f$  meets essentially one support, which is what hypothesis A quantifies — then each atom of the representation of  $f$  “sees” only one  $g_i$ , and the cost of  $(\sum_i g_i)f$  is governed by the *largest* relevant multiplier norm instead of their sum: the estimate behaves like an ultrametric bound, whence *non-Archimedean*. The level hypothesis B makes the separation scale-respecting: an atom may only meet a multiplier whose tail level has already been reached. The proof multiplies the representation of  $f$  atomwise, expands each product  $g_i a_Q$  by the restriction lemma on the induced grid of  $Q$ , transmutes the resulting  $\beta$ -smooth representations down to smoothness  $s$  (transmutation again — here `WeakGrid/Scales.lean` provides the  $\beta \rightarrow s$  rescaling), and reassembles; the bounded overlap of the supports is exactly what keeps the reassembled cost additive in the right way. The positive versions add the support and positivity conclusions of Remark 18.5, whose proofs rest on the support rigidity of Section 17; the infinite-index versions require, in addition, the pointwise theory: absolute convergence of  $\sum_i g_i(x)$  on  $\{f \neq 0\}$ , with the limit identified through the compactness of representations of Section 6.

If  $g_i \in M(\mathcal{B}_{p,q}^s)$ , the naive estimate is

$$\left| \left( \sum_i g_i \right) f \right|_{\mathcal{B}_{p,q}^s} \leq \left( \sum_i |g_i|_{M(\mathcal{B}_{p,q}^s)} \right) |f|_{\mathcal{B}_{p,q}^s}.$$

Remarkably, when the supports of the  $g_i$  are separated by the grid, one gets a far better estimate.

**Theorem 19.4** (Non-Archimedean property, finite version). *Let  $\beta > s$  and  $\tilde{q} \in [1, \infty]$ . There is  $C_{12} = C_{12}(\lambda_1, \lambda_2, C_1, s, \beta, p, q, \tilde{q})$  with the following*

property. Let  $\Lambda$  be finite,  $g_i$  with  $|g_i|_{\mathcal{B}_{p,\tilde{q},selfs}^{\beta,t_i}} < \infty$  for  $i \in \Lambda$  and  $t_i \in \mathbb{N}$ . Consider a function  $f$  with a  $\mathcal{B}_{p,q}^s$ -representation

$$f = \sum_k \sum_{Q \in \mathcal{P}^k} c_Q a_Q$$

by canonical Souza's atoms satisfying

A. We have

$$\sup_{\substack{Q \in \mathcal{P} \\ c_Q \neq 0}} \sum_{i: Q \cap \text{supp } g_i \neq \emptyset} |g_i|_{\mathcal{B}_{p,\tilde{q},selfs}^{\beta,t_i}} \leq N.$$

B. If  $Q \in \mathcal{P}^k$  satisfies  $c_Q \neq 0$  and  $Q \cap \text{supp } g_i \neq \emptyset$  then  $k \geq t_i$ .

Then  $(\sum_{i \in \Lambda} g_i)f \in \mathcal{B}_{p,q}^s$  and we can find a  $\mathcal{B}_{p,q}^s$ -representation

$$(7) \quad \left( \sum_{i \in \Lambda} g_i \right) f = \sum_k \sum_{Q \in \mathcal{P}^k} d_Q a_Q$$

by canonical Souza's atoms such that

$$(8) \quad \left( \sum_k \left( \sum_{Q \in \mathcal{P}^k} |d_Q|^p \right)^{q/p} \right)^{1/q} \leq C_{12} N \left( \sum_k \left( \sum_{Q \in \mathcal{P}^k} |c_Q|^p \right)^{q/p} \right)^{1/q}.$$

**Formalization.** The theorem `souzaNonArchimedeanPropertyLambdaFinite` in `GoodGrid/Multipliers/NonArchimedeanProperty.lean` (paper Proposition 18.4 for finite  $\Lambda$ ) produces, from the bounded-overlap hypothesis A and the level/scale hypothesis B, a representation of the product  $(\sum_i g_i)f$  of cost at most  $C_{12} N$  times the cost of the representation of  $f$ , with  $C_{12}$  independent of the multipliers, of  $N$ , and of  $\Lambda$ . The representations of  $f$  and of the product are by *canonical* Souza's atoms (`SouzaCanonicalRepresentation`), which is no loss of generality for Souza's atoms and is the precise form formalized.

```

theorem souzaNonArchimedeanPropertyLambdaFinite
  (G : GoodGridSpace (α := α))
  (s β : ℝ) (p q qtilde : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hβ : 0 < β) (hβs : s < β)
  (hβ_lt_inv : β < (p.toReal)⁻¹)
  (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)] [Fact (1 ≤ qtilde)] :
  ∃ Cgen : ℝ,
    0 ≤ Cgen ∧
    ∀ (Λ : Finset ℕ) (t : ℕ → ℕ) (g : ℕ → α → ℂ) (N : ℝ)
      (f : α → ℂ)
      (x : WeakGridSpace.BesovishSpace
        (souzaAtomFamily G s p hs hp hp_top) q)
      (R : WeakGridSpace.LpGridRepresentation
        (souzaAtomFamily G s p hs hp hp_top)
        (x : Lp ℂ p G.toWeakGridSpace.measure)),
      0 ≤ N →
      WeakGridSpace.RepresentsFunction
        (G := G.toWeakGridSpace) (p := p) f

```

```

(x : Lp C p G.toWeakGridSpace.measure) →
WeakGridSpace.LpGridRepresentation.FinitePQCost (q := q) R →
(∀ i ∈ Λ,
  ∃ C : ℝ,
    SouzaPointwiseSelfsTailBound
      G β p qtilde hβ hp hp_top (t i) (g i) C) →
(∀ k (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k),
  (R.block k).coeff Q ≠ 0 →
    nonArchimedeanRelevantTailSelfsSum
      G β p qtilde hβ hp hp_top ∧ t g Q ≤ N) →
(∀ k (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k) i,
  i ∈ Λ →
    (R.block k).coeff Q ≠ 0 →
      goodGridLevelCellMeetsSupport G Q (g i) →
        t i ≤ k) →
  ∃ y : WeakGridSpace.BesovishSpace
    (souzaAtomFamily G s p hs hp hp_top) q,
  ∃ S : WeakGridSpace.LpGridRepresentation
    (souzaAtomFamily G s p hs hp hp_top)
    (y : Lp C p G.toWeakGridSpace.measure),
  WeakGridSpace.RepresentsFunction
    (G := G.toWeakGridSpace) (p := p)
    (fun z => (∑ i ∈ Λ, g i z) * f z)
    (y : Lp C p G.toWeakGridSpace.measure) ∧
  WeakGridSpace.LpGridRepresentation.pqCost (q := q) S ≤
  Cgen * N *
  WeakGridSpace.LpGridRepresentation.pqCost (q := q) R := ...

```

**Theorem 19.5** (Non-Archimedean property, infinite version). *In the setting of Theorem 19.4 let  $\Lambda \subset \mathbb{N}$  be infinite, and replace hypothesis A. by the summability condition*

$$\sum_{i \in \Lambda} T_i \leq N, \quad T_i := \sup_{\substack{Q \in \mathcal{P}, c_Q \neq 0 \\ Q \cap \text{supp } g_i \neq \emptyset}} |g_i|_{\mathcal{B}_{p,q,\text{selfs}}^{\beta,t_i}},$$

*understood as a series with values in  $[0, \infty]$ . Then, for  $m$ -a.e.  $x$  in  $\{f \neq 0\}$ , the series  $\sum_{i \in \Lambda} g_i(x)$  converges absolutely, with*

$$\sum_{i \in \Lambda} |g_i(x)| \leq C_{12} N;$$

*the function  $h = (\sum_{i \in \Lambda} g_i)f$  (defined a.e. as this pointwise limit times  $f$ ) belongs to  $L^p$  and to  $\mathcal{B}_{p,q}^s$ , and it has a canonical representation (7) satisfying (8).*

**Formalization.** The theorem `souzaNonArchimedeanProperty` in `GoodGrid/Multipliers/NonArchimedeanProperty.lean` proves all the conclusions above: pointwise absolute summability of  $\sum_i g_i(x)$  with bound  $C_{12}N$  a.e. on  $\{f \neq 0\}$ , existence of the limit function  $h$  with  $\|h\| \leq C_{12}N\|f\|$  a.e., membership of  $h$  in  $L^p$ , and a canonical representation of  $h$  with the cost bound (8). The pointwise part is `exists_nonArchimedeanInfinite_pointwise_hasSum`; the identification of

the limit representation goes through the compactness of representations with uniform cost (`exists_strongly_convergent_subseq_of_uniform_pqCost`) applied to the truncations to finite initial segments of  $\Lambda$ . Paper Proposition 18.4 treats finite and infinite  $\Lambda$  together, reducing to the finite case; the formalization keeps them separate because the infinite case has genuinely more content.

```

theorem souzaNonArchimedeanProperty
  (G : GoodGridSpace ( $\alpha := \alpha$ ))
  (s  $\beta$  :  $\mathbb{R}$ ) (p q qtilde :  $\mathbb{R}_{\geq 0\infty}$ )
  (hs :  $0 < s$ ) (h $\beta$  :  $0 < \beta$ ) (h $\beta$ s :  $s < \beta$ )
  (h $\beta$ _lt_inv :  $\beta < (p.toReal)^{-1}$ )
  (hp :  $1 \leq p$ ) (hp_top :  $p \neq \infty$ )
  [Fact ( $1 \leq p$ )] [Fact ( $1 \leq q$ )] [Fact ( $1 \leq qtilde$ )] :
   $\exists$  Cgen :  $\mathbb{R}$ ,
     $0 \leq Cgen \wedge$ 
     $1 \leq Cgen \wedge$ 
     $\forall (\Lambda : \text{Set } \mathbb{N}) (t : \mathbb{N} \rightarrow \mathbb{N}) (g : \mathbb{N} \rightarrow \alpha \rightarrow \mathbb{C}) (N : \mathbb{R})$ 
    (f :  $\alpha \rightarrow \mathbb{C}$ )
    (x : WeakGridSpace.BesovishSpace
      (souzaAtomFamily G s p hs hp hp_top) q)
    (R : WeakGridSpace.LpGridRepresentation
      (souzaAtomFamily G s p hs hp hp_top)
      (x : Lp  $\mathbb{C}$  p G.toWeakGridSpace.measure)),
     $0 \leq N \rightarrow$ 
    WeakGridSpace.RepresentsFunction
      (G := G.toWeakGridSpace) (p := p) f
      (x : Lp  $\mathbb{C}$  p G.toWeakGridSpace.measure)  $\rightarrow$ 
    WeakGridSpace.LpGridRepresentation.FinitePQCost (q := q) R  $\rightarrow$ 
    ( $\forall i \in \Lambda$ ,
       $\exists C : \mathbb{R}$ ,
        SouzaPointwiseSelfsTailBound
          G  $\beta$  p qtilde h $\beta$  hp hp_top (t i) (g i) C)  $\rightarrow$ 
    ( $\forall k$  (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k),
      (R.block k).coeff Q  $\neq 0 \rightarrow$ 
         $\exists T : \mathbb{R}$ ,
          HasSum
            (fun i : {i // i  $\in \Lambda$ } =>
              nonArchimedeanRelevantTailSelfsInfiniteTerm
                G  $\beta$  p qtilde h $\beta$  hp hp_top  $\Lambda$  t g Q i)
            T  $\wedge$ 
            T  $\leq N$ )  $\rightarrow$ 
    ( $\forall k$  (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k) i,
      i  $\in \Lambda \rightarrow$ 
      (R.block k).coeff Q  $\neq 0 \rightarrow$ 
      goodGridLevelCellMeetsSupport G Q (g i)  $\rightarrow$ 
      t i  $\leq k$ )  $\rightarrow$ 
     $\exists h : \alpha \rightarrow \mathbb{C}$ ,
       $\exists$  absSum :  $\alpha \rightarrow \mathbb{R}$ ,
        ( $\forall^m z$   $\partial$ G.toWeakGridSpace.measure,
          f z  $\neq 0 \rightarrow$ 
          HasSum
            (fun i : {i // i  $\in \Lambda$ } => ||g i.1 z||)
            (absSum z)  $\wedge$ 

```

```

    absSum z ≤ Cgen * N) ∧
(∀m z ∂G.toWeakGridSpace.measure,
  HasSum
    (fun i : {i // i ∈ Λ} => g i.1 z * f z)
    (h z)) ∧
(∀m z ∂G.toWeakGridSpace.measure,
  ||h z|| ≤ Cgen * N * ||f z||) ∧
(∃ hmem : MemLp h p G.toWeakGridSpace.measure,
  ||MemLp.toLp h hmem|| ≤
    Cgen * N * ||(x : Lp C p G.toWeakGridSpace.measure)||) ∧
∃ y : WeakGridSpace.BesovishSpace
  (souzaAtomFamily G s p hs hp hp_top) q,
∃ S : WeakGridSpace.LpGridRepresentation
  (souzaAtomFamily G s p hs hp hp_top)
  (y : Lp C p G.toWeakGridSpace.measure),
WeakGridSpace.RepresentsFunction
  (G := G.toWeakGridSpace) (p := p) h
  (y : Lp C p G.toWeakGridSpace.measure) ∧
WeakGridSpace.LpGridRepresentation.FinitePQCost (q := q) S ∧
WeakGridSpace.LpGridRepresentation.pqCost (q := q) S ≤
  Cgen * N *
  WeakGridSpace.LpGridRepresentation.pqCost (q := q) R := ...

```

**Theorem 19.6** (Non-Archimedean property, positive version). *In the setting of Theorems 19.4 and 19.5 (finite or infinite  $\Lambda$ , with the respective hypotheses, the tail seminorms being replaced by their positive-cone counterparts*

$$|g|_{\mathcal{B}_{p,\vec{q},selfs}^{\beta+,t}} = \sup_{\substack{a_Q \in \mathcal{A}_{\beta,p}^{zz} \\ Q \in \mathcal{P}^j, j \geq t}} |ga_Q|_{\mathcal{B}_{p,\vec{q}}^{\beta+}}$$

for the functions  $g_i$ ), assume that each  $g_i$  is  $\mathcal{B}_{p,\vec{q}}^{\beta,t_i}$ -positive. Let  $R$  be the given canonical representation  $(c_Q)_Q$  of  $f$ . Then the representation  $S = (d_Q)_Q$  of  $(\sum_i g_i)f$  produced by Theorems 19.4/19.5 can be chosen so that, in addition:

- i. (positivity) if  $R$  is a positive representation ( $c_Q \geq 0$  for all  $Q$ ), then  $S$  is a cone-positive representation: all  $d_Q$  are nonnegative reals and the corresponding Souza's atoms are a.e. nonnegative;
- ii. (support) unconditionally — without assuming  $c_Q \geq 0$  — every cell  $Q$  active in  $S$  ( $d_Q \neq 0$ ) satisfies: there exists  $i \in \Lambda$  with  $g_i \neq 0$  a.e. on  $Q$ .

**Formalization.** In the file `NonArchimedeanPropertyPositiveStandalone.lean` of `GoodGrid/Multipliers/`, with cores in `NonArchimedeanProperty.lean` and `GoodGrid/PositiveCone.lean`. The finite case is the theorem `souzaNonArchimedeanPropertyPositiveCone`: it produces the representation  $S$  of the product with the cost bound of Theorem 19.4 and, in addition, the support conclusion ii. and the positivity implication i. The infinite case is `souzaNonArchimedeanPropertyPositiveConeInfinite` ( $\Lambda \subset \mathbb{N}$  infinite, with condition A as an  $[0, \infty]$ -valued series bound, no summability witness needed): it

proves all the conclusions of Theorem 19.5 plus i. and ii.; the limit arguments pass ii. through the convergence of the coefficients and i. through the closedness of the nonnegative real ray in  $\mathbb{C}$  and a.e. convergence of a subsequence of the atoms. The auxiliary product representation behind both is `exists_nonArchimedeanProductRepresentation_positive`, whose key local ingredient (`exists_nonArchimedeanLocalTransmutationData_pos`) assembles, one multiplier at a time, positive local transmutation data on the induced grids. Compared with paper Remark 18.5 the statement was made precise as follows: (a) the hypothesis on the representation  $R$  of  $f$  is only that it is canonical (arbitrary complex coefficients); (b) the two conclusions are separated according to their true strengths — the support conclusion ii. holds unconditionally, and is weakened to its a.e. form ( $g_i \neq 0$  a.e. on  $Q$  rather than  $Q \subset \text{supp } g_i$ ); (c) the positivity conclusion i. produces a *cone-positive* representation (nonnegative real coefficients, a.e. nonnegative Souza’s atoms), not necessarily one by canonical atoms. Axioms of all four non-Archimedean theorems checked: `propext`, `Classical.choice`, `Quot.sound`.

```

theorem souzaNonArchimedeanPropertyPositiveCone
  (G : GoodGridSpace (α := α))
  (s β : ℝ) (p q qtilde : ℝ≥0∞)
  (hs : 0 < s) (hβ : 0 < β) (hβs : s < β)
  (hβ_lt_inv : β < (p.toReal)-1)
  (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)] [Fact (1 ≤ qtilde)] :
  ∃ Cgen : ℝ,
    0 ≤ Cgen ∧
    ∀ (Λ : Finset ℕ) (t : ℕ → ℕ) (g : ℕ → α → ℂ) (N : ℝ)
      (f : α → ℂ)
      (x : WeakGridSpace.BesovishSpace
        (souzaAtomFamily G s p hs hp hp_top) q)
      (R : WeakGridSpace.LpGridRepresentation
        (souzaAtomFamily G s p hs hp hp_top)
        (x : Lp ℂ p G.toWeakGridSpace.measure)),
    0 ≤ N →
    WeakGridSpace.RepresentsFunction
      (G := G.toWeakGridSpace) (p := p) f
      (x : Lp ℂ p G.toWeakGridSpace.measure) →
    WeakGridSpace.LpGridRepresentation.FinitePQCost (q := q) R →
    SouzaCanonicalRepresentation G s p hs hp hp_top R →
    (∀ i ∈ Λ,
      SouzaPositiveFunction G β p qtilde hβ hp hp_top (g i)) →
    (∀ i ∈ Λ,
      ∃ C : ℝ≥0∞,
        SouzaPositivePointwiseSelfsTailBound
          G β p qtilde hβ hp hp_top (t i) (g i) C) →
    (∀ k (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k),
      (R.block k).coeff Q ≠ 0 →
        nonArchimedeanRelevantPositiveTailSelfsSum
          G β p qtilde hβ hp hp_top Λ t g Q ≤ ENNReal.ofReal N) →
    (∀ k (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k) i,
      i ∈ Λ →
        (R.block k).coeff Q ≠ 0 →

```

```

    goodGridLevelCellMeetsSupport G Q (g i) →
      t i ≤ k) →
  ∃ y : WeakGridSpace.BesovishSpace
    (souzaAtomFamily G s p hs hp hp_top) q,
  ∃ S : WeakGridSpace.LpGridRepresentation
    (souzaAtomFamily G s p hs hp hp_top)
    (y : Lp ℂ p G.toWeakGridSpace.measure),
  WeakGridSpace.RepresentsFunction
    (G := G.toWeakGridSpace) (p := p)
    (fun z => (∑ i ∈ Λ, g i z) * f z)
    (y : Lp ℂ p G.toWeakGridSpace.measure) ∧
  WeakGridSpace.LpGridRepresentation.FinitePQCost (q := q) S ∧
  WeakGridSpace.LpGridRepresentation.pqCost (q := q) S ≤
  Cgen * N *
  WeakGridSpace.LpGridRepresentation.pqCost (q := q) R ∧
  -- [ii] support: holds with only canonical atoms on 'R' (no sign on
  'c_Q').
  (∀ k (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k),
    (S.block k).coeff Q ≠ 0 →
      ∃ i ∈ Λ, ∀m z ∂(G.toWeakGridSpace.measure.restrict Q.1), g i z
      ≠ 0) ∧
  -- [i] positivity: only when 'R' is fully positive (canonical + 'c_Q
  ≥ 0').
  (SouzaPositiveRepresentation G s p hs hp hp_top R →
    SouzaConePositiveRepresentation G s p hs hp hp_top S) := ...

theorem souzaNonArchimedeanPropertyPositiveConeInfinite
  (G : GoodGridSpace (α := α))
  (s β : ℝ) (p q qtilde : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hβ : 0 < β) (hβs : s < β)
  (hβ_lt_inv : β < (p.toReal)-1)
  (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)] [Fact (1 ≤ qtilde)] :
  ∃ Cgen : ℝ,
    0 ≤ Cgen ∧
    1 ≤ Cgen ∧
    ∀ (Λ : Set ℕ) (t : ℕ → ℕ) (g : ℕ → α → ℂ) (N : ℝ)
      (f : α → ℂ)
      (x : WeakGridSpace.BesovishSpace
        (souzaAtomFamily G s p hs hp hp_top) q)
      (R : WeakGridSpace.LpGridRepresentation
        (souzaAtomFamily G s p hs hp hp_top)
        (x : Lp ℂ p G.toWeakGridSpace.measure)),
    0 ≤ N →
      WeakGridSpace.RepresentsFunction
        (G := G.toWeakGridSpace) (p := p) f
        (x : Lp ℂ p G.toWeakGridSpace.measure) →
      WeakGridSpace.LpGridRepresentation.FinitePQCost (q := q) R →
      SouzaCanonicalRepresentation G s p hs hp hp_top R →
      (∀ i ∈ Λ,
        SouzaPositiveFunction G β p qtilde hβ hp hp_top (g i)) →
      (∀ i ∈ Λ,
        ∃ C : ℝ ≥ 0 ∞,
          SouzaPositivePointwiseSelfsTailBound
            G β p qtilde hβ hp hp_top (t i) (g i) C) →

```

```

(∀ k (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k),
  (R.block k).coeff Q ≠ 0 →
    (∑' i : {i // i ∈ Λ},
      nonArchimedeanRelevantPositiveTailSelfInfiniteTerm
        G β p qtilde hβ hp hp_top Λ t g Q i) ≤ ENNReal.ofReal N) →
  (∀ k (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k) i,
    i ∈ Λ →
      (R.block k).coeff Q ≠ 0 →
        goodGridLevelCellMeetsSupport G Q (g i) →
          t i ≤ k) →
  ∃ h : α → ℂ,
  ∃ absSum : α → ℝ,
  (∀m z ∂G.toWeakGridSpace.measure,
    f z ≠ 0 →
      HasSum
        (fun i : {i // i ∈ Λ} => ||g i.1 z||)
        (absSum z) ∧
        absSum z ≤ Cgen * N) ∧
  (∀m z ∂G.toWeakGridSpace.measure,
    HasSum
      (fun i : {i // i ∈ Λ} => g i.1 z * f z)
      (h z)) ∧
  (∀m z ∂G.toWeakGridSpace.measure,
    ||h z|| ≤ Cgen * N * ||f z||) ∧
  (∃ hmem : MemLp h p G.toWeakGridSpace.measure,
    ||MemLp.toLp h hmem|| ≤
      Cgen * N * ||(x : Lp ℂ p G.toWeakGridSpace.measure)||) ∧
  ∃ y : WeakGridSpace.BesovishSpace
    (souzaAtomFamily G s p hs hp hp_top) q,
  ∃ S : WeakGridSpace.LpGridRepresentation
    (souzaAtomFamily G s p hs hp hp_top)
    (y : Lp ℂ p G.toWeakGridSpace.measure),
  WeakGridSpace.RepresentsFunction
    (G := G.toWeakGridSpace) (p := p) h
    (y : Lp ℂ p G.toWeakGridSpace.measure) ∧
  WeakGridSpace.LpGridRepresentation.FinitePQCost (q := q) S ∧
  WeakGridSpace.LpGridRepresentation.pqCost (q := q) S ≤
    Cgen * N *
      WeakGridSpace.LpGridRepresentation.pqCost (q := q) R ∧
  (∀ k (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k),
    (S.block k).coeff Q ≠ 0 →
      ∃ i ∈ Λ,
        ∀m z ∂(G.toWeakGridSpace.measure.restrict Q.1),
          g i z ≠ 0) ∧
  (SouzaPositiveRepresentation G s p hs hp hp_top R →
    SouzaConePositiveRepresentation G s p hs hp hp_top S) := ...

```

## 19.6. The tail selfs classes consist of multipliers.

**File:** GoodGrid/Multipliers/SelfsSubsetMultipliers.lean

**Imports (within the project):** GoodGrid/Multipliers/  
NonArchimedeanProperty.lean

The first consequence of the non-Archimedean property is the continuous inclusion of the tail `selfs` classes in the multiplier space.

**Corollary 19.7.** *Let  $0 < s < \beta < 1/p$  and  $q, \tilde{q} \in [1, \infty]$ , and fix a cutoff level  $t \in \mathbb{N}$ . Then*

$$\mathcal{B}_{p, \tilde{q}, \text{selfs}}^{\beta, t} \subset M(\mathcal{B}_{p, q}^s),$$

*and the inclusion is continuous:  $|g|_{M(\mathcal{B}_{p, q}^s)} \leq C |g|_{\mathcal{B}_{p, \tilde{q}, \text{selfs}}^{\beta, t}}$  with a constant  $C$  independent of  $g$ .*

**Formalization.** In `GoodGrid/Multipliers/SelfsSubsetMultipliers.lean` (paper Corollary 18.6). The quantitative form is `exists_souzaSelfsMultiplierConstant`: a tail `selfs` bound  $C$  for  $g$  produces the multiplier operator bound  $C_{\text{mult}} \cdot C$ . The inclusion is `souzaPointwiseMultiplier_of_souzaPointwiseSelfsTailClass` and the continuity at the level of (semi)norms is `souzaPointwiseMultiplierNorm.le_const_mul_selfsTailNorm`. The proof applies the non-Archimedean property (Theorem 19.4) to the one-element family  $\{g\}$ , after the level-lowering step `souzaPointwiseSelfsTailBound_levelZero`: a tail `selfs` bound from level  $t$  onward yields a full (level-0) bound with constant multiplied by the number of level- $t$  cells, because an atom on a shallow cell is the finite disjoint sum of its level- $t$  pieces, each a scaled atom on a tail cell. Axioms checked: `propext`, `Classical.choice`, `Quot.sound`.

```

theorem exists_souzaSelfsMultiplierConstant
  (G : GoodGridSpace (α := α)) (s β : ℝ) (p q qtilde : ℝ ≥ 0 ∞) (t : ℕ)
  (hs : 0 < s) (hβ : 0 < β) (hβs : s < β)
  (hβ_lt_inv : β < (p.toReal)-1)
  (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)] [Fact (1 ≤ qtilde)] :
  ∃ Cmult : ℝ,
    0 ≤ Cmult ∧
    ∀ (g : α → ℂ) (C : ℝ),
      SouzaPointwiseSelfsTailBound G β p qtilde hβ hp hp_top t g C →
      SouzaPointwiseMultiplierBound G s p q hs hp hp_top g (Cmult * C)
  := ...

theorem souzaPointwiseMultiplierNorm_le_const_mul_selfsTailNorm
  (G : GoodGridSpace (α := α)) (s β : ℝ) (p q qtilde : ℝ ≥ 0 ∞) (t : ℕ)
  (hs : 0 < s) (hβ : 0 < β) (hβs : s < β)
  (hβ_lt_inv : β < (p.toReal)-1)
  (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)] [Fact (1 ≤ qtilde)] :
  ∃ Cmult : ℝ,
    0 ≤ Cmult ∧
    ∀ g : α → ℂ,
      SouzaPointwiseSelfsTailClass G β p qtilde hβ hp hp_top t g →
      WeakGridSpace.pointwiseMultiplierNorm
        (A := souzaAtomFamily G s p hs hp hp_top) q g ≤

```

```

Cmult * souzaPointwiseSelfsTailNorm G β p qtilde hβ hp hp_top t
g := ...

```

### 19.7. Strongly regular domains and Pointwise Multipliers I.

**File:** GoodGrid/Multipliers/StronglyRegularDomains.lean  
**Imports (within the project):** GoodGrid/Multipliers/  
NonArchimedeanPropertyPositiveStandalone.lean

The positive-cone version of the non-Archimedean property was designed for indicators, and strongly regular domains are the class of sets whose indicators it handles.

**Definition 19.8.** Let  $a \in \mathbb{R}$ ,  $K \geq 0$  and  $k_1 \in \mathbb{N}$ . A measurable set  $\Omega$  is an  $(a, K, k_1)$ -**strongly regular domain** if for every grid cell  $Q$  of level at least  $k_1$  the intersection  $Q \cap \Omega$  is an exactly disjoint countable union of grid cells,

$$Q \cap \Omega = \bigcup_k \bigcup_{P \in \mathcal{F}^k} P,$$

with  $\mathcal{F}^k \subset \mathcal{P}^k$  and the level-by-level cost control

$$\sum_{P \in \mathcal{F}^k} m(P)^a \leq K m(Q)^a \quad \text{for every } k.$$

**Proposition 19.9.** *If  $\Omega$  is a  $(1 - \beta p, K, k_1)$ -strongly regular domain then*

$$|\mathbb{1}_\Omega|_{\mathcal{B}_{p,\infty, self}^{\beta+, k_1}} \leq K^{1/p} :$$

*for every grid cell  $Q$  of level at least  $k_1$ , the product of  $\mathbb{1}_\Omega$  with the canonical  $(\beta, p)$ -Souza's atom on  $Q$  admits a positive Souza representation of positive  $(p, \infty)$ -gauge at most  $K^{1/p}$ . In particular  $\mathbb{1}_\Omega \in \mathcal{B}_{p,\infty}^{\beta+}$ .*

**Formalization.** In GoodGrid/Multipliers/StronglyRegularDomains.lean (paper 18.7–18.8). The decomposition data inside one cell is the structure `StronglyRegularDecomposition` (families  $\mathcal{F}^k$  of genuine level- $k$  cells, exact disjoint cover of  $Q \cap \Omega$ , disjointness also across levels, and the cost bound), and the definition is `StronglyRegularDomain`. Proposition 19.9 is `souzaPositiveSelfsTailBound.of_stronglyRegularDomain`, with the weighted variant `souzaPositiveSelfsTailBound.smul_of_stronglyRegularDomain` ( $\Theta \cdot \mathbb{1}_\Omega$  with constant  $\Theta K^{1/p}$ ) and the membership `souzaPositiveFunction.of_stronglyRegularDomain`: the positive representation of  $\mathbb{1}_\Omega \cdot a_Q$  is read off directly from the decomposition families.

```

def StronglyRegularDomain
  (G : GoodGridSpace (α := α)) (Ω : Set α) (a K : ℝ) (k₁ : ℕ) : Prop :=
  MeasurableSet Ω ∧
  ∀ Q : GoodGridCell G, k₁ ≤ Q.level →
    Nonempty (StronglyRegularDecomposition G Ω a K Q)

```

```

theorem souzaPositiveSelfsTailBound_of_stronglyRegularDomain
  (G : GoodGridSpace (α := α)) (β : ℝ) (p : ℝ ≥ 0 ∞)
  (hβ : 0 < β) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)]
  {Ω : Set α} {K : ℝ} {k₁ : ℕ}
  (hΩ : StronglyRegularDomain G Ω (1 - β * p.toReal) K k₁) :
  SouzaPositivePointwiseSelfsTailBound G β p ∞ hβ hp hp_top k₁
  (Ω.indicator fun _ => (1 : ℂ))
  (ENNReal.ofReal (K ^ (p.toReal)⁻¹)) := ...

```

**Theorem 19.10** (Pointwise Multipliers I). *There is a constant  $C$  with the following property. Let  $\Omega_i$ , for  $i$  in a finite set  $\Lambda$ , be  $(1 - \beta p, K_i, t_i)$ -strongly regular domains, let  $\Theta_i > 0$  be weights, and let  $f$  be represented by a finite-cost canonical Souza representation  $R = (c_Q)_Q$  in  $\mathcal{B}_{p,q}^s$  such that*

- A. *on every active cell  $Q$  ( $c_Q \neq 0$ ), the total weighted cost  $\sum_{i: Q \cap \Omega_i \neq \emptyset} \Theta_i K_i^{1/p}$  is at most  $N$ ;*
- B. *every active cell meeting  $\Omega_i$  has level at least  $t_i$ .*

*Then  $g = \sum_{i \in \Lambda} \Theta_i \mathbb{1}_{\Omega_i}$  is a multiplier of the represented function: the product  $gf$  has a Souza representation  $S$  with finite cost and*

$$\text{cost}(S) \leq C N \text{cost}(R),$$

*and moreover (i) every active cell of  $S$  is, up to measure zero, contained in some  $\Omega_i$ , and (ii) if  $R$  is positive then  $S$  is cone-positive. An infinite-family version ( $\Lambda \subset \mathbb{N}$  countable) holds with condition A stated as an  $[0, \infty]$ -valued series bound.*

**Formalization.** In `GoodGrid/Multipliers/StronglyRegularDomains.lean` (paper Proposition 18.9). The finite case is `souzaPointwiseMultipliersI` and the countable case is `souzaPointwiseMultipliersIInfinite`; the weighted overlap costs of condition A are `stronglyRegularOverlapCost` and `stronglyRegularOverlapCostInfinite`. Both are direct applications of the positive-cone non-Archimedean theorems (Theorem 19.6) to the weighted indicators, whose positive tail `selfs` bounds come from Proposition 19.9; the support conclusion ii. of Theorem 19.6 (nonvanishing a.e.) here upgrades to a.e. *containment* in some  $\Omega_i$ , since the multipliers are indicators. Axioms checked: `propext`, `Classical.choice`, `Quot.sound`.

```

theorem souzaPointwiseMultipliersI
  (G : GoodGridSpace (α := α))
  (s β : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hβ : 0 < β) (hβs : s < β)
  (hβ_lt_inv : β < (p.toReal)⁻¹)
  (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)] :
  ∃ Cgen2 : ℝ,

```

```

0 ≤ Cgen2 ∧
∀ (Λ : Finset N) (t : N → N) (Ω : N → Set α) (K Θ : N → ℝ) (N : ℝ)
(f : α → ℂ)
(x : WeakGridSpace.BesovishSpace
  (souzaAtomFamily G s p hs hp hp_top) q)
(R : WeakGridSpace.LpGridRepresentation
  (souzaAtomFamily G s p hs hp hp_top)
  (x : Lp ℂ p G.toWeakGridSpace.measure)),
0 ≤ N →
(∀ i ∈ Λ,
  StronglyRegularDomain G (Ω i) (1 - β * p.toReal) (K i) (t i)) →
(∀ i ∈ Λ, 0 < Θ i) →
WeakGridSpace.RepresentsFunction
  (G := G.toWeakGridSpace) (p := p) f
  (x : Lp ℂ p G.toWeakGridSpace.measure) →
WeakGridSpace.LpGridRepresentation.FinitePQCost (q := q) R →
SouzaCanonicalRepresentation G s p hs hp hp_top R →
(∀ k (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k),
  (R.block k).coeff Q ≠ 0 →
    stronglyRegularOverlapCost G p Λ Ω K Θ Q ≤ N) →
(∀ k (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k) i,
  i ∈ Λ →
    (R.block k).coeff Q ≠ 0 →
      (Q.1 ∩ Ω i).Nonempty →
        t i ≤ k) →
  ∃ y : WeakGridSpace.BesovishSpace
    (souzaAtomFamily G s p hs hp hp_top) q,
  ∃ S : WeakGridSpace.LpGridRepresentation
    (souzaAtomFamily G s p hs hp hp_top)
    (y : Lp ℂ p G.toWeakGridSpace.measure),
  WeakGridSpace.RepresentsFunction
    (G := G.toWeakGridSpace) (p := p)
    (fun z => (∑ i ∈ Λ,
      (Θ i : ℂ) * (Ω i).indicator (fun _ => (1 : ℂ)) z) * f z)
    (y : Lp ℂ p G.toWeakGridSpace.measure) ∧
  WeakGridSpace.LpGridRepresentation.FinitePQCost (q := q) S ∧
  WeakGridSpace.LpGridRepresentation.pqCost (q := q) S ≤
    Cgen2 * N *
    WeakGridSpace.LpGridRepresentation.pqCost (q := q) R ∧
  (∀ k (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k),
    (S.block k).coeff Q ≠ 0 →
      ∃ i ∈ Λ,
        ∀m z ∂(G.toWeakGridSpace.measure.restrict Q.1),
          z ∈ Ω i) ∧
    (SouzaPositiveRepresentation G s p hs hp hp_top R →
      SouzaConePositiveRepresentation G s p hs hp hp_top S) := ...

```

### 19.8. $\mathcal{B}_{p,\infty}^{1/p} \cap L^\infty$ and Pointwise Multipliers II.

**File:** GoodGrid/Multipliers/Bp1overpinftyisMultiplier.lean  
**Imports (within the project):** GoodGrid/Multipliers/  
 NonArchimedeanProperty.lean, GoodGrid/DiracApproximations.lean,  
 GoodGrid/AlternativeRepresentationsAndNorms/  
 FiniteStandardNormimpliesBesov.lean

**Proposition 19.11** (Pointwise Multipliers II). *Let  $0 < s < 1/p$ . There is a constant  $C$  such that every  $g \in \mathcal{B}_{p,\infty}^{1/p} \cap L^\infty$  is a pointwise multiplier of  $\mathcal{B}_{p,q}^s$ , with*

$$|g|_{M(\mathcal{B}_{p,q}^s)} \leq C |g|_{\mathcal{B}_{p,\infty}^{1/p}} + |g|_\infty.$$

**Formalization.** In GoodGrid/Multipliers/Bp1overpinftyisMultiplier.lean (paper Proposition 18.10). The statement is `souzaPointwiseMultipliersII`, with the hypothesis  $g \in \mathcal{B}_{p,\infty}^{1/p} \cap L^\infty$  rendered as:  $g$  is represented by an element `xg` of the  $(1/p, p, \infty)$  Souza space and  $\|g\| \leq M$  a.e. The proof has three layers. (1) `exists_fouRepresentation` packages the input from Corollary `fou` (Section 16) and Proposition 17.1.B (Section 18): the canonical standard representation of  $g$  has  $(p, \infty)$ -cost controlled by  $|g|_{\mathcal{B}_{p,\infty}^{1/p}}$ , and all its ancestor-tower coefficient sums (`ancestorCoeffSum`) are bounded by  $M$ , because for  $s = 1/p$  the canonical atoms have value  $m(J)^{1/p-1/p} = 1$  on their cells, so the tower sums are exactly the partial standard sums, i.e. averages of  $g$  over cells. (2) The paper’s splitting  $gf = u_1 + u_2$  is `exists_mult_product_representation` (block-form core `exists_mult_product_blocks`): the block families `multU1Block` and `multU2Block` realize the atom algebra  $a_Q b_J$  ( $J \subsetneq Q$  and  $Q \subseteq J$ , respectively); the  $u_1$  cost is controlled, via the coefficient bound `multU1Block_coeff_norm_le` and single-ancestor counting, by a self-contained discrete Young inequality with the geometric kernel  $\lambda_2^{n(1/p-s)}$  (`geometric_conv_rpow_summable_and_tsum_le`), producing the constant  $C_{conv} = \lambda_2^\theta / (1 - \lambda_2^\theta)$  with  $\theta = 1/p - s$ ; the  $u_2$  cost is controlled by the tower-sum  $L^\infty$  bound (`multU2Block_levelCoeffPower_le`); both block series converge in  $L^p$  through `formalBlockSeq_hasRepresentation` (Section 6); and the identity  $gf = u_1 + u_2$  follows from the *exact* pointwise identity of truncated products (`mult_truncated_pointwise_identity`, proved along the tower of cells containing each point), passed to the limit along a.e.-convergent subsequences (`representsPointwiseProduct_of_tendsto_Lp_varying`) — the  $L^1/\text{comp1}$  route sketched in the paper was not needed. (3) The outer proof:  $\varepsilon$ -optimization over near-optimal representations of  $f$  and uniqueness of the product representative in  $L^p$ . Axioms of `souzaPointwiseMultipliersII`, `exists_mult_product_representation` and `exists_fouRepresentation` checked: `propext`, `Classical.choice`, `Quot.sound`.

```
theorem souzaPointwiseMultipliersII
  (G : GoodGridSpace (α := α)) (s : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hs_lt_inv : s < (p.toReal)-1)
  (h1p : 0 < (p.toReal)-1) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)] :
  ∃ Cmult : ℝ,
```

```

0 ≤ Cmult ∧
∀ (g : α → ℂ) (M : ℝ)
(xg : WeakGridSpace.BesovishSpace
  (souzaAtomFamily G (p.toReal)-1 p h1p hp hp_top) ∞),
WeakGridSpace.RepresentsFunction
  (G := G.toWeakGridSpace) (p := p) g
(xg : Lp ℂ p G.toWeakGridSpace.measure) →
(∀m z ∂G.toWeakGridSpace.measure, ‖g z‖ ≤ M) →
SouzaPointwiseMultiplierBound G s p q hs hp hp_top g
(Cmult * WeakGridSpace.BesovishSpace.Norm_Costpq
  (souzaAtomFamily G (p.toReal)-1 p h1p hp hp_top) ∞ xg
  + M) := ...

theorem exists_fouRepresentation
  (G : GoodGridSpace (α := α)) (p : ℝ ≥ 0 ∞)
  (h1p : 0 < (p.toReal)-1) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] :
  ∃ Cfou : ℝ,
  0 ≤ Cfou ∧
  ∀ (g : α → ℂ) (M : ℝ)
  (xg : WeakGridSpace.BesovishSpace
    (souzaAtomFamily G (p.toReal)-1 p h1p hp hp_top) ∞),
  WeakGridSpace.RepresentsFunction
    (G := G.toWeakGridSpace) (p := p) g
  (xg : Lp ℂ p G.toWeakGridSpace.measure) →
  (∀m z ∂G.toWeakGridSpace.measure, ‖g z‖ ≤ M) →
  ∃ Rg : WeakGridSpace.LpGridRepresentation
    (souzaAtomFamily G (p.toReal)-1 p h1p hp hp_top)
    (xg : Lp ℂ p G.toWeakGridSpace.measure),
  SouzaCanonicalRepresentation G (p.toReal)-1 p h1p hp hp_top Rg ∧
  WeakGridSpace.LpGridRepresentation.FinitePQCost (q := ∞) Rg ∧
  WeakGridSpace.LpGridRepresentation.pqCost (q := ∞) Rg ≤
  Cfou * WeakGridSpace.BesovishSpace.Norm_Costpq
    (souzaAtomFamily G (p.toReal)-1 p h1p hp hp_top) ∞ xg ∧
  ∀ (k : ℕ) (Q : WeakGridSpace.LevelCell G.toWeakGridSpace k),
  ‖ancestorCoeffSum G Rg Q‖ ≤ M := ...

```

**Remark 19.12** (Positive Pointwise Multipliers II). Replacing  $\mathcal{B}_{p,b}^a$  by the positive cones  $\mathcal{B}_{p,b}^{a+}$  everywhere in Proposition 19.11: if  $g$  is represented by an element  $x_g$  of the positive cone of  $\mathcal{B}_{p,\infty}^{1/p}$  with finite positive gauge and  $\|g\| \leq M$  a.e., then for every element  $x$  of the positive cone of  $\mathcal{B}_{p,q}^s$  with finite positive gauge the product  $gf$  has a *positive* Besov representative  $y$  with

$$|y|_{\mathcal{B}_{p,q}^{s+}} \leq (C |x_g|_{\mathcal{B}_{p,\infty}^{1/p,+}} + M) |x|_{\mathcal{B}_{p,q}^{s+}},$$

the positive gauges being the  $[0, \infty]$ -valued infima of the  $(p, q)$ -costs over positive representations (Section 17).

**Formalization.** In the same file (the paper’s positive-cone remark to Proposition 18.10, Remark pos3 of the source). The statement is

`souzaPointwiseMultipliersIIPositive`, with the bound stated for the positive gauges `souzaPositiveNorm`; the representation form is `exists_mult_product_representation_pos` (positive finite-cost representations of  $g$  and  $f$  give a positive representation of the product with cost bound  $(C_{conv} \cdot \text{cost}_{(p,\infty)} R_g + M) \cdot \text{cost}_{(p,q)} R_f$ ). The same block construction is reused: the canonical-atom hypothesis and the ancestor-tower bound of the non-positive version are *derived* from positivity (`souzaPositiveRepresentation.canonical` and `ancestorCoeffSum.norm_lessBound_of_positive`, the positive form of Proposition 17.1.B: the tower partial sums are nondecreasing with an a.e.-convergent subsequence, so they are bounded by  $M$  on each cell); positivity of the  $u_1/u_2$  blocks is inherited from the input representations (`multU1Block_positive`, `multU2Block_positive`); and the two pieces are summed inside the positive cone (`souzaPositiveRepresentationAdd`). Axioms checked: `propext`, `Classical.choice`, `Quot.sound`.

```

theorem souzaPointwiseMultipliersIIPositive
  (G : GoodGridSpace (α := α)) (s : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hs_lt_inv : s < (p.toReal)-1)
  (h1p : 0 < (p.toReal)-1) (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)] :
  ∃ Cmult : ℝ,
    0 ≤ Cmult ∧
    ∀ (g : α → ℂ) (M : ℝ)
      (xg : WeakGridSpace.BesovishSpace
        (souzaAtomFamily G (p.toReal)-1 p h1p hp hp_top) ∞),
      WeakGridSpace.RepresentsFunction
        (G := G.toWeakGridSpace) (p := p) g
        (xg : Lp ℂ p G.toWeakGridSpace.measure) →
        (∀m z ∂G.toWeakGridSpace.measure, ||g z|| ≤ M) →
        souzaPositiveNorm G (p.toReal)-1 p ∞ h1p hp hp_top xg ≠ ∞ →
        ∀ x : WeakGridSpace.BesovishSpace
          (souzaAtomFamily G s p hs hp hp_top) q,
          souzaPositiveNorm G s p q hs hp hp_top x ≠ ∞ →
          ∃ y : WeakGridSpace.BesovishSpace
            (souzaAtomFamily G s p hs hp hp_top) q,
            WeakGridSpace.RepresentsPointwiseProduct
              (G := G.toWeakGridSpace) (p := p) g
              (x : Lp ℂ p G.toWeakGridSpace.measure)
              (y : Lp ℂ p G.toWeakGridSpace.measure) ∧
              SouzaPositiveElement G s p q hs hp hp_top y ∧
              souzaPositiveNorm G s p q hs hp hp_top y ≤
              (ENNReal.ofReal Cmult *
                souzaPositiveNorm G (p.toReal)-1 p ∞ h1p hp hp_top xg +
                ENNReal.ofReal M) *
              souzaPositiveNorm G s p q hs hp hp_top x := ...

```

## 20. REGULAR DOMAINS

```

File: GoodGrid/RegularDomains.lean
Imports (within the project): GoodGrid/Multipliers/
StronglyRegularDomains.lean, GoodGrid/QuasiAlgebra.lean

```

Strongly regular domains control  $\mathbb{1}_\Omega$  through one cost constant per intersecting cell. The paper's *regular domains* (the subsection labelled `cf` in the source) relax this to a single geometric series across the whole family: the indicators are still bounded multipliers of  $\mathcal{B}_{p,q}^s \cap L^\infty$ , and the restrictions  $g \cdot \mathbb{1}_{\Omega_r}$  admit Souza representations whose coefficients are localized in  $\Omega_r$ . Throughout,  $k_0(\Omega)$  is the first level at which  $\Omega$  contains a grid cell (`firstContainedLevel`).

**Definition 20.1.** Let  $a \in \mathbb{R}$ ,  $C \geq 0$  and  $0 \leq c < 1$ . A countable family  $\{\Omega_r\}_{r \in \Lambda}$ ,  $\Lambda \subset \mathbb{N}$ , of pairwise disjoint measurable sets is an  $(a, C, c)$ -**regular family** if one can choose families  $\mathcal{F}^k(\Omega_r) \subset \mathcal{P}^k$  tiling each  $\Omega_r$  ( $\Omega_r = \bigcup_k \bigcup_{Q \in \mathcal{F}^k(\Omega_r)} Q$ , disjointly, with  $\mathcal{F}^k(\Omega_r) = \emptyset$  for  $k < k_0(\Omega_r)$ ) such that the level cost obeys

$$(\text{dom}) \quad \sum_{r \in \Lambda} \sum_{Q \in \mathcal{F}^k(\Omega_r)} m(Q)^a \leq C c^{k-k_0(\bigcup_r \Omega_r)} m\left(\bigcup_r \Omega_r\right)^a.$$

A measurable set  $\Omega$  is an  $(a, C, c)$ -**regular domain** if the singleton  $\{\Omega\}$  is an  $(a, C, c)$ -regular family.

**Formalization.** In `GoodGrid/RegularDomains.lean` (paper subsection `cf`). The two structures are `RegularFamily` and `RegularDomain`; both carry  $C \geq 0$  and  $0 \leq c < 1$  as fields and the cost `(dom)` as `cost`. The bridge from a domain to its singleton family is `RegularDomain.toRegularFamily_singleton`, and the reverse union construction (the family union is itself a regular domain) is `RegularFamily.regularDomain_union`, built from the per-level finite cell selection `unionFamily` and the summable active-family cost series `RegularFamily.cost_summable`.

**Proposition 20.2.** Let  $\beta > s$ . Every  $(1 - \beta p, K, 0)$ -strongly regular domain (that contains a grid cell) is a  $(1 - sp, C', \lambda_2^{(\beta-s)p})$ -regular domain for some  $C' \geq 0$ .

**Formalization.** In `GoodGrid/RegularDomains.lean` this is `regularDomain_of_stronglyRegularDomain`; the geometric ratio is the grid constant  $\lambda_2^{(\beta-s)p} < 1$ , and  $C' = K \lambda_1^{-k_0(\Omega)}$ . The candidate decomposition is assembled cell-by-cell over  $\mathcal{P}^{k_0(\Omega)}$  from the strong decompositions of  $Q \cap \Omega$ , and the quantitative step turns the strong  $1 - \beta p$  costs into the regular  $1 - sp$  geometric cost.

**Proposition 20.3.** *Let  $\Omega$  be a  $(1 - sp, C, c)$ -regular domain. Then its indicator is a Besov function: for finite  $q$ ,*

$$(estG) \quad |\mathbb{1}_\Omega|_{\mathcal{B}_{p,q}^s} \leq \frac{C^{1/p}}{(1 - c^{q/p})^{1/q}} m(\Omega)^{1/p-s},$$

*while at the endpoint  $q = \infty$  the geometric series collapses to a supremum (which is attained as  $c^{k-k_0} \leq 1$ ), so the factor  $(1 - c^{q/p})^{-1/q}$  disappears and*

$$(estG_\infty) \quad |\mathbb{1}_\Omega|_{\mathcal{B}_{p,\infty}^s} \leq C^{1/p} m(\Omega)^{1/p-s}.$$

*In both cases  $g \mapsto g \cdot \mathbb{1}_\Omega$  is a bounded multiplier of  $\mathcal{B}_{p,q}^s \cap L^\infty$ : there is  $C_{op} \geq 0$  with a representation of  $g \cdot \mathbb{1}_\Omega$  of cost at most  $C_{op} (|g|_{\mathcal{B}_{p,q}^s} + \|g\|_\infty)$ .*

**Formalization.** In `GoodGrid/RegularDomains.lean`. The finite- $q$  bound (estG) is `regularDomainIndicatorBesovNormBound` and the endpoint (estG $_\infty$ ) is `regularDomainIndicatorBesovNormBoundTop`; both are folded into the all- $q$  statement `regularDomainIndicatorBesovNormBoundAll`, whose explicit constant is `regularDomainIndicatorCost` (defined by case on  $q = \infty$ : the value is  $C^{1/p} m(\Omega)^{1/p-s}$  at the endpoint and  $C^{1/p} (1 - c^{q/p})^{-1/q} m(\Omega)^{1/p-s}$  for finite  $q$ ). The bounded multiplier is `regularDomainIndicatorMultiplierOnBoundedSouzaBesov` (finite  $q$ ) and `...top` ( $q = \infty$ , obtained by applying Pointwise Multipliers III at  $q = \infty$  to the indicator representation), unified as `...all`. The same four conclusions for the union of a regular family are `regularFamilyUnionIndicatorBesovNormBoundAll` and `regularFamilyUnionIndicatorMultiplierOnBoundedSouzaBesovAll` (each handling both finite  $q$  and the endpoint internally).

**Theorem 20.4** (Localized restriction representations). *If  $\{\Omega_r\}_{r \in \Lambda}$  is a  $(1 - ps, C, c)$ -regular family then there is a constant  $C_{rel} \geq 0$  such that for every  $g \in \mathcal{B}_{p,q}^s \cap L^\infty$  with  $\|g\|_\infty \leq M$  and each  $r \in \Lambda$  one finds a Souza representation*

$$(pdd) \quad g \cdot \mathbb{1}_{\Omega_r} = \sum_k \sum_{\substack{Q \in \mathcal{P}^k \\ Q \subset \Omega_r}} d_Q^r a_Q$$

*of finite cost whose nonzero coefficients are supported on cells contained in  $\Omega_r$ , with the mixed cost bound*

$$(hiip1) \quad \left( \sum_j \left( \sum_{r \in \Lambda} \sum_{\substack{Q \in \mathcal{P}^j \\ Q \subset \Omega_r}} |d_Q^r|^p \right)^{q/p} \right)^{1/q} \leq C_{rel} (|g|_{\mathcal{B}_{p,q}^s} + M).$$

*The statement is uniform in  $q \in [1, \infty]$ : at the endpoint  $q = \infty$  the outer  $\ell^q$  in (hiip1) is read as the supremum over levels  $j$  of the  $p$ -th roots  $(\sum_r \sum_{Q \subset \Omega_r} |d_Q^r|^p)^{1/p}$ , and  $C_{rel}$  uses the corresponding supremum form of the geometric reference cost.*

**Formalization.** In `GoodGrid/RegularDomains.lean` this is `regularFamilyRestrictionRepresentations`. The aggregated mixed cost of `(hiip1)` is `regularFamilyRestrictionCost` (its inner level term `regularFamilyRestrictionLevelCoeffPower` sums  $|d_Q^r|^p$  over both  $r \in \Lambda$  and cells  $Q$ ), and the geometric reference cost coming purely from regularity is `regularFamilyGeomRootCost`. Both `regularFamilyRestrictionCost` and `regularFamilyGeomRootCost` are defined by a case split on  $q = \infty$  (a supremum of the level roots at the endpoint, an  $\ell^q$  sum of those roots otherwise), so the single Lean theorem covers the whole range  $q \in [1, \infty]$ . The proof takes a weighted `fou` representation of  $g$  (`exists_weighted_fouRepresentation`), a localized Besov representation of each  $\mathbb{1}_{\Omega_r}$  (`regularFamilyIndicatorBesovRepresentation`), and combines them with the quasi-algebra  $u_1 + u_2$  product of Section 21 (`exists_quasi_product_of_tower_representations`); a convexity step  $|a + b|^p \leq 2^{p-1}(|a|^p + |b|^p)$  produces the explicit constant  $C_{rel} = (2^{p-1})^{1/p}(\text{regularFamilyGeomRootCost} + C_{fou})$ . Axioms checked: `propext`, `Classical.choice`, `Quot.sound`.

```

theorem regularFamilyRestrictionRepresentations
  (G : GoodGridSpace (α := α)) (Λ : Set ℕ) (Ω : ℕ → Set α)
  (s C c : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hs_lt_inv : s < (p.toReal)⁻¹)
  (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)]
  (hΩ : RegularFamily G Λ Ω (1 - p.toReal * s) C c) :
  ∃ Crel : ℝ,
    0 ≤ Crel ∧
    ∀ (g : α → ℂ) (M : ℝ)
      (xg : WeakGridSpace.BesovishSpace
        (souzaAtomFamily G s p hs hp hp_top) q),
      WeakGridSpace.RepresentsFunction
        (G := G.toWeakGridSpace) (p := p) g
        (xg : Lp ℂ p G.toWeakGridSpace.measure) →
        (∀m z ∂G.toWeakGridSpace.measure, ||g z|| ≤ M) →
      ∃ y : ℕ → WeakGridSpace.BesovishSpace
        (souzaAtomFamily G s p hs hp hp_top) q,
      ∃ R : (i : ℕ) →
        WeakGridSpace.LpGridRepresentation
          (souzaAtomFamily G s p hs hp hp_top)
          ((y i : ...) : Lp ℂ p G.toWeakGridSpace.measure),
      (∀ i ∈ Λ, WeakGridSpace.RepresentsFunction (p := p)
        (fun z => g z * (Ω i).indicator (fun _ => (1 : ℂ)) z)
        (y i : Lp ℂ p G.toWeakGridSpace.measure)) ∧
      (∀ i ∈ Λ,
        WeakGridSpace.LpGridRepresentation.FinitePQCost (q := q) (R i)) ∧
      (∀ i ∈ Λ, ∀ j (Q : WeakGridSpace.LevelCell G.toWeakGridSpace j),
        ((R i).block j).coeff Q ≠ 0 → Q.1 ⊆ Ω i) ∧
      regularFamilyRestrictionCost G s p q Λ y R ≤
      Crel *
      (WeakGridSpace.BesovishSpace.Norm_Costpq
        (souzaAtomFamily G s p hs hp hp_top) q xg + M) := ...

```

## 21. THE BOUNDED SOUZA–BESOV SPACE AS A QUASI-ALGEBRA

```

File: GoodGrid/QuasiAlgebra.lean
Imports (within the project): GoodGrid/Multipliers/
Bp1overpinfyisMultiplier.lean

```

This section documents the formalization of the quasi-algebra result of Section 19 of the published paper. Since a Besov-ish function need not be bounded,  $\mathcal{B}_{p,q}^s$  itself is not closed under products; but intersecting with  $L^\infty$  fixes this, and the proof is a two-sided run of the  $u_1 + u_2$  machinery of Section 19: each factor plays the role of the bounded multiplier  $g$  against the representation of the other, and the two mixed terms  $|f|_{\mathcal{B}} |g|_\infty + |g|_{\mathcal{B}} |f|_\infty$  appear.

**Proposition 21.1** (Pointwise Multipliers III). *Let  $0 < s < 1/p$ . The class  $\mathcal{B}_{p,q}^s \cap L^\infty$  is closed under pointwise multiplication: there is a constant  $C$ , depending only on the grid and the exponents, such that for all  $f, g \in \mathcal{B}_{p,q}^s \cap L^\infty$ ,*

$$|fg|_{\mathcal{B}_{p,q}^s} + |fg|_\infty \leq C (|f|_{\mathcal{B}_{p,q}^s} + |f|_\infty) (|g|_{\mathcal{B}_{p,q}^s} + |g|_\infty).$$

*In particular  $\mathcal{B}_{p,q}^s \cap L^\infty$ , with the norm  $|\cdot|_{\mathcal{B}_{p,q}^s} + |\cdot|_\infty$ , is a quasi-algebra.*

**Formalization.** In `GoodGrid/QuasiAlgebra.lean` (paper Proposition 19.1, named `mult33` in the source). The outer statement is `souzaPointwiseMultipliersIII`: concrete functions  $f, g$  represent Souza-Besov elements and carry a.e. bounds  $M_f, M_g$ , and the conclusion produces a representative  $y$  of the product with  $|y|_{\mathcal{B}} + M_f M_g \leq C (|f|_{\mathcal{B}} + M_f) (|g|_{\mathcal{B}} + M_g)$ , together with the a.e. bound  $\|fg\| \leq M_f M_g$  (the elementary  $L^\infty$  part, `ae_norm_mul_le_mul_bounds`). The technical core is `exists_quasiAlgebra_product_representation`, with the bilinear mixed bound  $C (|f|_{\mathcal{B}} M_g + |g|_{\mathcal{B}} M_f)$ : the  $u_1 + u_2$  construction of Section 19.8 is run for *two*  $(s, p)$  Souza representations — the role played in Proposition 19.11 by the tower sums of the  $(1/p, p)$  representation of  $g$  is now played by the *weighted* and *strict weighted* ancestor towers (`weightedAncestorCoeffSum`: every ancestor contributes its coefficient times the local value  $m(J)^{s-1/p}$  of the canonical atom), a weighted `fou` representation (`exists_weighted_fouRepresentation`) is extracted from the standard-representation machinery and Proposition 17.1.B, the two triangular block families are built directly with costs controlled by the towers, and the finite truncation identity is passed to the limit in  $L^p$  as in Section 19.8. Axioms of `souzaPointwiseMultipliersIII` and `exists_quasiAlgebra_product_representation` checked: `propext`, `Classical.choice`, `Quot.sound`.

```

theorem souzaPointwiseMultipliersIII
  (G : GoodGridSpace (α := α)) (s : ℝ) (p q : ℝ ≥ 0 ∞)
  (hs : 0 < s) (hs_lt_inv : s < (p.toReal)-1)
  (hp : 1 ≤ p) (hp_top : p ≠ ∞)
  [Fact (1 ≤ p)] [Fact (1 ≤ q)] :

```

```

    ∃ Cqa : ℝ,
    0 ≤ Cqa ∧
    ∀ (f g : α → ℂ) (Mf Mg : ℝ)
      (xf xg : WeakGridSpace.BesovishSpace
        (souzaAtomFamily G s p hs hp hp_top) q),
    WeakGridSpace.RepresentsFunction
      (G := G.toWeakGridSpace) (p := p) f
      (xf : Lp ℂ p G.toWeakGridSpace.measure) →
    WeakGridSpace.RepresentsFunction
      (G := G.toWeakGridSpace) (p := p) g
      (xg : Lp ℂ p G.toWeakGridSpace.measure) →
    (∀m z ∂G.toWeakGridSpace.measure, ‖f z‖ ≤ Mf) →
    (∀m z ∂G.toWeakGridSpace.measure, ‖g z‖ ≤ Mg) →
    ∃ y : WeakGridSpace.BesovishSpace
      (souzaAtomFamily G s p hs hp hp_top) q,
    WeakGridSpace.RepresentsFunction
      (G := G.toWeakGridSpace) (p := p)
      (fun z => f z * g z)
      (y : Lp ℂ p G.toWeakGridSpace.measure) ∧
    WeakGridSpace.BesovishSpace.Norm_Costpq
      (souzaAtomFamily G s p hs hp hp_top) q y + Mf * Mg ≤
    Cqa *
    ((WeakGridSpace.BesovishSpace.Norm_Costpq
      (souzaAtomFamily G s p hs hp hp_top) q xf + Mf) *
      (WeakGridSpace.BesovishSpace.Norm_Costpq
      (souzaAtomFamily G s p hs hp hp_top) q xg + Mg)) ∧
    (∀m z ∂G.toWeakGridSpace.measure, ‖f z * g z‖ ≤ Mf * Mg) := ...

```

## 22. A REMARKABLE DESCRIPTION OF $\mathcal{B}_{1,1}^s$

**File:** GoodGrid/AlternativeDescriptionBs11.lean

**Imports (within the project):** GoodGrid/RegularDomains.lean

This section documents the formalization of paper Section 20, labelled `rema` in the source. At the endpoint  $p = q = 1$ , and for  $0 < s < 1$ , the Souza–Besov space can be described without using arbitrary Souza atoms: one may use only normalized indicators of regular domains. Fix constants  $C \geq 0$  and  $0 \leq c < 1$ , let  $\mathcal{W}$  be the family of all  $(1 - s, C, c)$ -regular domains, and choose an admissible family  $\widehat{\mathcal{W}}$  satisfying

$$\mathcal{P} \subset \widehat{\mathcal{W}} \subset \mathcal{W}.$$

The alternative space consists of functions admitting a representation

$$(9) \quad f = \sum_{i=0}^{\infty} c_i \frac{\mathbb{1}_{\Omega_i}}{m(\Omega_i)^{1-s}}, \quad \Omega_i \in \widehat{\mathcal{W}}, \quad \sum_i |c_i| < \infty,$$

with the gauge given by the infimum of  $\sum_i |c_i|$  over all such representations. The formal statement uses concrete representatives  $f: I \rightarrow \mathbb{C}$  for the domain-indicator side and the existing subtype `SouzaBesov11` for the Souza side;

equality of spaces is therefore stated as a two-way representative theorem with two norm/gauge estimates.

**Proposition 22.1** (Regular-domain description of  $\mathcal{B}_{1,1}^s$ ). *The alternative regular-domain space defined by (9) and Souza’s space  $\mathcal{B}_{1,1}^s$  have the same concrete representatives, and their gauges are equivalent. More precisely, every regular-domain representation gives a Souza element with cost bounded by a constant multiple of the domain gauge, and every Souza element gives a regular-domain representation with domain gauge bounded by the Souza cost.*

**Formalization.** In `GoodGrid/AlternativeDescriptionBs11.lean`. The endpoint exponent  $1/(1-s)$  is `domainEndpointExponent`; the normalized indicator  $m(\Omega)^{s-1}\mathbb{1}_\Omega$  is `normalizedDomainIndicator`; the condition  $\mathcal{P} \subset \widehat{\mathcal{W}} \subset \mathcal{W}$  is `AdmissibleDomainClass`. A representation of the form (9) is `DomainAtomicRepresentation`; membership in the alternative space is `DomainBesovSpace`; and the infimum gauge is `domainBesovGauge`. The first inclusion is `domainBesovSpace_to_souzaBesov11`, using `regularDomainIndicator_besov_norm_bound_all` from Section 20. The reverse inclusion is `souzaBesov11_to_domainBesovSpace`: a Souza representation is flattened over the encodable set of grid cells, and the private coefficient estimate `souzaDomainFlattenedCoeff_summable_norm_and_tsum_le` keeps the  $\ell^1$  cost bounded by the original Souza representation cost. The final packaged equivalence is `domainBesovSpace_equiv_souzaBesov11`. Axioms checked for this theorem: `propext`, `Classical.choice`, `Quot.sound`.

```

structure DomainAtomicRepresentation
  (G : GoodGridSpace (α := α)) (s : ℝ) (What : Set (Set α))
  [Fact (1 ≤ domainEndpointExponent s)] (f : α → ℂ) where
  domain : ℕ → Set α
  domain_mem : ∀ i, domain i ∈ What
  coeff : ℕ → ℂ
  memLp : MemLp f (domainEndpointExponent s) G.grid.μ
  term_memLp :
    ∀ i, MemLp
      (fun z => coeff i * normalizedDomainIndicator G s (domain i) z)
      (domainEndpointExponent s) G.grid.μ
  hasSum :
    HasSum
      (fun i =>
        MemLp.toLp
          (fun z => coeff i * normalizedDomainIndicator G s (domain i) z)
          (term_memLp i))
      (MemLp.toLp f memLp)
  coeff_summable : Summable fun i => ||coeff i||

theorem domainBesovSpace_equiv_souzaBesov11
  (G : GoodGridSpace (α := α)) (s C c : ℝ) (What : Set (Set α))
  [Fact (1 ≤ domainEndpointExponent s)]
  (hs : 0 < s) (hs_lt_one : s < 1)
  (hW : AdmissibleDomainClass G s C c What) :
  (∃ C₀ : ℝ,

```

```

0 ≤ C₀ ∧
∀ f : α → ℂ,
  DomainBesovSpace G s What f →
    ∃ x : SouzaBesov11 G s hs,
      WeakGridSpace.RepresentsFunction (p := (1 : ℝ≥0∞))
        f (x : Lp ℂ (1 : ℝ≥0∞) G.toWeakGridSpace.measure) ∧
      WeakGridSpace.BesovishSpace.Norm_Costpq
        (souzaAtomFamily G s (1 : ℝ≥0∞) hs le_rfl ENNReal.one_ne_top)
        (1 : ℝ≥0∞) x ≤
        C₀ * domainBesovGauge G s What f) ∧
(∀ x : SouzaBesov11 G s hs,
  DomainBesovSpace G s What
    ((x : Lp ℂ (1 : ℝ≥0∞) G.toWeakGridSpace.measure) : α → ℂ) ∧
  domainBesovGauge G s What
    ((x : Lp ℂ (1 : ℝ≥0∞) G.toWeakGridSpace.measure) : α → ℂ) ≤
  WeakGridSpace.BesovishSpace.Norm_Costpq
    (souzaAtomFamily G s (1 : ℝ≥0∞) hs le_rfl ENNReal.one_ne_top)
    (1 : ℝ≥0∞) x) := ...

```

### 23. LEFT COMPOSITIONS

**File:** GoodGrid/LeftCompositions.lean  
**Imports (within the project):** GoodGrid/  
 AlternativeRepresentationsAndNorms/MeanOscillationNorm.lean

This section records the incorporated part of paper Section 21, labelled **expo** in the source. The paper's result says that a Lipschitz map  $\psi : \mathbb{C} \rightarrow \mathbb{C}$  fixing the origin acts on the left of the Besov space:  $f \mapsto \psi \circ f$  is bounded. The Lean file currently formalizes the analytic estimate in the mean-oscillation model. If  $\psi$  is  $K$ -Lipschitz and  $\psi(0) = 0$ , then the  $L^p$  term and the oscillation term of  $\psi \circ f$  are each bounded by  $K$  times the corresponding term for  $f$ . The proof follows the short argument in the manuscript: for a cell  $Q$ , test the oscillation of  $\psi \circ f$  against the constants  $\psi(c)$ , and use

$$|\psi(f(x)) - \psi(c)| \leq K|f(x) - c|.$$

**Proposition 23.1** (Left composition, mean-oscillation form). *Let  $G$  be a good grid, let  $\psi : \mathbb{C} \rightarrow \mathbb{C}$  be  $K$ -Lipschitz with  $\psi(0) = 0$ , and assume  $p > 0$  and either  $q = \infty$  or  $q > 0$ . Then*

$$\|\psi \circ f\|_{\text{MO}(s,p,q)} \leq K \|f\|_{\text{MO}(s,p,q)}.$$

*In particular, every local oscillation, every level block, the oscillation seminorm, and the full mean-oscillation gauge are stable under left composition by  $\psi$ .*

**Formalization.** In GoodGrid/LeftCompositions.lean. The elementary  $L^p$  estimate is `eLpNorm_comp_le_of_lipschitzWith`; the restricted estimate around a constant is `eLpNorm_comp_sub_const_le_of_lipschitzWith`. These give the cell

oscillation estimate `osc_comp_le_of_lipschitzWith`. Summing over all level- $k$  cells gives `levelOscillationBlock_comp_le_of_lipschitzWith`; taking the  $1/p$ -root gives `levelOscillationBlock_root_comp_le_of_lipschitzWith`. The finite  $q$  and  $q = \infty$  cases are packaged together in `oscillationSeminorm_comp_le_of_lipschitzWith`, and the final  $L^p$ -plus-oscillation estimate is `meanOscillationNorm_comp_le_of_lipschitzWith`. This is marked as a partial formalization of paper Proposition `expo`: the remaining work is to transport/package the mean-oscillation estimate as the full Souza–Besov left-composition theorem.

```

theorem osc_comp_le_of_lipschitzWith
  (G : GoodGridSpace (α := α)) {K : NNReal} {g : ℂ → ℂ}
  (hg : LipschitzWith K g) (p : ℝ ≥ 0∞) (f : α → ℂ) (Q : GoodGridCell G) :
  osc G p (fun x => g (f x)) Q ≤ (K : ℝ ≥ 0∞) * osc G p f Q := ...

theorem oscillationSeminorm_comp_le_of_lipschitzWith
  (G : GoodGridSpace (α := α)) {K : NNReal} {g : ℂ → ℂ}
  (hg : LipschitzWith K g) (s : ℝ) (p q : ℝ ≥ 0∞)
  (hp : 0 < p.toReal) (hq : q ≠ ∞ → 0 < q.toReal) (f : α → ℂ) :
  oscillationSeminorm G s p q (fun x => g (f x)) ≤
    (K : ℝ ≥ 0∞) * oscillationSeminorm G s p q f := ...

theorem meanOscillationNorm_comp_le_of_lipschitzWith
  (G : GoodGridSpace (α := α)) {K : NNReal} {g : ℂ → ℂ}
  (hg : LipschitzWith K g) (hg_zero : g 0 = 0)
  (s : ℝ) (p q : ℝ ≥ 0∞)
  (hp : 0 < p.toReal) (hq : q ≠ ∞ → 0 < q.toReal) (f : α → ℂ) :
  meanOscillationNorm G s p q (fun x => g (f x)) ≤
    (K : ℝ ≥ 0∞) * meanOscillationNorm G s p q f := ...

```

### III. GUIDE TO THE REPOSITORY

#### 24. GUIDE TO THE REPOSITORY FILES

All paths below are relative to the library root `BesovSpacesGoodGrid/`; the aggregate file `BesovSpacesGoodGrid.lean` is the main entry point and imports the whole pipeline.

##### 24.1. Weak-grid layer (Part I).

- `WeakGrid/Definition.lean`: weak-grid structures (`WeakGrid`, `WeakGridSpace`) and the overlap estimates of Section 3.
- `WeakGrid/Atoms.lean`: weak-grid cells, local spaces, and atom families (Section 4).
- `WeakGrid/BesovishSpaces.lean`: level blocks, representations, costs, Besov-ish spaces, the cost gauge `Norm_Costpq`, the  $L^t$  embedding and the normed structure (Section 5).

- `WeakGrid/Scales.lean`: scaled atom families and the smoothness-scale inclusion (Section 7).
- `WeakGrid/Completeness.lean`: representation limits, compactness of cost balls, and completeness (Section 6).
- `WeakGrid/InducedGrid.lean`: induced weak grids on a fixed cell and the contraction into the ambient Besov-ish space (Section 8).
- `WeakGrid/Transmutation.lean`: transmutation objects and Claims I, II, III, including the explicit Claim C embedding constants (Section 9).
- `WeakGrid/Multipliers.lean`: the abstract multiplier and self-class predicates on weak grids (Section 10).
- `Sums.lean`: block-index types (`BlockIndex`, `ChildIndex`, `ParentIndex`) and block-sum notation used throughout (Section 10).

#### 24.2. Good-grid layer (Part II).

- `GoodGrid/Definition.lean`: good-grid structures (Section 11).
- `GoodGrid/BesovSpace.lean`: Souza’s atoms and good-grid Besov spaces, with completeness, compactness, and density (Sections 12 and 13).
- `GoodGrid/PositiveCone.lean`: the positive cone, positive and cone-positive representations, decompositions, density, and the support theorems (Section 17).
- `GoodGrid/DiracApproximations.lean`: the grid Dirac kernels, the evaluation of partial standard sums as cell averages, and the  $L^\infty$  bounds on ancestor coefficient sums of paper Proposition 17.1 (Section 18).
- `GoodGrid/BesovAtoms.lean`: Besov’s atoms on good grids and the Souza/Besov sandwich theorem (Section 14).
- `GoodGrid/AlternativeRepresentationsAndNorms.lean` and `GoodGrid/AlternativeRepresentationsAndNorms/`: the norm-comparison layer of Section 16 — `HaarRepresentationNorm.lean` (normalized Haar coefficients and the Haar gauge), `HaarParametrizedRepresentation.lean` and `ComparingHaarRepresentations1.lean` (Haar expansions as parametrized representations and their comparison), `standardRepresentation.lean` (standard atomic coefficients and gauge), `standardNormleqHaarRepresentenstionNorm.lean` (Haar controls standard), `StandarRepresentationNormleqBesovNorm.lean` (the Besov norm controls the standard norm), `FiniteStandardNormimpliesBesov.lean` (finite standard norm endpoint), `FiniteHaarNormimpliesLp.lean` (finite Haar norm endpoint), `MeanOscillationNorm.lean`

(mean-oscillation definitions and local oscillation lemmas), `OscillationNormleqBesovNorm.lean` (Besov controls mean oscillation), and `HaarNormleqOscillationNorm.lean` (mean oscillation controls Haar).

- `GoodGrid/Distribution.lean`: cell indicators (`atomIndicator`), the test-function module (`TestFunctions`), and its algebraic dual (`Distributions`).

### 24.3. Applications and multiplier layer (Part II).

- `GoodGrid/Multipliers.lean` and `GoodGrid/Multipliers/`: the pointwise-multiplier layer of Section 19 — `Definition.lean` (the `selfs` classes and seminorms), `Besovspq.lean` (induced-cell representations, restriction to a cell, and the general inclusion  $M(\mathcal{B}_{p,q}^s \subset \mathcal{B}_{p,q, selfs}^s)$ ), `Besovs11.lean` (the  $\mathcal{B}_{1,1}^s$  characterization), `MultipliersareBounded.lean` (boundedness of `selfs` multipliers, uniform in the tail level, and the canonical tail classes), `NonArchimedeanProperty.lean` (the finite and infinite non-Archimedean estimates and the cores of their positive versions), `NonArchimedeanPropertyPositiveStandalone.lean` (the user-facing positive-cone statements, finite and infinite), `SelfsSubsetMultipliers.lean` (the continuous inclusion of the tail `selfs` classes in the multiplier space, paper Corollary 18.6), `StronglyRegularDomains.lean` (strongly regular domains and Pointwise Multipliers I, paper 18.7–18.9), and `Bp1overpinfityisMultiplier.lean` (Pointwise Multipliers II, paper Proposition 18.10, together with its positive-cone version, Remark 19.12).
- `GoodGrid/QuasiAlgebra.lean`: the quasi-algebra property of  $\mathcal{B}_{p,q}^s \cap L^\infty$ , Pointwise Multipliers III (Section 21).
- `GoodGrid/RegularDomains.lean`: regular families and regular domains (paper subsection cf, Section 20) — the definitions and the union construction, the strongly-regular  $\Rightarrow$  regular comparison, the indicator Besov/multiplier bounds `estG`, and the localized restriction representation estimate `pdd/hiip1` (`regularFamily_restriction_representations`).
- `GoodGrid/AlternativeDescriptionBs11.lean`: the regular-domain indicator-series description of  $\mathcal{B}_{1,1}^s$  from paper Section 20, with the final packaged equivalence `domainBesovSpace_equiv_souzaBesov11` (Section 22).
- `GoodGrid/LeftCompositions.lean`: partial Section 21 infrastructure for left compositions; it proves the  $L^p$ , restricted distance-to-constants, cell oscillation, level-block, oscillation seminorm, and full

mean-oscillation gauge estimates for Lipschitz maps fixing zero (Section 23).

**24.4. External dependencies.** The repository depends on the external Lean repositories `UnbalancedHaarWavelet` (grids and unbalanced Haar systems), `UnconditionalSchauderBasis`, `LaminarFamiliesMaximalBinaryTrees` and `Burkholder` (all by SmaniaD), which provide the Girardi–Sweldens construction and the unconditional basis property used in Section 15.

**24.5. Repository status.** At the date of this document (June 17, 2026) a full lake build succeeds (3462 jobs) and the project Lean files contain no `sorry`, no `admit`, and no project-local `axiom` or `constant` declarations; every project module, including the multiplier and quasi-algebra files, `AlternativeDescriptionBs11.lean`, and `LeftCompositions.lean`, is imported by the aggregate root `BesovSpacesGoodGrid.lean`. The axioms of the main theorems — the finite and infinite, non-positive and positive non-Archimedean properties and their consequences, namely

```
souzaNonArchimedeanProperty,
souzaNonArchimedeanPropertyLambdaFinite,
souzaNonArchimedeanPropertyPositiveCone,
souzaNonArchimedeanPropertyPositiveConeInfinite,
exists_souzaSelfsMultiplierConstant, souzaPointwiseMultipliersI,
souzaPointwiseMultipliersIInfinite, claimA_standard, claimA_
positive,
claimB, exists_fouRepresentation, souzaPointwiseMultipliersII,
souzaPointwiseMultipliersIIPositive, souzaPointwiseMultipliersIII,
regularFamily_restriction_representations, domainBesovSpace_equiv_
souzaBesov11
```

— were checked with `#print axioms`: only the standard axioms `propext`, `Classical.choice`, `Quot.sound` appear (no `sorryAx`). The running verification log is kept in `status.md` at the repository root.

## REFERENCES

- [1] D. Smania, *Besov-ish spaces through atomic decomposition*, Analysis & PDE **15** (2022), no. 1, 123–174. A prerelease version of the published paper is included in this repository as `preprint-source/besovish-2020-07-18.tex` (with PDF); it is the source from which the present document was produced.
- [2] M. Girardi and W. Sweldens, *A new class of unbalanced Haar wavelets that form an unconditional basis for  $L_p$  on general measure spaces*, J. Fourier Anal. Appl. **3** (1997), 457–474.
- [3] G. S. de Souza, *Spaces formed by special atoms*, PhD thesis, SUNY Albany, 1980.
- [4] D. Smania, *BesovSpacesGoodGrid: a Lean 4 formalization*, git repository, 2026. Dependencies: `UnbalancedHaarWavelet`, `UnconditionalSchauderBasis`, `LaminarFamiliesMaximalBinaryTrees`, `Burkholder` (SmaniaD).

DEPARTAMENTO DE MATEMÁTICA, INSTITUTO DE CIÊNCIAS MATEMÁTICAS E DE  
COMPUTAÇÃO-UNIVERSIDADE DE SÃO PAULO (ICMC/USP) - SÃO CARLOS, CAIXA  
POSTAL 668, SÃO CARLOS-SP, CEP 13560-970, BRAZIL.

*Email address:* `smania@icmc.usp.br`

*URL:* <https://sites.icmc.usp.br/smania/>