

DEMO-001 — Submission

LaTeX response — synthetic student

Q1 — Calculus

5 / 5

Let $u = x^2$, so $du = 2x \, dx$. Therefore

$$I = \frac{1}{2} \int_0^1 e^u \, du = \frac{1}{2} [e^u]_0^1 = \frac{e - 1}{2}.$$

Q2 — Mechanics

8 / 8

The normal force is $N = mg \cos 30^\circ$. Along the slope,

$$ma = mg \sin 30^\circ - \mu_k N, \quad a = 3.20 \, \text{m s}^{-2}.$$

Starting from rest, $v = \sqrt{2as} = 4.38 \, \text{m s}^{-1}$.

DEMO-001 — Submission

LaTeX response — synthetic student

Q3 — Linear algebra

7 / 7

If $Ax = 0$, then $A^T Ax = 0$. Conversely,

$$x^T A^T Ax = \|Ax\|^2 = 0,$$

hence $Ax = 0$. Rank-nullity completes the argument.

Q4 — Programming

6 / 10

```
import math
def stable_softmax(xs):
    if not xs: return []
    exps = [math.exp(x) for x in xs]
    return [value / sum(exps) for value in exps]
```

REVIEW SIGNAL — extreme values overflow because $\max(xs)$ is not subtracted.