

# CTRM STATISTICS

ABSTRACT. We fit a max-renewal process, (aka 'CTRM', Continuous Time Random Maxima) to data.

## 1. INTRODUCTION

## 2. THE MODEL

The stochastic model  $X(t)$  below has been studied under the various names “Shock process” [EM73, SS83, SS84, SS85, And87, Gut99], “max-renewal process”, [Sil04, ST04, BŠ14] and “CTRM (continuous time random maxima) [BSM07, MS09, Hee14, HS15]: Assume i.i.d. pairs of random variables  $(W_k, J_k)$ ,  $k = 1, 2, \dots$  where  $W_k > 0$  represents an inter-arrival time of certain events and  $J_k \in \mathbb{R}$  the corresponding event magnitude. Write  $N(t) = \max\{n \in \mathbb{N} : W_1 + \dots + W_n \leq t\}$  for the renewal process associated with the  $W_k$  (where the maximum of the empty set is set to 0), and define

$$X(t) = \bigvee_{k=1}^{N(t)} J_k = \max\{J_k : k = 1, \dots, N(t)\}.$$

If  $W_k$  is interpreted as the time leading up to the event with magnitude  $J_k$ , then  $X(t)$  is the largest magnitude up to time  $t$ . The alternative case where  $W_k$  is the inter-arrival time *following*  $J_k$  is termed “second type” (in the shock model literature) or OCTRM (overshooting CTRM), and the largest magnitude up to time  $t$  is then given by

$$Y(t) = \bigvee_{k=1}^{N(t)+1} J_k.$$

Finally, the model is called coupled when  $W_k$  and  $J_k$  are not independent; in this article we focus on the uncoupled case. For this case,  $X(t)$  and  $Y(t)$  have the same limiting behaviour for large times and magnitudes [HS15] and hence we focus on the CTRM  $X(t)$ .

Any probability distribution with cumulative distribution function (CDF)  $F$  on  $\mathbb{R}$  lies in the max-domain of attraction of a generalized extreme value (GEV) distribution with CDF  $\Lambda$ . That is, there exist sequences  $a_n \in \mathbb{R}^+$  and  $b_n \in \mathbb{R}$  such that

$$F(a_n x + b_n) \rightarrow \Lambda(x), \quad x \in (x_{\min}, x_{\max}).$$

## REFERENCES

- [And87] KK Anderson. Limit Theorems for General Shock Models with Infinite Mean Intershock Times. *J. Appl. Probab.*, 24(2):449–456, 1987.

- [BŠ14] Bojan Basrak and Drago Špoljarić. Extremal behaviour of random variables observed in renewal times. 2014.
- [BSM07] David a. Benson, Rina Schumer, and Mark M. Meerschaert. Recurrence of extreme events with power-law interarrival times. *Geophys. Res. Lett.*, 34(16):1–5, aug 2007.
- [EM73] J. D. Esary and A. W. Marshall. Shock Models and Wear Processes, 1973.
- [Gut99] Allan Gut. Extreme Shock Models. *Extremes*, (1983):295–307, 1999.
- [Hee14] Katharina Hees. *Gemeinsame Konvergenz von Summe und Maximum und Extremwertverhalten von gekoppelten Continuous Time Random Walks*. PhD thesis, Universität Siegen, 2014.
- [HS15] Katharina Hees and Hans-peter Scheffler. Coupled continuous time random maxima. pages 1–24, 2015.
- [MS09] MM Meerschaert and SA Stoev. Extremal Limit Theorems for observations separated by random waiting times. *J. Stat. Plan. Inference*, 139(7):2175–2188, 2009.
- [Sil04] Dmitrii S Silvestrov. *Limit Theorems for Randomly Stopped Stochastic Processes*. Springer (Berlin, Heidelberg), 2004.
- [SS83] JG Shanthikumar and U Sumita. General shock models associated with correlated renewal sequences. *J. Appl. Probab.*, 20(3):600–614, 1983.
- [SS84] JG Shanthikumar and Ushio Sumita. Distribution Properties of the System Failure Time in a General Shock Model. *Adv. Appl. Probab.*, 16(2):363–377, 1984.
- [SS85] J G Shanthikumar and U Sumita. A class of correlated cumulative shock models. *Adv. Appl. Probab.*, 17(2):347–366, 1985.
- [ST04] Dmitrii S. Silvestrov and Jozef L. Teugels. Limit theorems for mixed max-sum processes with renewal stopping. *Ann. Appl. Probab.*, 14(4):1838–1868, nov 2004.

PETER STRAKA, SCHOOL OF MATHEMATICS AND STATISTICS, UNSW AUSTRALIA, SYDNEY, NSW 2052, AUSTRALIA

*E-mail address:* p.straka@unsw.edu.au