

Thesis

Smarak Nayak - z3415339

July 2016

Distribution of duration

Let F be the cdf of a gev random variable (write this properly). Let a be the threshold level, then define T_a as the duration between jumps that have been thresholded(?) at the a -level. From (insert reference here) we know that (explain proof here?)

$$T_a \sim (-\log F(a))^{-\frac{1}{\beta}} X^{\frac{1}{\beta}} D(1).$$

Where $D(1)$ is a random variable of the stable distribution with stability parameter β and skewness parameter 1 and where X is a standard exponential random variable. After taking the logarithms of both sides we arrive at

$$\log T_a \sim \frac{1}{\beta} \log X + \log D(1) - \frac{1}{\beta} \log(-\log F(a)).$$

In order to find the distribution for $\log T_a$ we will first need to find the distributions of $\frac{1}{\beta} \log X$ and $\log D$. We have that

$$\begin{aligned} f_{\frac{1}{\beta} \log X}(x) &= \frac{d}{dx} \mathbb{P}\left(\frac{1}{\beta} \log X \leq x\right) \\ &= \frac{d}{dx} \mathbb{P}(X \leq e^{x\beta}) \\ &= f_X(e^{x\beta}) \beta e^{x\beta}. \end{aligned} \tag{a}$$

Similarly we have

$$\begin{aligned} f_{\log D}(x) &= \frac{d}{dx} \mathbb{P}(\log D \leq x) \\ &= \frac{d}{dx} \mathbb{P}(D \leq e^x) \\ &= f_D(e^x) e^x. \end{aligned} \tag{b}$$

Using convolution we know that

$$f_{\frac{1}{\beta} \log X + \log D}(x) = \int_{-\infty}^{\infty} f_{\frac{1}{\beta} \log X}(x-y) f_{\log D}(y) dy$$

and thus

$$f_{\log T_a}(x) = \int_{-\infty}^{\infty} f_{\frac{1}{\beta} \log X}\left(x-y + \frac{1}{\beta} \log(-\log F(a))\right) f_{\log D}(y) dy \tag{1}$$

as $\frac{1}{\beta} \log(-F(a))$ is a constant with respect to x . Substituting (a) and (b) into (1) we get

$$f_{\log T_a}(x) = \int_{-\infty}^{\infty} f_X(e^{\beta(x-y+\frac{1}{\beta} \log(-\log F(a)))}) \beta e^{\beta(x-y+\frac{1}{\beta} \log(-\log F(a)))} f_D(e^y) e^y dy$$

which simplifies to

$$f_{\log T_a}(x) = \int_{-\infty}^{\infty} -\log F(a) f_X(-\log F(a) e^{\beta(x-y)}) \beta e^{\beta(x-y)} f_D(e^y) e^y dy.$$