

GrapeMaster: A Crop-Season-Centered Digital Platform for Traceable Vineyard Disease Management

Author information to be completed before submission

Affiliations, author order, and corresponding author details will be completed before journal submission.

Abstract

Grapevine is a high-value fruit crop. In humid subtropical vineyards, epidemic diseases, particularly downy mildew, remain major constraints on yield, fruit quality, and production cost. Disease simulation, warning, diagnosis, and advisory assistance have each advanced substantially, but their outputs are often developed, validated, and adopted as separate functions, constraining their practical value for disease management. This limitation arises because vineyard disease management must respond to interactions among phenological susceptibility, infection risk, symptom diagnosis, epidemic development, fungicide application, and protection status throughout the growing season, rather than to isolated single-event outputs. To address this limitation, we developed GrapeMaster, a vineyard-level disease-management platform that uses the crop season as the core organizational unit to link daily disease pressure, plant-protection operations, and task feedback within a dynamic seasonal process. A backend-record audit showed that 126 crop seasons retained linked weather, phenology, risk, notification, task, fungicide-feedback, protection-state updating, advisory assistance, and message records. A representative season replay demonstrated the platform's capacity to construct a prediction–notification–task–treatment–feedback–review workflow. Module-level validation further supported the analytical readiness of downy mildew warning, broad-stage phenology display, and disease image recognition for the Guangxi context. These results indicate that GrapeMaster offers a deployable platform architecture for converting isolated digital tools into a closed-loop and traceable management system, with potential value for addressing complex production environments and improving the precision, standardization, and digital transformation of vineyard management.

Keywords: grapevine; downy mildew; decision support system; crop-season management; digital agriculture; advisory support; record traceability

1 Introduction

Grapevine is an important economic crop for fresh consumption, wine production, and processing. In humid subtropical vineyards, high humidity, rainfall, dew formation, and heterogeneous field microclimates readily favor epidemic diseases represented by downy mildew (Gessler et al., 2011; Yu et al., 2017, 2022; Peng et al., 2024). In severe cases, this disease can cause more than 75% yield loss (Koledenkova et al., 2022; Peng et al., 2024). However, conventional plant-protection practice often depends on fragmented field scouting, empirical spray calendars, manual symptom diagnosis, informal advisory sources with uneven reliability, and retrospective paper records. This non-integrated and weakly standardized decision process can lead to mismatches between spray timing and disease-risk periods, inappropriate pesticide concentration or mixing, and control outcomes that are difficult to trace and quantitatively evaluate (Pertot et al., 2017; Rossi et al., 2014; Oliver et al., 2023; Wise et al., 2010).

Several analytical technologies relevant to vineyard disease management have advanced substantially. Grapevine downy mildew models have progressed from empirical warning rules to mechanistic or semi-mechanistic descriptions of pathogen development and infection processes (Orlandini et al., 1993; Park et al., 1997; Caffi et al., 2011b), and warning systems can improve treatment timing when meteorological inputs, local calibration, and implementation context are adequate (Caffi et al., 2010; Caffi and Rossi, 2018; Carisse, 2016). Grapevine phenology models provide crop-stage information for interpreting host susceptibility, warning windows, canopy development, and operation timing (Caffarra and Eccel, 2010; Parker et al., 2011; García de Cortázar-Atauri et al., 2017; Wang et al., 2020). Image-recognition and advisory systems can convert visible symptoms into structured diagnostic information and management-oriented explanation (Yan et al., 2024, 2025; Shafay et al., 2026; Rai et al., 2026). These technologies have substantial value in different settings. The challenge for extension and industrial deployment is how to coordinate their separate

53 outputs within a shared seasonal management context.

54 More broadly, digital agriculture and decision support system (DSS) implementation studies in-
55 creasingly show that production deployment is not determined by model performance alone. De-
56 ployed agricultural DSS platforms have begun to combine monitoring, analytical models, advisory
57 delivery, and user-facing workflows. In viticulture, vite.net addressed the implementation problem
58 in vineyard DSSs through integrated vineyard monitoring, model-based decision support, and two-
59 way communication (Rossi et al., 2014), whereas MISFITS-DSS harmonized regional grapevine
60 downy mildew risk assessment for public plant-protection services (Bregaglio et al., 2022). Be-
61 yond viticulture, Fuxi Brain illustrated a whole-cycle intelligent management platform based on IoT
62 perception, multimodal fusion, and generative decision making in maize production (Chen et al.,
63 2026). Together, these examples indicate that DSSs intended for production use must move beyond
64 model output alone and require stable data ingestion, modular architecture, reliable service opera-
65 tion, user-facing task processes, feedback mechanisms, record tracking, workflow evaluation, and
66 traceability. However, for perennial fruit-crop production contexts, they also expose a platform-
67 level architectural problem: how analytical outputs can be organized into a crop-season-centered
68 workflow so that disease-management decisions can be delivered, recorded, executed, updated, and
69 reviewed within a traceable seasonal frame.

70 Therefore, for grape production, the unresolved problem is not only whether individual models or
71 diagnostic tools can perform accurately, but whether their outputs can be organized into a verifiable
72 seasonal decision chain. Otherwise, after a warning is generated, the system may not be able to trace
73 whether the warning was delivered, reviewed, converted into an operation, reflected in subsequent
74 protection status, and used as the updated management context for subsequent risk assessment and
75 evaluation.

76 To address this gap, we developed and deployed GrapeMaster, a closed-loop and traceable vineyard
77 disease-management platform that uses the crop season as the core organizational unit. Here, the
78 crop season denotes the annual production cycle from budburst to fruit maturity and harvest, and
79 is used as a shared unit for storing vineyard and crop context and linking crop development, risk

interpretation, warning delivery, task execution and feedback, diagnostic and advisory records, and message storage. The central research question was whether this platform-implemented architecture can convert independent analytical and modular functions into a traceable workflow linking prediction, action, feedback, symptom or advisory records, and review across the crop season. We evaluated this question through three methodological layers: the crop-season data structure and database relationships that implement this management unit, workflow constraints and state-transition logic, and backend-record audit using linkage checks and representative season replay. GrapeMaster provides a platform-level pathway and methodological basis for consistent, standardized, traceable, and evaluable disease management in perennial fruit production.

2 Materials and methods

2.1 Study design and evaluation data

The methodological objective of this study was to construct a traceability data architecture that uses the crop season as the core organizational unit, implement a closed workflow linking risk notification, plant-protection task creation, fungicide protection, task feedback, and risk updating, embed platform-compatible analytical modules, and check whether the deployed backend retained the record objects and unique identifiers required for workflow reconstruction.

The study used two complementary data inputs. Backend records from the deployed platform were used for the backend-record audit and representative replay. Field and image data were used to configure and evaluate the platform-compatible analytical modules. The two data sources were analyzed separately. The module-evaluation datasets are listed in Table 1, and additional module-level technical details are provided in Supplementary Table S1.

Table 1. Field and image datasets used for platform-compatible module evaluation.

Site or dataset	Cultivar or scope	Years	Data types
Mingyang, Nanning	Jufeng	2023, 2024	Phenology observations, downy mildew monitoring, airborne sporangia monitoring, and management records
Du'an, Hechi	Maoputao Yenang No. 2	2022, 2023, 2024	Phenology observations, downy mildew monitoring, airborne sporangia monitoring, and management records
Xing'an, Guilin	Wenke	2022	Phenology observations, downy mildew monitoring, airborne sporangia monitoring, and management records
Grape disease image dataset	Multiple grape disease categories	Not site-year based	Symptom images

2.2 Platform architecture and service layers

GrapeMaster uses a layered architecture in which the mobile client, backend services, analytical model services, persistent storage, and external data sources are separated into logical service layers. The service layers are listed in Table 2. Detailed software dependencies and implementation components are listed in Supplementary Table S6.

Table 2. Logical architecture layers of GrapeMaster used in the current study.

Layer	Main components	Main responsibilities
Application and presentation layer	Mobile interfaces for field setup, crop-season workspace, warnings, tasks, image upload, consultation, map display, and notifications	User input, crop-season state display, warning review, plant-protection task operation, symptom-image submission, and advisory interaction
Backend service and storage layer	API services, authentication, persistent storage, scheduled jobs, master-data tables, and business objects	API routing, access control, data persistence, scheduled updates, field and crop-season records, and operational record storage
Analytical input and risk-processing layer	Weather preprocessing, phenology workflow, spraying-suitability workflow, and disease-risk interface	Weather transformation, crop-stage simulation, disease-risk request construction, treatment-window generation, and risk-state interpretation
Symptom diagnosis and advisory-record layer	Image-upload pathway, visual disease-recognition service, consultation interface, and advisory-response records	Symptom-image processing, disease-category output, visual-record storage, advisory-message storage, and management-oriented explanation
Risk delivery, task, and feedback layer	Notification service, task service, fungicide and pesticide records, treatment windows, and protection-state feedback	Warning delivery, task creation, task completion, overdue handling, fungicide-history conversion, and protection-state or risk-state updating after completed operations

2.3 Crop-season management unit and traceability structure

2.3.1 Field and crop-season records

The data structure was organized around two foundational management objects: the vineyard field and the crop season. The field record represents the persistent spatial and ownership unit, including field identity, area, boundary, location, administrative attributes, sharing state, and management status. The crop-season record links the field with grape cultivar, cultivation method, vine age, field coordinates, field size, and cultivar metadata. Detailed field-record attributes and related crop-season attributes are listed in Supplementary Table S7.

At the implementation level, weather records, phenology-simulation outputs, disease-risk outputs, plant-protection task schedules, disease-risk updates, pesticide or fungicide application ledgers, irrigation tasks, notes, warning notifications, disease images, and user-facing statistical summaries are grouped under the same crop-season management unit. Standardized background databases, including grape variety, cultivation method, BBCH stage, disease type, fungicide list, pesticide list, and reminder-rule tables, provide reference values across the functional steps.

2.3.2 Traceability object and minimum linkage chain

For one crop season c , the implementation-level traceability representation was formalized as:

$$T_c = (F_c, M_c, W_c, P_c, R_c, N_c, O_c, B_c, E_c, A_c), \quad (1)$$

where F_c is the linked field record, M_c is crop-season metadata, W_c is weather input, P_c is phenology state, R_c is disease-risk and request-response output, N_c is notification or event delivery, O_c is task or field-operation state, B_c is fungicide or pesticide feedback, E_c is symptom or image records, and A_c is advisory or message records. The minimum traceability chain evaluated in this study was:

$$F_c \rightarrow M_c \rightarrow (W_c, P_c) \rightarrow R_c \rightarrow N_c \rightarrow O_c \rightarrow B_c, \quad (2)$$

with E_c and A_c treated as additional symptom or advisory records when present. A crop season was considered analytically linked when W_c , P_c , and R_c shared the same crop-season key. Risk-to-operation linkage was present when N_c or O_c could be retrieved under the same retained user, field, or crop-season context. Treatment-feedback linkage was present when B_c was connected to O_c through task identifiers. The formal crop-season object, database relationship rules, workflow state-transition rules, and linkage-check rules are specified in Supplementary Tables S2–S5. Figure 1 provides a schematic overview of the crop-season-centered traceability architecture used to operationalize the formal object and linkage chain defined above.




Figure 1. Crop-season-centered traceability architecture implemented in GrapeMaster. The crop season serves as the platform management unit linking field setup, environmental and crop state, disease-risk interpretation, event delivery, field operation, treatment feedback, optional symptom and advisory records, and replay/review records. The feedback path indicates that completed plant-protection records can update later protection-state or risk interpretation, while replay/review records support reconstruction of the prediction–action–feedback chain.

2.4 Risk-task-protection-feedback workflow

2.4.1 Risk delivery and task generation

The risk-task-protection-feedback workflow starts from the daily disease-risk output generated for an active crop season. The platform translates model output into user-facing risk states, such as protected, low-risk, favourable, or high-risk periods. When weather, phenology, and disease-risk

outputs indicate a favourable infection window, the crop-season page and notification interface display the corresponding risk state and treatment window.

The notification is linked to the crop season rather than to a stand-alone warning message. From the warning page, the user can enter the plant-protection workflow, inspect the suggested treatment timing, and create a plant-protection task if intervention is required. This rule connects risk viewing, decision review, and task creation within the same crop-season context.

2.4.2 Treatment feedback and protection-state updating

After a plant-protection task is created, the user can keep it pending, complete it after field operation, or leave it unexecuted. A completed task records the execution date and pesticide or fungicide information and is then used by the platform as management feedback. This rule follows the practical plant-protection assumption that fungicide applications can provide a temporary protection window whose duration depends on product properties, application timing, host growth, rainfall, and disease pressure (Caffi and Rossi, 2018; Bleyer et al., 2020; Oliver et al., 2023; Warneke et al., 2020). In this workflow, a completed fungicide-related task activates a rule-based protection period or low-risk management state. During this period, the crop-season risk display is updated to a protected or low-risk state.

When the protection period ends, the platform resumes dynamic risk prediction using the updated weather, phenology, disease-risk input, and management history. If a new favourable or high-risk window appears, the system again displays the risk state and prompts the user to review the need for a new plant-protection task. The resulting sequence is risk prediction, risk-state display, notification, task creation, task completion, protected-state updating, and renewed risk prediction.

2.4.3 Symptom-image and advisory branches

The image-recognition and LLM-assisted advisory modules create additional record branches when users observe or report symptoms during the crop season. A user can upload a field image for disease identification, and the platform returns a grape-disease diagnosis result linked to the current crop-season context. For downy mildew, the image record can be stored alongside the risk-

prediction workflow. For other supported grape diseases, such as powdery mildew, the image record remains a symptom-level diagnostic branch rather than a disease-risk forecasting record.

The LLM-assisted function is implemented as a supplementary advisory-record channel within the crop-season context. LLM messages and advisory outputs are stored as advisory records and linked to the crop-season context when a crop-season identifier is available.

2.5 Platform-compatible analytical modules

2.5.1 Weather-data preprocessing for analytical modules

For each active crop season, scheduled workflows retrieve historical and forecast weather using the field location. In the current implementation, weather data were obtained from the Open-Meteo weather API (<https://open-meteo.com/>) using field coordinates. The weather workflow combines historical observations with forecast records where available. Hourly and daily variables include temperature, humidity, dew-point or wetness-related variables, precipitation, wind conditions, and daily summary variables. Missing values in date-aligned weather series were filled by linear interpolation before analytical use so that downstream workflows received complete input series.

The backend transforms weather data into three working schemas. Daily mean temperature is converted into phenology-model input. Hourly weather is converted into the disease-model schema, including datetime, dew point, precipitation, relative humidity, air temperature, and wind speed. Short-term hourly records are used to evaluate spraying suitability. This preprocessing belongs to the GrapeMaster platform workflow and defines the input schema required by the platform-compatible analytical modules.

2.5.2 Phenology simulation and Guangxi calibration

The phenology workflow estimates grapevine growth-stage progression from daily temperature and crop-season metadata. The backend calculates daily thermal accumulation and maps accumulated development to broad growth-stage classes related to the BBCH (Biologische Bundesanstalt, BUNDessortenamt and Chemical industry) scale. The output provides a crop-development timeline for mobile display and a daily crop-stage sequence for disease-risk interpretation, warning windows,

and management-task scheduling.

For regional evaluation, the phenology observations summarized in Table 1 were mapped to broad major-stage classes: early season or budburst, leaf development, inflorescence emergence, flowering, berry development, ripening, and post-harvest or senescence. Daily mean temperature was converted to cumulative growing degree days (GDD) after a meteorological biofix:

$$\text{GDD}_t = \sum_{d=1}^t f_{\text{WE}}(T_d), \quad (3)$$

where T_d is daily mean temperature and $f_{\text{WE}}(\cdot)$ is the Wang–Engel temperature-response function.

The accumulated value was mapped to the broad-stage sequence using cultivar-specific major-stage thresholds. For mobile display, the predicted major stage was supplemented with within-stage progress and a transition label when the thermal state was close to an adjacent stage boundary.

The Guangxi observations were evaluated using grouped validation by site-year and cultivar-stratified grouping where possible. The primary metrics were exact major-stage agreement, user-visible agreement after transition-window display, adjacent-stage agreement, severe misclassification rate, and mean absolute stage distance. The cultivar Wenke was represented by one sparse site-year and was reported separately.

2.5.3 FSIM-S downy mildew first-infection risk engine

The disease-risk service receives weather, crop-stage sequence, cultivar susceptibility, target disease information, and fungicide-history inputs through the platform interface. It returns date-aligned disease-risk states, field-risk states, protection states, recommendations, and treatment windows. For GrapeMaster, the disease-risk module was configured with FSIM-S for humid subtropical grape-growing areas in Guangxi. The module predicts the first seasonal infection of grapevine downy mildew under conditions where early-season infection may be driven by recurrent airborne asexual sporangia rather than by a discrete oospore-driven primary infection event. It was calibrated and evaluated using the downy mildew monitoring records summarized in Table 1.

FSIM-S uses a mechanistic infection chain adapted from Rossi primary-infection modelling and

related sporangia-based infection modelling work (Rossi et al., 2008; Caffi et al., 2009, 2011a; Brischetto et al., 2021). In the subtropical setting, airborne sporangia are treated as the relevant inoculum source for the first seasonal infection, and the model represents the main steps from sporangia availability and deposition to infection establishment, incubation, and symptom expression. The FSIM-S configuration evaluated in this study uses field-derived spore pressure to regulate the infection-triggering threshold. A daily sporangia pressure field $S(d)$ is constructed from field spore monitoring data after interpolation, normalization, and smoothing. Hourly pressure is represented as a three-day rolling mean:

$$P_t = \frac{1}{3} \sum_{k=0}^2 S(d_t - k), \quad (4)$$

where P_t is the spore-pressure index at hour t , and d_t is the corresponding calendar day.

Infection is triggered when the wetness-duration term and its temperature response exceed a spore-pressure-adjusted effective threshold:

$$\begin{aligned} \text{IC}_t^* &= \begin{cases} \text{IC} \exp(-\alpha_{\text{pos}} P_t), & P_t \geq 0, \\ \text{IC} \exp[\alpha_{\text{neg}} (-P_t)], & P_t < 0, \end{cases} \\ \text{infection}_t &= \mathbb{I}(\text{WD}_t \text{TWD}_t \geq \text{IC}_t^*), \end{aligned} \quad (5)$$

where WD_t is the wetness-duration component, TWD_t is the temperature response during the wetness period, IC is the base infection threshold, α_{pos} and α_{neg} are calibration parameters for positive and negative spore-pressure effects, and $\mathbb{I}(\cdot)$ is an indicator function. Positive spore pressure lowers the effective infection threshold, whereas low pressure raises it.

The disease-risk workflow was limited to downy mildew because FSIM-S was calibrated for this target disease. Other grape diseases were represented in image-recognition or advisory functions but were not disease-risk prediction targets in this study. Weather inputs were transformed into the platform disease-model schema, and the spore-pressure term was used as an FSIM-S model-level calibration input rather than as a required live input for routine platform operation. Supplementary Figure S1 presents the platform-level inputs, transformations, and user-facing outputs required for

integrating the disease-risk engine with the GrapeMaster workflow.

2.5.4 Grape disease recognition and LLM-assisted advisory

The image-based workflow records symptom-level information after disease signs are observed or suspected. Users can upload field images through the mobile client, and the image-recognition service returns disease or healthy categories with associated diagnostic or management text for display. Image outputs can be linked to crop-season notes, disease reports, and advisory records, keeping visual records within the same crop-season context as other management records.

The image-recognition and advisory module follows the published CDIP-ChatGLM3 dual-model framework, which combines a deep-learning image classifier with a fine-tuned ChatGLM3-based advisory model for crop disease identification and management guidance (Yan et al., 2025). In this study, the framework was adapted to the grape scenario by expanding the grape disease image dataset summarized in Table 1 with additional field photographs and two grape disease categories. The platform-compatible visual model was selected according to grape-focused validation performance.

For image recognition, each candidate model outputs class probabilities using a softmax layer and is trained with cross-entropy loss:

$$\begin{aligned} p(y = k | x) &= \frac{\exp(z_k)}{\sum_{j=1}^K \exp(z_j)}, \\ \mathcal{L}_{\text{CE}} &= - \sum_{k=1}^K y_k \log p(y = k | x), \end{aligned} \tag{6}$$

where x is the input image, z_k is the logit for disease class k , K is the number of grape disease or healthy categories, and y_k is the one-hot class label. Model selection is based on grape-focused validation metrics, including accuracy, precision, recall, and F1-score:

$$m^* = \arg \max_{m \in \mathcal{M}} \text{Score}_{\text{val}}(m), \tag{7}$$

where $\mathcal{M}$ is the set of candidate vision models and $\text{Score}_{\text{val}}$ is the selected validation criterion, such

as accuracy or macro-F1. The selected classifier provides the disease label and confidence score used by the advisory interface. Supplementary Figure S2 summarizes the image-recognition and advisory workflow.

Table 3. Provenance, platform roles, and evaluation basis of the main GrapeMaster components. Backend records and module-level validation results are listed separately because they were generated from different data sources.

Module	Source or provenance	Platform role	Evaluation basis
Platform workflow	Deployed GrapeMaster back-end, mobile client, database, scheduled jobs, and notification/task services	Links field, crop season, weather, phenology, risk, task, fungicide, image, advisory, and record objects	Backend-record export and linkage checks
Phenology module	Weather-driven phenology workflow calibrated with Guangxi field observations	Supplies crop-stage timing for risk interpretation, advisory timing, and task scheduling	Offline Guangxi phenology validation
FSIM-S risk engine	Team-developed downy mildew first-infection model calibrated and validated with Guangxi field data	Supplies first seasonal infection-period warning for task triggering and protection-state interpretation	Offline Guangxi downy mildew validation
Grape disease recognition	Platform-compatible grape disease visual model based on the published CDIP-ChatGLM3 framework	Supplies post-symptomatic image category output and confidence information	Offline grape-image validation
LLM advisory	Fine-tuned ChatGLM3-based advisory model from the CDIP-ChatGLM3 framework	Stores consultation, explanation, and advisory-message records	Platform message records and published advisory framework

2.6 Backend-record audit and representative replay

2.6.1 Backend export and account screening

Backend database exports from GrapeMaster were used to check whether the platform had generated the record types required by the workflow. The exported tables included user and profile tables, field records, crop-season records, notifications, plant-protection tasks, irrigation tasks, fungicide and pesticide records, weather-related records, phenology records, disease-risk records, image-recognition and advisory records, disease-report records, note records, and message records. The export represented the operational records available at the time of data extraction and was used as a cross-sectional record audit.

Account-level screening was applied before summarizing GrapeMaster records. Accounts without business records or without both field and crop-season records were excluded from the workflow-audit set. One non-target-region account was also removed from the vineyard evaluation set. De-

tailed screening rules are listed in Supplementary Table S8.

2.6.2 Linkage checks and workflow reconstruction

The screened records were used for four workflow checks: field and crop-season creation and linkage, attachment of weather, phenology, risk, and notification records to crop seasons, creation and status updating of plant-protection tasks and pesticide/fungicide records, and linkage of image, advisory, disease-report, and note records to management contexts.

The representative replay anchor was selected as a chain-complete audit case from retained crop seasons that had the minimum traceability chain required for risk-to-feedback reconstruction: linked weather, phenology, disease-risk or request-response payloads, notification records, completed plant-protection tasks, and fungicide-product records. The replay was used to test whether the expected workflow objects could be reconstructed through shared crop-season or task identifiers and whether optional symptom or incidence records were present or absent under the selected anchor.

The backend-record audit and representative replay assessed record retention, identifier linkage, and workflow reconstructability, rather than agronomic efficacy, pesticide reduction, user adoption, or economic outcomes.

3 Results

3.1 Backend-record audit of crop-season-centered linkage

The cleaned GrapeMaster backend export contained 95 deduplicated registered accounts. Among them, 47 accounts contained both field records and crop-season records, whereas the remaining 48 accounts did not contain field-crop-season linkage data.

Within the 47 accounts with field-crop-season linkage, the backend contained 139 vineyard fields and 128 crop seasons. Analytical-state records were linked at the crop-season level, including 126 weather payloads, 126 phenology payloads, and 252 disease-risk or model request-response payloads. Event and operation records were also retained, including 3590 notifications, 151 plant-protection or irrigation tasks, and 234 fungicide or pesticide records.

The optional symptom and advisory branches contained 94 field-note or disease-report records, 2808 image-analysis records, 1360 advisory or message records, and 27 forum records. Table 4 summarizes the retained record classes, linkage basis, and interpretation scope. The backend linkage topology is provided in Supplementary Figure S4.

Among the retained backend records, 124 field records from the 47 retained accounts had valid coordinates and crop-season linkage and were mapped as 27 regional aggregates (Figure 2). The same map also locates the independent experimental sites used for module-level validation.




Figure 2. Spatial context of mappable platform field records and independent experimental module-validation sites. Teal bubbles show mappable field records aggregated by region, with bubble size indicating the number of records per region. Red points indicate experimental sites used for module-level validation and are not part of the backend-record denominator.

Table 4. Backend-record audit of crop-season-centered linkage in the retained GrapeMaster export.

Audit level	Retained records	Linkage basis	Result summary
Screened workflow set	47 of 95 deduplicated accounts retained after screening	At least one field and one crop season	Accounts available for crop-season workflow evaluation
Field-season denominator	139 fields and 128 crop seasons	Retained account, field UUID, and crop-season UUID	Field and crop-season records established as the audit base
Crop-season state records	126 crop seasons with linked weather, phenology, risk, and request-response payloads	Shared crop-season UUID	Environmental, crop-stage, and disease-risk states reconstructable by crop season
Event and operation records	3590 notifications and 151 plant-protection or irrigation tasks	Retained user, field, or crop-season context	Event-delivery and field-operation records retained
Treatment-feedback records	234 fungicide or pesticide records linked to 141 plant-protection tasks	Task UUID	Product records connected to completed or recorded task contexts
Symptom and advisory records	94 field-note or disease-report records, 2808 image-analysis records, 1360 advisory or message records, and 27 forum records	Account, field, crop-season, image, or message identifiers	Optional symptom/advisory branches retained, with completeness varying by season
Interpretation scope	No controlled production comparison in the backend export	Not applicable	No efficacy, pesticide-reduction, adoption, or economic outcome inferred

Abbreviation: UUID, universally unique identifier.

3.2 Crop-season workspace entry and state display

The mobile client displayed vineyard fields and crop seasons as the entry points for crop-season management functions. Field-management pages provided access to field setup, crop-season selection, weather, phenology, disease-risk warning, notifications, tasks, notes, and image-related records (Supplementary Figure S3).

Within this workspace, weather, operation-timing, growth-stage, and summary field-risk outputs were displayed under the selected crop-season context (Figure 3). The displayed records retained field-specific environmental and phenology states within the crop-season workspace.

The weather and phenology interfaces corresponded to the linked weather and phenology payloads reported in Table 4. Disease-risk delivery, task feedback, image recognition, and advisory interaction records are reported in Section 3.3.




Figure 3. Crop-season workspace for weather and phenology state in GrapeMaster. (a) The weather page displays near-term field-specific weather information. (b) The operation-timing page displays hourly weather and spray-suitability information. (c) The crop-season workspace displays field-risk and growth-stage summaries under the selected field context.

3.3 Risk delivery, operation feedback, and record capture

Disease-risk outputs were displayed as date-aligned mobile timelines for crop-season review (Figure 4). Across the displayed cases, downy mildew risk, field-level risk, and fungicide-feedback-dependent protection states were retained in a shared temporal display. The warning and notification interfaces displayed crop-season and disease-risk events as notification records (Figure 5), and these records corresponded to the retained event-delivery records summarized in Table 4. Task interfaces retained field-operation and product-feedback information after crop-season events were reviewed (Figure 6). The symptom-image and advisory interfaces provided the optional diagnostic and consultation branch of the workflow (Figure 7), corresponding to the image-analysis and advisory/message records reported in Table 4. Across these interfaces, the displayed record sequence

324 followed the configured workflow order from risk-state display to event delivery, task recording,
 325 treatment-feedback capture, and optional symptom-image or advisory-record entry.







Figure 4. Disease-risk timeline in the GrapeMaster mobile workflow. (a) Mingyang 2024 shows date-aligned downy mildew and field-risk states. (b) A Guangxi crop-season case without fungicide-feedback records shows the untreated risk-state sequence. (c) The same Guangxi crop-season case with fungicide-feedback records shows the corresponding protected-state updates.



(a)



(b)



(c)

Figure 5. Alert and notification records for crop-season events in GrapeMaster. (a) The disease-alert dialog displays a crop-season disease-risk event. (b) The management notice dialog provides a task-related prompt. (c) The notification list archives reviewable crop-season events.




Figure 6. Task and product-feedback interfaces in GrapeMaster. (a) The task list records field operations and task status. (b) The product-selection page stores fungicide or pesticide entries linked to plant-protection tasks. (c) The in-tank mixing page records sprayer capacity, product dosage, plot area, water volume, and planned date.




Figure 7. Symptom-image and advisory-record interfaces in GrapeMaster. (a) The image-upload page records post-symptom visual information. (b) The recognition-result page displays the predicted grape disease category. (c) The advisory interface stores consultation text linked to the crop-season workspace.

3.4 Record-level reconstruction of the risk-to-feedback loop

The selected loop-complete crop-season replay reconstructed the retained record chain from state updating to risk interpretation, event delivery, task execution, and treatment feedback (Figure 8 and Table 5). Under the same crop-season UUID or linked task identifiers, the replay retrieved all record groups required by the minimum risk-to-feedback chain.

Weather, phenology, and disease-risk payloads shared the crop-season key. Notifications and completed tasks were recorded under the same crop-season context, and fungicide-product records carried task identifiers.

No field-note or disease-incidence note was present under this replay anchor, although symptom-image records were present in the broader retained account set. This absence was retained as an explicit entry for the optional field-note branch.




Figure 8. Record-level replay of the risk-to-feedback loop in GrapeMaster. Each node summarizes a retained record group under the selected crop-season management unit, and node labels report the corresponding record counts in this loop-complete replay.

Table 5. Record-level reconstruction results for a loop-complete crop-season replay in GrapeMaster.

Record group	Retained record set	Identifier or status	Result note
Replay record set	Weather, phenology, risk, notification, task, and fungicide records	One crop-season UUID	Loop-complete crop-season replay
State and risk records	Weather, phenology, and disease-risk payloads	Shared crop-season key	Shared seasonal record set
Operation records	Notifications and completed plant-protection tasks	Same crop-season context	Notification and task records
Treatment-feedback records	Fungicide-product records	Task identifiers	Product-task record set
Optional field-note branch	No field-note or incidence-note record	Explicit absence	No field/incidence note in this replay

3.5 Platform-compatible analytical module results

The module-level result set included downy mildew first-infection timing, field-observed grapevine growth stages, and grape disease image-recognition metrics.

The FSIM-S configuration evaluated for GrapeMaster produced a validation mean absolute error of 6.3 days and a validation root mean square error of 6.5 days for downy mildew first-infection

timing (Figure 9). The image-recognition module produced class-wise F1-scores from 0.926 to 0.998 across seven grape disease categories. Downy mildew, the disease target linked to FSIM-S risk prediction, reached an F1-score of 0.986.

The phenology dataset contained 151 valid field observations from six Guangxi site-years. In grouped three-fold validation, the broad-stage configuration reached 58.0% exact major-stage agreement and 76.0% user-visible agreement after transition-window display, with a mean absolute stage distance of 0.45 major stages when the sparsely observed post-harvest class was excluded from the core metrics. Most non-exact cases were transition-window or adjacent-stage cases, and severe mismatches represented a small fraction of observations (Figure 10). Fruit development and ripening stages had lower mean stage distance than early-season stages. The Wenke cultivar subset contained one sparse site-year and is reported separately from the two larger cultivar subsets.




Figure 9. Module-level results for disease-risk and image-recognition components. (a) The FSIM-S panel reports validation error for downy mildew first-infection timing. (b) The image-recognition panel reports class-wise F1-scores for grape disease categories, with downy mildew highlighted.




Figure 10. Validation results for the platform-compatible broad-stage phenology module. (a) Outcome proportions are summarized by cultivar. (b) Outcome proportions are summarized by observed major stage. (c) Mean absolute stage distance is summarized by observed major stage. BD, bud development; LD, leaf development; IE, inflorescence emergence; FL, flowering; FD, fruit development; RM, ripening of berries.

4 Discussion

4.1 Crop season as the platform integration unit

The vineyard field remains spatially stable, whereas weather exposure, phenological stage, inoculum pressure, protection status, observations, and management decisions are season-specific and must be reinterpreted within each crop season. Therefore, in GrapeMaster, the crop season is positioned as the core organizational unit of the vineyard disease-management workflow, rather than as a descriptive time label or database field. This need for a season-specific management unit is consistent with previous plant disease DSS studies showing that weather-driven disease-risk models and advisory systems can improve decision timing only when their outputs are usable within operational production contexts (Seem, 2004; Rossi et al., 2014; Bregaglio et al., 2022). It is also consistent with grapevine phenology studies showing that cultivar and regional climatic context constrain crop-stage interpretation, making seasonal anchoring necessary for operational disease

management rather than merely convenient for data storage (Parker et al., 2011; Wang et al., 2020). This architecture enables cross-module retrieval through a shared seasonal unit, so related environmental, risk, operation, feedback, and observation records can be queried without manually reconstructing links. It also makes season replay possible by keeping the risk-to-feedback sequence within a single temporal frame. For perennial grapevine production systems, repeated crop-season-centered records from the same field can support longitudinal comparison rather than only producing fragmented summaries of isolated records (Nogueira Júnior et al., 2018). Architecture-level linkage results are informative not because of record volume alone, but because retained records remained interpretable within a shared temporal and relational frame. These functions distinguish GrapeMaster from field-centered or user-centered record systems in which the crop season is used mainly as a filtering label or descriptive attribute.

4.2 Traceability from risk display to field-operation feedback

In vineyard disease management, task feedback is essential because fungicide application can alter short-term infection risk, epidemic development, and disease severity by creating a temporary protection period (Caffi and Rossi, 2018). This feedback-based adjustment may need to occur repeatedly within a crop season as weather, host growth, disease pressure, and protection status change. Although disease-warning models have advanced substantially, a warning output alone does not indicate whether the information was reviewed, whether an operation was executed, or whether previous fungicide management has been incorporated into the current decision context. Disease-warning studies similarly indicate that prediction value should be interpreted through action-relevant probability and decision context rather than through risk classification alone (Yuen and Hughes, 2002; Zhao et al., 2011). The GrapeMaster workflow addresses this gap by organizing isolated disease-risk warnings into a dynamic within-season risk–reminder–task–protection–feedback process.

The representative replay shows this workflow logic at the record level (Figure 8 and Table 5). Weather and phenology inputs were linked to risk display and notification, warning states were con-

nected with plant-protection tasks, completed tasks generated fungicide-use records with product and mixing information, and these records supported subsequent protection-state interpretation and updating. Although this study did not measure agronomic efficacy, pesticide reduction, or grower adoption, it established the traceable record chain and architectural pattern required for evaluating these outcomes in future production studies.

Disease diagnosis and agronomic advisory interaction form auxiliary decision-support and supplementary record branches within the same crop-season frame. They do not replace the core workflow chain, but, when present, allow observed symptoms, diagnostic outputs, consultation messages, vineyard context, and operation history to be stored and reviewed together. This design gives growers a structured entry point for symptom-level interpretation and advisory interaction, can reduce the delay between symptom observation and access to management-relevant information, and reduces dependence on informal information sources of uneven reliability while preserving the consultation context for later review (Yao et al., 2025; Yan et al., 2025).

GrapeMaster creates a structured record basis for evaluating treatment timing, unnecessary fungicide use, disease-control outcomes, and advisory consistency in future production studies. This is particularly important because fungicides are a major pesticide class that affects production costs and raises concerns about environmental exposure (Zubrod et al., 2019). Thus, the value of the platform is not only that it delivers warnings, but that it converts warnings into management sequences that can be stored, traced, and audited under a shared management unit, thereby supporting outcome-oriented evaluation.

4.3 Lightweight deployment and regional adaptation

Agricultural DSSs are increasingly expected to function as deployable platform services rather than isolated analytical tools. In viticulture, vite.net integrated vineyard monitoring, pest and disease models, web-based decision support, and two-way user communication to support operational decision making in grape production (Rossi et al., 2014), whereas MISFITS-DSS demonstrated how public research institutions and regional plant-protection services can coordinate grapevine downy

mildew risk assessment across regions (Bregaglio et al., 2022). More recently, Fuxi Brain combined IoT-based perception, multimodal data fusion, and generative-model decision making for whole-cycle maize production management (Chen et al., 2026). These examples indicate that deployment requires more than accurate model output. It also requires workable data sources, user-facing workflows, feedback handling, and adaptation to local production conditions. In this deployment context, GrapeMaster offers a lightweight pathway because the crop-season workflow can operate with routine field records and public weather data before additional sensors or local calibration are introduced.

The same architecture can subsequently be strengthened in two directions. First, when field-scale management requires finer microclimate, crop-stage, or pathogen information, the platform can support optional integration of orchard-scale weather stations, phenology cameras, spore samplers, or other IoT sensors, including for complex mountain or hilly microclimates (Bakir et al., 2026; Nkwocha and Chandel, 2025; Brischetto et al., 2020; Rai et al., 2026). Second, the platform can be extended to support grower-reported phenological stages, disease occurrence, treatment feedback, and other field records for local checking and adjustment of model and advisory outputs. This is a requirement emphasized in DSS implementation studies because local observations and advisory workflows determine how DSS outputs are interpreted, checked, and acted upon (Rossi et al., 2014; Seem, 2004; González-Domínguez et al., 2023). This deployment pathway is consistent with biovigilance frameworks that connect observations, risk interpretation, and management decisions under changing production conditions (Carisse et al., 2017). It allows the platform to remain lightweight at initial deployment while progressively improving adaptation to local conditions as sensor data and user-confirmed records accumulate. However, field-scale validation remains necessary before such adaptations are used as operational recommendations.

4.4 Analytical modules and platform readiness

Independent analytical modules provide decision context for GrapeMaster. The FSIM-S disease-risk module supports warning of the first seasonal infection period of grapevine downy mildew, the

phenology module provides crop-stage context for interpreting susceptibility and operation timing, and image recognition adds visual diagnostic records when orchard disease symptoms become visible. Although the disease-management workflow evaluated here is centred on downy mildew, the image-recognition and LLM-assisted advisory modules provide broader support for diagnosis and agronomic advice when other disease symptoms are observed. The LLM-assisted advisory component provides season-long consultation and management-oriented explanation within the crop-season context, but it remains distinct from validated disease models, expert diagnosis, pesticide labels, and regulated plant-protection decisions. These modules are positioned at different decision moments within the season, so their results should be judged by their contribution to the workflow rather than only by standalone algorithmic performance.

The design value of modular embedding also lies in the ability to update, calibrate, and locally adapt analytical components without relying on a single fixed model. For example, humid subtropical vineyards in Guangxi differ from many temperate or Mediterranean grape-growing regions in rainfall regime, disease pressure, canopy growth, cultivation calendars, and control practices. These differences can change the timing, intensity, and frequency of the same disease and limit direct transfer of warning thresholds or management rules (Seem, 2004; Yu et al., 2017; Wang et al., 2020). FSIM-S is therefore best interpreted as a locally calibrated first-infection prediction component: it represents a Guangxi context in which the first seasonal downy mildew infection is associated with recurrent asexual sporangia under humid conditions, rather than a temperate pattern driven only by oospore-based primary infection (Rossi et al., 2008; Caffi et al., 2009, 2011a; Brischetto et al., 2021). The same principle applies to phenology simulation, image recognition, and advisory support. When the platform is transferred to other cultivars, training systems, or climatic regions, the GrapeMaster architecture supports local recalibration and validation of individual modules rather than reuse as fixed experience-based assumptions.

4.5 Evaluation scope and future validation

The results of this study are best interpreted as a record-based assessment of a traceable and closed-loop workflow and of module readiness, rather than as a final agronomic-impact evaluation. Architecture-level linkage summaries, user-facing workflow records, representative replay results, and module-level tests together show that the platform can generate the expected data objects around the crop-season unit, preserve linkage rules, support feedback closure, and host platform-compatible analytical modules.

Because a single grapevine crop season spans a long annual production cycle, this study did not yet obtain large-scale deployment data across repeated seasons for evaluating production outcomes such as fungicide-use efficiency, disease-control performance, and economic return. Such production-level data will need to be accumulated through broader regional use, repeated crop-season operation, and monitoring across years, cultivars, and vineyard microclimates. Wider user studies should also quantify interaction rate, task-completion rate, recommendation adoption, and suitability across farms of different sizes.

Future work should evaluate GrapeMaster through repeated crop-season use under production conditions. Priorities include multi-region deployment, quantitative comparison with existing grower plant-protection practice, task-completion analysis, and user-centered evaluation of decision adoption. Cross-regional use will also require local checks of weather-data quality, cultivar-specific phenology calibration, disease-risk parameterization, product-rule adjustment, and advisory-content validation.

5 Conclusion

This study developed and evaluated GrapeMaster, a digital platform that uses the crop season as the core organizational unit for closed-loop and traceable vineyard disease management. The platform connects vineyard field identity, weather and phenology state, downy mildew risk interpretation, warning delivery, plant-protection tasks, treatment feedback, symptom-image records, advisory interactions, and archived management records under the crop-season management unit. The rep-

representative season replay showed that the retained backend records linked one active crop season with weather and phenology inputs, one risk-interpretation payload, 53 notifications, 45 risk or spray reminders, two completed plant-protection tasks, four fungicide-product records, and subsequent protection-state review. This indicates that GrapeMaster can convert disease-risk information into a dynamic prediction–notification–task–treatment–feedback–review sequence. Embedded module-level tests further supported the readiness of platform-compatible disease-risk, phenology, and grape disease-recognition modules for the Guangxi humid subtropical vineyard context. By turning separate analytical, advisory, and record-keeping functions into a traceable seasonal workflow, GrapeMaster extends isolated disease-management tools into an auditable, locally adaptable, and scalable management system, providing a digital and methodological foundation for more standardized vineyard-level decision processes, advisory coordination, diagnostic support, and future production-outcome evaluation.

CRediT authorship contribution statement

To be completed before submission.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

The de-identified derived datasets and analysis scripts supporting the figures, tables, and summary statistics reported in this article are available from the corresponding author upon reasonable request. These materials include backend-record coverage summaries, representative crop-season replay summaries, phenology validation outputs, disease-risk timeline data, figure source data, and the code used for data summarization and visualization.

Raw operational backend exports from the deployed GrapeMaster platform are not publicly

available because they contain user account records, field locations, uploaded images, messages, task records, and management information subject to privacy and confidentiality restrictions. De-identified data will be provided after confirmation to protect user privacy.

Acknowledgements

To be completed before submission.

Ethics and user data privacy statement

The backend records used in this study were analyzed in aggregated or de-identified form, and no personally identifiable user information is reported.

Supplementary material

Supplementary material associated with this article is provided as a separate file.

References

- Bakir, M.E., Perea, A.R., Funk, M., Rahman, S., Spetter, M.J., Macon, L., Cox, A., Estell, R.E., Cao, H., Cibils, A.F., Spiegel, S.A., Bestelmeyer, B.T., Utsumi, S.A., 2026. A scalable, end-to-end IoT and remote sensing platform for precision rangeland and livestock management. *Computers and Electronics in Agriculture* 247, 111615. doi: 10.1016/j.compag.2026.111615.
- Bleyer, G., Lösch, F., Schumacher, S., Fuchs, R., 2020. Together for the Better: Improvement of a Model Based Strategy for Grapevine Downy Mildew Control by Addition of Potassium Phosphonates. *Plants* 9, 710. doi: 10.3390/plants9060710.
- Bregaglio, S., Savian, F., Raparelli, E., Morelli, D., Epifani, R., Pietrangeli, F., Nigro, C., Bugiani, R., Pini, S., Culatti, P., Tognetti, D., Spanna, F., Gerardi, M., Delillo, I., Bajocco, S., Fanchini, D., Fila, G., Ginaldi, F., Manici, L.M., 2022. A public decision support system for the assessment

of plant disease infection risk shared by Italian regions. *Journal of Environmental Management* 317, 115365. doi: 10.1016/j.jenvman.2022.115365.

Brischetto, C., Bove, F., Fedele, G., Rossi, V., 2021. A Weather-Driven Model for Predicting Infections of Grapevines by Sporangia of *Plasmopara viticola*. *Frontiers in Plant Science* 12, 636607. doi: 10.3389/fpls.2021.636607.

Brischetto, C., Bove, F., Languasco, L., Rossi, V., 2020. Can Spore Sampler Data Be Used to Predict *Plasmopara viticola* Infection in Vineyards? *Frontiers in Plant Science* 11, 1187. doi: 10.3389/fpls.2020.01187.

Caffarra, A., Eccel, E., 2010. Increasing the robustness of phenological models for *Vitis vinifera* cv. Chardonnay. *International Journal of Biometeorology* 54, 255–267. doi: 10.1007/s00484-009-0277-5.

Caffi, T., Rossi, V., 2018. Fungicide models are key components of multiple modelling approaches for decision-making in crop protection. *Phytopathologia Mediterranea* 57. doi: 10.14601/Phytopathol_Mediterr-22471.

Caffi, T., Rossi, V., Bugiani, R., 2010. Evaluation of a Warning System for Controlling Primary Infections of Grapevine Downy Mildew. *Plant Disease* 94, 709–716. doi: 10.1094/PDIS-94-6-0709.

Caffi, T., Rossi, V., Bugiani, R., Spanna, F., Flamini, L., Cossu, A., Nigro, C., 2009. A model predicting primary infections of *plasmopara viticola* in different grapevine-growing areas of Italy. *Journal of Plant Pathology* , 535–548.

Caffi, T., Rossi, V., Carisse, O., 2011a. Evaluation of a Dynamic Model for Primary Infections Caused by *Plasmopara viticola* on Grapevine in Quebec. *Plant Health Progress* 12, 22. doi: 10.1094/PHP-2011-0126-01-RS.

- Caffi, T., Rossi, V., Legler, S.E., Bugiani, R., 2011b. A mechanistic model simulating ascospore infections by *erysiphe necator*, the powdery mildew fungus of grapevine. *Plant Pathology* 60, 522–531. doi: 10.1111/j.1365-3059.2010.02395.x.
- Carisse, O., 2016. Development of grape downy mildew (*Plasmopara viticola*) under northern viticulture conditions: Influence of fall disease incidence. *European Journal of Plant Pathology* 144, 773–783. doi: 10.1007/s10658-015-0748-y.
- Carisse, O., Fall, M.L., Vincent, C., 2017. Using a biovigilance approach for pest and disease management in quebec vineyards. *Canadian Journal of Plant Pathology* 39, 393–404. doi: 10.1080/07060661.2017.1366368.
- Chen, H., Hou, G., Hua, C., Wang, S., Chen, Z., Zhang, Y., 2026. Agricultural autonomous decision-making system "fuxi brain" based on generative large model fusion internet of things. *Computers and Electronics in Agriculture* 244, 111454. doi: 10.1016/j.compag.2026.111454.
- García de Cortázar-Atauri, I., Duchêne, E., Destrac-Irvine, A., Barbeau, G., De Ressaigui, L., Lacombe, T., Parker, A.K., Saurin, N., Van Leeuwen, C., 2017. Grapevine phenology in France: From past observations to future evolutions in the context of climate change. *OENO One* 51, 115. doi: 10.20870/oeno-one.2016.0.0.1622.
- Gessler, C., Pertot, I., Perazzolli, M., 2011. *Plasmopara viticola*: A review of knowledge on downy mildew of grapevine and effective disease management. *Phytopathologia Mediterranea* 50, 3–44.
- González-Domínguez, E., Caffi, T., Rossi, V., Salotti, I., Fedele, G., 2023. Plant disease models and forecasting: Changes in principles and applications over the last 50 years. *Phytopathology*® 113, 678–693.
- Koledenkova, K., Esmaeel, Q., Jacquard, C., Nowak, J., Clément, C., Ait Barka, E., 2022. *Plasmopara viticola* the Causal Agent of Downy Mildew of Grapevine: From Its Taxonomy to Disease Management. *Frontiers in Microbiology* 13, 889472. doi: 10.3389/fmicb.2022.889472.

- Nkwocha, C.L., Chandel, A.K., 2025. Towards an end-to-end digital framework for precision crop disease diagnosis and management based on emerging sensing and computing technologies: State over past decade and prospects. *Computers* doi: 10.3390/computers14100443.
- Nogueira Júnior, A.F., Amorim, L., Savary, S., Willocquet, L., 2018. Modelling the dynamics of grapevine growth over years. *Ecological Modelling* 369, 77–87. doi: 10.1016/j.ecolmodel.2017.12.016.
- Oliver, C., Cooper, M., Lewis Ivey, M.L., Brannen, P., Miles, T.D., Lowder, S.R., Mahaffee, W.F., Moyer, M., 2023. Fungicide Use Patterns in Select United States Wine Grape Production Regions. *Plant Disease* , PDIS–04–23–0798–REdoi: 10.1094/PDIS-04-23-0798-RE.
- Orlandini, S., Gozzini, B., Rosa, M., Egger, E., Storchi, P., Maracchi, G., Miglietta, F., 1993. PLASMO: A simulation model for control of *Plasmopara viticola* on grapevine. *EPPO Bulletin* 23, 619–626. doi: 10.1111/j.1365-2338.1993.tb00559.x.
- Park, E.W., Seem, R.C., Gadoury, D.M., Pearson, R.C., 1997. DMCAST: A prediction model for grape downy mildew development. *Wein-Wissenschaft* 52, 182–189.
- Parker, A., De Cortázar-Atauri, I., Van Leeuwen, C., Chuine, I., 2011. General phenological model to characterise the timing of flowering and veraison of *Vitis vinifera* L.: Grapevine flowering and veraison model. *Australian Journal of Grape and Wine Research* 17, 206–216. doi: 10.1111/j.1755-0238.2011.00140.x.
- Peng, J., Wang, X., Wang, H., Li, X., Zhang, Q., Wang, M., Yan, J., 2024. Advances in understanding grapevine downy mildew: From pathogen infection to disease management. *Molecular Plant Pathology* 25, e13401. doi: 10.1111/mpp.13401.
- Pertot, I., Caffi, T., Rossi, V., Mugnai, L., Hoffmann, C., Grando, M., Gary, C., Lafond, D., Duso, C., Thiery, D., Mazzoni, V., Anfora, G., 2017. A critical review of plant protection tools for reducing pesticide use on grapevine and new perspectives for the implementation of IPM in viticulture. *Crop Protection* 97, 70–84. doi: 10.1016/j.cropro.2016.11.025.

- Rai, N., (Dana) Choi, D., Boyd, N.S., Schumann, A.W., 2026. Advancing site-specific disease and pest management in precision agriculture: From reasoning-driven foundation models to adaptive, feedback-based learning. *Computers and Electronics in Agriculture* 250, 111859. doi: 10.1016/j.compag.2026.111859.
- Rossi, V., Caffi, T., Giosuè, S., Bugiani, R., 2008. A mechanistic model simulating primary infections of downy mildew in grapevine. *Ecological Modelling* 212, 480–491. doi: 10.1016/j.ecolmodel.2007.10.046.
- Rossi, V., Salinari, F., Poni, S., Caffi, T., Bettati, T., 2014. Addressing the implementation problem in agricultural decision support systems: The example of vite.net®. *Computers and Electronics in Agriculture* 100, 88–99. doi: 10.1016/j.compag.2013.10.011.
- Seem, R.C., 2004. Forecasting plant disease in a changing climate: A question of scale. *Canadian Journal of Plant Pathology* 26, 274–283. doi: 10.1080/07060660409507144.
- Shafay, M., Velayudhan, D., Owais, M., Hassan, T., Seneviratne, L., Hussain, I., Werghi, N., 2026. An end-to-end vision–language framework with LLM-based reasoning for decision-oriented tomato leaf disease management in greenhouse environments. *Computers and Electronics in Agriculture* 247, 111692. doi: 10.1016/j.compag.2026.111692.
- Wang, X., Li, H., García De Cortázar Atauri, I., 2020. Assessing grapevine phenological models under Chinese climatic conditions. *OENO One* doi: 10.20870/oeno-one.2020.54.3.3195.
- Warneke, B., Thiessen, L.D., Mahaffee, W.F., 2020. Effect of Fungicide Mobility and Application Timing on the Management of Grape Powdery Mildew. *Plant Disease* 104, 1167–1174. doi: 10.1094/PDIS-06-19-1285-RE.
- Wise, J.C., Jenkins, P.E., Schilder, A.M., Vandervoort, C., Isaacs, R., 2010. Sprayer type and water volume influence pesticide deposition and control of insect pests and diseases in juice grapes. *Crop Protection* 29, 378–385. doi: 10.1016/j.cropro.2009.11.014.

- Yan, C., Liang, Z., Cheng, H., Li, S., Yang, G., Li, Z., Yin, L., Qu, J., Wang, J., Wu, G., Tian, Q., Yu, Q., Zhao, G., 2025. CDIP-ChatGLM3: A dual-model approach integrating computer vision and language modeling for crop disease identification and prescription. *Computers and Electronics in Agriculture* 236, 110442. doi: 10.1016/j.compag.2025.110442.
- Yan, C., Liang, Z., Yin, L., Wei, S., Tian, Q., Li, Y., Cheng, H., Liu, J., Yu, Q., Zhao, G., et al., 2024. AFM-YOLOv8s: An accurate, fast, and highly robust model for detection of sporangia of *plasmopara viticola* with various morphological variants. *Plant Phenomics* 6, 0246.
- Yao, L., Zhao, G., Yan, C., Srivastava, A.K., Tian, Q., Jin, N., Qu, J., Yin, L., Yao, N., Webber, H., Luedeling, E., Yu, Q., 2025. Influence of Information Sources on Technology Adoption in Apple Production in China. *Agriculture* 15, 1785. doi: 10.3390/agriculture15161785.
- Yu, S., Li, B., Guan, T., Liu, L., Wang, H., Liu, C., Zang, C., Huang, Y., Liang, C., 2022. A comparison of three types of “vineyard management” and their effects on the structure of *plasmopara viticola* populations and epidemic dynamics of grape downy mildew. *Plants* 11, 2175. doi: 10.3390/plants11162175.
- Yu, S., Liu, C., Liang, C., Zang, C., Liu, L., Wang, H., Guan, T., 2017. Effects of Rain-shelter Cultivation on the Temporal Dynamics of Grape Downy Mildew Epidemics. *Journal of Phytopathology* 165, 331–341. doi: 10.1111/jph.12566.
- Yuen, J.E., Hughes, G., 2002. Bayesian analysis of plant disease prediction. *Plant Pathology* 51, 407–412. doi: 10.1046/j.0032-0862.2002.00741.x.
- Zhao, C.J., Li, M., Yang, X.T., Sun, C.H., Qian, J.P., Ji, Z.T., 2011. A data-driven model simulating primary infection probabilities of cucumber downy mildew for use in early warning systems in solar greenhouses. *Computers and Electronics in Agriculture* 76, 306–315. doi: 10.1016/j.compag.2011.02.009.
- Zubrod, J.P., Bundschuh, M., Arts, G., Brühl, C.A., Imfeld, G., Knäbel, A., Payraudeau, S., Ras-

658 mussen, J.J., Rohr, J., Scharmüller, A., 2019. Fungicides: An overlooked pesticide class? Envi-
659 ronmental science & technology 53, 3347–3365.