

1 Irrigation strategies under contrasting water-storage and rainfall

2 conditions in the North China Plain

3 Linjia Yao^{1,†}, Bin Chen^{2,†}, Linchao Li³, Hao Feng², Qiang Yu^{2,*}, Gang Zhao^{2,*}

4 ¹ College of Natural Resources and Environment, Northwest A&F University, Yangling 712100, China

5 ² State Key Laboratory of Soil and Water Conservation and Desertification Control, College of Soil and Water
6 Conservation Science and Engineering, Northwest A&F University, Yangling, Shaanxi 712100, China

7 ³ College of Agronomy, Inner Mongolia Agricultural University, Hohhot, 010019, China

8 † Linjia Yao and Bin Chen contributed equally.

9 * Corresponding authors: Qiang Yu (yuq@nwfufu.edu.cn) and Gang Zhao (gang.zhao@nwfufu.edu.cn).

10 Cover letter

11 Dear Editor,

12 Please consider the research article “Irrigation strategies under contrasting water-storage and rainfall conditions in
13 the North China Plain” for publication in Agricultural Water Management.

14 Reducing irrigation while sustaining grain production requires balancing water consumption and the distribution of
15 production losses. Using a process-based crop model, the study compares uniform and spatially targeted irrigation
16 under equal water budgets and tests whether satellite water-storage information improves rainfall-based irrigation
17 decisions.

18 At the same 50% irrigation reduction, targeting retained 4.87% more simulated grain than uniform cuts, but
19 increased evapotranspiration and reduced grain production in some locations. Adding regional storage information
20 did not change irrigation quotas or outcomes relative to matched rainfall-only strategies within the tested design.

21 These model-based findings support irrigation strategies that combine production goals with limits on consumptive
22 use and safeguards for local production. Their focus on irrigation allocation and crop water use makes the study
23 well suited to Agricultural Water Management.

24 Thank you for considering this manuscript.

25 Sincerely,

26 Qiang Yu and Gang Zhao

27 State Key Laboratory of Soil and Water Conservation and Desertification Control, College of Soil and Water
28 Conservation Science and Engineering, Northwest A&F University, Yangling, Shaanxi 712100, China
29 yuq@nwfau.edu.cn; gang.zhao@nwfau.edu.cn

30 **Highlights**

- 31 • Irrigation cuts require balancing grain retention, water use and local losses.
- 32 • Targeting added 0.732 t ha⁻¹ dry grain and 20.76 mm ET yearly over uniform 50% cuts.
- 33 • At 50% cuts, mean grain was below uniform allocation on 37.60% of the area.
- 34 • Over nine GRACE years, grain retention was 96.27% (95% target) and 99.63% (98%).
- 35 • Matched rainfall-only controls differed in grain production by 0% at both targets.

36 **Manuscript**

37 **Abstract**

38 Efficient irrigation allocation is essential for sustaining grain production while limiting water consumption in the
39 North China Plain wheat–maize rotation. How spatial targeting redistributes production losses, and whether
40 regional water-storage information improves rainfall-based irrigation decisions, remain unresolved. Continuous
41 process-based crop simulations compared uniform and targeted allocations under equal irrigation budgets and
42 evaluated annual storage–rainfall strategies against matched rainfall-only controls during 2014–2025. Growth
43 parameters were fitted to multisite observations; water-use parameters were calibrated against Wuqiao treatments in
44 2016–2018 and retrospectively tested in 2019. At a 50% irrigation reduction, targeting retained 0.73 t ha⁻¹ yr⁻¹
45 more simulated combined dry grain than uniform cuts across 9.87 million rotation hectares, equivalent to 7.22
46 million t yr⁻¹ or 4.87% of uniform production. Mean production was lower than under uniform cuts on 37.60% of
47 mapped area, while crop-plus-fallow evapotranspiration increased by 20.76 mm yr⁻¹ (2.05 km³ yr⁻¹). Across nine
48 years with eligible Gravity Recovery and Climate Experiment (GRACE) observations, strategies selected at 95%
49 and 98% grain targets retained 96.27% and 99.63% of conventional production, respectively. The 95% strategy

50 reduced irrigation by 130.72 mm and evapotranspiration by 36.84 mm per rotation. Matched storage–rainfall and
51 rainfall-only strategies selected identical quotas and produced identical outcomes. These conditional results support
52 joint production and consumptive-use constraints and local production safeguards, while showing that additional
53 storage information provides no decision benefit within the tested quota design.

54 **Keywords**

55 winter wheat–summer maize rotation; irrigation management; crop modeling; GRACE; water availability;
56 consumptive water use

57 **1. Introduction**

58 Irrigation management in the North China Plain must reconcile grain production with water-resource constraints
59 across the winter wheat–summer maize rotation (Yang et al., 2022). Wheat grows through a relatively dry season,
60 whereas maize receives more summer rainfall; soil-water reserves connect their responses to irrigation (Yang et al.,
61 2024). Long-term well-watered observations show that rising grain yield has accompanied increasing actual
62 evapotranspiration (ET), despite comparatively stable reference ET (Zhang et al., 2011). Maintaining production
63 with less irrigation therefore requires attention to both the water delivered to fields and the water consumed over the
64 complete rotation.

65 The timing and location of irrigation supply determine its marginal production return. A sixteen-year experiment in
66 2003–2018 compared seven wheat–maize irrigation schedules and found diminishing production gains as irrigation
67 increased (Yang et al., 2022). A four-year Wuqiao experiment also showed that precipitation conditions influenced
68 the preferred wheat schedule and grain production in the subsequent maize crop (Yang et al., 2024). These
69 responses reflect the sequence of crop demand, rainfall and soil-water carry-over across both seasons (Yang et al.,
70 2024; Yang et al., 2022). At regional scale, irrigation requirements vary among locations and hydrological years
71 (Ren et al., 2022; Wu et al., 2019). Crop-model optimization and irrigation-allocation studies provide approaches
72 for using this heterogeneity to guide management (Jiang et al., 2016; Xiao et al., 2024). Under a common water
73 budget, targeting locations with larger marginal grain returns can improve regional production retention, while
74 changing the distribution of production losses.

75 Assessing these allocation consequences requires consistent crop and water accounting. Field delivery, ET, return
76 flows and storage changes describe different quantities, so a reduction in irrigation does not directly establish an
77 equivalent reduction in consumption (Grafton et al., 2018; Perry et al., 2009). Process-based applications of APSIM

78 and RZ-SHAW in the North China Plain connect crop production with water productivity, field water balance and
79 ET (Chen et al., 2010b; Fang et al., 2014). However, transpiration formulation remains a major source of variation
80 in simulated wheat water use (Cammarano et al., 2016). Phenology, canopy, biomass, grain and ET observations
81 therefore constrain complementary processes, while documented irrigation treatments assess their combined
82 response to water supply. Continuous simulation retains the soil-water carry-over needed to compare complete
83 rotation histories.

84 Spatial allocation determines where irrigation is supplied, while antecedent hydrological conditions can inform the
85 annual irrigation budget. Recent rainfall describes water input at the crop-system scale, whereas GRACE total water
86 storage integrates several terrestrial stores over a regional footprint (Save et al., 2016; Scanlon et al., 2016). Recent
87 aquifer recovery under water transfers, abstraction management and changing climate illustrates the broader context
88 of regional storage variation (Long et al., 2025; Xu et al., 2026). Storage information may thus characterize
89 conditions beyond local rainfall. Its practical value for irrigation nevertheless depends on whether it changes the
90 selected action or improves the resulting grain–water outcome relative to rainfall-based decisions.

91 Two connected management questions remain unresolved. First, it is unclear how much grain spatial targeting can
92 retain under an identical irrigation budget, and what consumptive-water costs and persistent local production losses
93 accompany that regional benefit. Rotation-wide accounting must retain soil-water carry-over to assess these
94 outcomes together. Second, the additional decision value of regional storage information beyond antecedent rainfall
95 remains uncertain. A storage indicator can distinguish hydrological conditions without changing an irrigation
96 choice. Comparisons using the same production targets, candidate quotas and monitoring availability are needed to
97 isolate its contribution to management outcomes.

98 The objectives were to (i) quantify rotation grain production, crop-plus-fallow ET and profile drainage under
99 uniform and spatially targeted irrigation allocations with common regional water budgets; (ii) characterize the
100 spatial distribution, magnitude and annual persistence of allocation-related grain gains and losses; and (iii) assess
101 whether antecedent regional storage information changes annual irrigation quotas and grain–water outcomes
102 relative to rainfall-only controls under matched selection and monitoring conditions. Continuous Open Crop Model
103 simulations of the North China Plain wheat–maize rotation, parameterized using multisite crop observations and
104 documented irrigation treatments, provide the basis for these conditional comparisons during 2014–2025.

2. Materials and methods

2.1. Overall workflow

The workflow links crop-model calibration and evaluation to regional irrigation-management comparisons through four steps (Figure 1). First, crop development and growth are calibrated and evaluated, and crop water-use parameters are fitted to irrigation-treatment responses and tested retrospectively. The resulting parameters remain fixed for the management comparisons. Second, continuous winter wheat–summer maize simulations quantify crop responses to five irrigation levels, with soil water carried through successive crops and intervening fallows.




Figure 1. Study workflow. Crop growth and water-use calibration and evaluation precede continuous regional rotation simulations. Historical crop responses support spatial irrigation allocation under equal water budgets and annual quota selection from antecedent water availability. The strategies are compared during 2014–2025 using rotation grain production, field irrigation, crop-plus-fallow ET, profile drainage and the spatial distribution of production gains and losses.

Third, historical simulated responses support two complementary irrigation decisions: where to allocate a regional water budget and how much irrigation to supply in a given year. Spatial targeting and uniform reductions share identical regional irrigation budgets. Annual quota selection compares antecedent storage–rainfall classes with rainfall-only controls under matched production targets, candidate quotas and monitoring availability. Finally, the

selected strategies are compared during 2014–2025 using rotation grain production, field irrigation, crop-plus-fallow evapotranspiration (ET), profile drainage and the spatial distribution and persistence of production gains and losses.

2.2. Study area

The study area encompasses the winter-wheat–summer-maize lowlands of the North China Plain within 112–121.5°E and 32–41°N, extending from Beijing, Tianjin and Hebei through Shandong and Henan to the northern Anhui and Jiangsu plains (Figure 2) (Wu et al., 2019). The landscape consists of extensive alluvial lowlands, with piedmont plains along the Taihang and Yanshan mountains in the west and north and lower coastal plains toward the east (Long et al., 2025; Xu et al., 2026). Its low relief supports intensive agricultural production, while climatic and soil-water differences contribute to spatial variation in irrigation requirements (Ren et al., 2022).




Figure 2. Study area, rotation area, observation sites and regional water storage. (a) Mapped 2020 wheat–maize rotation area. (b) Five crop-observation sites, the Wuqiao irrigation experiment and 32 regional simulation units. (c) Observed CSR monthly terrestrial-water-storage anomalies and a 12-calendar-month mean requiring at least nine observations. Missing months remain

gaps. Thin grey lines show provincial boundaries; black outlines delimit the operational physiographic study domain. Maps use an
equiarectangular projection (36°N, 117°E). Provincial boundaries: Natural Earth Admin 1 v5.1.1 (Natural Earth, n.d.).

The region has a continental monsoon climate, with cold, dry winters and warm, rainy summers; regional
assessments report mean annual temperatures of approximately 10–15°C (Ren et al., 2022). Rainfall is concentrated
in the summer monsoon, whereas spring is frequently dry and precipitation varies substantially among years and
locations (Wu et al., 2019; Xu et al., 2026). Winter wheat is sown in autumn and harvested in early summer,
followed by summer maize harvested in autumn. Wheat therefore grows largely outside the rainy season and
commonly requires supplementary irrigation before sowing and during spring growth. Maize overlaps the rainy
season, with irrigation demand depending more strongly on rainfall distribution and dry-year conditions. The two
crops share a soil-water reserve, so irrigation in wheat can affect the water available to subsequent maize (Yang et
al., 2024; Yang et al., 2022).

Groundwater has historically supplied much of the irrigation deficit, particularly in the northern plain, where
intensive pumping for agriculture and urban demand caused prolonged groundwater decline (Xu et al., 2026). The
aquifer system includes both unconfined and deeper confined units, with substantial spatial differences in
groundwater depth and depletion (Long et al., 2025). Irrigation water availability also reflects access to river
diversions and the allocation of transferred surface water. Since 2020, groundwater recovery has been documented
across much of the northern North China Plain following the expansion of South-to-North Water Diversion
supplies, reduced abstraction and water-saving measures, with wet years reinforcing recovery (Long et al., 2025;
Xu et al., 2026). This transition places irrigation management within a changing water-supply setting: maintaining
recovery requires attention to consumptive use and seasonal demand as well as the volume delivered to fields.

2.3. Field observations and environmental data

2.3.1. Multisite crop observations

Winter-wheat and summer-maize observations supported crop-growth parameter estimation and comparisons across
five North China Plain sites. Long-term monitoring at Fengqiu, Luancheng, Shangqiu and Yucheng (Fengqiu
Station, 2020a; Luancheng Station, 2020; Shangqiu Station, 2021; Yucheng Station, 2025a; Yucheng Station, 2021)
was complemented by the Gucheng maize sowing-date experiment (Geng et al., 2023). The National Ecosystem
Science Data Center archives contain intermittent crop seasons during 1998–2022; exact site–crop–year coverage
and dataset identifiers are given in Tables S1 and S2.

Whole site-years retained their calibration or retrospective-evaluation assignments (Figure 3). The archive comprised 132 crop–plot–season trajectories, of which 127 contained eligible LAI, biomass, ET or harvest observations (Table 1). Daily ET and antecedent soil moisture came from the Fengqiu soil-water archive (Fengqiu Station, 2020b); ET was also available from the Yucheng water-environment archive (Yucheng Station, 2025b). Grain and harvest-biomass samples were confined to Yucheng. Within-season measurements and quadrats describe temporal dynamics and sampling variation rather than independent growing seasons.




Figure 3. Evidence partitions. (a) Whole station site-years retain their original calibration and retrospective-evaluation partitions; dates denote observation-group years. All five eligible sites are included. (b) Four Wuqiao treatment means per crop-year provide 12 calibration seasons in 2016–2018 and four retrospective testing seasons in 2019. Station comparisons retain earlier growth-fitting history and unconfirmed complete irrigation management.

Observations were matched to predictions by crop, plot, quantity and sampling date or averaging window. Screening excluded wheat LAI above $7 \text{ m}^2 \text{ m}^{-2}$ and duplicated maize harvest groups of unresolved authenticity; raw values and exclusion reasons were retained. Complete plot-specific irrigation histories remain unconfirmed, so station comparisons are conditional on archived management assumptions and have no weight in the water-response fit based on documented Wuqiao treatments (Section 2.3.2). Measurement definitions, screening details and antecedent soil-moisture initialization are specified in Supplementary Section S1.

177 **Table 1. Eligible station-observation records by site and variable.**

Site	LAI	Biomass	Daily ET	Yield	Harvest biomass
Fengqiu	159	163	962	0	0
Gucheng	216	0	0	0	0
Luancheng	123	165	0	0	0
Shangqiu	114	148	0	0	0
Yucheng	183	67	4713	192	192

178 Counts denote individual observations, not independent experiments. Station comparisons retain inherited growth-
179 fitting history and unconfirmed complete management. Yucheng grain and harvest biomass contain six dry-matter
180 samples per crop-year; the duplicated 2012 and 2017 maize groups remain excluded.

181 **2.3.2. Field irrigation experiment**

182 The Wujiao winter-wheat–summer-maize experiment (37.630°N, 116.444°E) comprised four wheat irrigation
183 regimes during 2015–2019 (Yang et al., 2024). W0 received no wheat irrigation; W1 received 75 mm before
184 sowing; W2 received an additional 75 mm at jointing; and W3 received a further 75 mm at anthesis. Total wheat
185 irrigation was therefore 0, 75, 150 and 225 mm, respectively. Treatments had three field replicates and a common
186 nitrogen input of 158.4 kg ha⁻¹ per crop.

187 Maize irrigation was identical across treatments within each year: no irrigation in 2016 and 2018, and 75 mm on 18
188 June in 2017 and 2019. Maize W0–W3 labels consequently denote preceding wheat irrigation rather than different
189 maize irrigation schedules. The treatment structure distinguishes direct wheat irrigation contrasts from maize
190 responses associated with preceding wheat management.

191 Crop–treatment–year observations comprised seasonal ET, grain yield and dry aboveground biomass. ET was
192 estimated from rainfall, irrigation and depletion of 0–2 m soil-water storage, assuming negligible runoff and
193 drainage; grain yields were standardized to 13% moisture (Yang et al., 2024). Observations from 2016–2018
194 constrained water-response coefficients, while 2019 remained outside fitting and parameter selection. Prior
195 inspection during model development makes the latter comparison retrospective (Figure 3). Simulation
196 initialization, ET accounting and grain digitization are detailed in Supplementary Section S4.

197 **2.3.3. Meteorological, soil, crop-rotation and terrain data**

198 Daily AgERA5 v2.0 weather data on the common 0.1° grid supplied minimum and maximum air
199 temperature, precipitation, solar radiation, reference evapotranspiration, vapor pressure and 10-m wind
200 speed for 1996–2025 (Copernicus Climate Change Service, 2020). Reference evapotranspiration (ET_0)
201 entered the water-balance module directly.

202 Native 250 m SoilGrids v2.0 maps of texture, bulk density, organic carbon and coarse fragments were averaged to
203 0.01° , then aggregated over mapped rotation pixels within 0.1° meteorological cells, for six depth intervals between
204 0 and 2 m (ISRIC — World Soil Information, 2020). Hydraulic properties were derived using Saxton–Rawls
205 equations with bulk-density and gravel adjustments (Saxton and Rawls, 2006). Organic matter was approximated
206 using an assumed organic-matter/organic-carbon ratio of 1.724, and air-dry water content was assumed to equal half
207 the wilting-point value. Hydraulic derivation is detailed in Supplementary Section S2.

208 The 500 m ChinaCP map for 2020 identified winter-wheat–summer-maize rotation pixels, which were paired with
209 the corresponding CPM cropland-percentage map (Qiu et al., 2022). Physical classified-cell areas were corrected
210 for projection scale and multiplied by cropland percentages to obtain rotation-area weights. CPM values were
211 sampled at ChinaCP pixel centers after coordinate transformation because the source grids had different origins.
212 Pixel centers assigned these weights to the study area and 0.1° meteorological cells. The rotation footprint was
213 9,866,733 ha before soil screening; three cells with wholly missing soil profiles, totaling 131 ha, were excluded,
214 leaving 9,866,602 ha in 3,641 meteorological cells. The fixed 2020 footprint supplies a common area basis across
215 climate years, so modeled totals describe climatic and management responses rather than historical production
216 trends. The 2015–2021 area series and common-extent comparison with CCD2020 quantify mapping sensitivity in
217 Supplementary Section S3.

218 SRTMGL1 v003 elevation was bilinearly resampled to 0.001° , and slope was derived at that support (NASA JPL,
219 2013). Aligned 10×10 cell means provided elevation and slope on a 0.01° grid. Catchment screening, elevation
220 ≤ 200 m and slope $\leq 2^\circ$ delineated a connected lowland study area of 404,658 km² within the geographical extent
221 described in Section 2.2. Geographical and terrain-threshold sensitivities are reported in Supplementary Section S3.
222 Crop-area weights and aggregated soil profiles were linked to daily weather through common 0.1° cell identifiers.

223 **2.3.4. Terrestrial-water-storage observations**

224 CSR RL06.3 monthly GRACE/GRACE-FO total-water-storage anomalies were aggregated using physical cell
225 areas and the official CSR land mask (Save, 2020; Save et al., 2016). The land-mask overlap covers 395,969 km²
226 within the 404,658-km² study area; excluded coastal supports account for 2.15% of its area. The archive contains
227 259 observed monthly solutions from April 2002 to July 2026 relative to the 2004–2009 mean. Source values in
228 centimeters were converted to millimeters, and missing months remain missing. A 12-calendar-month moving mean
229 requires at least nine observations. The 0.25° product grid is finer than the effective GRACE observation scale, so
230 one regional storage indicator is shared across all simulation units (Save, 2020; Save et al., 2016). Soil, surface and
231 other stores were not subtracted; the indicator represents total water storage rather than groundwater alone.
232 The corresponding aggregation without the CSR land mask is retained as a sensitivity analysis. Regional storage
233 changes provide hydrological context for water-management pressures, with no causal attribution to particular
234 irrigation treatments, water transfers or groundwater regulations (Long et al., 2025; Long et al., 2020).

235 **2.4. Crop model description**

236 Open Crop Model (OCM) (Zhao et al., submitted) couples crop growth and a layered soil-water balance at a daily
237 time step (Figure 4). Wheat development combines temperature, vernalization and photoperiod responses; maize
238 development follows thermal time. Development stage (DVS) is anchored at emergence, flowering and maturity.
239 Germination begins when water above the wilting-point storage is available in the represented layer containing the
240 seed; subsequent thermal development starts from that date. Stage coefficients remain fixed, while the soil-water
241 state can alter establishment timing. Recorded or prescribed harvest dates define the management endpoint
242 separately. Soil-water storage passes continuously through crops and fallows.




Figure 4. Crop growth and soil-water coupling in Open Crop Model. Gold denotes photosynthetically active radiation (PAR) input; blue solid arrows denote water fluxes; green solid arrows denote dry-matter transfers; dashed arrows denote process controls and feedbacks. Nutrition is prescribed, the lower boundary uses free drainage, and soil water carries through crop/fallow segments. Plant and soil sketches are not to scale.

Canopy interception of radiation follows Beer–Lambert attenuation. Aboveground production depends on intercepted PAR, crop-specific RUE, temperature, prescribed nutrition and an empirical power response to realized transpiration supply. PAR is 0.45 of incident shortwave radiation (Kiniry et al., 1989). Stage-specific SLA determines new leaf area, while established cohorts retain their birth area-to-mass ratio (de Wit and Berghuijs, n.d.). Stage-dependent thermal senescence and a separate daily stress-loss fraction determine leaf survival (Tan et al., 2020). LAI emerges from these carbon and area balances without a maximum-LAI parameter. Simulated biomass at wheat flowering or maize silking determines grain number; fixed individual grain mass defines sink capacity, and assimilate supply and wheat stem reserves constrain grain filling.

The water balance represents interception, ponding, infiltration, evaporation, root uptake, runoff and bottom drainage. Atmospheric demands are partitioned from reference ET using canopy cover and effective transpiration and evaporation coefficients. Soil evaporation follows a storage-based two-stage reduction (Allen et al., 1998).

258 Root uptake depends on rooted-layer water, local depletion stress and a bounded compensation fraction that
259 redirects unmet demand toward better-supplied rooted layers. Vertical transport uses van Genuchten–Mualem
260 hydraulic functions (Mualem, 1976; van Genuchten, 1980), signed gravity–matric gradients and conservative
261 storage bounds. Relative-arithmetic interface mobility and adaptive internal steps permit wetting and redistribution
262 within 24 daily outer steps. The lower boundary is free drainage with no capillary-rise supply. This explicit layered
263 approximation differs from a full implicit Richards-equation solution. Profile depths follow the supplied inputs,
264 including 2 m for Wuqiao and regional simulations.

265 **2.5. Model parameterization, calibration and evaluation**

266 Phenological coefficients, stage-specific specific leaf area and senescence, initial LAI, stem-reserve coefficients and
267 nutrition modifiers follow the crop parameterization described in Section S2. RUE, leaf allocation and grain-
268 number parameters were estimated from 480 wheat and 433 maize growth and harvest observations across four
269 wheat sites and five maize sites. Whole site-years retain their original partitions, with balanced contributions from
270 variables, sites, site-years and crop cases. Station simulations retain their recorded site and cultivar
271 parameterization; Wuqiao and regional simulations use the shared crop parameters (Table S3). Individual grain
272 masses are fixed at 0.045 g for wheat and 0.300 g for maize.

273 Eight water-use parameters per crop are fitted to the documented Wuqiao treatments in 2016–2018: the
274 transpiration coefficient, soil evaporation coefficient, readily available water fraction, assimilation water-stress
275 exponent, root-density decay, maximum rooting depth, root water-uptake coefficient and root compensation
276 fraction. Each crop contributes 12 seasonal ET, 12 dry-grain and 12 dry-biomass targets, with three annual W3–W0
277 ET contrasts derived from the same treatment seasons. RUE, leaf allocation and grain-number parameters remain
278 fixed. Bounded least squares combines scaled observation residuals and weak parameter priors; objective weights,
279 bounds and convergence settings are specified in Section S2.4.

280 The fitted water parameters are fixed before comparison with the four 2019 treatment means per crop. Previous
281 inspection of these outcomes makes the test retrospective. Station observations have zero weight in this water fit
282 and assess transfer under unconfirmed complete irrigation histories. Regional strategy selection uses historical
283 weather responses; the crop-calibration years overlap the 2014–2025 regional comparison period, so those
284 outcomes are conditional scenarios rather than time-independent crop validation. Observed and simulated quantities
285 retain matched sampling windows and grain-moisture bases.

286 **2.6. Regional upscaling and irrigation scenarios**

287 Thirty-two area-weighted climate–soil representatives were selected from actual source cells using deterministic
288 clustering of latitude, longitude, plant-available water, and 1997–2013 mean precipitation and reference ET.
289 Features were standardized with rotation-area weights. Each cluster uses the actual cell nearest its centroid; the full
290 source-cell mapping and represented areas are retained. Climate years used for policy evaluation do not enter the
291 climate features used for selection. Shared crop parameter sets are transferred regionally without extrapolating site-
292 specific adjustments.

293 Each representative has five continuous histories at irrigation fractions 0, 0.25, 0.50, 0.75 and 1.00. The
294 conventional wheat schedule is 60 mm at sowing on 12 October, 90 mm on 25 March, 90 mm on 15 April and 60
295 mm on 10 May, totaling 300 mm. Maize receives 80 mm at sowing on 15 June. These common schedules are
296 declared scenarios and are not observed universal practice across the plain. Every event amount is multiplied by the
297 irrigation fraction, with its timing fixed. Wheat is simulated through 10 June and maize through 5 October;
298 intervening fallows carry the soil state forward. Initial plant-available water is 80% at the start of the 1996 spin-up,
299 with no subsequent seasonal storage reset. All crop and fallow ET, drainage and runoff are included. The final 2025
300 cycle ends on 5 October because the next fallow lies outside the declared endpoint. Nutrition is represented by
301 prescribed fertilized-crop modifiers; nitrogen cycling and spatially varying nutrient limitations are not dynamically
302 simulated.

303 Eight purposively selected non-representative cells provide paired spatial aggregation checks under all five
304 irrigation fractions. Two cells from each latitude quartile have the largest area-times-feature-distance score. Actual-
305 cell weather and profiles use the same continuous model and crop parameters. Their differences from
306 representative-cell predictions measure crop-model aggregation error rather than field-observation accuracy
307 (Supplementary Section S3).


308 **2.7. Irrigation strategies and comparisons**

309 **2.7.1. Spatial allocation under common irrigation budgets**

310 Uniform allocation assigns the same irrigation fraction to every represented hectare. Targeted allocation
311 chooses area shares x_{jk} for representative j and simulated irrigation fraction k by linear programming. The
312 objective maximizes $\sum_j A_j \sum_k x_{jk} \bar{Y}_{jk}$, where $\bar{Y}_{jk}$ is mean annual wheat-plus-maize grain in 1997–2013.
313 Constraints require $\sum_k x_{jk} = 1$, $0 \leq x_{jk} \leq 1$ and $\sum_j A_j \sum_k x_{jk} I_k = (1 - r) \sum_j A_j I_1$, with reductions

314 $r = 0.25, 0.50$ or 0.75 . Within-representative mixtures describe shares of area receiving actual simulated
 315 irrigation treatments; crop responses are not interpolated between irrigation fractions. Allocations remain
 316 fixed in 2014–2025. Both policies deliver exactly the same total field-irrigation water in every year, so
 317 their differences reflect allocation under the same volume constraint.

318 Spatial contrasts at the common 50% irrigation cut use frozen shares of complete continuous irrigation histories.
 319 Their area-weighted grain, ET and drainage differences map from 32 representatives to 3,641 source cells. Fixed
 320 southern ($32\text{--}36^\circ\text{N}$), central ($36\text{--}38^\circ\text{N}$) and northern ($38\text{--}41^\circ\text{N}$) bands summarize the distribution. Counts of
 321 positive annual contrasts describe consistency within the modeled record (Figure 5; Table S9).



322 **Figure 5. Spatial redistribution at a common 50% regional irrigation reduction, 2014–2025. (a) Targeted fraction of the conventional**
 323 **quota; uniform allocation retains 50% everywhere. (b,c) Targeted-minus-uniform combined rotation dry grain and crop-plus-fallow**
 324 **ET. (d) Mapped rotation-area shares with positive, negative or unchanged mean grain contrasts. The 3,641 cells inherit 32**
 325 **representative responses. Thin grey provincial lines and black study-domain outlines follow Figure 2; unmapped areas remain blank.**

326 **2.7.2. Annual quotas based on antecedent water availability**

327 Relative water-availability classes combine two antecedent indicators: regional total-water-storage status and local
328 rainfall before wheat sowing. The storage indicator is the mean of available August and September GRACE
329 observations in the calendar year preceding harvest, with at least one monthly solution required. Precipitation is
330 accumulated from 1 July to 30 September at each simulation unit, covering 92 days before the prescribed 12
331 October wheat sowing. The corresponding precipitation/reference-ET ratio is retained as a diagnostic of climatic
332 water availability. These antecedent indicators do not use rainfall or crop outcomes from the subsequent growing
333 season.

334 Thresholds use 2003–2013 harvest years, the common selection period with antecedent GRACE coverage. The
335 regional storage median is 13.38 mm, and the rotation-area-weighted precipitation median is 431.74 mm. Values
336 below their respective thresholds define lower storage and lower rainfall; values at or above define higher storage
337 and higher rainfall. Crossing the two indicators gives four hydroclimatic classes. The classes describe relative
338 antecedent water availability, rather than absolute water scarcity derived from water demand, allocation rights or
339 access. Spatial differences within a year arise from precipitation; GRACE does not supply independently resolved
340 field-scale storage classes.

341 For each class, five continuous irrigation histories provide paired grain and ET responses relative to conventional
342 irrigation, weighted by rotation area and year. The primary candidate is the lowest fraction retaining at least 95% of
343 class-mean conventional grain in 2003–2013; a 98% target assesses a stricter production criterion. Selected
344 fractions and class thresholds remain frozen in 2014–2025. Higher-storage classes have no evaluation exposure.
345 Required GRACE observations are missing for 2014, 2018 and 2019.

346 Two annual strategies apply the irrigation fractions selected from fixed-fraction responses at grain-retention targets
347 of 95% and 98%. At each simulation unit, the antecedent storage–rainfall class determines the fraction before wheat
348 sowing. This fraction scales all four wheat applications and the subsequent maize application in the same rotation
349 year. Application dates, crop parameters, weather, hydraulic profiles and the free-drainage lower boundary are
350 identical to those in the fixed-fraction experiment. Conventional irrigation applies before 2003 and whenever an
351 eligible storage class is unavailable, including evaluation harvest years 2014, 2018 and 2019.

352 Each strategy is simulated continuously from the 1996 spin-up through 2025. Soil-water states pass between crops
353 and fallows and across annual changes in irrigation fraction. Conventional irrigation supplies the paired continuous
354 reference; replays at two simulation units verify the scenario driver against conventional outputs. Class thresholds
355 and irrigation fractions remain fixed after selection. Outcomes cover all twelve comparison years and, separately,

the nine years with eligible classes. Production retention is the ratio of period-mean regional grain production to its corresponding conventional mean; annual ratios describe variation within the period. The strategies assume access to antecedent observation periods, while release availability of GRACE before sowing remains unverified.

2.7.3. Rainfall-only controls and matched monitoring coverage

Rainfall-only controls use the same 2003–2013 simulation-unit years with eligible GRACE observations, the same rotation-area weights, irrigation candidates and grain-retention targets. Selection groups responses by antecedent rainfall alone and chooses the lowest fraction meeting the relevant target. The rainfall threshold is identical to that in the joint classification. Matched controls receive conventional irrigation whenever the storage–rainfall strategy lacks an eligible class, preserving both the selection cohort and monitoring coverage. Continuous differences between these paired strategies consequently assess the incremental effect of storage-based grouping.

Separate rainfall-available controls apply the selected rainfall-only fractions whenever an eligible rainfall group exists, including years without GRACE coverage. Their complete histories assess the combined effect of rainfall-based management and greater monitoring availability. Irrigation changes in earlier gap years can alter the soil-water state of later crops, so these controls are reported separately from the matched comparison.

2.8. Evaluation metrics, spatial statistics and water accounting

Performance is summarized by RMSE, mean bias error, nRMSE relative to observed mean, Nash–Sutcliffe efficiency (NSE) (Nash and Sutcliffe, 1970) and squared Pearson correlation (R^2). Station scores give equal weight to available sites, site-years within sites, crop cases within site-years and records within cases. Wuqiao treatment means receive equal weights within each crop and partition. Field calibration and retrospective testing remain separate. Grain and harvest biomass in station comparisons are dry matter; Wuqiao grain results retain the published 13% moisture basis, with a single 0.87 conversion between dry and reported grain. Biomass remains dry matter throughout.

Seasonal curves group matching observations and predictions into 15-day wheat and 7-day maize bins at observation-window midpoints in days after sowing. Case means receive equal weight within site-years, site-year means within sites and available sites within bins. Standard errors describe variation among contributing site-year means rather than model uncertainty. Site-specific curves use equal site-year weights. Coverage varies among bins, and gaps remain uninterpolated. Accuracy scores use unbinned records. Yucheng harvest error bars describe quadrat sampling variation within crop-years.


ET sums use identical observed and simulated measurement days within each case and retain the station balancing weights. They exclude unmeasured days and do not represent complete-season observed totals. Regional grain production uses dry mass and rotation-area weights; actual ET includes crops and fallows. Paired policy differences cover 2014–2025. Daily water and carbon residuals and continuity of carried soil-water storage assess numerical conservation. Field irrigation, ET, drainage below 2 m and groundwater storage have distinct accounting boundaries.

Spatial distribution statistics describe targeted-minus-uniform policy contrasts in mean annual combined wheat–maize dry grain and water fluxes over 2014–2025. The 3,641 mapped 0.1° source cells carry outcomes from 32 representative response units. Cell weights are the static 2020 mapped rotation areas after cropland-fraction correction; they represent mapped rotation area rather than observed harvested area. Gains and losses exceed $+10^{-6}$ and fall below -10^{-6} kg ha⁻¹, respectively; contrasts within these bounds are classified as unchanged. Group means use rotation-area weights, regional contributions sum area multiplied by the corresponding contrast, and weighted quantiles use the inverse empirical cumulative area distribution. Annual loss frequencies use the same sign threshold across the 12 years. Connected loss footprints comprise occupied cells sharing a horizontal or vertical grid edge (rook adjacency). The statistics describe the conditional modeled allocation footprint; mapped cells do not provide independent replication or cell-specific validation.

3. Results

3.1. Conditional multisite crop comparisons

Retrospective-evaluation LAI RMSE was 1.843 m² m⁻² for wheat and 0.995 m² m⁻² for maize, with NSE of 0.184 and 0.689, respectively (Figure 6). LAI RMSE in the original calibration partition was 1.727 and 1.138 m² m⁻². In-season biomass RMSE in retrospective evaluation was 4.413 and 2.789 t ha⁻¹, with biases of +1.314 and +1.158 t ha⁻¹ and NSE of 0.349 and 0.796. Site-specific seasonal comparisons retain their available observations and original partitions (Figures S6–S11).



407 **Figure 6. Multisite seasonal crop comparisons. Wheat and maize occupy the top and bottom rows; columns show LAI, aboveground**
408 **dry biomass and daily actual ET. Filled and open observations distinguish the original calibration and retrospective-evaluation**
409 **partitions; solid blue and dashed orange lines show the corresponding simulations. Error bars describe standard errors among site-**
410 **year means. Balanced means use 15-day wheat and seven-day maize bins, and unobserved bins remain gaps. Accuracy scores use**
411 **unbinned records. Complete station irrigation management remains unconfirmed.**

412 Dry-grain RMSE was 1.256 t ha^{-1} for wheat and 1.402 t ha^{-1} for maize in the original calibration partition,
413 compared with 0.834 and 2.286 t ha^{-1} in retrospective evaluation (Figure S5). Evaluation grain biases were $+0.485$
414 and -1.586 t ha^{-1} , and NSE was 0.099 and -1.566 . Evaluation harvest-biomass RMSE was 2.736 t ha^{-1} for wheat
415 and 3.719 t ha^{-1} for maize; the corresponding observation counts were 54 in nine wheat years and 48 in eight maize
416 years.

417 Daily ET RMSE in retrospective evaluation was 1.809 mm d^{-1} for wheat and 2.572 mm d^{-1} for maize, with biases
418 of $+0.141$ and -0.734 mm d^{-1} and NSE of 0.121 and 0.012 (Figure 6). ET summed over matched measurement days
419 had evaluation RMSE of 51.29 mm for wheat and 74.17 mm for maize, biases of $+25.94$ and -55.58 mm , and NSE
420 of 0.826 and 0.118 (Table 2). These sums exclude days without ET measurements.

421 **Table 2. Conditional station ET sums on matched measurement days. Observed and simulated sums use identical measurement days**
 422 **within cases, with balanced site, site-year and case weights. Unmeasured days are excluded. Original calibration partitions retain an**
 423 **earlier growth-fitting history; evaluation is retrospective. Complete irrigation-log matching remains unconfirmed.**

Crop	Dataset	Seasons	NSE	RMSE (mm)	MBE (mm)
Maize	Prior calibration partition	11	0.511	45.60	-34.92
Maize	Retrospective evaluation	10	0.118	74.17	-55.58
Wheat	Prior calibration partition	11	-0.300	84.94	-23.53
Wheat	Retrospective evaluation	11	0.826	51.29	25.94

424 **3.2. Wuqiao calibration and retrospective testing**

425 Seasonal ET calibration RMSE was 40.19 mm for wheat and 37.44 mm for maize across 12 treatment seasons per
 426 crop, with biases of −12.65 and +3.60 mm and nRMSE of 8.97% and 12.69% (Table 3; Figure 7). Across the four
 427 2019 testing means per crop, ET RMSE was 73.80 and 55.64 mm, bias was +48.93 and −35.90 mm, nRMSE was
 428 21.98% and 14.13%, and NSE was 0.503 and −0.346, respectively.




Figure 7. Wuqiao calibration and retrospective testing. (a,b) Seasonal ET; (c,d) annual treatment-mean grain at 13% moisture. Wheat occupies the left column and maize the right. Circles show 2016–2018 treatments used to fit water-use parameters; squares show 2019, withheld from the fit and selection. Dashed lines denote 1:1 agreement. Crop-specific grain axes focus on their observed ranges. Horizontal grain error bars are the published SE from three field replicates. Observed ET follows the published 0–2 m water balance; simulated dry grain is converted once by division by 0.87. Maize labels retain preceding wheat management with identical maize irrigation within each year.

At 13% grain moisture, calibration RMSE was 1.249 t ha⁻¹ for wheat and 2.267 t ha⁻¹ for maize; retrospective-testing RMSE was 3.161 and 1.700 t ha⁻¹. Testing grain biases were +3.108 and +1.403 t ha⁻¹, with nRMSE of 83.27% and 15.97% and NSE of -4.654 and -3.641 (Table 3; Figure 7). Dry aboveground-biomass RMSE was 4.222 and 4.688 t ha⁻¹ in calibration and 5.427 and 2.038 t ha⁻¹ in testing, respectively. Testing biomass nRMSE was 55.66% for wheat and 10.44% for maize.

W3–W0 seasonal ET-contrast RMSE was 45.79 mm for wheat and 66.84 mm for maize across three calibration years; the absolute 2019 contrast errors were 116.37 and 52.74 mm. For grain contrasts of W1–W3 relative to same-year W0, calibration RMSE was 0.519 and 1.669 t ha⁻¹ and retrospective-testing RMSE was 1.316 and 1.323 t ha⁻¹, respectively, at 13% moisture (Table S5; Figure S2). Maize treatment labels denote preceding wheat irrigation; maize irrigation was identical within each year.

Table 3. Wuqiao water calibration and retrospective testing.


Crop	Period	Target	n	Unit	RMSE	MBE	nRMSE (%)	NSE	R ²
Maize	2016–2018	Seasonal ET	12	mm	37.444	3.604	12.69	0.533	0.619
Maize	2016–2018	Grain	12	t ha ⁻¹	2.267	-1.738	18.78	-2.940	0.128
Maize	2016–2018	Biomass	12	t ha ⁻¹	4.688	-4.188	21.79	-5.625	0.073
Maize	2019	Seasonal ET	4	mm	55.637	-35.902	14.13	-0.346	0.343
Maize	2019	Grain	4	t ha ⁻¹	1.700	1.403	15.97	-3.641	0.972
Maize	2019	Biomass	4	t ha ⁻¹	2.038	0.726	10.44	-0.313	0.123
Wheat	2016–2018	Seasonal ET	12	mm	40.188	-12.649	8.97	0.689	0.758
Wheat	2016–2018	Grain	12	t ha ⁻¹	1.249	-0.445	16.74	0.746	0.817
Wheat	2016–2018	Biomass	12	t ha ⁻¹	4.222	-2.480	25.48	0.496	0.781
Wheat	2019	Seasonal ET	4	mm	73.805	48.929	21.98	0.503	0.915
Wheat	2019	Grain	4	t ha ⁻¹	3.161	3.108	83.27	-4.654	0.870
Wheat	2019	Biomass	4	t ha ⁻¹	5.427	4.908	55.66	-0.752	0.924

Calibration uses 12 treatment seasons per crop in 2016–2018; retrospective testing uses four in 2019. Counts are treatment means rather than independent replicate seasons. ET is a 0–2 m water-balance estimate under negligible runoff and drainage assumptions. Grain is at 13% moisture and aboveground biomass is dry matter. Annual grain means and SE are digitized from source Supplementary Figure S2. Initial total storage is observed; layer distribution and transferred hydraulics remain conditional.

3.3. Grain production and water balance under irrigation allocation

At a 50% regional irrigation reduction, targeted allocation gave a simulated annual rotation dry-grain yield of 15.761 t ha⁻¹ yr⁻¹ for wheat and maize combined, compared with 15.029 under uniform cuts. Both yields remained below the conventional yield of 16.756 t ha⁻¹ yr⁻¹. Targeting therefore retained 0.732 t ha⁻¹ yr⁻¹ more grain than

455 uniform allocation (4.87% of uniform production), equivalent to 7.219 million t yr⁻¹ across the fixed 9.867 million
 456 rotation hectares (Table 4; Figure 8). Targeted production remained 9.813 million t yr⁻¹ below conventional
 457 production.



458 **Figure 8. Irrigation allocation and grain–water trade-offs during 2014–2025. (a,b) Rotation-area-weighted annual combined wheat–**
 459 **maize dry grain and crop-plus-fallow ET; error bars are between-year SD. (c) Annual targeted-minus-uniform grain at identical**
 460 **irrigation budgets. (d) Period means; connectors join policies with the same budget. Values per hectare use the same 9.867 million**
 461 **mapped rotation hectares, rather than separate crop areas. Targeting retains more grain than uniform cuts while remaining below**
 462 **conventional production.**

463 Both allocations supplied 190 mm yr⁻¹ of field irrigation. Targeted and uniform crop-plus-fallow ET averaged
 464 827.70 and 806.94 mm yr⁻¹, respectively. The grain advantage was accompanied by 20.76 mm yr⁻¹ (2.048 km³ yr⁻¹)
 465 greater ET. Profile drainage was 79.04 versus 96.34 mm yr⁻¹, and runoff was 14.43 versus 18.99 mm yr⁻¹ (Table 4).
 466 The targeted-minus-uniform grain contrast at the 50% cut was positive in all twelve comparison years (Figure 8c).
 467 The positive regional contrast was accompanied by spatially contrasting yield responses (Section 3.4).

468 **Table 4. Simulated annual rotation production and water balance under equal irrigation reductions, 2014–2025.**

Strategy	Dry grain (t ha ⁻¹ yr ⁻¹)	Irrigation (mm yr ⁻¹)	ET (mm yr ⁻¹)	Drainage (mm yr ⁻¹)	Runoff (mm yr ⁻¹)
Conventional	16.756	380.00	887.43	185.26	39.32
Uniform, 25% cut	16.177	285.00	856.41	135.94	25.71
Targeted, 25% cut	16.562	285.00	870.81	125.90	19.86
Uniform, 50% cut	15.029	190.00	806.94	96.34	18.99
Targeted, 50% cut	15.761	190.00	827.70	79.04	14.43
Uniform, 75% cut	13.347	95.00	741.57	68.72	15.83
Targeted, 75% cut	14.018	95.00	752.30	60.01	13.68

469 All per-hectare means use the same 9.866602 million mapped rotation hectares. Dry grain combines wheat and
 470 maize harvests; ET includes crops and fallows. Regional grain in million tonnes equals rotation yield multiplied by
 471 mapped area in million hectares. Regional water in km³ equals depth in mm multiplied by hectares and 10, divided
 472 by 10⁹. The conventional schedule supplies 300 mm to wheat and 80 mm to maize.

473 **3.4. Spatial distribution and persistence of yield contrasts**

474 Relative to uniform allocation, the central and northern latitude bands contributed grain gains of 7.475 and 1.886
 475 million t yr⁻¹, while the southern band contributed a loss of 2.142 million t yr⁻¹. The southern band occupied 60.7%
 476 of mapped rotation area. Mean rotation-yield contrasts were +2.415, +2.420 and -0.357 t ha⁻¹ yr⁻¹ in the central,
 477 northern and southern bands, respectively (Figure 5; Table S9).

478 At the 50% regional irrigation reduction, targeted allocation increased mean annual combined wheat–maize dry
 479 grain relative to uniform reduction on 43.92% of the mapped rotation area (4.333 Mha), decreased it on 37.60%
 480 (3.710 Mha), and left it unchanged on 18.48% (1.823 Mha). Mean contrasts within the gain and loss areas were
 481 +2.439 and -0.903 t ha⁻¹ yr⁻¹, respectively. Their regional contributions were +10.568 and -3.349 Mt yr⁻¹, giving a
 482 net regional grain difference of 7.219 Mt yr⁻¹ (0.732 t ha⁻¹ yr⁻¹ across the full mapped area). Local losses offset
 483 31.69% of the gross positive contribution (Figure 5d).

484 The area-weighted median grain contrast was 0.000 t ha⁻¹ yr⁻¹, and the 10th and 90th percentiles were -1.182 and
 485 +3.553 t ha⁻¹ yr⁻¹. Relative to uniform reduction, mean annual field irrigation increased by 119.0 mm yr⁻¹ in the
 486 gain area and decreased by 139.0 mm yr⁻¹ in the loss area. The corresponding crop-plus-fallow ET contrasts were

487 +94.4 and -55.1 mm yr⁻¹. Irrigation changes balanced at regional scale, whereas the area-weighted mean ET
488 contrast was +20.8 mm yr⁻¹.

489 Negative yield contrasts persisted over 2014–2025: 21.34% of the mapped rotation area had lower yields than under
490 uniform allocation in all 12 years, and 37.60% experienced a loss in at least one year. Every response unit with a
491 negative period-mean contrast recorded losses in at least 9 years; all response units with a positive period-mean
492 contrast recorded gains in all 12 years. Across years, the mean annual area exposed to a loss was 35.37%. Mapped
493 cells with negative period-mean contrasts formed 8 connected components under rook adjacency. The largest
494 contained 3.700 Mha of mapped rotation area (99.73% of all loss area), with occupied cell centers extending from
495 32.0 to 36.3°N and 112.8 to 121.5°E.

496 Negative period-mean yield contrasts occupied 32.80%, 37.60%, and 60.08% of mapped rotation area at reductions
497 of 25%, 50%, and 75%, respectively. At the 75% reduction, the area-weighted median contrast became negative
498 (-0.788 t ha⁻¹ yr⁻¹), although gains on 37.04% of the area produced a positive regional balance. Gross positive and
499 negative contributions at that reduction were +14.100 and -7.487 Mt yr⁻¹, leaving a net gain of 6.614 Mt yr⁻¹, below
500 the net gain at the 50% reduction.

501 **3.5. Antecedent water availability and annual irrigation strategies**

502 Four antecedent storage–rainfall classes occurred during 2003–2013. Nine comparison years had eligible GRACE
503 observations, all in the lower-storage category; three years without eligible GRACE observations received
504 conventional irrigation quotas. At the 95% grain target, both storage strata selected an irrigation fraction of 0.75
505 under lower rainfall and 0.50 under higher rainfall. The 98% target selected 1.00 and 0.75, respectively (Table 5).

506 Continuous strategies selected at the 95% and 98% targets retained 96.27% and 99.63% of conventional grain
507 production over the nine years with eligible GRACE observations. Field-irrigation reductions were 130.72 and
508 35.72 mm per rotation; corresponding ET reductions were 36.84 and 6.17 mm (Figure 9; Table S7). The 95%
509 strategy retained at least 95% in 8 of nine years, with a minimum of 92.12%.




Figure 9. Continuous annual irrigation strategies during 2014–2025. (a) Field irrigation; (b) grain retained relative to conventional irrigation; (c) crop-plus-fallow ET reduction; (d) annual production–irrigation combinations. Blue circles and orange squares identify 95% and 98% selection targets. Shading and open markers identify missing-GRACE years with conventional quotas. Grain and ET can still differ in those years because preceding strategy decisions alter carried soil-water states. Matched rainfall-only histories coincide with the storage–rainfall histories.

Storage–rainfall and matched rainfall-only strategies selected identical irrigation fractions and produced identical grain, ET, drainage and irrigation histories at both targets. Rainfall-only strategies that applied decisions during GRACE gaps followed different irrigation histories and carried different soil-water states into subsequent crops (Table S8).

519 **Table 5. Selected irrigation candidates by relative water-availability class.**

Class	Irrigation fraction	Wheat irrigation (mm)	Maize irrigation (mm)	Selection grain retention (%)
Higher storage dry	0.750	225.000	60.000	95.537
Higher storage wet	0.500	150.000	40.000	97.093
Lower storage dry	0.750	225.000	60.000	96.617
Lower storage wet	0.500	150.000	40.000	97.431

520 Fractions meet the 95% class-mean grain target during selection years 2003–2013. Current shared crop parameters
521 apply throughout. Both storage strata select identical fractions within each rainfall group; higher-storage classes
522 have no later-period exposure.

523 **4. Discussion**

524 We used a calibrated crop model to simulate continuous winter wheat–summer maize rotations across the North
525 China Plain, compare uniform and spatially targeted irrigation under equal water budgets, and test whether
526 antecedent regional water storage improved decisions based on rainfall. Sustaining grain production under restricted
527 irrigation requires consideration of both consumptive water use and the distribution of production losses across
528 locations. The conditional results show that higher regional grain retention can accompany greater
529 evapotranspiration and lower production in some locations, whereas storage information did not change the selected
530 irrigation quotas within the tested design. The management value of these strategies therefore depends on
531 accounting for the complete rotation water balance, soil-water carry-over across crops and fallows, and reliable crop
532 responses to irrigation.

533 **4.1. Marginal irrigation returns and spatial allocation**

534 Spatial targeting exploits differences in marginal grain response to irrigation. Diminishing production gains across
535 field schedules (Yang et al., 2022) and spatial variation in irrigation requirements (Wu et al., 2019) provide a basis
536 for redistributing a limited supply, consistent with regional allocation studies (Jiang et al., 2016). The equal-budget
537 comparison attributes the simulated production difference to delivery allocation. Complete continuous histories
538 retain the effects of earlier irrigation and rainfall on subsequent crop responses.

539 At the 50% delivery reduction, targeting retained $0.732 \text{ t ha}^{-1} \text{ yr}^{-1}$ more combined wheat–maize dry grain than
540 uniform cuts. Across 9.8666 million rotation hectares, the corresponding reduction in grain loss was 7.219 million t
541 yr^{-1} , or 4.87% of the uniform total. Targeted production remained below conventional production. The positive
542 contrast in all twelve years indicates consistent regional gains within the simulated record. Its magnitude remains
543 dependent on the crop-response functions and declared management schedules (Table 4; Figure 8).

544 The spatial distribution sharpens this distinction between regional and local outcomes. Relative to uniform
545 allocation, mean losses occupy 37.60% of mapped rotation area, and losses persist in every comparison year on
546 21.34%. Gross gains of $10.568 \text{ Mt yr}^{-1}$ are partly offset by losses of 3.349 Mt yr^{-1} ; the area-weighted median is
547 zero despite a positive regional mean. The regional grain objective balances gains and losses among locations.
548 Local production floors constitute an additional allocation constraint.

549 Regional gain has a pronounced distributional cost. Central and northern band-mean rotation yields increased by
550 2.415 and $2.420 \text{ t ha}^{-1} \text{ yr}^{-1}$, while the southern mean decreased by $0.357 \text{ t ha}^{-1} \text{ yr}^{-1}$. The southern band covers
551 60.7% of the rotation area, so its losses substantially offset gains elsewhere (Figure 5; Table S9). The production
552 objective maximizes the area-weighted regional total and supplies no minimum yield for an individual location.
553 Local production floors and constraints on the distribution of irrigation reductions are therefore separate
554 requirements for allocation that protects affected farms.

555 **4.2. Rotation carry-over and the timing of water supply**

556 The spatial production response integrates water effects across both crops. Wheat irrigation can support canopy
557 growth and grain formation, replenish water available to maize or increase drainage. Reduced wheat delivery can
558 increase subsequent dependence on summer rainfall, producing different annual grain responses under different
559 rainfall sequences. The sixteen-year irrigation experiment shows diminishing production gains, and the Wuqiao
560 experiment links precipitation conditions with wheat scheduling and subsequent maize production (Yang et al.,
561 2024; Yang et al., 2022). These coupled responses make the complete rotation the relevant unit for irrigation
562 allocation.

563 Continuous soil-water histories preserve the delayed benefits and costs of allocation and annual quota changes
564 (Figure S3). Production can be maintained by drawing down profile storage, with later rainfall or irrigation
565 determining its replenishment. A finite-period delivery reduction can therefore include a contribution from stored

566 soil water. Crop-plus-fallow accounting also captures evaporation and drainage between harvest and sowing. Their
567 inclusion connects apparent production retention with the water remaining for the next crop and the complete
568 rotation water balance.

569 The field evidence constrains only part of this carry-over mechanism. Wuqiao simulations use reported total profile
570 storage with an assumed layer distribution. Wheat crop-start storage follows the pre-sowing simulation, whereas
571 maize is independently initialized from treatment-specific observations. Maize irrigation is identical among
572 treatments within each year, and its treatment labels identify preceding wheat schedules (Yang et al., 2024). Maize
573 comparisons thus evaluate crop responses to the initial state associated with wheat management (Figure 7; Figure
574 S2). Regional carry-over additionally depends on the model's continuous propagation and vertical redistribution of
575 water between crops.

576 **4.3. Production, consumptive use and groundwater accounting**

577 The production benefit of targeting carries a consumptive-use trade-off. At the common 50% delivery reduction,
578 targeting increased ET by $2.048 \text{ km}^3 \text{ yr}^{-1}$ and reduced profile drainage by $1.707 \text{ km}^3 \text{ yr}^{-1}$ relative to uniform
579 allocation. With rainfall and total irrigation held constant, differences in ET are balanced by drainage, runoff and
580 soil-storage change (Table 4). A shared delivery constraint therefore permits different consumptive outcomes.
581 Irrigation productivity and consumptive-water savings require distinct objectives (Grafton et al., 2018; Perry et al.,
582 2009).

583 The soil-profile balance also has a different boundary from the aquifer balance. Drainage below 2 m enters a deeper
584 vadose zone, with storage and transit outside the modeled profile (Wu et al., 2023). Reduced field delivery or
585 profile drainage therefore provides insufficient information to quantify changes in pumping, aquifer recharge or
586 groundwater storage. Production and ET comparisons can identify agronomic trade-offs, while groundwater
587 outcomes require the corresponding subsurface and abstraction accounting.

588 GRACE offers regional storage context at a broader scale. Its anomalies integrate groundwater with other terrestrial
589 stores, and its effective observation scale exceeds the displayed grid resolution (Save et al., 2016; Scanlon et al.,
590 2016). Recent aquifer recovery reflects water transfers, abstraction management and hydroclimatic conditions
591 (Long et al., 2025; Xu et al., 2026). The crop-model scenarios quantify changes in field delivery, consumption and
592 profile outflow without attributing those observed storage changes to individual irrigation strategies.

593 **4.4. When storage information changes an irrigation decision**

594 A storage indicator adds decision value when its inclusion changes the selected action or improves the grain–water
595 outcome under common constraints. GRACE represents regional storage conditions, while antecedent rainfall
596 describes more local water input (Save et al., 2016; Scanlon et al., 2016). Their different physical support motivates
597 joint classification, but decision value also depends on how the resulting classes distinguish among the available
598 irrigation fractions.

599 Both storage strata selected the same fraction for a given rainfall group at the 95% and 98% grain targets. Storage–
600 rainfall and matched rainfall-only strategies consequently produced identical complete histories of irrigation, crop
601 states, grain, ET and drainage. The storage split supplied additional environmental description without changing an
602 irrigation action in this five-level quota design. All nine later years with eligible observations fell in the lower-
603 storage category, so the higher-storage choices also lacked later-period exposure (Table 5; Figure 9). These features
604 delimit the configuration and conditions represented by the equal outcomes.

605 Monitoring coverage must remain distinct from the grouping criterion. Rainfall-available controls can change
606 irrigation during GRACE gaps, and those earlier decisions alter the soil-water state inherited in later years. Their
607 outcomes therefore combine rainfall-based selection with additional years of management. Matched coverage
608 preserves the appropriate comparison for the incremental effect of storage information. Finer quota choices,
609 alternative thresholds and verified release availability would define different decision experiments with their own
610 selection and evaluation requirements.

611 **4.5. Crop-response constraints and management reliability**

612 Management selection depends on the modeled marginal response to water, in addition to absolute grain and canopy
613 agreement. Intercepted-PAR radiation-use efficiency (Kiniry et al., 1989), cohort-specific leaf area and thermal
614 senescence connect canopy development with carbon supply; grain number and fixed individual grain mass
615 constrain the sink. Root uptake and the assimilation stress exponent then determine how water shortage changes
616 production. Variation among wheat-model transpiration formulations demonstrates the sensitivity of water-use
617 predictions to these processes (Cammarano et al., 2016). Water coefficients near their prescribed bounds remain
618 effective empirical estimates whose physiological interpretation and uniqueness are limited.

619 The estimation and evaluation data provide different evidence for these responses. Earlier growth estimates use 480
620 wheat and 433 maize observations across four wheat sites and five maize sites, with whole site-years and balanced
621 site contributions. Incomplete station irrigation histories limit transfer comparisons, and station observations have

no weight in the fitted water-response objective. Wuqiao 2016–2018 supplies the documented treatment seasons for that fit. Wuqiao 2019 supplies a single test year with reported total storage, assumed layer distributions and transferred soil hydraulics; prior inspection of its outcomes makes the test retrospective (Figures 3, 6 and 7). Regional comparison years also overlap crop-fitting years, so regional production differences characterize model scenarios rather than time-forward crop validation.

The absolute regional production reference also depends on the prescribed management and crop parameters. Conventional simulations produce 8.065 t ha⁻¹ of wheat dry grain and 8.691 t ha⁻¹ of maize dry grain per year. These values represent fixed irrigation dates, crop calendars and nutrition modifiers rather than an independently validated regional harvest inventory. The 0.732 t ha⁻¹ yr⁻¹ allocation advantage depends on the same simulated marginal responses; its regional scaling does not remove field-response uncertainty.

Substantial field-test errors constrain confidence in irrigation-response transfer. Wheat grain nRMSE was 83.27%, with a bias of +3.108 t ha⁻¹ at the published 13% moisture basis; the absolute W3–W0 ET-contrast error was 116.37 mm. Maize ET NSE was negative, with an ET-contrast error of 52.74 mm (Table 3; Table S5). These discrepancies affect the response functions used to select regional allocations. Numerical water-balance closure demonstrates conservation but supplies no measure of response accuracy. The eight purposive actual-cell checks address a separate uncertainty, spatial aggregation, and provide no regional field-validation sample. Depth-resolved soil water and complete dated management would better constrain uptake and redistribution; canopy and soil-moisture assimilation also requires treatment of retrieval errors and observation operators (Ines et al., 2013).

Production-retention targets introduce a further management distinction between period means and annual reliability. Mean retention of 96.27% and 99.63% coincided with annual minima of 92.12% and 98.58%; the 95% strategy met that threshold in eight of nine class-available years (Figure 9). A mean target consequently supplies no shortfall probability or guarantee for an individual farm. Local production floors, acceptable shortfall frequency and a reserve entering the next crop can complement regional delivery and consumption constraints. Forecast-based irrigation studies illustrate explicit treatment of water-stress risk (Roy et al., 2021). Field scheduling additionally requires current soil-water information, weather forecasts and application timing.

4.6. Limitations and future directions

The principal limitation is uncertainty in the crop responses used for strategy selection. Field testing covers a single, previously inspected year, while incomplete station management and assumed initial water profiles constrain transfer. Remaining errors in ET and grain responses prevent the simulated policy advantages from being

651 interpreted as independently validated field benefits. Further evaluation requires new site-years with complete
652 management records, withheld from both model development and strategy selection. Concurrent measurements of
653 soil water by depth, ET, canopy development and harvest would allow separate assessment of water redistribution,
654 crop stress and irrigation-treatment contrasts.

655 Regional results also depend on a static rotation map, representative climate–soil units, prescribed irrigation dates
656 and fixed nutrition modifiers. Checks at purposively selected source cells cannot bound regional uncertainty.

657 Dynamic nitrogen cycling is absent, and changes in crop area or actual farm practices are not reconstructed. Future
658 analyses should expand evaluation at individual source cells and propagate uncertainty in crop parameters, soil
659 hydraulics, weather and spatial aggregation through paired policy histories. Allocation stability and the magnitude
660 and persistence of local losses require uncertainty assessment alongside the regional mean.

661 The storage comparison is specific to the tested quota grid, classification thresholds and lower-storage conditions
662 represented during later comparison years. GRACE release availability before sowing remains unverified, so the
663 scenarios do not establish operational scheduling performance. Prospective decision experiments should use
664 information available on each decision date, maintain matched rainfall-only controls and evaluate both storage
665 states. Finer quotas and application timing informed by soil water could be tested with constraints on consumptive
666 use, local production and the risk of production shortfalls.

667 Finally, the profile uses free drainage at its lower boundary, with no capillary supply or deeper subsurface storage,
668 and field irrigation is not linked to measured abstraction. Groundwater applications require joint accounting of soil
669 water and aquifer flows, constrained by observations of pumping, recharge and storage change. Delivery capacity,
670 irrigation costs and farmer participation also require assessment before the agronomic scenarios can support
671 operational management.

672 **5. Conclusions**

673 At an identical 50% irrigation reduction, targeted allocation retained $0.732 \text{ t ha}^{-1} \text{ yr}^{-1}$ more simulated combined
674 wheat–maize dry grain than uniform cuts during 2014–2025. This reduced regional grain loss by 7.219 million t
675 yr^{-1} (4.87% of uniform production), while increasing crop-plus-fallow ET by 20.76 mm yr^{-1} ($2.048 \text{ km}^3 \text{ yr}^{-1}$).

676 Relative to uniform allocation, mean grain losses occurred on 37.60% of mapped rotation area and persisted in all
677 twelve years on 21.34%. Regional production retention therefore requires separate consumption and local
678 production constraints.

679 Annual strategies selected at 95% and 98% grain-retention targets retained 96.27% and 99.63% of conventional
680 production over nine years with eligible GRACE observations. The 95% strategy reduced irrigation by 130.72 mm
681 and ET by 36.84 mm per rotation. Matched rainfall-only controls selected identical quotas and produced identical
682 outcomes, so storage classification provided no additional decision benefit within the tested quota design and
683 monitoring coverage.

684 Continuous rotation accounting preserves the influence of preceding management on subsequent production and
685 water use. The field-test errors and incomplete station management delimit confidence in transferring the estimated
686 crop responses. Translation of these regional scenarios into irrigation management depends on observed response
687 accuracy, explicit production-risk constraints and field scheduling informed by soil water and forecasts.
688 Groundwater effects additionally require pumping, recharge and deeper subsurface storage beyond the crop-model
689 boundary.

690 **Data and code availability**

691 Model code, frozen calibration and regional inputs, observation subsets and quality exclusions, weather, calibrated
692 parameters, predictions, evaluation metrics and figure-data tables are available at [https://github.com/SmartAG-](https://github.com/SmartAG-Team/agri_water_management)
693 [Team/agri_water_management](https://github.com/SmartAG-Team/agri_water_management). The repository includes a standalone Open Crop Model package, pinned
694 dependencies, reproduction commands and SHA-256 provenance. Reproduction uses the included inputs without
695 another model checkout or access to the historical workspace. Original datasets retain their provider licences and
696 attribution requirements, documented in the repository source catalogue.

697 **Declaration of generative AI and AI-assisted technologies**

698 AI assisted with grammar checking.

699 **Supplementary material**

700 **Supplementary methods and results**

701 **S1. Observation sources and eligibility**

702 Fengqiu, Luancheng and Shangqiu crop phenology, leaf area and biomass records were obtained from the station
703 growth-monitoring datasets (Fengqiu Station, 2020a; Luancheng Station, 2020; Shangqiu Station, 2021). The
704 Gucheng sowing-date dataset supplies maize development and LAI observations (Geng et al., 2023). Yucheng
705 records combine the 1998–2006 crop archive and the 2005–2022 biological and water-environment archives
706 (Yucheng Station, 2025b; Yucheng Station, 2025a; Yucheng Station, 2021). Fengqiu ET and antecedent soil-
707 moisture profiles come from the farmland soil-water physical-properties archive (Fengqiu Station, 2020b).

708 Published crop-model evaluations used Fengqiu, Luancheng and Yucheng observations (Chen et al., 2010a; Yu et
709 al., 2006). Regional crop-water-use evaluation also used CERN monitoring records from Fengqiu, Luancheng and
710 Shangqiu (Wang and Shi, 2024). These publications document earlier use of the station monitoring programs; the
711 exact records, plots and measurement periods retained here are identified by the source datasets and case inventory
712 (Tables S1 and S2).

713 The station archive preserves 132 crop–plot–season trajectories, whole site-year partitions and source metadata. The
714 7,397 eligible records occur in 127 cases and retain all five observation sites for their available variables.

715 Calibration and retrospective-evaluation trajectory counts remain 71 and 61. Variable-specific availability and
716 missing combinations are retained in the accompanying case and site tables; missing measurements do not become
717 zero observations.

718 Wheat LAI above $7 \text{ m}^2 \text{ m}^{-2}$ remains excluded under the study-specific rule. Duplicated six-replicate maize harvest
719 vectors in Yucheng 2012 and 2017 account for 24 excluded grain and biomass records, divided equally between the
720 original calibration and evaluation partitions. These source-based exclusions retain raw values and do not depend
721 on prediction residuals.

722 Earlier multisite growth estimates retain 480 wheat and 433 maize calibration observations. Current station growth
723 and ET comparisons are conditional on archived management whose full irrigation-log completeness is unresolved.
724 They have zero weight in the water-parameter fit. Wuqiao 2016–2018 supplies 12 documented treatment seasons
725 per crop for that fit (Yang et al., 2024); 2019 supplies four retrospective testing seasons per crop. Seasonal ET, grain
726 and biomass targets refer to the same treatment-season population, while ET contrasts are derived from those
727 seasonal targets.

728 Twenty-six Fengqiu cases retain antecedent moisture initialization from the preceding calendar month (Fengqiu
729 Station, 2020b). Volumetric percentages from fifteen 100-mm source layers spanning 0–1.5 m map conservatively
730 to the represented model layers, preserving depth-integrated storage without depth extrapolation or clipping. The
731 identical eligibility rule applies to 11 original calibration and 15 evaluation cases. Monthly timing, aggregation and
732 subplot support remain unresolved relative to the exact sowing-day state. Original cases remain available when a
733 measured antecedent profile is absent.

734 Observation eligibility requires valid dates, compatible crop and plot identities, consistent stage sequences and
735 documented quantity units. Measurements and predictions share recorded sampling dates or documented averaging
736 windows. Stage-indexed Gucheng LAI measurements use observed stage dates as sampling coordinates without
737 prescribing simulated development. Fengqiu ET identifies the lysimeter measurement plot, whereas Yucheng ET
738 retains its source measurement definitions. Provider grain and aboveground dry masses in g m^{-2} are multiplied by
739 10 to obtain kg ha^{-1} .

740 **Table S1. Crop seasons and calibration/evaluation case counts by site.**

Site	Crop	Years	Calibration cases	Evaluation cases
Fengqiu	Maize	2004, 2005, 2006, 2007, 2008	11	9
Fengqiu	Wheat	2004, 2005, 2006, 2007, 2008	11	10
Gucheng	Maize	2018, 2019, 2020	8	4
Luancheng	Maize	2004, 2005, 2006, 2007, 2008	2	3
Luancheng	Wheat	2004, 2005, 2006, 2007, 2008	2	3
Shangqiu	Maize	2002, 2004, 2005, 2006, 2007	2	6
Shangqiu	Wheat	2001, 2002, 2003, 2004, 2005, 2006, 2007	9	8
Yucheng	Maize	1998, 2000, 2001, 2002, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022	14	9
Yucheng	Wheat	1999, 2000, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022	12	9

741 Years denote archived crop-season groups, which retain all their plot cases. Coverage is intermittent; years absent
742 from the table are not assumed to have observations.

743 **Table S2. Source datasets for evaluated field observations.**

Site	Dataset	DOI	Source
Fengqiu	2004-2008年河南封丘站	10.12199/nesdc.ecodb.mon.	National Ecosystem
	作物生育及生长动态监测	2020.dp2011.fqa.002	Science Data Center /
	数据		Science Data Bank
Fengqiu	2005-2008年河南封丘站	10.12199/nesdc.ecodb.mon.	National Ecosystem
	农田土壤水分物理性质监	2020.dp2011.fqa.006	Science Data Center /
	测数据		Science Data Bank
Gucheng	2018-2021年华北北部不	10.57760/sciencedb.o00119	National Ecosystem
	同播期对玉米生长发育的	.00017	Science Data Center /
	影响数据集		Science Data Bank
Luancheng	2000-2008年河北栾城站	10.12199/nesdc.ecodb.mon.	National Ecosystem
	作物生育及生长动态监测	2020.dp2011.lca.002	Science Data Center /
	数据		Science Data Bank
Shangqiu	2000-2007年河南商丘站	10.12199/nesdc.ecodb.mon.	National Ecosystem
	作物生育及生长动态监测	2020.dp2011.sqa.003	Science Data Center /
	数据		Science Data Bank
Yucheng	1998-2006年山东禹城站	10.12199/nesdc.ecodb.mon.	National Ecosystem
	作物生育及生长动态监测	2020.dp2011.yca.003	Science Data Center /
	数据		Science Data Bank
Yucheng	禹城站2005-2022年生物	10.12199/nesdc.ecodb.2021	National Ecosystem
	要素观测数据集	YFF0703900.yca.2025.2	Science Data Center /
			Science Data Bank
Yucheng	禹城站2005-2022年水环	10.12199/nesdc.ecodb.2021	National Ecosystem
	境要素观测数据集	YFF0703900.yca.2025.3	Science Data Center /
			Science Data Bank

744 Dataset DOIs identify official source records; original metadata and observation-row provenance are retained in the
745 reproducibility archive.

746 **S2. Crop processes, parameter definitions and water accounting**

747 **S2.1. Model structure and phenological development**

748 Open Crop Model simulates phenology, green leaf area, live and dead aboveground biomass, grain dry matter,
749 rooting depth and layered soil-water storage. Inputs comprise weather, soil profiles, crop parameters, dated sowing
750 and harvest, and irrigation events. Soil water carries continuously between crop and fallow segments in regional
751 histories; Wuqiao crop starts retain their separately specified storage conditioning.

752 Wheat stage development combines temperature, vernalization and photoperiod modifiers with inherited stage
753 requirements. Maize uses a subdaily temperature response reconstructed from daily extrema, with base, optimum
754 and maximum temperatures of 6, 30 and 44 °C. The shared medium-maturity maize card requires approximately
755 1,909 °C d to reach R6. DVS anchors are wheat BBCH 09, 61 and 89 and maize VE, R1 and R6 (Meier, 2018).

756 Wheat dough stage and full ripeness retain their distinct definitions.

757 Seed-layer water above wilting-point storage permits germination. The represented layer is selected by sowing
758 depth, with a 50-mm prior when no depth is supplied. Thermal development and wheat vernalization begin from the
759 first eligible germination day rather than accumulating during dry-seed days. This layer-average availability is an
760 effective establishment condition, not a measured seed-scale water potential. Fixed stage coefficients therefore do
761 not imply fixed calendar-stage predictions across water states. New biomass production ends at wheat full ripeness
762 or maize physiological maturity; green maize area can persist under the terminal senescence rate until harvest.
763 Harvest is a separate management endpoint and does not substitute for an observed maturity date.

764 **S2.2. Canopy development, biomass accumulation and grain yield**

765 Green cover is $f_c = 1 - \exp(-k_L \text{LAI})$. Daily aboveground dry-matter production is
766 $\Delta B = 10R_s p_{\text{PAR}} f_c \varepsilon f_T f_W^\gamma f_N$, where R_s is incident shortwave radiation, $p_{\text{PAR}} = 0.45$, ε is RUE per MJ of
767 intercepted PAR, f_T is the temperature modifier, f_N is the prescribed nutrition factor and f_W is realized
768 transpiration divided by remaining potential transpiration demand after wet-canopy evaporation. The
769 factor 10 converts g m^{-2} to kg ha^{-1} . PAR enters once (Kiniry et al., 1989). The fitted empirical
770 assimilation exponents are 1.85985 for wheat and 0.25000 for maize. They govern carbon response
771 separately from leaf-expansion and leaf-survival sensitivity.

SLA is linearly interpolated through DVS 0, 0.5, 1 and 2, with the DVS-2 value equal to DVS 1. New leaf dry matter acquires area using current SLA; established cohorts retain their birth area-to-mass ratio (de Wit and Berghuijs, n.d.). Thermal senescence hazard $h(\text{DVS})$ is interpolated through DVS 0, 1, 1.4, 1.7 and 2, with a constant preflowering rate and increasing grain-filling rates (Tan et al., 2020). Daily survival is $\exp[-h(\text{DVS}) \max(T_{\text{mean}} - T_{\text{base}}, 0)](1 - q_s)$, with $q_s = 0.025(1 - f_w f_N)$. The terminal hazard continues at maize R6, whereas wheat BBCH 89 senesces remaining green leaves. LAI emerges from the cohort area balance; no maximum-LAI parameter or cap enters this formulation.

Leaf allocation follows crop development, with separate early and late vegetative multipliers for maize and additional temperature, water-supply and nutrition limits on leaf expansion. Initial LAI is an emergence condition. Stage-specific SLA and senescence coefficients retain their inherited estimates and conditional trait priors (APSIM Initiative, 2026; de Wit, 2022; de Wit, 2020); their values do not represent universal physiological bounds.

Grain number is fixed at the first grain-set day as $N_g = 0.1B_f c_g$, where B_f is simulated aboveground dry mass immediately before grain set and c_g is the grain-number coefficient. Grain set starts at wheat BBCH 61 and maize R1. Maximum grain dry mass is $Q_g = 10N_g m_g$, with fixed m_g of 0.045 g grain⁻¹ for wheat and 0.300 g grain⁻¹ for maize. Assimilate supply and development-dependent filling capacity constrain grain accumulation. Wheat remobilization transfers an explicit subset of stem mass to grain while conserving total aboveground mass. Live and dead leaves, stems and grain remain in the aboveground carbon pool; observed yield or harvest index does not initialize grain.

Prescribed nutrition factors distinguish archived fertilized and unfertilized management classes. Wuqiao and regional simulations retain the fertilized class. These factors describe effective growth limitations; the model does not resolve a mechanistic nitrogen balance.

S2.3. Soil-water balance and actual evapotranspiration

Potential transpiration and exposed-soil evaporation are $T_p = K_T \text{ET}_0 f_c$ and $E_p = K_E \text{ET}_0 (1 - f_c)$, with ET_0 supplied by AgERA5. Rainfall interception fills a canopy store of capacity 1 mm $\times f_c$. Wet-canopy evaporation reduces the transpiration demand remaining within each outer water step. Actual ET is $T_a + E_s + E_c$, the sum of transpiration, soil evaporation and canopy evaporation. Wuqiao residue cover is unmeasured and its exposed-soil boundary is retained; prescribed organic mulch, where supplied, reduces potential soil evaporation using the FAO-56 cover approximation (Allen et al., 1998).

Two-stage evaporation depends on actual surface-layer storage. Total evaporable water is
 $TEW = (\theta_{FC} - \theta_{AD})\Delta z_1$ and readily evaporable water is $REW = 0.30TEW$. Surface depletion is
 $D_e = (\theta_{FC} - \theta_1)\Delta z_1$. The evaporation reduction is $K_r = \min \left[1, \max \left(0, \frac{TEW - D_e}{TEW - REW} \right) \right]$, with actual
extraction bounded by storage above air dry. Ponded water evaporates before soil storage. This storage
closure follows the two-stage drying principle (Allen et al., 1998) without resolving a complete dual-crop-
coefficient surface balance.

Root activity integrates an exponential depth distribution over rooted parts of each layer and is normalized
over the represented rooted volume. Local depletion stress is $s_i = \min \left[1, \frac{\max(0, W_i - \theta_{WP,i}\Delta z_i)}{(1-p)(\theta_{FC,i} - \theta_{WP,i})\Delta z_i} \right]$, where p
is the fitted readily available water fraction. Base uptake combines root activity, local stress, atmospheric
demand and the daily extraction coefficient; accessible storage and extraction capacity bound each
contribution. The compensation fraction allocates a bounded part of unmet demand to rooted layers whose
stress factors equal or exceed the activity-weighted mean. Total uptake remains bounded by demand times
the best rooted-layer stress factor and by available extraction capacity. Compensation redistributes
demand and does not eliminate uniformly dry-profile stress.

Effective saturation is $S_e = \frac{\theta - \theta_{AD}}{\theta_{SAT} - \theta_{AD}}$. With $m = 1 - \frac{1}{n}$, the retention relation is $S_e = [1 + (\alpha |h|)^n]^{-m}$
(van Genuchten, 1980), and conductivity is $K = K_{sat} S_e^{1/2} \left[1 - (1 - S_e^{1/m})^m \right]^2$ (Mualem, 1976; van
Genuchten, 1980). Adjacent half-layer saturated resistances define $K_{sat, face} = \frac{d}{\frac{\Delta z_i}{2K_{sat,i}} + \frac{\Delta z_{i+1}}{2K_{sat,i+1}}}$, where
 $d = \frac{\Delta z_i + \Delta z_{i+1}}{2}$. Relative-arithmetic face mobility is $r_{face} = \frac{\Delta z_{i+1}r_i + \Delta z_i r_{i+1}}{\Delta z_i + \Delta z_{i+1}}$, with $r_i = \frac{K_i}{K_{sat,i}}$. Thus
 $K_{face} = K_{sat, face} r_{face}$. This interface approximation permits a wet layer to supply a dry neighbor while
retaining the resistance of low-conductivity material.

Signed interface flux is $q_i = K_{face} \left[1 + \frac{h_i - h_{i+1}}{d} \right]$, positive downward. Transfers have equal and opposite
contributions in neighboring layer balances and are bounded by donor storage and receiving pore capacity.
Twenty-four outer steps partition daily forcing, evaporation and uptake; adaptive internal transport steps
limit net θ increments to 0.005. The free-drainage bottom flux equals the current unsaturated conductivity
under unit hydraulic gradient. Upward redistribution is an internal profile flux, with no external capillary-
rise supply. This bounded explicit layered approximation does not resolve the full implicit Richards
system or saturated positive pressure.

827 For a field of profile storage S , the accounting identity is $\Delta S = P + I - ET_a - D - Q$. Applied irrigation I
828 is a field-boundary input. Flood-irrigation scenarios wet the entire field, with application, drift and
829 conveyance losses set to zero at that boundary. Drainage D leaves the modeled profile and is distinct from
830 water reaching an aquifer after deeper vadose-zone transport (Wu et al., 2023).

831 **S2.4. Hydraulic derivation, shared crop parameters and numerical checks**

832 Saxton–Rawls equations use normalized sand and clay fractions and organic matter percent (Saxton and
833 Rawls, 2006). SoilGrids carbon in dg kg^{-1} is converted to organic matter percent by $\frac{1.724 \times \text{SOC}}{100}$; bulk
834 density in cg cm^{-3} is divided by 100; coarse fragments in permille are divided by 1,000 (ISRIC — World
835 Soil Information, 2020). Regional porosity uses $1 - \frac{\text{BD}}{2.65}$, with prescribed density and gravel adjustments.
836 Air-dry water content retains its declared prior. Field-capacity, wilting-point and saturation ordering is
837 checked before simulation; unavailable complete profiles remain excluded under the spatial-source rule.

838 The van Genuchten α and n values are conditional hydraulic priors reconstructed from supplied field-
839 capacity and wilting-point endpoints at pressure heads of $-3,300$ and $-150,000$ mm, with θ_{AD} as residual
840 water and θ_{SAT} as saturated content. Endpoint reconstruction preserves those storage thresholds rather
841 than fitting curves to crop outcomes. Neither depth-resolved Wuqiao retention measurements nor exact
842 seasonal initial layer distributions are available. The separate Wuqiao particle-density prior is 2.75 g cm^{-3} .

843 Table S3 contains effective shared crop values after applying the frozen fit multipliers. Regional and Wuqiao inputs
844 use identical shared cards within each crop. Maximum root depth is limited by the supplied profile depth. Stage-
845 specific SLA and senescence, fixed grain mass and inherited growth coefficients remain distinct from the eight
846 water coefficients fitted in the current run. Near-zero root-density decay yields approximately uniform activity over
847 represented rooted volume; it does not identify an observed root-mass profile.

848 The 164 selected station and field trajectories conserve daily water and aboveground carbon, with maximum
849 absolute residuals of 1.00×10^{-12} mm and 6.71×10^{-12} kg ha $^{-1}$. Exact inputs, observation identities, frozen
850 parameters, source snapshots and SHA-256 provenance accompany the predictions. Conservation establishes
851 arithmetic consistency; hydraulic and crop-response accuracy depend on the conditional observations and
852 retrospective field test.

853 Eight existing water-response coefficients per crop are estimated from documented Wuqiao management in 2016–
854 2018: the transpiration multiplier, soil evaporation coefficient, readily available water fraction, assimilation water-
855 stress exponent, root-density decay, maximum rooting depth, $\log_{10}$ daily root extraction coefficient and root
856 compensation fraction. RUE, leaf allocation and grain-number multipliers remain fixed. Each crop contributes 12
857 seasonal ET, 12 dry-grain and 12 dry-biomass targets, together with three W3–W0 ET contrasts derived from the
858 same seasonal targets. The criterion retains weights of 0.325, 0.175, 0.15 and 0.10 for those components. ET
859 residual scales are $\max(50 \text{ mm}, 0.15 \times \text{observed ET})$; grain and biomass scales are $\max(1,000 \text{ kg ha}^{-1}, 0.20 \times$
860 $\text{observed dry mass})$. Contrast residuals use 50 mm. Eight coefficient-prior terms have total weight 0.01, with
861 inherited bounds, centres and scales.

862 Bounded least squares starts from the inherited water vector, with coefficients scaled to their prescribed intervals, a
863 40-evaluation limit and convergence tolerances of 8×10^{-5} . Reference and refitted coefficients use the identical
864 field-only criterion; the equal-crop mean calibration data loss is 0.592580 for the reference and 0.577934 for the
865 selected refit. Selection excludes prior penalties from this comparison and precedes 2019 testing. Station
866 observations, daily ET and phase ET with unresolved sampling windows have zero weight in the water-parameter
867 fit. Source observations, weather, soil priors, initialization rules and native process equations remain fixed during
868 this refit.

869 **Table S3. Shared crop parameters transferred to regional simulations.**

Parameter	Unit	Wheat	Maize
Radiation-use efficiency	g MJ ⁻¹ intercepted PAR	3.082080	3.379974
Specific leaf area at DVS 0	m ² g ⁻¹	0.028482	0.024045
Specific leaf area at DVS 0.5	m ² g ⁻¹	0.027578	0.029960
Specific leaf area at DVS 1	m ² g ⁻¹	0.023462	0.024000
Leaf senescence rate before flowering	(°C d) ⁻¹	0.000536	0.000010
Leaf senescence rate at DVS 1.4	(°C d) ⁻¹	0.001907	0.000023
Leaf senescence rate at DVS 1.7	(°C d) ⁻¹	0.005327	0.001650
Leaf area index at emergence	m ² m ⁻²	0.141900	0.070844
Grain number per biomass at grain set	grains g ⁻¹	26.272389	4.242740
Maximum individual grain mass	g grain ⁻¹	0.045000	0.300000
Transpiration coefficient	dimensionless	0.838813	1.023273
Soil evaporation coefficient	dimensionless	1.295161	0.100000
Root water uptake coefficient	d ⁻¹	0.042671	0.033543
Leaf biomass partitioning multiplier	dimensionless	1.010797	—
Early vegetative leaf partitioning multiplier	dimensionless	—	0.912989
Late vegetative leaf partitioning multiplier	dimensionless	—	1.779539
Stem reserve fraction	dimensionless	0.270612	—
Stem reserve release fraction	dimensionless	0.067334	—
Assimilation water-stress exponent	dimensionless	1.859851	0.250000
Readily available water fraction	dimensionless	0.543149	0.674994
Root density decay	m ⁻¹	3.41 × 10 ⁻¹⁰	2.27 × 10 ⁻¹³
Maximum rooting depth	mm	1888.654910	1699.999998
Root compensation fraction	dimensionless	1.000000	6.62 × 10 ⁻⁸
Readily evaporable water fraction	dimensionless	0.300000	0.300000

870 Effective shared crop parameters from the selected current cards. RUE uses intercepted PAR with PAR fraction
871 0.45. DVS-2 SLA equals its DVS-1 value; senescence is a thermal hazard with a separate daily stress loss. Wheat
872 BBCH 89 terminates remaining green leaf area; maize retains its terminal hazard at R6. Grain mass is fixed. Water

873 coefficients are estimated from Wuqiao 2016–2018; growth coefficients retain inherited multisite estimates. Root
874 depth is limited by profile depth. Dashes denote crop-specific processes. No maximum-LAI parameter is fitted.

875 **S3. Study-area and spatial aggregation checks**

876 The SRTM-derived terrain grid (NASA JPL, 2013) contains 720 eligible lowland components; the connected
877 component retained by the operational definition covers 404,658 km². Disconnected eligible terrain totals 6,465 km²
878 and contains approximately 7,068 ha of CPM-corrected rotation area. The main definition reaches the southern and
879 eastern bounding-box edges, so geographic sensitivity remains relevant to regional interpretation. Nine elevation–
880 slope alternatives combine elevations 100, 200 and 300 m with slopes 1°, 2° and 3°. Seven annual mapped-area
881 layers cover 2015–2021.

882 A common-extent comparison between ChinaCP plus CPM (Qiu et al., 2022) and the 2020 China Crop Dataset
883 layer (Fu et al., 2026; Fu et al., 2025) yields 9.694 and 7.841 million ha, respectively; CCD2020 is 19.11% lower on
884 that common support. This disagreement is map uncertainty rather than proof of either product’s accuracy. The
885 modeled 2020 area is held fixed to isolate climate and irrigation effects. Historical crop expansion, rotation
886 switching and double-counting of harvested crop area are not estimated.

887 Thirty-two actual-cell representatives use rotation-area weights and selection-period climate means. Eight
888 purposively selected non-representative cells test approximation under all five continuous irrigation fractions. Each
889 crop–fraction comparison retains 96 evaluation cell-years in 2014–2025. Absolute grain, ET and management-
890 response differences compare actual-cell with representative-cell model predictions. The sample measures
891 aggregation sensitivity rather than observed regional accuracy or a probabilistic error bound (Table S4; Figure S1).


892 **Table S4. Crop-model spatial aggregation errors in eight purposively selected simulation cells.**

Crop	Irrigation fraction	Variable	Unit	N	RMSE	Bias
Maize	0.000	Grain yield	t ha ⁻¹	96	1.17	0.18
Maize	0.000	ET	mm	96	35.21	6.41
Maize	0.250	Grain yield	t ha ⁻¹	96	0.72	0.13
Maize	0.250	ET	mm	96	27.39	5.17
Maize	0.500	Grain yield	t ha ⁻¹	96	0.50	0.08
Maize	0.500	ET	mm	96	22.19	3.82
Maize	0.750	Grain yield	t ha ⁻¹	96	0.39	0.08
Maize	0.750	ET	mm	96	17.85	4.38
Maize	1.000	Grain yield	t ha ⁻¹	96	0.35	0.09
Maize	1.000	ET	mm	96	14.97	5.69
Wheat	0.000	Grain yield	t ha ⁻¹	96	1.49	0.19
Wheat	0.000	ET	mm	96	51.66	-3.18
Wheat	0.250	Grain yield	t ha ⁻¹	96	1.39	0.07
Wheat	0.250	ET	mm	96	43.32	-4.26
Wheat	0.500	Grain yield	t ha ⁻¹	96	1.19	-7.41 × 10 ⁻³
Wheat	0.500	ET	mm	96	37.97	-4.56
Wheat	0.750	Grain yield	t ha ⁻¹	96	0.89	-0.01
Wheat	0.750	ET	mm	96	31.90	-2.05
Wheat	1.000	Grain yield	t ha ⁻¹	96	0.64	0.11
Wheat	1.000	ET	mm	96	24.70	3.21

893 Current model aggregation discrepancies against eight actual-cell simulations under all five irrigation fractions.

894 Grain is converted from kg ha⁻¹ to t ha⁻¹ in every displayed grain-valued column. Exactly zero irrigation

895 discrepancies are omitted from display; full inputs and metrics remain in the current-results product.



896 **Figure S1. Current crop-model spatial aggregation errors under all five irrigation fractions in eight purposively selected cells during**
 897 **2014–2025. Actual-cell and representative predictions share the current native model and crop parameters. These discrepancies**
 898 **assess spatial representation rather than field-observation accuracy.**

899 **S4. Annual field yields and irrigation-treatment contrasts**

900 Annual Wujiao grain targets comprise sixteen treatment-year means per crop in 2016–2019, digitized with standard
 901 errors from the primary Supplementary Figure S2. Grain retains its 13% moisture basis in comparisons. The twelve
 902 2016–2018 means per crop enter calibration after a single multiplication by 0.87 to dry matter. The four 2019
 903 means remain outside fitting and selection but had been inspected during earlier development. Four-year means
 904 summarize the same years and supply no additional independent targets.

905 Annual grain calibration RMSE is 1.249 t ha⁻¹ for wheat and 2.267 t ha⁻¹ for maize; retrospective-testing RMSE is
 906 3.161 and 1.700 t ha⁻¹ at 13% moisture. Within-year W1–W3 minus W0 grain-contrast RMSE is 0.519 and 1.669 t
 907 ha⁻¹ in calibration and 1.316 and 1.323 t ha⁻¹ in 2019 (Table S5; Figure S2). Maize treatment labels retain
 908 preceding wheat management because maize irrigation is identical within each year.

909 All printed seasonal ET targets remain eligible after matching the 2016 maize rainfall period. Maize-season rainfall
 910 of 499.1 mm excludes the 100.5 mm during wheat from the 599.6-mm rotation total. Reconstructed balance
 911 residuals for W0–W3 are +0.1, +0.1, -0.9 and +0.1 mm. Source ET and depletion values remain unchanged.
 912 Storage-depletion diagnostics share the source ET balance and its unconfirmed exact sampling dates; they are
 913 conditional accounting comparisons rather than independent soil-water validation.

914 Field simulations retain the recorded irrigation, sowing and harvest dates and archived daily weather. Wheat crop-
 915 start storage follows the pre-sowing simulation, whereas maize is initialized separately from observed treatment-
 916 specific total storage in 2 m; its distribution among layers is assumed. Field capacity, wilting point and bulk density


917 are transferred from the published Luancheng profile, with the bottom 10 cm extrapolated (Sun et al., 2006).
918 Conductivity follows the SoilGrids–Saxton–Rawls prior (ISRIC — World Soil Information, 2020; Saxton and
919 Rawls, 2006), with a particle-density prior of 2.75 g cm^{-3} . Reported wheat-straw mulch has no measured cover
920 fraction, so maize retains the exposed-soil evaporation boundary (Yang et al., 2024). The model represents runoff
921 and drainage, which the published ET balance assumes negligible.

922 Harvest aboveground biomass comprises grain and residues cut at ground level and oven-dried to constant mass,
923 requiring no moisture adjustment. Annual grain means and standard errors digitized from source Supplementary
924 Figure S2 are reconciled with its eight printed four-year means (Yang et al., 2024). Pixel tolerances are
925 approximately 0.043 t ha^{-1} for wheat and 0.056 t ha^{-1} for maize. Observed grain is multiplied once by 0.87 for dry-
926 matter fitting, and simulated dry grain is divided by 0.87 for comparison at 13% moisture. Annual and four-year
927 means overlap and provide no additional independent observations.

928 **Table S5. Wuqiao annual grain and within-year treatment contrasts by fitting and testing partition.**

Crop	Dataset	Comparison	N	RMSE (t/ha)	Bias (t/ha)	NSE	R ²
Maize	Calibration (2016–2018)	Annual grain yield	12	2.267	-1.738	-2.940	0.128
Maize	Calibration (2016–2018)	Within-year W1–W3 minus W0	9	1.669	0.296	-0.788	0.076
Maize	Retrospective evaluation (2019)	Annual grain yield	4	1.700	1.403	-3.641	0.972
Maize	Retrospective evaluation (2019)	Within-year W1–W3 minus W0	3	1.323	-0.836	-1.403	1.000
Wheat	Calibration (2016–2018)	Annual grain yield	12	1.249	-0.445	0.746	0.817
Wheat	Calibration (2016–2018)	Within-year W1–W3 minus W0	9	0.519	0.064	0.857	0.882
Wheat	Retrospective evaluation (2019)	Annual grain yield	4	3.161	3.108	-4.654	0.870
Wheat	Retrospective evaluation (2019)	Within-year W1–W3 minus W0	3	1.316	-1.312	-0.712	0.994

929 Grain is at 13% moisture. Calibration comprises 2016–2018 and retrospective testing comprises 2019. Source
930 means and standard errors are digitized from Supplementary Figure S2, with pixel tolerances near 0.043 t ha⁻¹ for
931 wheat and 0.056 for maize. Contrasts compare W1–W3 with same-year W0. Four-year means overlap these annual
932 targets; maize contrasts retain preceding wheat histories.



933 **Figure S2.** Annual Wuqiao grain at 13% moisture under the selected current parameters. Panels (a–d) show wheat and (e–h) maize;
 934 columns from left to right show 2016, 2017, 2018 and 2019. Black points and error bars are published means $\pm$ standard error of
 935 three replicates; blue squares show current predictions. The 2016–2018 treatment means contribute to water calibration after
 936 conversion to dry grain; 2019 is retrospective testing, excluded from fitting and selection. Maize irrigation is identical within each
 937 year and labels retain preceding wheat management. Source means and SE are digitized from Supplementary Figure S2 of the
 938 primary experiment.

939 **S5. Class exposure, continuous states and rainfall-only controls**

940 All nine common GRACE-available years occupy lower-storage classes; three other years use conventional
 941 fallback. The selected current quotas are identical across storage strata within each rainfall group. Matched rainfall-
 942 only and storage–rainfall histories have identical actions, crop states and outcomes throughout the modeled record.
 943 Available-rainfall controls apply reduced quotas during monitoring gaps and therefore create different carried water
 944 states before later common years. These history effects are separate from the value of the storage classification.
 945 Table S7 reports all-year and common-year retention and water use; Table S8 reports the matched and available-
 946 rainfall controls.

947 **Table S6. Class exposure and missing-storage coverage by period.**

Period	Class	Unit-years	Years	Area-years (ha yr)	Area-year fraction
evaluation	Lower storage / lower rainfall	174	9	55410278.787	0.468
evaluation	Lower storage / higher rainfall	114	9	33389136.355	0.282
evaluation	Missing storage	96	3	29599805.048	0.250
selection	Higher storage / lower rainfall	68	6	19466918.675	0.179
selection	Higher storage / higher rainfall	124	6	39732691.420	0.366
selection	Lower storage / lower rainfall	109	5	34747673.179	0.320
selection	Lower storage / higher rainfall	51	5	14585335.233	0.134

948 **Table S7. Continuous class-adaptive irrigation evaluation during 2014–2025.**

Strategy	Period	Years	Grain retained (%)	Irrigation cut (mm)	ET cut (mm)	Minimum grain (%)
Storage– rainfall 95%	Nine GRACE years	9	96.270	130.721	36.842	92.115
Storage– rainfall 95%	All twelve years	12	96.818	98.040	29.733	92.115
Storage– rainfall 98%	Nine GRACE years	9	99.626	35.721	6.171	98.579
Storage– rainfall 98%	All twelve years	12	99.622	26.790	5.208	98.579
Conventional	Nine GRACE years	9	100.000	0.000	0.000	100.000
Conventional	All twelve years	12	100.000	0.000	0.000	100.000

949 Current continuous histories. Grain retention is a ratio of period-mean regional dry-grain production to conventional
 950 irrigation. ET includes crops and fallows. All-year summaries include conventional fallback; common-year
 951 summaries use the nine GRACE-available years.

952 **Table S8. Continuous GRACE–rainfall and rainfall-only strategy comparison during 2014–2025.**

Strategy	Period	Grain retained (%)	Irrigation (mm)	ET (mm)
Storage–rainfall 95%	Nine GRACE years	96.27	249.279	848.527
Storage–rainfall 95%	All twelve years	96.82	281.960	857.698
Storage–rainfall 98%	Nine GRACE years	99.63	344.279	879.197
Storage–rainfall 98%	All twelve years	99.62	353.210	882.223
Rainfall 95%, available	Nine GRACE years	95.40	249.279	844.203
Rainfall 95%, available	All twelve years	95.04	248.565	842.945
Rainfall 95%, matched	Nine GRACE years	96.27	249.279	848.527
Rainfall 95%, matched	All twelve years	96.82	281.960	857.698
Rainfall 98%, available	Nine GRACE years	99.41	344.279	878.151
Rainfall 98%, available	All twelve years	99.36	343.565	879.669
Rainfall 98%, matched	Nine GRACE years	99.63	344.279	879.197
Rainfall 98%, matched	All twelve years	99.62	353.210	882.223


953 Matched controls share the complete GRACE-availability and fallback calendar. Available-rainfall controls act
 954 during missing-storage years and alter subsequent soil states. All values use current parameters; differences do not
 955 identify causal groundwater responses.



956 **Figure S3. Simulated 0–2 m soil-water storage in representative units (a) 0 and (b) 31. State carries continuously through wheat,**
 957 **maize, fallows and annual irrigation choices. Displayed histories compare conventional irrigation with selected current strategies;**
 958 **they are model states rather than observed groundwater levels.**

959 **S6. Harvest data quality and crop model recalibration**

960 Original Yucheng harvest sources contain identical six-replicate maize vectors in 2012 and 2017 under different
 961 date and cultivar labels. Both groups are retained as raw evidence and excluded from scored grain and harvest
 962 biomass. Eligibility depends on source authenticity rather than model residuals. Current plots preserve observed
 963 quadrat variation and the original whole site-year partitions (Figure S4).



964 **Figure S4. Original Yucheng harvest observations at YCAZH01ABC_01. Panels (a, b) show wheat aboveground biomass and grain;**
 965 **panels (c, d) show maize aboveground biomass and grain. Faint points are six quadrat observations; dark markers and error bars**
 966 **are means $\pm$ replicate standard error. Magenta diamonds identify the identical full maize source vectors in 2012 and 2017. Raw**
 967 **values remain displayed for source inspection; both groups are excluded from comparisons with quality-controlled observations.**
 968 **Error bars describe sampling variation, rather than model or policy uncertainty.**

969 **S7. Spatial decomposition under the common irrigation budget**

970 Frozen area shares weight complete current irrigation histories before spatial aggregation. Dry-grain differences in
 971 kg ha⁻¹ multiply represented hectares and divide by 1,000 for tonnes; water depths multiply hectares by 10 for cubic
 972 metres. Independent sums reproduce regional contrasts. Latitude bands describe the distribution of gains and losses,
 973 while the 3,641 mapped cells inherit 32 simulated representative responses (Table S9).

974 **Table S9. Spatial decomposition at a common 50% regional irrigation reduction during 2014–2025.**


Latitude band	Area share (%)	Targeted irrigation fraction	Grain change (t ha ⁻¹)	ET change (mm)	Grain contribution (Mt)
Southern (32–36°N)	60.733	0.305	-0.357	-25.599	-2.142
Central (36–38°N)	31.369	0.799	2.415	93.497	7.475
Northern (38–41°N)	7.897	0.813	2.420	88.358	1.886

975 Current targeted-minus-uniform means at the same 50% regional irrigation cut. Latitude bands are reporting
976 partitions. Source cells inherit representative responses and are not independent observations.


977 **S8. Seasonal crop trajectories and harvest outcomes**

978 Current seasonal curves use the selected frozen parameters and all sites with eligible observations for each variable.
979 Observation and prediction channels share windows, case identities and aggregation weights. Bins span 15 days for
980 wheat and 7 days for maize; site-year contributions and populations can change among bins. Standard errors
981 describe variation across site-year means and are omitted for single-year bins. Missing bins remain gaps. Unbinned
982 records determine performance scores (Figures S6–S11).


983 The original station calibration partitions retain earlier growth-fitting history; all station comparisons are
984 conditional on unconfirmed complete irrigation management. Station observations have no weight in the latest
985 water fit. Yucheng has no eligible in-season biomass evaluation observations, while its harvest-biomass endpoints
986 remain separately available. Figure S5 shows dry-grain and aboveground harvest means with quadrat SE. Six
987 samples within a crop-year describe sampling variation, and the duplicated 2012 and 2017 maize groups remain
988 excluded. Exact case availability, site-specific metrics, bin contributions and harvest source rows accompany the
989 figures.




990 **Figure S5. Conditional Yucheng harvest comparisons. Wheat occupies the left column and maize the right; the top row shows**
 991 **aboveground biomass and the bottom row grain. Black points show observed means $\pm$ standard errors among six quadrat samples**
 992 **per crop-year; blue points show current coupled-model predictions at the same sampling windows. Circles and squares distinguish**
 993 **the original calibration and retrospective-evaluation partitions. Grain and total aboveground biomass are dry matter. The duplicated**
 994 **maize 2012 and 2017 source vectors remain excluded. Within-year replicates describe spatial sampling variation and are not**
 995 **independent crop seasons. Management completeness remains unconfirmed and current water coefficients use Wuqiao fitting targets,**
 996 **while growth coefficients retain their earlier multisite fitting history.**




997 **Figure S6. Site-specific wheat LAI comparisons: (a) Fengqiu; (b) Luancheng; (c) Shangqiu; (d) Yucheng. Calibration and**
 998 **retrospective-evaluation observations and simulations share the same axes within each site. Error bars describe site-year**
 999 **errors; gaps are retained. The observations, weighting and conditional management interpretation follow Figure 6.**



1000 **Figure S7. Site-specific wheat aboveground biomass comparisons: (a) Fengqiu; (b) Luancheng; (c) Shangqiu; (d) Yucheng.**
 1001 **Calibration and retrospective-evaluation observations and simulations share the same axes within each site. Error bars describe site-**
 1002 **year standard errors; gaps are retained. The observations, weighting and conditional management interpretation follow Figure 6.**



1003 **Figure S8. Site-specific wheat ET comparisons: (a) Fengqiu; (b) Yucheng. Calibration and retrospective-evaluation observations and**
 1004 **simulations share the same axes within each site. Error bars describe site-year standard errors; gaps are retained. The observations,**
 1005 **weighting and conditional management interpretation follow Figure 6.**



1006 **Figure S9. Site-specific maize LAI comparisons: (a) Fengqiu; (b) Gucheng; (c) Luancheng; (d) Shangqiu; (e) Yucheng. Calibration**
 1007 **and retrospective-evaluation observations and simulations share the same axes within each site. Error bars describe site-year**
 1008 **standard errors; gaps are retained. The observations, weighting and conditional management interpretation follow Figure 6.**




Figure S10. Site-specific maize aboveground biomass comparisons: (a) Fengqiu; (b) Luancheng; (c) Shangqiu; (d) Yucheng. Calibration and retrospective-evaluation observations and simulations share the same axes within each site. Error bars describe site-year standard errors; gaps are retained. The observations, weighting and conditional management interpretation follow Figure 6.




Figure S11. Site-specific maize ET comparisons: (a) Fengqiu; (b) Yucheng. Calibration and retrospective-evaluation observations and simulations share the same axes within each site. Error bars describe site-year standard errors; gaps are retained. The observations, weighting and conditional management interpretation follow Figure 6.

References

- Yang, X., Wang, G., Chen, Y., Sui, P., Pacenka, S., Steenhuis, T.S., Siddique, K.H.M., 2022. Reduced groundwater use and increased grain production by optimized irrigation scheduling in winter wheat–summer maize double cropping system—A 16-year field study in North China Plain. *Field Crops Res.* 275, 108364.

1019 <https://doi.org/10.1016/j.fcr.2021.108364>

1020 Yang, L., Fang, X., Zhou, J., Zhao, J., Hou, X., Yang, Y., Zang, H., Zeng, Z., 2024. Optimal irrigation for wheat-
1021 maize rotation depending on precipitation in the North China Plain: Evidence from a four-year experiment.
1022 *Agric. Water Manag.* 294, 108726. <https://doi.org/10.1016/j.agwat.2024.108726>

1023 Zhang, X., Chen, S., Sun, H., Shao, L., Wang, Y., 2011. Changes in evapotranspiration over irrigated winter wheat
1024 and maize in North China Plain over three decades. *Agric. Water Manag.* 98, 1097–1104.
1025 <https://doi.org/10.1016/j.agwat.2011.02.003>

1026 Wu, D., Fang, S., Li, X., He, D., Zhu, Y., Yang, Z., Xu, J., Wu, Y., 2019. Spatial-temporal variation in irrigation
1027 water requirement for the winter wheat-summer maize rotation system since the 1980s on the North China Plain.
1028 *Agric. Water Manag.* 214, 78–86. <https://doi.org/10.1016/j.agwat.2019.01.004>

1029 Ren, P., Huang, F., Li, B., 2022. Spatiotemporal patterns of water consumption and irrigation requirements of
1030 wheat-maize in the Huang-Huai-Hai Plain, China and options of their reduction. *Agric. Water Manag.* 263,
1031 107468. <https://doi.org/10.1016/j.agwat.2022.107468>

1032 Xiao, L., Wang, G., Wang, E., Liu, S., Chang, J., Zhang, P., Zhou, H., Wei, Y., Zhang, H., Zhu, Y., Shi, Z., Luo, Z.,
1033 2024. Spatiotemporal co-optimization of agricultural management practices towards climate-smart crop
1034 production. *Nature Food* 5, 59–71. <https://doi.org/10.1038/s43016-023-00891-x>

1035 Jiang, Y., Xu, X., Huang, Q., Huo, Z., Huang, G., 2016. Optimizing regional irrigation water use by integrating a
1036 two-level optimization model and an agro-hydrological model. *Agricultural Water Management* 178, 76–88.
1037 <https://doi.org/10.1016/j.agwat.2016.08.035>

1038 Perry, C., Steduto, P., Allen, R.G., Burt, C.M., 2009. Increasing productivity in irrigated agriculture: Agronomic
1039 constraints and hydrological realities. *Agric. Water Manag.* 96, 1517–1524.
1040 <https://doi.org/10.1016/j.agwat.2009.05.005>

1041 Grafton, R.Q., Williams, J., Perry, C.J., Molle, F., Ringler, C., Steduto, P., Udall, B., Wheeler, S.A., Wang, Y.,
1042 Garrick, D., Allen, R.G., 2018. The paradox of irrigation efficiency. *Science* 361, 748–750.
1043 <https://doi.org/10.1126/science.aat9314>

1044 Chen, C., Wang, E., Yu, Q., 2010b. Modelling the effects of climate variability and water management on crop
1045 water productivity and water balance in the North China Plain. *Agric. Water Manag.* 97, 1175–1184.
1046 <https://doi.org/10.1016/j.agwat.2008.11.012>

1047 Fang, Q.X., Ma, L., Flerchinger, G.N., Qi, Z., Ahuja, L.R., Xing, H.T., Li, J., Yu, Q., 2014. Modeling
 1048 evapotranspiration and energy balance in a wheat–maize cropping system using the revised RZ-SHAW model.
 1049 *Agricultural and Forest Meteorology* 194, 218–229. <https://doi.org/10.1016/j.agrformet.2014.04.009>

1050 Cammarano, D., Rötter, R.P., Asseng, S., Ewert, F., Wallach, D., Martre, P., Hatfield, J.L., Jones, J.W., Rosenzweig,
 1051 C., Ruane, A.C., Boote, K.J., Thorburn, P.J., Kersebaum, K.C., Aggarwal, P.K., Angulo, C., Basso, B., Bertuzzi,
 1052 P., Biernath, C., Brisson, N., Challinor, A.J., Doltra, J., Gayler, S., Goldberg, R., Heng, L., Hooker, J., Hunt,
 1053 L.A., Ingwersen, J., Izaurralde, R.C., Müller, C., Kumar, S.N., Nendel, C., O’Leary, G.J., Olesen, J.E., Osborne,
 1054 T.M., Palosuo, T., Priesack, E., Ripoche, D., Semenov, M.A., Shcherbak, I., Steduto, P., Stöckle, C.O.,
 1055 Stratonovitch, P., Streck, T., Supit, I., Tao, F., Travasso, M., Waha, K., White, J.W., Wolf, J., 2016. Uncertainty
 1056 of wheat water use: Simulated patterns and sensitivity to temperature and CO₂. *Field Crops Research* 198, 80–
 1057 92. <https://doi.org/10.1016/j.fcr.2016.08.015>

1058 Save, H., Bettadpur, S., Tapley, B.D., 2016. High-resolution CSR GRACE RL05 mascons. *Journal of Geophysical*
 1059 *Research: Solid Earth* 121, 7547–7569. <https://doi.org/10.1002/2016jb013007>

1060 Scanlon, B.R., Zhang, Z., Save, H., Wiese, D.N., Landerer, F.W., Long, D., Longuevergne, L., Chen, J., 2016.
 1061 Global evaluation of new GRACE mascon products for hydrologic applications. *Water Resources Research* 52,
 1062 9412–9429. <https://doi.org/10.1002/2016wr019494>

1063 Long, D., Xu, Y., Cui, Y., Cui, Y., Butler, J.J., Dong, L., Wang, L., Liu, D., Wada, Y., Hu, L., Bai, G., Li, B., Wang,
 1064 S., Nong, X., Cai, Y., Cheng, C., Mu, Y., Qiao, Y., Wang, J., Wang, H., Scanlon, B.R., 2025. Unprecedented
 1065 large-scale aquifer recovery through human intervention. *Nat. Commun.* 16, 7296.
 1066 <https://doi.org/10.1038/s41467-025-62719-5>

1067 Xu, Y., Long, D., Jiang, W., Cui, Y., Chen, F., Wang, J., Wada, Y., 2026. Quantifying Human and Climatic
 1068 Contributions of Large-Scale Groundwater Recovery in the North China Plain. *Water Resour. Res.* 62,
 1069 e2025WR043052. <https://doi.org/10.1029/2025wr043052>

1070 Natural Earth, n.d. Natural Earth, 1:10m Admin 1 – States, Provinces (version 5.1.1).

1071 Fengqiu Station, 2020a. Crop phenology and growth dynamics at Fengqiu Station, Henan Province, 2004–2008.
 1072 <https://doi.org/10.12199/nesdc.ecodb.mon.2020.dp2011.fqa.002>

1073 Luancheng Station, 2020. Crop phenology and growth dynamics at Luancheng Station, Hebei Province, 2000–
 1074 2008. <https://doi.org/10.12199/nesdc.ecodb.mon.2020.dp2011.lca.002>

1075 Shangqiu Station, 2021. Crop phenology and growth dynamics at Shangqiu Station, Henan Province, 2000–2007.
 1076 <https://doi.org/10.12199/nesdc.ecodb.mon.2020.dp2011.sqa.003>

1077 Yucheng Station, 2021. Crop phenology and growth dynamics at Yucheng Station, Shandong Province, 1998–2006.
 1078 <https://doi.org/10.12199/nesdc.ecodb.mon.2020.dp2011.yca.003>

1079 Yucheng Station, 2025a. Biological observations at Yucheng Station, 2005–2022.
 1080 <https://doi.org/10.12199/nesdc.ecodb.2021YFF0703900.yca.2025.2>

1081 Geng, J., Zhou, G., Song, Y., Ren, S., Zhao, H., Zhou, H., Song, X., Tian, X., 2023. A dataset of effects of different
 1082 sowing dates on maize growth and development in the northern part of North China from 2018 to 2021.
 1083 <https://doi.org/10.57760/sciencedb.o00119.00017>

1084 Fengqiu Station, 2020b. Farmland soil-water physical properties at Fengqiu Station, Henan Province, 2005–2008.
 1085 <https://doi.org/10.12199/nesdc.ecodb.mon.2020.dp2011.fqa.006>

1086 Yucheng Station, 2025b. Water-environment observations at Yucheng Station, 2005–2022.
 1087 <https://doi.org/10.12199/nesdc.ecodb.2021YFF0703900.yca.2025.3>

1088 Copernicus Climate Change Service, 2020. Agrometeorological indicators from 1979 to present derived from
 1089 reanalysis. <https://doi.org/10.24381/CDS.6C68C9BB>

1090 ISRIC — World Soil Information, 2020. SoilGrids250m version 2.0.

1091 Saxton, K.E., Rawls, W.J., 2006. Soil Water Characteristic Estimates by Texture and Organic Matter for Hydrologic
 1092 Solutions. *Soil Science Society of America Journal* 70, 1569–1578. <https://doi.org/10.2136/sssaj2005.0117>

1093 Qiu, B., Hu, X., Chen, C., Tang, Z., Yang, P., Zhu, X., Yan, C., Jian, Z., 2022. Maps of cropping patterns in China
 1094 during 2015–2021. *Scientific Data*. <https://doi.org/10.1038/s41597-022-01589-8>

1095 NASA JPL, 2013. NASA Shuttle Radar Topography Mission Global 1 arc second.
 1096 <https://doi.org/10.5067/MEASURES/SRTM/SRTMGL1.003>

1097 Save, H., 2020. CSR GRACE and GRACE-FO RL06 Mascon Solutions v02. <https://doi.org/10.15781/cgq9-nh24>

1098 Long, D., Yang, W., Scanlon, B.R., Zhao, J., Liu, D., Burek, P., Pan, Y., You, L., Wada, Y., 2020. South-to-North
 1099 Water Diversion stabilizing Beijing’s groundwater levels. *Nat. Commun.* 11, 3665.
 1100 <https://doi.org/10.1038/s41467-020-17428-6>

1101 Zhao, G., Yao, L., Chen, B., submitted. OpenCropModel: A process-based framework for integrated nutrient, water
1102 and crop protection recommendations. Submitted to European Journal of Agronomy.

1103 Kiniry, J.R., Jones, C.A., O'toole, J.C., Blanchet, R., Cabelguenne, M., Spanel, D.A., 1989. Radiation-use efficiency
1104 in biomass accumulation prior to grain-filling for five grain-crop species. Field Crops Research.
1105 [https://doi.org/10.1016/0378-4290\(89\)90023-3](https://doi.org/10.1016/0378-4290(89)90023-3)

1106 de Wit; Berghuijs, n.d.. Python Crop Simulation Environment: WOFOST leaf dynamics implementation [software].
1107 Version Archived master-branch source snapshot.
1108 https://github.com/ajwdewit/pcse/blob/master/pcse/crop/leaf_dynamics.py (accessed 4 October 2026).

1109 Tan, M., Gou, F., Stomph, T.J., Wang, J., Yin, W., Zhang, L., Chai, Q., van der Werf, W., 2020. Dynamic process-
1110 based modelling of crop growth and competitive water extraction in relay strip intercropping: Model
1111 development and application to wheat-maize intercropping. Field Crops Research 246, 107613.
1112 <https://doi.org/10.1016/j.fcr.2019.107613>

1113 Allen, R.G., Pereira, L.S., Raes, D., Smith, M., 1998. Crop evapotranspiration: Guidelines for computing crop
1114 water requirements, FAO Irrigation and Drainage Paper. Food and Agriculture Organization of the United
1115 Nations, Rome.

1116 van Genuchten, M.T., 1980. A Closed-form Equation for Predicting the Hydraulic Conductivity of Unsaturated
1117 Soils. Soil Science Society of America Journal 44, 892–898.
1118 <https://doi.org/10.2136/sssaj1980.03615995004400050002x>

1119 Mualem, Y., 1976. A new model for predicting the hydraulic conductivity of unsaturated porous media. Water
1120 Resources Research 12, 513–522. <https://doi.org/10.1029/wr012i003p00513>

1121 Nash, J.E., Sutcliffe, J.V., 1970. River flow forecasting through conceptual models part I — A discussion of
1122 principles. Journal of Hydrology 10, 282–290. [https://doi.org/10.1016/0022-1694\(70\)90255-6](https://doi.org/10.1016/0022-1694(70)90255-6)

1123 Wu, L., Min, L., Liu, M., Zhang, Y., Pei, H., Li, H., Zhang, G., Wang, S., Shen, Y., 2023. Monitoring of Thick
1124 Vadose Zone Water Dynamics Under Irrigation Using a 48 m Deep Caisson at the Luancheng Critical Zone
1125 Observatory. Water Resour. Res. 59, e2022WR032965. <https://doi.org/10.1029/2022wr032965>

1126 Ines, A.V.M., Das, N.N., Hansen, J.W., Njoku, E.G., 2013. Assimilation of remotely sensed soil moisture and
1127 vegetation with a crop simulation model for maize yield prediction. Remote Sens. Environ. 138, 149–164.
1128 <https://doi.org/10.1016/j.rse.2013.07.018>

1129 Roy, A., Narvekar, P., Murtugudde, R., Shinde, V., Ghosh, S., 2021. Short and Medium Range Irrigation Scheduling
 1130 Using Stochastic Simulation-Optimization Framework With Farm-Scale Ecohydrological Model and Weather
 1131 Forecasts. *Water Resources Research* 57, e2020WR029004. <https://doi.org/10.1029/2020wr029004>
 1132 Chen, C., Wang, E., Yu, Q., 2010a. Modeling Wheat and Maize Productivity as Affected by Climate Variation and
 1133 Irrigation Supply in North China Plain. *Agronomy Journal* 102, 1037–1049.
 1134 <https://doi.org/10.2134/agronj2009.0505>
 1135 Yu, Q., Saseendran, S.A., Ma, L., Flerchinger, G.N., Green, T.R., Ahuja, L.R., 2006. Modeling a wheat–maize
 1136 double cropping system in China using two plant growth modules in RZWQM. *Agricultural Systems* 89, 457–
 1137 477. <https://doi.org/10.1016/j.agsy.2005.10.009>
 1138 Wang, M., Shi, W., 2024. The annual dynamic dataset of high-resolution crop water use in China from 1991 to
 1139 2019. *Scientific Data* 11, 1373. <https://doi.org/10.1038/s41597-024-04185-0>
 1140 Meier, U. (Ed.), 2018. Growth stages of mono- and dicotyledonous plants: BBCH Monograph. Julius Kühn-Institut,
 1141 Quedlinburg. <https://doi.org/10.5073/20180906-074619>
 1142 de Wit, A., 2020. Parameter sets for the WOFOST 7.2/7.3 cropping system model for wheat.
 1143 de Wit, A., 2022. Parameter sets for the WOFOST 7.2/7.3 model for simulating maize.
 1144 APSIM Initiative, 2026. APSIM Next Generation (ApsimX): wheat and maize model definitions [software].
 1145 Version 074d15f.
 1146 [https://github.com/APSIMInitiative/ApsimX/tree/074d15f51889a386d3667acfed184d03e4e8f65f/Models/Resou](https://github.com/APSIMInitiative/ApsimX/tree/074d15f51889a386d3667acfed184d03e4e8f65f/Models/Resources)
 1147 [rces](https://github.com/APSIMInitiative/ApsimX/tree/074d15f51889a386d3667acfed184d03e4e8f65f/Models/Resources) (accessed 4 October 2026).
 1148 Fu, Y., Dong, J., Shen, R., Peng, Q., Li, X., Zheng, Y., Pan, B., Dong, J., Qiu, B., Huang, J., Yuan, W., 2025. China
 1149 Crop Dataset: A long-term and high-resolution crop distribution dataset in China.
 1150 <https://doi.org/10.57760/sciencedb.32361>
 1151 Fu, Y., Dong, J., Shen, R., Peng, Q., Li, X., Zheng, Y., Pan, B., Dong, J., Qiu, B., Huang, J., Huang, X., Yuan, W.,
 1152 2026. A 30-m annual distribution dataset of major crops in China from 2001-2024. *Scientific Data* 13, 1074.
 1153 <https://doi.org/10.1038/s41597-026-07370-5>
 1154 Sun, H.-Y., Liu, C.-M., Zhang, X.-Y., Shen, Y.-J., Zhang, Y.-Q., 2006. Effects of irrigation on water balance, yield
 1155 and WUE of winter wheat in the North China Plain. *Agric. Water Manag.* 85, 211–218.
 1156 <https://doi.org/10.1016/j.agwat.2006.04.008>