

Math 220 A - Lecture 13

November 9, 2020

Logistics

Wed, Nov 11 - holiday - no lecture

Office Hour - Wed 4-5 PM (discuss homework, midterm)

Questions about the midterm?

Last time - Laurent expansion

Theorem Let $f: \Delta(a; r, R) \rightarrow \mathbb{C}$ holomorphic. Then

$$f(z) = \sum_{k=-\infty}^{\infty} a_k (z - a)^k \quad \text{can be represented as}$$

Laurent series, converging absolutely & uniformly on compact subsets of

$\Delta(a; r, R)$.

Today - classification of singularities Conway v. 1.

- characterization of singularities.

Types of singularities

$f: \Delta^*(a, R) \rightarrow \mathbb{C}$, holomorphic.

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z-a)^n \quad \text{Laurent series.}$$

Terminology [I] the coefficient of $(z-a)^{-1}$:

$$a_{-1} = \operatorname{Res}_{z=a} f = \text{residue}$$

$$[II] \quad \sum_{n=-\infty}^{-1} a_n (z-a)^n = \text{principal part.}$$

Three cases

$$[A] \quad a_k = 0 \quad \forall k < 0 \Leftrightarrow \text{Taylor expansion}$$

$\Leftrightarrow f$ extends holomorphically across a

Removable singularity

$$[B] \quad a_k = 0 \quad \forall k < -N, \quad a_{-N} \neq 0$$

Pole of order N .

$$[C] \quad a_k \neq 0, \quad k < 0 \text{ happens infinitely often}$$

Essential singularity

Case A a removable singularity

Theorem A TFAE $f: \Delta^*(a, R) \rightarrow \mathbb{C}$ holomorphic

i f extends holomorphically across a

ii f extends continuously across a

iii f bounded near a

iv $\lim_{z \rightarrow a} f(z) \cdot (z-a) = 0.$

Proof i \Rightarrow ii \Rightarrow iii \Rightarrow iv is obvious

iv \Rightarrow i. WLOG $a=0$, else work with $f(z+a)$.

We show $a_k = 0 \ \forall k < 0$. Fix $\varepsilon > 0$. Since

$$\lim_{z \rightarrow 0} z f(z) = 0 \Rightarrow |f(z)| < \frac{\varepsilon}{|z|} \text{ if } |z| < \delta.$$

We have for $0 < r < \delta < R$:

$$\begin{aligned} |a_k| &= \left| \frac{1}{2\pi i} \int_{|z|=r} \frac{f(z)}{z^{k+1}} dz \right| \leq \frac{1}{2\pi} \cdot \frac{\varepsilon}{r} \cdot \frac{1}{r^{k+1}} \cdot 2\pi r \\ &= \frac{\varepsilon}{r^{k+1}}. \end{aligned}$$

If $k = -1$: $|a_{-1}| < \varepsilon \quad \forall \varepsilon > 0 \Rightarrow a_{-1} = 0.$

If $k < -1$, take $\varepsilon = 1$, $|a_k| < \frac{1}{r^{k+1}}$. Make $r \rightarrow 0$

to obtain $a_k = 0$. since $k < -1$.

Example $f: U \rightarrow \mathbb{C}$ holomorphic

$$g(z) = \begin{cases} \frac{f(z) - f(a)}{z - a}, & z \neq a \\ f'(a), & z = a \end{cases} \quad \text{is holomorphic}$$

Indeed $\frac{f(z) - f(a)}{z - a}$ has a **removable** singularity at a

by item iv. & g is the continuous / holomorphic extension across a .

Case B a pole of order N .

$$f(z) = (z-a)^{-N} g(z) \quad \leftarrow \text{holomorphic}$$

$$g(z) = \sum_{k=0}^{\infty} a_{k-N} (z-a)^k, \quad g(a) = a_{-N} \neq 0.$$

Zeros versus Poles

$$\frac{1}{f(z)} = (z-a)^N \cdot \frac{1}{g(z)}, \quad \frac{1}{g} \text{ holomorphic near } a$$

f pole of order N at $a \iff \frac{1}{f}$ zero of order N at a

Lemma 8 $f: \Delta^*(a, R) \rightarrow \mathbb{C}$ holomorphic TFAE

(i) a is a pole

(ii) $\lim_{z \rightarrow a} f(z) = \infty$.

Proof (i) \Rightarrow (ii) Write $f(z) = (z-a)^{-N} g(z)$.

Since $g(a) \neq 0 \Rightarrow |g(z)| \geq M > 0$ in $|z-a| < \delta$.

$$\Rightarrow |f(z)| = \frac{|g(z)|}{|z-a|^N} \geq \frac{M}{|z-a|^N}. \text{ Make } z \rightarrow a \text{ to}$$

conclude $\lim_{z \rightarrow a} f(z) = \infty$.

$$\boxed{ii} \Rightarrow \boxed{i} \text{ Note } \lim_{z \rightarrow a} f(z) = \infty \Rightarrow \lim_{z \rightarrow a} \frac{1}{f(z)} = 0$$

$\Rightarrow \frac{1}{f}$ bounded near $a \Rightarrow \frac{1}{f}$ can be extended across a holomorphically. Note the extension vanishes at a , say of order $N \Rightarrow f$ has a pole at a of order N .

Definition $S \subseteq U$ **discrete**. A function f holomorphic in $U \setminus S$, with at worst poles at S is called **meromorphic**.

Example

$$\boxed{i} \quad f(z) = \frac{P(z)}{Q(z)}, \quad u = \mathbb{C} \text{ meromorphic.}$$

polynomials ↙

$$\boxed{ii} \quad f(z) = \frac{1}{\sin \frac{1}{z}}, \quad u = \mathbb{C}^\times$$

check $z = \frac{1}{n\pi}, n \in \mathbb{Z}^\times$ are poles. These do not

accumulate in $U = \mathbb{C} \setminus \{0\}$. Thus f meromorphic in
 $U = \mathbb{C}^*$.

Case C $f: \Delta^*(a, R) \rightarrow \mathbb{C}$ holomorphic

a essential singularity e.g. $f(z) = e^{\frac{1}{z}}$.

Example $f(z) = e^{\frac{1}{z}} = \sum_{k=0}^{\infty} \frac{1}{k!} \cdot \frac{1}{z^k} \Rightarrow$

$\Rightarrow a=0$ is essential singularity.


Remark f cannot be bounded or go to ∞ .

(see cases A & B)

Question How does f behave near a ?

Theorem (Big Picard Theorem) - 220 C.

$$\forall \Delta^*(a, \varepsilon) \subseteq \Delta^*(a, R), \quad f(\Delta^*(a, \varepsilon)) = \mathbb{C} \text{ or } \mathbb{C} \setminus \{\text{point}\}.$$



Example $f(z) = e^{\frac{1}{z}}$, $a = 0$.

Claim $f(\Delta^*(0, \varepsilon)) = \mathbb{C} \setminus \{0\}$, $\forall \varepsilon > 0$

Proof $y \neq 0$: $y = e^{\frac{1}{z}}$, $z \in \Delta^*(0, \varepsilon)$.

$$\Leftrightarrow \frac{1}{z} = \log y + 2n\pi i \text{ for any choice of } \log$$

$$\Leftrightarrow z = \frac{1}{\log y + 2n\pi i} \in \Delta^*(0, \varepsilon) \text{ if } n \gg 0.$$

Theorem C (Casorati - Weierstraß) $f: \Delta^*(a, R) \rightarrow \mathbb{C}$

(i) f has essential singularity at a

(ii) $\forall \Delta^*(a, \varepsilon) \subseteq \Delta^*(a, R)$, $f(\Delta^*(a, \varepsilon))$ is

dense in \mathbb{C} .

Proof $\boxed{11} \Rightarrow \boxed{12}$ Assume for some $\varepsilon > 0$, the set

$f(\Delta^*(a, \varepsilon))$ is not dense in \mathbb{C} . Then $\exists \Delta(\lambda, R)$

$$(*). \quad f(\Delta^*(a, \varepsilon)) \cap \Delta(\lambda, R) = \emptyset.$$

Define $g = \frac{1}{f - \lambda}$ in $\Delta^*(a, \varepsilon)$. By (*) we know

$$|f - \lambda| \geq R \text{ in } \Delta^*(a, \varepsilon) \Rightarrow |g| \leq \frac{1}{R} \text{ in } \Delta^*(a, \varepsilon)$$

Thm A

$\Rightarrow a$ is removable singularity for g . But

$$f = \lambda + \frac{1}{g}. \quad (+)$$

If a is not a zero for $g \Rightarrow \frac{1}{g}$ holomorphic \Rightarrow (+)

f extends holomorphically across $a \Rightarrow$ removable singularity.

If a is a zero for $g \Rightarrow \frac{1}{g}$ has a pole at $a \Rightarrow$ (+)
 $\Rightarrow f$ has pole at a .

Both cases are impossible.

16 \Rightarrow 6 Assume a *removable* singularity \Rightarrow

Thm A

$\Rightarrow f$ bounded near $a \Rightarrow \exists M > 0, \varepsilon > 0$ with

$$|f(z)| < M \text{ in } \Delta^*(a, \varepsilon)$$

$\Rightarrow f(\Delta^*(a, \varepsilon))$ cannot be dense.

Lemma B

Assume a *pole* $\Rightarrow \lim_{z \rightarrow a} f(z) = \infty \Rightarrow$

$\Rightarrow \exists \varepsilon > 0, |f(z)| \geq 1$ in $\Delta^*(a, \varepsilon) \Rightarrow$

$\Rightarrow f(\Delta^*(a, \varepsilon))$ cannot be dense.

Thus a is *essential* singularity.



Felice Casorati
1835 - 1890



Karl Weierstraß
1815 - 1897



Émile Picard
1856 - 1941