

Pieri Rules for Schur functions in superspace

Miles Eli Jones
joint work with Luc Lapointe

Universidad de Talca, Chile

July 9, 2015



Miles Eli Jones joint work with Luc Lapointe

Pieri Rules for Schur functions in superspace

Overview

Symmetric function theory

Overview

Symmetric function theory

Symmetric function theory IN SUPER SPACE!!!!




Symmetric Function Theory

$$\mathbb{K}[z_1, \dots, z_N]^{S_N}$$

1. partitions

$(5, 4, 2, 2, 1)$



2. simple bases $(m_\lambda, p_\lambda, e_\lambda, h_\lambda, s_\lambda, \dots)$

3. other bases: Macdonald polynomials, Jack polynomials



Miles Eli Jones joint work with Luc Lapointe


Pieri Rules for Schur functions in superspace

Symmetric Function Theory

$$\mathbb{K}[z_1, \dots, z_N]^{S_N}$$

1. partitions

$(5, 4, 2, 2, 1)$



2. simple bases $(m_\lambda, p_\lambda, e_\lambda, h_\lambda, s_\lambda, \dots)$

3. other bases: Macdonald polynomials, Jack polynomials



Miles Eli Jones joint work with Luc Lapointe


Pieri Rules for Schur functions in superspace

Symmetric Function Theory

$$\mathbb{K}[z_1, \dots, z_N]^{S_N}$$

1. partitions

$(5, 4, 2, 2, 1)$



2. simple bases $(m_\lambda, p_\lambda, e_\lambda, h_\lambda, s_\lambda, \dots)$

3. other bases: Macdonald polynomials, Jack polynomials




Symmetric Function Theory

$$\mathbb{K}[z_1, \dots, z_N]^{S_N}$$

1. partitions

$(5, 4, 2, 2, 1)$



2. simple bases $(m_\lambda, p_\lambda, e_\lambda, h_\lambda, s_\lambda, \dots)$

3. other bases: Macdonald polynomials, Jack polynomials



Symmetric function theory

$$M_{\lambda}^{(q,t)}$$

Symmetric function theory

$$\begin{array}{c} M_{\lambda}^{(q,t)} \\ \downarrow \begin{array}{l} q=t^{\alpha} \\ t \rightarrow 1 \end{array} \end{array}$$

Symmetric function theory

$$\begin{array}{c} M_{\lambda}^{(q,t)} \\ \downarrow \begin{array}{l} q=t^{\alpha} \\ t \rightarrow 1 \end{array} \\ J_{\lambda}^{(\alpha)} \end{array}$$

Symmetric function theory

$$\begin{array}{c} M_{\lambda}^{(q,t)} \\ \downarrow \begin{array}{l} q=t^{\alpha} \\ t \rightarrow 1 \end{array} \\ J_{\lambda}^{(\alpha)} \\ \downarrow \alpha \rightarrow 1 \end{array}$$

Symmetric function theory

$$\begin{array}{c} M_{\lambda}^{(q,t)} \\ \downarrow \begin{array}{l} q=t^{\alpha} \\ t \rightarrow 1 \end{array} \\ J_{\lambda}^{(\alpha)} \\ \downarrow \alpha \rightarrow 1 \\ s_{\lambda} \end{array}$$

Symmetric function theory

$$M_{\lambda}^{(q,t)}$$

$$\begin{array}{c} \downarrow \\ q=t^{\alpha} \\ t \rightarrow 1 \end{array}$$

$$J_{\lambda}^{(\alpha)}$$

$$\begin{array}{c} \downarrow \\ \alpha \rightarrow 1 \end{array}$$

$$s_{\lambda}$$

COMBINATORICS

PHYSICS

combinatorics

Symmetric function theory

Common characterization:

- ▶ $M_{\lambda}^{(q,t)} = m_{\lambda} + \text{smaller terms}$ (triangularity)
- ▶ $\langle M_{\lambda}^{(q,t)}, M_{\mu}^{(q,t)} \rangle = 0$ if $\lambda \neq \mu$ (orthogonality)

Scalar product:

$$\langle p_{\lambda}, p_{\mu} \rangle = \delta_{\lambda\mu} z_{\lambda} \prod_i \frac{1 - q^{\lambda_i}}{1 - t^{\lambda_i}}$$

Symmetric function theory

Common characterization:

- ▶ $M_{\lambda}^{(q,t)} = m_{\lambda} + \text{smaller terms}$ (triangularity)
- ▶ $\langle M_{\lambda}^{(q,t)}, M_{\mu}^{(q,t)} \rangle = 0$ if $\lambda \neq \mu$ (orthogonality)

Scalar product:

$$\langle p_{\lambda}, p_{\mu} \rangle = \delta_{\lambda\mu} z_{\lambda} \prod_i \frac{1 - q^{\lambda_i}}{1 - t^{\lambda_i}}$$

Macdonald positivity

$$\tilde{M}_{\lambda}^{(q,t)} = \sum_{\mu} K_{\mu\lambda}(q, t) s_{\mu} \quad \text{with} \quad K_{\mu\lambda}(q, t) \in \mathbb{N}[q, t]$$

$K_{\mu\lambda}(1, 1)$ = number of standard tableaux of shape μ

Macdonald positivity

$$\tilde{M}_{\lambda}^{(q,t)} = \sum_{\mu} K_{\mu\lambda}(q,t) s_{\mu} \quad \text{with} \quad K_{\mu\lambda}(q,t) \in \mathbb{N}[q,t]$$

$K_{\mu\lambda}(1,1)$ = number of standard tableaux of shape μ

Macdonald positivity

$$\tilde{M}^{(q,t)}_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}} = t s_{\begin{smallmatrix} \square & \square & \square \end{smallmatrix}} + (1 + qt) s_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}} + q s_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}}$$

1 2 3

1 2
3

1 3
2

1
2
3

Macdonald positivity

$$\tilde{M}^{(q,t)}_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \\ \hline \end{array}} = t s_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}} + (1 + qt) s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \\ \hline \end{array}} + q s_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}}$$

1 2 3

1 2
3

1 3
2

1
2
3

Supersymmetry

2 types of particles in nature

bosons (integer spin: $0, 1, 2, \dots$)

fermions (half integer spin: $1/2, 3/2, \dots$)

$$\Psi \longrightarrow \Psi$$

exchange of two **bosons**

$$\Psi \longrightarrow -\Psi$$

exchange of two **fermions**
(*Pauli's exclusion principle*)

Supersymmetry

2 types of particles in nature

bosons (integer spin: $0, 1, 2, \dots$)

fermions (half integer spin: $1/2, 3/2, \dots$)

$$\Psi \longrightarrow \Psi$$

exchange of two **bosons**

$$\Psi \longrightarrow -\Psi$$

exchange of two **fermions**
(*Pauli's exclusion principle*)

Supersymmetry

2 types of particles in nature

bosons (integer spin: $0, 1, 2, \dots$)

fermions (half integer spin: $1/2, 3/2, \dots$)

$$\Psi \longrightarrow \Psi$$

exchange of two **bosons**

$$\Psi \longrightarrow -\Psi$$

exchange of two **fermions**
(*Pauli's exclusion principle*)

Supersymmetry

2 types of particles in nature

bosons (integer spin: $0, 1, 2, \dots$)

fermions (half integer spin: $1/2, 3/2, \dots$)

$$\Psi \longrightarrow \Psi$$

exchange of two **bosons**

$$\Psi \longrightarrow -\Psi$$

exchange of two **fermions**
(*Pauli's exclusion principle*)

A symmetric function theory in superspace

$$\mathbb{K}[z_1, \dots, z_N, \theta_1, \dots, \theta_N]^{S_N} \quad \text{with} \quad (\theta_i \theta_j = -\theta_j \theta_i \quad \text{and} \quad \theta_i^2 = 0)$$

$$N = 2: \quad (z_1 - z_2) \theta_1 \theta_2$$

Power sums: $p_1, p_2, \dots, \tilde{p}_0, \tilde{p}_1, \dots$

$$p_r = z_1^r + z_2^r + \dots \quad \text{and} \quad \tilde{p}_k = \theta_1 z_1^k + \theta_2 z_2^k + \dots$$

$$\tilde{p}_i \tilde{p}_j = -\tilde{p}_j \tilde{p}_i \quad \text{and} \quad \tilde{p}_i^2 = 0$$

A symmetric function theory in superspace

$$\mathbb{K}[z_1, \dots, z_N, \theta_1, \dots, \theta_N]^{S_N} \quad \text{with} \quad (\theta_i \theta_j = -\theta_j \theta_i \quad \text{and} \quad \theta_i^2 = 0)$$

$$\underline{N = 2}: \quad (z_1 - z_2) \theta_1 \theta_2$$

Power sums: $p_1, p_2, \dots, \tilde{p}_0, \tilde{p}_1, \dots$

$$p_r = z_1^r + z_2^r + \dots \quad \text{and} \quad \tilde{p}_k = \theta_1 z_1^k + \theta_2 z_2^k + \dots$$

$$\tilde{p}_i \tilde{p}_j = -\tilde{p}_j \tilde{p}_i \quad \text{and} \quad \tilde{p}_i^2 = 0$$

A symmetric function theory in superspace

$$\mathbb{K}[z_1, \dots, z_N, \theta_1, \dots, \theta_N]^{S_N} \quad \text{with} \quad (\theta_i \theta_j = -\theta_j \theta_i \quad \text{and} \quad \theta_i^2 = 0)$$

$$\underline{N = 2}: \quad (z_1 - z_2) \theta_1 \theta_2$$

Power sums: $p_1, p_2, \dots, \tilde{p}_0, \tilde{p}_1, \dots$

$$p_r = z_1^r + z_2^r + \dots \quad \text{and} \quad \tilde{p}_k = \theta_1 z_1^k + \theta_2 z_2^k + \dots$$

$$\tilde{p}_i \tilde{p}_j = -\tilde{p}_j \tilde{p}_i \quad \text{and} \quad \tilde{p}_i^2 = 0$$

A symmetric function theory in superspace

$$\mathbb{K}[z_1, \dots, z_N, \theta_1, \dots, \theta_N]^{S_N} \quad \text{with} \quad (\theta_i \theta_j = -\theta_j \theta_i \quad \text{and} \quad \theta_i^2 = 0)$$

$$\underline{N = 2}: \quad (z_1 - z_2) \theta_1 \theta_2$$

Power sums: $p_1, p_2, \dots, \tilde{p}_0, \tilde{p}_1, \dots$


$$p_r = z_1^r + z_2^r + \dots \quad \text{and} \quad \tilde{p}_k = \theta_1 z_1^k + \theta_2 z_2^k + \dots$$

$$\tilde{p}_i \tilde{p}_j = -\tilde{p}_j \tilde{p}_i \quad \text{and} \quad \tilde{p}_i^2 = 0$$

Superpartitions

$$\Lambda = (\Lambda^a; \Lambda^s) \quad \begin{cases} \Lambda^s \text{ is usual partition} \\ \Lambda^a \text{ has no repeated parts} \end{cases}$$

$$(4, 2, 0; 3, 2, 1, 1) \longleftrightarrow (4, 3, 2, 2, 1, 1, 0) \longleftrightarrow$$




simple bases $(m_\Lambda, p_\Lambda, e_\Lambda, h_\Lambda, s_\Lambda, \dots)$

other bases: Macdonald polynomials and Jack polynomials in superspace.

Superpartitions

$$\Lambda = (\Lambda^a; \Lambda^s) \quad \begin{cases} \Lambda^s \text{ is usual partition} \\ \Lambda^a \text{ has no repeated parts} \end{cases}$$

$$(4, 2, 0; 3, 2, 1, 1) \longleftrightarrow (4, 3, 2, 2, 1, 1, 0) \longleftrightarrow$$



simple bases $(m_\Lambda, p_\Lambda, e_\Lambda, h_\Lambda, s_\Lambda, \dots)$

other bases: Macdonald polynomials and Jack polynomials in superspace.

Superpartitions

$$\Lambda = (\Lambda^a; \Lambda^s) \quad \begin{cases} \Lambda^s \text{ is usual partition} \\ \Lambda^a \text{ has no repeated parts} \end{cases}$$

$$(4, 2, 0; 3, 2, 1, 1) \longleftrightarrow (4, 3, 2, 2, 1, 1, 0) \longleftrightarrow$$



simple bases $(m_\Lambda, p_\Lambda, e_\Lambda, h_\Lambda, s_\Lambda, \dots)$

other bases: Macdonald polynomials and Jack polynomials in superspace.

Symmetric function theory in superspace

Common characterization:

- ▶ $M_{\Lambda}^{(q,t)} = m_{\Lambda} + \text{smaller terms}$ (triangularity)
- ▶ $\langle M_{\Lambda}^{(q,t)}, M_{\Omega}^{(q,t)} \rangle = 0$ if $\Lambda \neq \Omega$ (orthogonality)

Scalar product:

$$\langle p_{\Lambda}, p_{\Omega} \rangle_{qt} = \delta_{\Lambda\Omega} q^{|\Lambda^a|} z_{\Lambda^s} \prod_i \frac{1 - q^{\Lambda_i^s}}{1 - t^{\Lambda_i^s}}$$

$$\Lambda = (\Lambda^a; \Lambda^s) \quad \text{and} \quad p_{\Lambda} = \tilde{p}_{\Lambda^a} p_{\Lambda^s}$$

Symmetric function theory in superspace

Common characterization:


- ▶ $M_{\Lambda}^{(q,t)} = m_{\Lambda} + \text{smaller terms}$ (triangularity)
- ▶ $\langle M_{\Lambda}^{(q,t)}, M_{\Omega}^{(q,t)} \rangle = 0$ if $\Lambda \neq \Omega$ (orthogonality)

Scalar product:

$$\langle p_{\Lambda}, p_{\Omega} \rangle_{qt} = \delta_{\Lambda\Omega} q^{|\Lambda^a|} z_{\Lambda^s} \prod_i \frac{1 - q^{\Lambda_i^s}}{1 - t^{\Lambda_i^s}}$$

$$\Lambda = (\Lambda^a; \Lambda^s) \quad \text{and} \quad p_{\Lambda} = \tilde{p}_{\Lambda^a} p_{\Lambda^s}$$

Symmetric function theory in superspace

$$M_{\Lambda}^{(q,t)}$$



Symmetric function theory in superspace

$$\begin{array}{c} M_{\Lambda}^{(q,t)} \\ \downarrow \begin{array}{l} q=t^{\alpha} \\ t \rightarrow 1 \end{array} \\ J_{\Lambda}^{(\alpha)} \end{array}$$


Symmetric function theory in superspace

$$\begin{array}{c} M_{\Lambda}^{(q,t)} \\ \downarrow \begin{array}{l} q=t^{\alpha} \\ t \rightarrow 1 \end{array} \\ J_{\Lambda}^{(\alpha)} \\ \downarrow \alpha \rightarrow 1 \\ s_{\Lambda}^1 \end{array}$$


Symmetric function theory in **superspace**



Symmetric function theory in superspace



Symmetric function theory in **superspace**



$$\langle p_\Lambda, p_\Omega \rangle_{qt} = \delta_{\Lambda\Omega} q^{|\Lambda^a|} z_{\Lambda^s}(q, t)$$

Symmetric function theory in superspace

$$\begin{array}{c} M_{\Lambda}^{(q,t)} \\ \vdots \\ q=t=0 \\ \vdots \\ s_{\Lambda}^0 \end{array}$$

Macdonald positivity conjecture in superspace!!

$$\tilde{M}_{\Lambda}^{(q,t)} = \sum_{\Omega} K_{\Omega\Lambda}(q, t) s_{\Omega}^0 \quad \text{with} \quad K_{\Omega\Lambda}(q, t) \in \mathbb{N}[q, t]???$$

Macdonald positivity conjecture in superspace

$$\tilde{M}_{\begin{smallmatrix} \square \\ \square \end{smallmatrix} \bullet}^{(q,t)} = t s_{\begin{smallmatrix} \square & \square \end{smallmatrix} \bullet}^0 + s_{\begin{smallmatrix} \square \\ \square \end{smallmatrix} \bullet}^0 + qt s_{\begin{smallmatrix} \square & \square \\ \bullet \end{smallmatrix}}^0 + q s_{\begin{smallmatrix} \square \\ \square \\ \bullet \end{smallmatrix}}^0$$

$$\tilde{M}_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}}^{(q,t)} = t s_{\begin{smallmatrix} \square & \square & \square \end{smallmatrix}} + (1 + qt) s_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}} + q s_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}}$$

Refinement of the original problem!!

Macdonald positivity conjecture in superspace

$$\tilde{M}_{\begin{smallmatrix} \square \\ \square \end{smallmatrix} \bullet}^{(q,t)} = t s_{\begin{smallmatrix} \square & \square \end{smallmatrix} \bullet}^0 + s_{\begin{smallmatrix} \square \\ \square \end{smallmatrix} \bullet}^0 + qt s_{\begin{smallmatrix} \square & \square \\ \bullet \end{smallmatrix}}^0 + q s_{\begin{smallmatrix} \square \\ \square \\ \bullet \end{smallmatrix}}^0$$

$$\tilde{M}_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}}^{(q,t)} = t s_{\begin{smallmatrix} \square & \square & \square \end{smallmatrix}} + (1 + qt) s_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}} + q s_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}}$$

Refinement of the original problem!!

Macdonald positivity conjecture in superspace

$$\tilde{M}_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \bullet}^{(q,t)} = t s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \bullet} + s_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \bullet} + qt s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \bullet & \end{array}} + q s_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} \bullet}$$

$$\tilde{M}_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \end{array}}^{(q,t)} = t s_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \end{array}} + qt s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \end{array}} + q s_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}}$$


Refinement of the original problem!!

Symmetric function theory in superspace

$$\begin{array}{c}
 M_{\Lambda}^{(q,t)} \\
 \vdots \\
 q=t=0 \\
 \vdots \\
 \nearrow \\
 s_{\Lambda}^0
 \end{array}$$


$$\langle p_{\Lambda}, p_{\Omega} \rangle_{qt} = \delta_{\Lambda\Omega} q^{|\Lambda^a|} z_{\Lambda^s}(q, t)$$

Symmetric function theory in **superspace**



Symmetric function theory in superspace

Pieri rules, tableaux generating functions (monomial expansions),
Cauchy identities (RSK).




Symmetric function theory in superspace


Mathieu

Blondeau-Fournier


Pieri Rule




Pieri Rule for $e_r \cdot s_{\Lambda}^0$




Pieri Rule for $e_r \cdot s_{\Lambda}^0$



Pieri Rule for $\tilde{e}_r \cdot s_\Lambda^0$



Pieri Rule for $\tilde{e}_r \cdot s_\Lambda^0$



Monomial expansions

$$s_{\Lambda}^0 \xleftarrow{\text{essentially}} \xrightarrow{\text{dual}} s_{\Lambda}^{\infty}$$

$$s_{\Lambda}^{\infty} = \sum_{T \in \text{Tab}_{\Lambda}^0} (x^{\theta})^T$$


Symmetric function theory in superspace

Lapointe


Preville-Ratelle

MJ


Pieri Rule for s_{Λ}^{∞} .



Pieri Rule for $e_r \cdot s_\Lambda^\infty$



Pieri Rule for $\tilde{e}_r \cdot s_\Lambda^\infty$



Monomial expansions

$$s_{\Lambda}^0 \xleftarrow{\text{essentially}} \xrightarrow{\text{dual}} s_{\Lambda}^{\infty}$$

$$s_{\Lambda}^{\infty} = \sum_{T \in \text{Tab}_{\Lambda}^0} (x\theta)^T$$

$$s_{\Lambda}^0 = \sum_{T \in \text{Tab}_{\Lambda}^{\infty}} (x\theta)^T$$

Cauchy Identity

$$\sum_{\Lambda} s_{\Lambda}^0(x, \theta) s_{\Lambda'}^{\infty}(y, \phi) = \prod_{i,j} (1 + x_i y_j + \theta_i \phi_j)$$


Pieri Rules proofs

$$M_{\Lambda}^{qt} = \mathcal{O}^{qt} E_{\eta(\Lambda)}^{qt} \theta_1 \dots \theta_m + \text{other terms}_{(\text{symmetrization})}$$

$\eta(\Lambda)$ is a composition based on Λ

$E_{\eta(\Lambda)}^{qt}$ is a non-symmetric Macdonald polynomial.

\mathcal{O}^{qt} is some operator.



Key polynomials




Pieri Rules proofs

$$M_{\Lambda}^{qt} = \mathcal{O}^{qt} E_{\eta(\Lambda)}^{qt} \theta_1 \dots \theta_m + \text{other terms}_{(\text{symmetrization})}$$

$\eta(\Lambda)$ is a composition based on Λ

$E_{\eta(\Lambda)}^{qt}$ is a non-symmetric Macdonald polynomial.

\mathcal{O}^{qt} is some operator.




Pieri Rules proofs

$$M_{\Lambda}^{qt} = \mathcal{O}^{qt} E_{\eta(\Lambda)}^{qt} \theta_1 \dots \theta_m + \text{other terms}_{(\text{symmetrization})}$$


$\eta(\Lambda)$ is a composition based on Λ

$E_{\eta(\Lambda)}^{qt}$ is a non-symmetric Macdonald polynomial.

\mathcal{O}^{qt} is some operator.



Symmetric function theory in superspace



$$J_{\Lambda}^{(\alpha)}$$


The Pieri Rules for $e_1 J_{\Lambda}^{(\alpha)}$ are quotients of linear factors in α .



$$\frac{3\alpha(5\alpha + 2)}{(3\alpha + 2)^2(5\alpha + 3)}$$

$$J_{\Lambda}^{(\alpha)}$$


The Pieri Rules for $e_1 J_{\Lambda}^{(\alpha)}$ are quotients of linear factors in α .



$$\frac{3\alpha(5\alpha + 2)}{(3\alpha + 2)^2(5\alpha + 3)}$$

$J_{\Lambda}^{(\alpha)}$

The Pieri Rules for $e_2 J_{\Lambda}^{(\alpha)}$ are quotients of linear factors in α .
 Sometimes there are quadratic factors




$$-\frac{2\alpha^3(3\alpha^2 + \alpha - 1)}{(6\alpha + 5)(7\alpha + 5)(\alpha + 1)(\alpha + 2)(3\alpha + 1)(2\alpha + 1)}$$

Sum of 2 terms

$J_{\Lambda}^{(\alpha)}$

The Pieri Rules for $e_3 J_{\Lambda}^{(\alpha)}$ are quotients of linear factors in α .
 Sometimes there are degree 6 factors!!!!!!!




$$\frac{1}{1152} \frac{\alpha^4 (2\alpha + 3)(3\alpha + 4)(416\alpha^6 + 2000\alpha^5 + 3484\alpha^4 + 2608\alpha^3 + 559\alpha^2 - 256\alpha - 108)}{(4\alpha + 3)(5\alpha + 4)(7\alpha + 6)(2\alpha + 1)(\alpha + 1)^{10}}$$

Sum of 6 terms???????

$J_{\Lambda}^{(\alpha)}$

The Pieri Rules for $e_3 J_{\Lambda}^{(\alpha)}$ are quotients of linear factors in α .
Sometimes there are degree 6 factors!!!!!!!



$$\frac{1}{1152} \frac{\alpha^4(2\alpha + 3)(3\alpha + 4)(416\alpha^6 + 2000\alpha^5 + 3484\alpha^4 + 2608\alpha^3 + 559\alpha^2 - 256\alpha - 108)}{(4\alpha + 3)(5\alpha + 4)(7\alpha + 6)(2\alpha + 1)(\alpha + 1)^{10}}$$


Sum of 7 terms!!!!!!!!!!!!!!

Alternating Sign Matrices!!!!

1, 1, 2, 7, 42, 429, ...

Square Ice!!!!

Sum corresponds to partition function of square ice!!!!!!!



Thank you