

Loss on Ignition and Kjeldahl Digestion for Estimating Organic Carbon and Total Nitrogen in Estuarine Marsh Soils: Calibration with Dry Combustion¹

C. B. CRAFT

Duke Wetland Center

School of Forestry and Environmental Studies

Duke University

Durham, North Carolina 27706

E. D. SENECA

Department of Botany

North Carolina State University

Raleigh, North Carolina 27695-7612

S. W. BROOME

Department of Soil Science

North Carolina State University

Raleigh, North Carolina 27695-7619

ABSTRACT: Soils ($n = 250$) were collected from ten salt and brackish-water marshes of North Carolina and analyzed for organic matter content by loss on ignition (LOI) and Kjeldahl nitrogen (KN). Total organic carbon and total nitrogen were determined on the same samples using an elemental CHN analyzer. Regression analyses indicated that LOI and KN were excellent estimators of organic C ($R^2 = 0.990$) and total N ($R^2 = 0.986$), respectively, in low clay content (0–11%) marsh soils containing a wide range of soil organic C (0.1–28%) and total N (0–1.6%). A quadratic equation best described the relationship between organic C and organic matter (Organic C = $0.40 [\text{LOI}] + 0.0025 [\text{LOI}]^2$) while a linear model accurately described the relationship between total N and Kjeldahl N (Total N = $1.048 [\text{KN}] - 0.010$). The proportion of organic C in organic matter (C/OM) increased with increasing soil organic matter content, probably as a result of aging. Young marshes, which are characterized by low soil organic content contain C/OM ratios similar to emergent vegetation (40–45%). In old organic soils (70–80% organic matter), C/OM increased to 57–60% due to accumulation of reduced organic materials.

Introduction

Marsh soils are a major reservoir of organic carbon (C) and total nitrogen (N) in estuarine ecosystems (Friedman and DeWitt 1978; Craft et al. 1988b) and are important sinks in the global C cycle (Feijtel et al. 1985; Armentano and Menges 1986). Studies of estuarine nutrient cycles and food webs often rely on the measurement of soil organic matter content by loss on ignition (LOI) (Nixon

and Oviatt 1973; Lord 1980; Morris and Bowden 1986; Craft et al. 1988b) or by dichromate oxidation (Cammen 1975; Lindau and Hossner 1981; Broome et al. 1986). Both LOI and dichromate methods require correction factors for estimating soil organic C.

The amount of organic C recovered by dichromate oxidation is highly variable, and ranges from 44% to 92% of the soil organic matter (Nelson and Sommers 1982). Other disadvantages of the dichromate method include interference from chloride and ferrous ions (both of which may be abundant in reduced, estuarine marsh soils; Nelson and Sommers 1982) and the problem of safe disposal of the dichromate-sulfuric acid solution.

Loss on ignition (LOI) is a rapid, safe, and in-

¹ Paper No. 12228 of the Journal Series of the North Carolina Agricultural Research Service, Raleigh, North Carolina 27695-7601. The use of trade names in this publication does not imply endorsement by the North Carolina Agricultural Research Service of the products named, nor criticism of similar ones not mentioned.

expensive method for measuring soil organic matter. The presence of carbonates does not affect the magnitude of the ignition loss (Davies 1974). However, loss of structural water during ignition may overestimate organic matter content in soils containing large amounts of clay (Mook and Hoskin 1982).

Factors ranging from 0.22 to 0.52 have been used to covert organic matter to organic C in marsh soils (Lord 1980; Nixon 1980; Bowden 1984; Morris and Whiting 1986; Craft et al. 1988b). This wide range suggests that, in many studies, LOI may yield inaccurate estimates of organic C.

The Kjeldahl procedure is widely used for estimating total N (KN) in marsh soils (Lindau and Hossner 1981; Bowden 1984; Morris and Bowden 1986; Craft et al. 1988b). The standard Kjeldahl procedure recovers ammonium and much of the organic N. Without appropriate modifications, however, it does not quantitatively recover nitrogen in compounds containing N-O or N-N linkages, or nitrogen in refractory heterocyclic compounds (Bremner and Mulvaney 1982).

We have compared LOI and KN with the dry combustion technique for measuring organic C and total N. Ten estuarine marshes containing a wide range of soil organic C and total N were used. Our goal was to generate regression equations for converting soil organic matter as determined by LOI to organic C and Kjeldahl N to total N.

Methods

Soils were collected from ten estuarine marshes in eastern North Carolina. The sites included both mineral (Carteret series; mixed, thermic typic psammaquents) and organic (Lafitte series; euic, thermic typic medisaprists) soils (Daniels et al. 1984). Soil organic matter content ranged from 0% to 72%. A detailed site description is presented by Craft et al. (1988b). Soils were sampled by taking 10 to 20 cores (8.5 cm diameter by 30 cm deep) from each marsh. Cores were subdivided into 0–10 and 10–30 cm sections and the soil was air-dried, ground, and passed through a 2 mm-mesh diameter sieve.

Samples were tested for carbonates by treating with 6 N HCl and observing effervescence. Carbonates were present in soils collected from two of the ten marshes. Soils taken from these two marshes were treated with 6 N HCl before analyzing for organic C. The clay content of the marsh soils ranged from 0% to 11% (Craft et al. 1986; Sacco 1989). Soils containing appreciable amounts (6–11%) of clay also had high organic matter levels (19–51%) (Craft et al. 1986; Sacco 1989). Mook and Hoskin (1982) observed that an estuarine sed-

iment containing 87% clay, 13% silt, and 0% organic matter lost 10% of its mass after ignition at 450°C. Based on this estimate, we calculate that structural water accounts for only 2% of the ignition loss in our high clay content (11%) soil. For this reason, we believe the contribution of structural water to ignition loss in our soils was negligible.

Two hundred and fifty samples were analyzed for organic C and total N. The LOI and Kjeldahl N data were published previously by Craft et al. (1988b). Organic matter content was measured on weighed 5 cm or 10 cm³ samples by loss on ignition at 450°C for 8 h. Kjeldahl N was determined on 0.25–1.0 g soil samples according to Bremner and Mulvaney (1982). Organic C and total N were measured on 5–75 mg samples using a Perkin-Elmer 2400 CHN analyzer (Nelson and Sommers 1982). Organic matter, organic C, total N, and Kjeldahl N values were corrected to oven-dried weight by drying a subsample at 105°C overnight.

The CHN analyzer was calibrated using acetanilide (71.09% C, 10.36% N). Carbon and nitrogen values for acetanilide standards (n = 17) were $70.99 \pm 0.07\%$ (± 1 standard error) and $10.36 \pm 0.04\%$, respectively. Analysis of soybean leaf (nitrogen = 4.32%) standards (n = 20) yielded values of $43.66 \pm 0.04\%$ for C and $4.33 \pm 0.02\%$ for N.

Fifty soil samples were randomly selected for duplicate analysis. Mean coefficients of variation (CV) were 15.75% and 6.00% for LOI and organic C, respectively. Likewise, the dry combustion technique for total N (mean CV = 9.80%) was more precise than the Kjeldahl method (mean CV = 11.15%).

Regression analysis was used to describe the relationships between the dependent variables, organic C and total N, and the independent variables, Kjeldahl N and organic matter. The data were square-root transformed to improve homogeneity of variance. Residuals were plotted against the independent variable to determine whether a linear or quadratic equation was more appropriate. Once a general model was chosen, the best model (regression through the origin versus simple regression) was determined by an F test ($\alpha = 0.05$) of the difference in the sum of squares error (SSE) and degrees of freedom of the error term (df) between the reduced and full models according to the expression (Steel and Torrie 1980):

$$\frac{\text{SSE (reduced model)} - \text{SSE (full model)}}{\text{df (reduced model)} - \text{df (full model)}} \bigg/ \frac{\text{Mean Square Error (full model)}}{\text{df (full model)}}$$




Fig. 1. The relationship between organic carbon and loss on ignition (LOI) in marsh soils ($n = 250$). The regression equation (± 1 standard error) is Organic C = $(0.40 \pm 0.01)\text{LOI} + (0.0025 \pm 0.0003)\text{LOI}^2$.

Results and Discussion

REGRESSION ANALYSES

The relationship between organic C and organic matter was described by the equation:

$$\text{Organic C} = (0.40 \pm 0.01) \text{LOI} + (0.0025 \pm 0.0003) \text{LOI}^2.$$

The quadratic form (Fig. 1) suggests that the proportion of organic C in soil organic matter (C/OM) increases with soil organic matter content. This does not appear to be due to different organic C sources. Estuarine marshes accumulate soil organic matter by deposition of residues of net primary production (NPP) from emergent vegetation (Craft et al. 1988a). Because marsh emergents such




Fig. 2. The relationship between total nitrogen and Kjeldahl nitrogen (KN) in marsh soils ($n = 250$). The regression equation (± 1 standard error) is Total N = $(1.048 \pm 0.008)\text{KN} - (0.010 \pm 0.005)$.

as *Spartina*, *Juncus*, and *Cladium* have similar tissue organic C content (42–45%) (Craft et al. 1986), different C sources do not explain the increase in C/OM with increasing soil organic matter content.

Differences in the ages of the marshes could account for the increase in C/OM with increasing soil organic content. Young marshes are characterized by low soil organic content (Craft et al. 1988b). The organic matter accumulating in these young soils is recently deposited plant material and the C/OM values reflect those of emergent vegetation. As marshes age, the interaction of water-saturated soils and deposition of NPP result in continued accumulation of soil organic matter. Over time, anaerobic decomposition produces reduced compounds, such as refractory organic compounds, fatty acids, and methane (Stevenson 1986; Reddy and Graetz 1988), containing a higher C content than "fresh" plant tissue. Thus, C/OM increases with increasing soil organic matter content (and marsh age) through accumulation of reduced organic compounds.

With the exception of Morris and Whiting (1986), published conversion factors for estimating organic C from LOI data in wetland and upland soils are in close agreement (Table 1). High clay content or the presence of carbonates in their low organic C (1–4%) soils could explain the low conversion factor used by these researchers.

Regression analysis indicated that the Kjeldahl method recovered approximately 95% of the total soil N (Fig. 2). It is likely that the shortfall is in the

TABLE 1. Factors for converting soil organic matter to organic carbon in wetland and upland soils.

	Reference	Conversion Factor
Wetland:		
(marsh)	Morris and Whiting (1986) ^a	0.225
(marsh)	Craft et al. (this study) ^a	0.400–0.600
Upland:		
(litter)	Lunt (1931) ^{a,b}	0.529–0.556
	Goldin (1987) ^a	0.417
	David (1988) ^a	0.51–0.53
(mineral soil)	Goldin (1987) ^a	0.405
	David (1988) ^a	0.520
(litter and mineral soil)	Ball (1964) ^b	0.458–0.476
	Ranney (1969) ^a	0.556

^a Calibrated by dry combustion.

^b Calibrated by chromic acid oxidation.




Fig. 3. The relationship between total nitrogen and loss on ignition (LOI) in marsh soils ($n = 250$). The regression equation (± 1 standard error) is $\text{Total N} = (0.020 \pm 0.001)\text{LOI} + (0.00005 \pm 0.00002)\text{LOI}^2 - (0.018 \pm 0.008)$.

form of refractory heterocyclic compounds and compounds containing N-N and N-O linkages (Bremner and Mulvaney 1982).

The relationship between total N and organic matter was best described by the quadratic equation:

$$\begin{aligned} \text{Total N} = & (0.020 \pm 0.001)\text{LOI} \\ & + (0.00005 \pm 0.00002)\text{LOI}^2 \\ & - (0.018 \pm 0.008) \end{aligned}$$

The proportion of N in organic matter (N/OM) increased with increasing soil organic content (Fig. 3). The large coefficient of determination ($R^2 = 0.971$, $n = 250$) for this model indicates that soil organic matter content as measured by LOI is useful also for estimating total N in marsh soils.

MARSH SOIL ORGANIC CARBON AND TOTAL NITROGEN

Marsh soil organic C, organic matter content (LOI), and total N ranged from 0.11% to 28.09%, 0.2% to 51%, and 0% to 1.63%, respectively (Table 2). Three of the marshes (N.C. Phosphate #2, Snow's cut #2, Texasgulf #2) were underlain by organic soils (Typic Medisaprists). Percent organic C was greater in the 10–30 cm depth compared to the 0–10 cm depth in these three marshes (Table 2). Likewise, total N increased with depth in two of the three organic soils.

Carbonates were present in soils collected from the Pine Knoll Shores marshes (Table 2) and accounted for 10–31% and 29–45% of the total C

TABLE 2. Mean organic carbon, loss on ignition (LOI), total nitrogen and elemental organic carbon : nitrogen (C:N) ratios in the ten marshes. (Values in parentheses are the number of samples collected from each depth within a given marsh).

Site	Depth (cm)	Organic C (%)	LOI ^a (%)	Total N (%)	C:N (wt : wt)
N.C. Phosphate #1 (15)	0–10	0.21	1.1	0.01	24.5
	10–30	0.17	1.0	0.01	19.8
N.C. Phosphate #2 (20)	0–10	24.92	48.0	1.63	17.8
	10–30	27.50	50.9	1.52	21.1
Oregon inlet #1 (10)	0–10	1.40	2.4	0.10	16.3
	10–30	0.11	0.2	0	—
Oregon inlet #2 (10)	0–10	2.46	5.5	0.18	15.9
	10–30	0.60	1.6	0.03	23.3
Pine Knoll Shores #1 (10)	0–10	0.29	1.2	0.02	16.9
	10–30	[0.13] ^b 0.11 [0.09]	0.4	0	—
Pine Knoll Shores #2 (10)	0–10	0.36	1.2	0.03	14.0
	10–30	[0.04] 0.20 [0.08]	0.6	0.01	23.3
Snow's cut #1 (10)	0–10	1.42	3.9	0.08	20.7
	10–30	0.27	1.2	0.01	31.5
Snow's cut #2 (10)	0–10	7.41	19.3	0.37	23.4
	10–30	10.60	23.5	0.51	24.2
Texasgulf #1 (15)	0–10	2.22	5.0	0.07	37.0
	10–30	1.89	3.8	0.05	44.1
Texasgulf #2 (15)	0–10	22.79	44.0	1.32	20.1
	10–30	28.09	48.9	1.52	21.6

^a From Craft et al. 1988b.

^b Percent carbonate-carbon.

present in the 0–10 and 10–30 cm depths, respectively.

Elemental organic C:N ratios ranged from 14.0 to 44.1 (Table 2). In nine of the marshes, the C:N ratios indicated net mineralization ($\text{C:N} < 20$) of organic N or an equilibrium ($\text{C:N} = 20\text{--}30$) between mineralization and immobilization (Stevenson 1986). The high C:N ratios of the Texasgulf #2 marsh suggested microbial immobilization ($\text{C:N} > 30$) of N in the soil of this marsh (Stevenson 1986). Organic C:N ratios generally increased from the 0–10 cm depth to the 10–30 cm depth (Table 2).

In summary, LOI is a reliable predictor of organic C and Kjeldahl N is a reliable predictor of total N in low clay content (0–11%) marsh soils containing a wide range of soil organic C (0.11–28.09%) and total N (0–1.63%). The amount of organic C in organic matter (C/OM) increases with increasing soil organic matter content. In young, low (1%) organic matter soils, C/OM is similar to marsh emergent vegetation ($\text{C/OM} = 40\%$). Accumulation of reduced organic C compounds in

old, organic (60–80% organic matter) marsh soils produces a C/OM of 55–60%.

ACKNOWLEDGMENTS

We thank W. Robarge, N. Blair, and J. Burkholder for their thorough and constructive reviews of the manuscript.

LITERATURE CITED

- ARMENTANO, T. V. AND E. S. MENGES. 1986. Patterns of change in the carbon balance of organic soil-wetlands of the temperate zone. *Journal of Ecology* 74:755–774.
- BALL, D. F. 1964. Loss-on-ignition as an estimate of organic matter and organic carbon in non-calcareous soils. *Journal of Soil Science* 15:84–92.
- BOWDEN, W. B. 1984. Nitrogen and phosphorus in the sediments of a tidal, freshwater marsh in Massachusetts. *Estuaries* 7:108–118.
- BREMNER, J. M. AND C. S. MULVANEY. 1982. Nitrogen-total. In A. L. Page, R. H. Miller, and D. R. Keeney (eds.), *Methods of Soil Analysis, Part 2. Agronomy* 9:595–624.
- BROOME, S. W., E. D. SENECA, AND W. W. WOODHOUSE, JR. 1986. Long-term growth and development of transplants of the salt-marsh grass *Spartina alterniflora*. *Estuaries* 9:63–74.
- CAMMEN, L. M. 1975. Accumulation rate and turnover time of organic carbon in a salt marsh sediment. *Limnology and Oceanography* 20:1012–1015.
- CRAFT, C. B., S. W. BROOME, AND E. D. SENECA. 1986. Carbon, nitrogen and phosphorus accumulation in man-initiated marsh soils, p. 117–131. In A. Amoozegar (ed.), *Proceedings of the 29th Annual Meeting of the Soil Science Society of North Carolina*. January 20–21, Raleigh, North Carolina.
- CRAFT, C. B., S. W. BROOME, AND E. D. SENECA. 1988a. Estimating sources of soil organic matter in natural and transplanted estuarine marshes using stable isotopes of carbon and nitrogen. *Estuarine Coastal and Shelf Science* 26:633–641.
- CRAFT, C. B., S. W. BROOME, AND E. D. SENECA. 1988b. Nitrogen, phosphorus and organic carbon pools in natural and transplanted marsh soils. *Estuaries* 11:272–280.
- DANIELS, R. B., H. J. KLEISS, S. W. BUOL, H. J. BYRD, AND J. A. PHILLIPS. 1984. *Soil Systems in North Carolina*. North Carolina Agricultural Research Service, bulletin no. 467, Raleigh, North Carolina. 77 p.
- DAVID, M. B. 1988. Use of loss-on-ignition to assess soil organic carbon in forest soils. *Communications in Soil Science and Plant Analysis* 19:1593–1599.
- DAVIES, B. E. 1974. Loss-on-ignition as an estimate of soil organic matter. *Soil Science Society of America Proceedings* 38:150–151.
- FEIJTEL, T. C., R. D. DELAUNE, AND W. H. PATRICK, JR. 1985. Carbon flow in coastal Louisiana. *Marine Ecology Progress Series* 24:255–260.
- FRIEDMAN, R. M. AND C. B. DEWITT. 1978. Wetlands as carbon and nutrient reservoirs: A spatial, historical and societal perspective, p. 175–185. In P. E. Greeson, J. R. Clark, and J. E. Clark (eds.), *Wetlands Functions and Values: The State of our Understanding*. American Water Resources Association, Minneapolis, Minnesota.
- GOLDIN, A. 1987. Reassessing the use of loss-on-ignition for estimating organic matter content in non-calcareous soils. *Communications in Soil Science and Plant Analysis* 18:1111–1116.
- LINDAU, C. W. AND L. R. HOSSNER. 1981. Substrate characterization of an experimental marsh and three natural marshes. *Soil Science Society of America Journal* 45:1171–1176.
- LORD, C. J., III. 1980. The chemistry and cycling of iron, manganese and sulfur in salt marsh sediments. Ph.D. Dissertation, University of Delaware, Newark, Delaware. 177 p.
- LUNT, H. A. 1931. The carbon-organic matter factor in forest soil humus. *Soil Science* 32:27–33.
- MOOK, D. H. AND C. M. HOSKIN. 1982. Organic determinations by ignition: Caution advised. *Estuarine Coastal and Shelf Science* 15:697–699.
- MORRIS, J. T. AND W. B. BOWDEN. 1986. A mechanistic, numerical model of sedimentation, mineralization and decomposition for marsh sediments. *Soil Science Society of America Journal* 50:96–105.
- MORRIS, J. T. AND G. J. WHITING. 1986. Emission of gaseous carbon dioxide from salt marsh sediments and its relation to other carbon losses. *Estuaries* 9:9–19.
- NELSON, D. W. AND L. E. SOMMERS. 1982. Total carbon, organic carbon and organic matter. In A. L. Page, R. H. Miller, and D. R. Keeney (eds.), *Methods of Soil Analysis, Part 2. Agronomy* 9:539–579.
- NIXON, S. W. 1980. Between coastal marshes and coastal waters—A review of twenty years of speculation and research on the role of salt marshes in estuarine productivity and water chemistry, p. 437–525. In P. Hamilton and K. B. Macdonald (eds.), *Estuarine and Wetland Processes with Emphasis on Modeling*. Plenum Publishing, New York.
- NIXON, S. W. AND C. A. OVIATT. 1973. Ecology of a New England salt marsh. *Ecological Monographs* 43:463–498.
- RANNEY, R. W. 1969. An organic carbon-organic matter conversion equation for Pennsylvania surface soils. *Soil Science Society of America Proceedings* 33:809–811.
- REDDY, K. R. AND D. A. GRAETZ. 1988. Carbon and nitrogen dynamics in wetland soils, p. 307–318. In D. D. Hook (ed.), *The Ecology and Management of Wetlands*. Vol. 1. Croom Helm Ltd., London.
- SACCO, J. N. 1989. Infaunal community development of artificially established salt marshes in North Carolina. M.S. Thesis, North Carolina State University, Raleigh, North Carolina.
- STEEL, R. G. D. AND J. H. TORRIE. 1980. *Principles and Procedures of Statistics*. McGraw-Hill Book Company, New York. 633 p.
- STEVENSON, F. J. 1986. *Cycles of Soil*. John Wiley and Sons, New York. 380 p.

Received for consideration, May 14, 1990
Accepted for publication, September 20, 1990