



16S V4 metabarcoding: Illumina library preparation using two-step PCR

Contributors: Katherine R. Murphy

Date of last update: 20-Aug-2026

BACKGROUND:

The 16S metabarcoding protocol detailed here is designed to prepare 16S fragments for sequencing on Illumina platforms using a two-step amplification process. The first step (which we call the amplification PCR) will amplify your region of interest and add an iTru tail. The tail acts as the priming site for the second step (which we call the indexing PCR) that will add iTru indices with the Illumina-required sequencing adapter to your product. The resulting libraries are dual-indexed and can be sequenced on any Illumina sequencer.

This protocol is designed such that it can be modified for any organism or locus by using custom primers in place of the 16S primers for the amplification PCR, as long as the primers also contain the appropriate iTru tail and the amplicon is an appropriate size for Illumina sequencing. To modify, replace the 16S forward and 16S reverse portions of the primers (in the case below, 515-F and 806-R, respectively) with your specific forward and reverse primers. Please note – if you use alternative locus-specific primers that include inosines, you must use regular Taq (such as GoTaq) instead of a high fidelity, proof-reading DNA polymerase. If your primers do not contain inosine, then we recommend using a high fidelity enzyme such as KAPA HiFi. Please make sure to adjust PCR conditions accordingly if using a different locus and/or polymerase.

For running these libraries on an Illumina sequencer, we recommend spiking 10-20% PhiX for MiSeq runs or 20-40% for NextSeq runs to the low complexity of amplicons if your primers did not include heterogeneity spacers. If your locus-specific primers include heterogeneity spacers and/or you're pooling libraries from multiple loci, then the PhiX spike-in can be decreased to 5-20% (or whatever the chosen sequencing core recommends for medium diversity libraries).

For Smithsonian researchers: LAB stocks four plates of unique iTru i5 indices and four plates of unique iTru i7 indices; this is enough to uniquely dual index 384 samples (4 plates), such that no i5 or i7 is repeated within the set. These iTru indexing primers are available for purchase to our LAB users; contact Katie Murphy or a member of LAB staff for details.

Please feel free to reach out to the LAB genomics specialist, Katie Murphy for assistance.



16S PRIMERS (AMPLIFICATION PCR)

- Primer design (three fields separated by spaces below)
 - 1. TruSeq tail 1 → this must be on any custom primers for this protocol; based on TruSeq chemistry. Do NOT edit this portion of the primer sequences.
 - Forward i5: ACACCTCTTTCCCTACACGACGCTCTTCCGATCT
 - Reverse i7: GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT
 - 2. Heterogeneity Spacer (0-8bp sequence) – optional, but highly recommended. This allows for decreased PhiX spike-in and typically improves sequencing run performance.
 - For ease of use, we suggest using heterogeneity spacers designed by Glenn et al. (2019) (<https://baddna.uga.edu/>).
 - 3. 16S primer sequence (or other locus-specific sequence) → can be altered to fit your needs.
 - Forward (515F-Parada): GTGYCAGCMGCCGCGGTA*A
 - Reverse (806R-Apprill): GGACTACNVGGGTWTCTAA*T
 - Make sure to include a phosphorothioate bond at the 3' end (indicated by * above) if you'll be using a high fidelity enzyme.
- These primer examples show the full iTru-tailed 16S primer design. They use the updated EMP 16S V4 primer sequences that have been used for many years in the Earth Microbiome protocol (515F-Parada & 806R-Apprill) and include heterogeneity spacers based on Glenn et al (2019). These are the primers that LAB typically uses; contact Katie Murphy for more information.
 - Product size: approx. 368-374 bp (~291 bp 16S + 67 bp tails + 10-16 bp het. spacer)
 - **iTru Forward: i5 (F) 16S_515FB_iTru_HS** – a mix of the 4 primers below
ACACCTCTTTCCCTACACGACGCTCTTCCGATCT GGTAC GTGYCAGCMGCCGCGGTA*A
ACACCTCTTTCCCTACACGACGCTCTTCCGATCT cAACAC GTGYCAGCMGCCGCGGTA*A
ACACCTCTTTCCCTACACGACGCTCTTCCGATCT atcGGTT GTGYCAGCMGCCGCGGTA*A
ACACCTCTTTCCCTACACGACGCTCTTCCGATCT tcgGTCAA GTGYCAGCMGCCGCGGTA*A
 - **iTru Reverse: i7 (R) 16S_806RB_iTru_HS** – a mix of the 4 primers below
GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT AGGAA GGACTACNVGGGTWTCTAA*T
GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT gAGTGG GGACTACNVGGGTWTCTAA*T
GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT ccACGTC GGACTACNVGGGTWTCTAA*T
GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT ttcTCAGC GGACTACNVGGGTWTCTAA*T



INDEX PRIMERS (INDEXING PCR):

- For LAB users: LAB has iTru indices available in 8 sets of pre-aliquoted plates (10 µl/well at 10 µM). There are 4 plates of i5 and 4 plates of i7. This is enough for 384 unique dual indices, or 147,456 combinatorial pairs. *We highly recommend using unique dual indices over combinatorial as this will eliminate issues due to index hopping.*
- Each sample needs one i7 and one i5 index. While each index can be used for multiple samples if you use combinatorial indexing, the combination of i7 and i5 must be unique to each sample you plan to pool. However, it's even better to use unique dual indices (UDIs) when possible, such as the iTru set described below, as this mitigates any complications due to index hopping.
- This PCR will increase your fragment size by 69 bp. The final library size using this protocol's 16S V4 primers is 437-443 bp.
- Index primer design (fields separated by spaces below)
 - 1. Illumina adapter sequence
 - 2. i7 or i5 index sequence (shown in **bold**)
 - 3. iTru tail 2

- Example Index Primer sequences:

Type	i5 or i7	Name	Sequence
iTru	i7	iTru7_101_01	CAAGCAGAAGACGGCATACGAGAT GGTAACGT GTGACTGGAGTTCA*G
iTru	i5	iTru5_01_A	AATGATACGGCGACCACCGAGATCTACAC ACCGACAA ACACTCTTTCCCTA*C

* indicates a phosphorothioate bond.

- Please refer to the "LAB Indexing Primers 2024" excel file for a full list of available iTru indexing sequences.

SUPPLIES:

- 10 µM iTru-tailed locus-specific primers (forward & reverse)
- 10 µM iTru indexing primers (i5 & i7)
- 2X KAPA HiFi master mix
 - Please remember to adjust your PCR protocols accordingly if you choose to use a different polymerase and/or use an alternative locus.*
- Sterile 0.2 mL PCR tubes or plates
- Sterile silicone sealing mats or films (if using PCR plates)
- KAPA Pure beads (and/or ExoSap Express)
- Ethanol
- 10mM Tris, pH 8.0
 - Alternatively, you can use TLE (10 mM Tris-HCl, 0.1 mM EDTA). *Make sure to NOT use standard 1X TE, as the higher level of EDTA can interfere with sequencing.*
- Nuclease-free water
- HS Qubit or Quant-iT assay
- Filter pipette tips



PROTOCOL:

Make sure to thoroughly clean your bench with 10% bleach (or lab disinfectant) followed by 70% EtOH prior to beginning work each day. Also, make sure to use filter tips.

Step 1 (Optional) – Quantification & dilution of DNA

Quantification and dilution is not mandatory, especially if you are amplifying a large number of samples. However, not quantifying may affect amplification success and product concentration.

- 1.1 Quantify your DNA sample(s) using Qubit or Quant-iT.
- 1.2 Normalize your DNA to desired concentration.

Step 2 – 16S PCR

This protocol describes the use of triplicate reactions. Please note that your project may require more replicates than this. We strongly suggest you review recent literature when planning your project, in case a different number of replicates is applicable.

- 2.1 Retrieve reagents from the freezer and allow them to thaw at room temperature. Vortex and spin down, then place on ice.
- 2.2 Perform the initial PCR **in triplicate** using 10 µL reactions. *Make sure to include negative PCR controls (PCR blanks).* Follow the recipe below to make your master mix, scaling up accordingly. Volumes are given in µL. Vortex the final master mix thoroughly, then spin briefly.

Component	Per rxn	Your master mix:	Final concentration
Nuclease-free water	3.4 µL		
2X KAPA HiFi mix	5.0 µL		1X
10 µM 16S_515F (Parada) with tail	0.3 µL		0.3 µM
10 µM 16S_806R (Aprill) with tail	0.3 µL		0.3 µM
template DNA	1.0 µL		varies
total	10 µL		

- 2.3 Aliquot 9.0 µL of master mix into each PCR tube or well, then add 1.0 µL of template DNA. Close tubes or cover plate with a sterile silicone mat, then spin to collect liquid to the bottom of wells.
- 2.4 Place your tubes/plate into a thermocycler and run the following program with a heated lid:
95°C – 3 min
35 cycles of:
98°C – 20 sec
60°C – 15 sec
72°C – 30 sec
72°C – 1 min
12°C – hold
- 2.5 After the PCR finishes, go directly to Step 3, or samples may be stored at 4°C for up to 3 days (or at -20°C for longer periods).

Step 3 – Gel verification & Pooling

- 3.1 Prepare a 1.5% agarose gel using 1X SB for each plate of PCRs. Each gel should have 8 tiers if you're running PCR plates.
- 3.2 Mix 2 µl PCR product from Step 2 with 2 µl 2X loading dye/10X GelRed. Run gels for a 6 minutes at 125V.
 - a. If you need to make more 2X loading dye/10X GelRed: add 1 µl GelRed stock into 1 mL of 2X loading dye. Mix thoroughly.
- 3.3 Visualize and/or image the gels to determine the presence/absence of PCR products. Record wells that show incorrect or absent bands.
 - a. Product size should be 368-374 bp for 16S V4 primers with heterogeneity spacers.
- 3.4 Pool PCR replicates for each sample (24 uL total per sample), omitting any that had incorrect bands present.



- 3.5 Proceed directly to Step 4, or samples may be stored at 4°C for up to 3 days (or at -20°C for longer periods).

Step 4 – PCR Cleanup – choose 4A or 4B

Clean up your 16S PCR products using either 1.2X KAPA Pure beads (Step 4A) or ExoSap Express (Step 4B). If you have less than 24 µL per sample, please scale the amount of beads or ExoSap accordingly. If you are using a different locus, you may need to adjust the bead ratio based on your amplicon size; refer to bead manufacturer manual for suggestions.

Step 4A – Cleanup with KAPA Pure Beads

- 4A.1 Remove KAPA Pure beads from the fridge and allow to come to room temperature. While you wait, make fresh 80% Ethanol (you need 450 µl per sample).
- 4A.2 Vortex the beads until fully resuspended (~30 seconds).
- 4A.3 To 24 µL of PCR product, add 28.8 µL of KAPA Pure beads (1.2X bead-to-sample ratio).
- 4A.4 Mix thoroughly by pipetting up/down at least 10 times.
- 4A.5 Incubate at room temperature for 5 minutes to bind the DNA to the beads.
- 4A.6 Place the plate/tubes on a magnet for ~3 minutes, or until the supernatant is clear and beads are pelleted.
- 4A.7 Carefully remove and discard the supernatant, making sure to not disturb the pellet.
- 4A.8 With the plate/tubes still on the magnet, wash the beads by adding 200µL of 80% ethanol. Do not mix.
- 4A.9 Incubate at room temp for 30 seconds.
- 4A.10 Remove and discard the supernatant.
- 4A.11 Repeat the ethanol wash (steps 4A.8-10).
- 4A.12 Briefly remove the plate or tubes from the magnet and spin down for a few seconds, then return to the magnetic rack.
- 4A.13 Carefully remove any residual ethanol using a P20 pipette.
- 4A.14 Allow beads to airdry for 3 minutes (not longer!).
- 4A.15 Remove the plate from the magnet.
- 4A.16 Add 22 µL of 10 mM Tris-HCl (pH 8.0) (or TLE: 10 mM Tris-HCl, 0.1 mM EDTA) to each well and use a pipette to fully resuspend the beads.
- 4A.17 Incubate the plate at room temperature for 5 minutes.
- 4A.18 Place the plate back on the magnet for ~3 minutes, or until the supernatant clears.
- 4A.19 Carefully transfer 20 µL of the clear supernatant to a new plate/tube, making sure to not disturb the bead pellet.
- 4A.20 After the cleanup, go directly to Step 5, or samples may be stored at 4°C for up to 3 days (or at -20°C for longer periods).

Step 4B – Cleanup with ExoSap Express

Traditional ExoSap can somewhat degrade your product when you used high fidelity polymerase in your PCR. If you used HiFi or other high fidelity polymerase, use either ExoSap express or a bead clean instead of traditional ExoSap. Please note, LAB no longer stocks traditional ExoSap (we only have Express).

- 4B.1 Dilute your ExoSap stock five-fold with water (eg. 1 µL ExoSap + 4 µL water). Make sure to make enough for all of your reactions; you'll need 6 µL per 24 µL PCR product.
- 4B.2 Add 6 µL of diluted ExoSap to each pooled PCR product. Your final volume in each well should be 30 µL.
- 4B.3 Pipette to mix, seal the plate or tubes, and spin briefly.
- 4B.4 Place into the thermocycler and run the following program with a heated lid:
 - 37°C – 4 min
 - 80°C – 1 min
 - 12°C – hold
- 4B.5 After the cleanup, go directly to Step 5, or samples may be stored at 4°C for up to 3 days (or at -20°C for longer periods).



Step 5 (Optional) – Quantification & Dilution

Step 5 is not mandatory, especially if you are amplifying a large number of samples, and we often omit this step at LAB. However, not quantifying may affect amplification success and product concentration. To compare results between samples, it's always ideal to normalize concentrations.

- 5.1 Use Qubit or Quant-iT to determine the concentration of each sample.
- 5.2 If needed, dilute your samples to 20 ng/uL or lower in 10mM Tris-HCl (pH 8.0).
 - a. You can also normalize all samples to the same concentration, though we find it's often not necessary for this protocol.
- 5.3 After quantification & dilution, go directly to Step 6, or samples may be stored at 4°C for up to 3 days (or at -20°C for longer periods).

Step 6 – Indexing PCR

- 6.1 Determine which indices will be used for which samples; make sure to record these exactly. Each sample needs a unique i7 and i5.
- 6.2 Retrieve the required PCR reagents from the freezer and allow them to thaw at room temperature. Once thawed, briefly vortex the master mix. Invert the indexing primers ~10 times to mix. Spin briefly.
- 6.3 Set up the indexing PCR using 25 μ L reactions; only use one reaction per sample. Make sure you remember to include a new indexing blank as well as continue with the 16S PCR blank. Follow the recipe below to make your master mix, scaling up accordingly. Mix thoroughly then spin briefly.

*a. Remember, your indices are sample-specific; do **NOT** include them in your master mix.*

Component	Per rxn	Your master mix	Final conc.
Nuclease-free water	9.0 μ L		
2X KAPA HiFi mix	12.5 μ L		1X
10 μ M i7 index	0.75 μ L	-----	0.3 μ M
10 μ M i5 index	0.75 μ L	-----	0.3 μ M
template (16S tailed product)	2.0 μ L	-----	varies
total	25 μ L		

- 6.4 After making your master mix that includes only 2X HiFi mix and water, pipette 21.5 μ L of master mix into each well of a new plate.
- 6.5 Add 0.75 μ L of the required i5 index primer to each well. Next, add 0.75 μ L of the required i7 index primer to each well.
 - a. Note – if your i5 and i7 were pre-combined, simply add 1.5 μ L total.
- 6.6 Lastly, add 2 μ L of your template DNA (16S PCR product). Your total reaction volume in each well should be 25 μ L.
- 6.7 Using a P20 or P200 pipette, set to ~15 μ L, mix up/down 5 times. *Avoid making bubbles!*
- 6.8 Seal the plate with a silicone mat or plate seal and spin briefly to collect liquid at the bottom of the wells.
- 6.9 Place the plate (or tubes) into a thermocycler and run the following indexing program with 5-10 cycles, using a heated lid:
 - 95°C – 3 min
 - 5-10 cycles of: (6 cycles is a good starting point)
 - 98°C – 30 sec
 - 65°C – 15 sec
 - 72°C – 30 sec
 - 72°C – 1 min
 - 12°C – hold
- 6.10 After the PCR finishes, go directly to Step 7, or samples may be stored at 4°C for up to 3 days (or at -20°C for longer periods).



Step 7 – Gel verification

- 7.1 Prepare a 1.5% agarose gel with 1X SB.
- 7.2 Mix 2 µl PCR product from Step 6 with 2 µl 2X loading dye/10X GelRed. Run gels for 6-8 minutes at 125V.
- 7.3 Visualize and/or image the gels to determine the presence/absence of indexed PCR products. Record wells that show incorrect or absent bands.
 - a. Product size should be 437-443 bp for correctly indexed 16S V4 libraries. A second band at ~368 bp indicates that not all of your 16S PCR product was indexed.
- 7.4 Go directly to Step 8, or store samples at 4°C for up to 3 days (or -20°C for longer periods).

Step 8 – PCR Cleanup – choose 8A or 8B

*Clean up your PCR products using either KAPA Pure beads (Step 8A) or ExoSap (Step 8B). If you have less than 23 µL per sample, please scale the amount of beads or ExoSap accordingly. Note – if you use ExoSap (Step 8B) at this stage, you **must** clean-up your final pool with Kapa Pure beads (Step 10) before submitting it for sequencing; the sequencing run will fail if there has been no bead cleanup step. If you follow Step 8A, then you will not need to clean the final pool with beads.*

Step 8A – Cleanup with Kapa Pure Beads

If you are using this protocol for amplicons other than 16S, you may need to adjust the bead-to-sample ratio.

- 8A.1 Remove KAPA Pure beads from the fridge and allow to come to room temperature. While you wait, make fresh 80% Ethanol (450 µl per sample).
- 8A.2 Vortex the beads until fully resuspended (~30 seconds).
- 8A.3 In your indexing PCR plate or tubes, add 18 µl of KAPA Pure beads to the ~23 µL of PCR product. This represents a ~0.8X bead-to-sample ratio.
 - a. This ratio is dependent upon the size of the product. You can use less for larger products (refer to KAPA Pure manual).
- 8A.4 Mix thoroughly by pipetting at least 10 times, until the mixture is homogenous.
- 8A.5 Incubate at room temperature for 5 minutes to bind the DNA to the beads.
- 8A.6 Place the plate/tubes on a magnet for ~3 minutes, or until the supernatant is clear and beads are pelleted.
- 8A.7 Carefully remove and discard the supernatant, making sure to not disturb the pellet.
- 8A.8 With the plate/tubes still on the magnet, wash the beads by adding 200 µL of 80% ethanol. Do not mix.
- 8A.9 Incubate at room temp for 30 seconds.
- 8A.10 Remove and discard the supernatant.
- 8A.11 Repeat the ethanol wash (steps 8A.8-10).
- 8A.12 Briefly remove the plate or tubes from the magnet and spin down for a few seconds, then return to the magnetic rack.
- 8A.13 Carefully remove any residual ethanol using a P20 pipette.
- 8A.14 Let the plate dry on the magnetic for 3 minutes (*not longer!*).
- 8A.15 Remove the plate from the magnet.
- 8A.16 Add 22 µL of 10 mM Tris-HCl (pH 8.0) (or TLE: 10 mM Tris-HCl, 0.1 mM EDTA) to each well and use a pipette to fully resuspend the beads.
- 8A.17 Incubate the plate at room temperature for 5 minutes.
- 8A.18 Place the plate back on the magnet for ~3 minutes, or until the supernatant clears.
- 8A.19 Carefully transfer 20 µL of the clear supernatant to a new plate/tube.
- 8A.20 After the cleanup, go directly to Step 9, or samples may be stored at 4°C for up to 3 days (or at -20°C for longer periods).

Step 8B – Cleanup with ExoSap Express

Traditional ExoSap may somewhat degrade your product when you used high fidelity polymerase in your PCR. If you used HiFi or other high fidelity polymerase, use ExoSap express or a bead clean instead of traditional ExoSap. Please note, LAB no longer stocks traditional ExoSap (we only have Express).

- 8B.1 Dilute your ExoSap Express stock five-fold with water (eg. 1 µL ExoSap + 4 µL water). Make sure to make enough for all of your reactions.



- a. You'll need 5.75 μ L per 23 μ L PCR product.
- 8B.2 Add 5.75 μ L of diluted ExoSap to each pooled PCR product. Your final volume in each well should now be ~ 28.75 μ L.
- 8B.3 Pipette to mix, seal the plate, and spin briefly.
- 8B.4 Place into the thermocycler and run the following program with a heated lid:
 - 37°C – 4 min
 - 80°C – 1 min
 - 12°C – hold
- 8B.5 After the cleanup, go directly to Step 9, or samples may be stored at 4°C for up to 3 days (or at -20°C for longer periods).

Step 9 – Library Quantification & Pooling

- 9.1 Quantify your libraries using the Qubit or Quant-iT HS dsDNA assay kit, following manufacturer's instructions.
- 9.2 Pool each library to be sequenced in equal ng or equimolar amounts in a new 1.5 mL tube:
 - a. If you are using a single gene and all samples will be approximately the same size (such as with 16S V4) you can pool using ng.
 - b. If you are using multiple amplicons that vary in size, you should instead pool in equimolar amounts.

Step 10 – Final Pool Cleanup

This protocol is written for cleaning up a 50 μ L aliquot of your pool, but you can adjust this to whatever volume you need. Just make sure to scale the beads proportionally.

*This step is **ONLY** required if you followed Step 8B (ExoSap Cleanup). **Omit** this step if you followed Step 8A (Bead Cleanup), unless you have adapter-dimer concerns. For 16S libraries, we recommend a 0.8X bead-to-sample ratio.*

- 10.1 Remove KAPA Pure beads from the fridge and allow to come to room temperature. While you wait, make 1mL of fresh 80% Ethanol.
- 10.2 Vortex the beads until fully resuspended (~30 seconds).
- 10.3 To a 50 μ L aliquot of your pool in a 1.5 mL tube, add 40 μ L of KAPA Pure beads. This represents a 0.8X bead-to-sample ratio.
 - a. If you wish to clean up a different volume of your pool, make sure to adjust the bead volume accordingly.
 - b. This ratio is dependent upon the size of the product (refer to KAPA Pure manual).
- 10.4 Mix thoroughly by pipetting or vortexing, until the mixture is homogenous.
- 10.5 Incubate at room temperature for 5-15 minutes to bind the DNA to the beads.
- 10.6 Place the tube on a magnetic stand for ~3 minutes, or until the supernatant is clear and beads are pelleted.
- 10.7 Carefully remove and discard the supernatant, making sure to not disturb the pellet.
- 10.8 With the plate/tubes still on the magnet, wash the beads by adding 500 μ L of 80% ethanol. Do not mix.
- 10.9 Incubate at room temp for 30 seconds.
- 10.10 Remove and discard the supernatant.
- 10.11 Repeat the ethanol wash (steps 10.8-10).
- 10.12 Briefly remove the tube from the magnet and spin down for a few seconds, then return to the magnetic rack.
- 10.13 Carefully remove any residual ethanol using a P20 pipette.
- 10.14 Let the plate dry on the magnetic for 3 minutes.
- 10.15 Remove the plate from the magnet.
- 10.16 Add 50 μ L of 10 mM Tris-HCl (pH 8.0) (or TLE: 10 mM Tris-HCl, 0.1 mM EDTA) to the tube and pipette to fully resuspend the beads.
 - a. Adjust the volume of 10 mM Tris if you wish to concentrate or dilute your pool.
- 10.17 While off the magnet, incubate the tube at room temperature for 5 minutes.
- 10.18 Place the tube back on the magnet for ~3 minutes, or until the supernatant clears.
- 10.19 Carefully transfer 48 μ L of the clear supernatant to a new tube.



Step 11 – Final Pool QC

- 11.1 Use HS dsDNA Qubit or Quant-iT to measure the concentration (ng/μL) of your library pool.
- 11.2 Run a 1:10 dilution of your pool on a HS D1000 tape or on the Bioanalyzer, to ensure the product is the correct size and no adapter-dimer is present.
 - a. Alternatively, run 2 uL of your undiluted pool on a 1.5% agarose gel with an appropriate DNA ladder. Suggested settings for mini-gels are 125V for approximately 30 minutes; this will need to be adjusted for different gel rigs. Visualize the gel to determine if an adapter-dimer band is present. Please note that this will not be as sensitive as the Tapestation or Bioanalyzer.
 - b. If adapter-dimer is present, we suggest performing an additional 0.8X bead cleanup. Repeat gel and Qubit quantification after any additional cleanups.
- 11.3 Determine the molarity (nM) of your pool using the equation given in the “Additional Information” section of this guide.
- 11.4 Your pool should be stored at -20°C until submitted for sequencing.

You're now ready to submit an aliquot of your pool to an Illumina sequencing facility, along with any required documentation.

If you need to ship your pool to a sequencing facility, we recommend shipping it on blue or dry ice.

If the pool will be sequenced at NMNH-LAB, directions for order submission can be found on LAB's website. Reach out to Katie Murphy with any questions.

ADDITIONAL INFORMATION:

- If you need to convert from ng/ul to nM, please use the following equation.

$$\frac{(\text{concentration in ng}/\mu\text{L})}{(660 \text{ g/mol}) \times (\text{average library size in bp})} \times 10^6 = \text{concentration in nM}$$

Average library size should be the total length (including index/adapters), not just your insert length.

USEFUL REFERENCES:

1. Glenn, T. C., R. Nilsen, T. J. Kieran, J. W. Finger Jr., T. W. Pierson, K. E. Bentley, S. L. Hoffberg, S. Louha, F. J. Garcia-De Leon, M. A. D. R. Portilla, K. D. Reed, J. L. Anderson, J. K. Meece, S. E. Aggrey, R. Rekaya, M. Alabady, M. Belanger, K. Winker, and B. C. Faircloth. Adapterama I: Universal stubs and primers for thousands of dual-indexed Illumina libraries (iTru & iNext). Preprint at <http://biorxiv.org/content/early/2016/06/15/049114>
2. Glenn TC, Pierson TW, Bayona-Vásquez NJ, Kieran TJ, Hoffberg SL, Thomas IV JC, Lefever DE, Finger JW, Gao B, Bian X, Louha S, Kolli RT, Bentley KE, Rushmore J, Wong K, Shaw TI, Rothrock Jr MJ, McKee AM, Guo TL, Mauricio R, Molina M, Cummings BS, Lash LH, Lu K, Gilbert GS, Hubbell SP, Faircloth BC. 2019. Adapterama II: universal amplicon sequencing on Illumina plat- forms (TaggiMatrix). PeerJ 7:e7786 <http://doi.org/10.7717/peerj.7786>.
3. Bell, J. (2011). Overview of tailed amplicon sequencing approach with MiSeq. Illumina.
4. KAPA Pure Beads technical data sheet, KR1245 – v3.16. July 2016.
5. Herbold, C.W. et al. (2015). A flexible and economical barcoding approach for highly multiplexed amplicon sequencing of diverse target genes. Front Microbiol. <https://doi.org/10.3389/fmicb.2015.00731>
6. 16S Metagenomic sequencing library preparation. Illumina part # 15044223 Rev. A.
7. Smithsonian Global ARMS Program – 18S Metabarcoding protocol.



8. Earth Microbiome 16S metabarcoding protocol - <https://earthmicrobiome.org/protocols-and-standards/16s/>
9. Apprill, A., McNally, S., Parsons, R., & Weber, L. (2015). Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquatic Microbial Ecology*, 75(2), 129–137. <http://doi.org/10.3354/ame01753>
10. Parada, A. E., Needham, D. M., & Fuhrman, J. A. (2016). Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environmental Microbiology*, 18(5), 1403–1414.
11. Beaudry MS, Wang J, Kieran TJ, Thomas J, Bayona-Vasquez NJ, Gao B, Devault A, Brunelle B, Lu K, Wang J-S, Rhodes OE, Glenn TC. 2021. Improved microbial community characterization of 16S rRNA via metagenome hybridization capture enrichment. *Frontiers in Microbiology*, 12:6444662.