



SPECIFICATIONS



TESTS		REQUIREMENTS
Description		White crystalline powder
Identification		The Infrared absorption spectrum of the sample should be concordant with the spectrum obtain from standard
• By FTIR		
Identification		The retention time of the major peak in the chromatogram of the assay preparation should correspond to that in the chromatogram of the standard preparation, as obtained in the assay.
• By HPLC		
Aroma		Very light odour
Assay by HPLC (%w/w on anhydrous basis)		Not less than 90.0 %
Water content by KF (%w/w)		Not more than 0.5 %
Emulsifier food grade		Not more than 10.0%
pH		Between 3.0 to 7.0
Melting range		Between 98.0°C to 102.0°C
Residual Solvents (Class III)		Not more than 5000 ppm
Related Substances (By HPLC)		
a. Palmitic acid		Not more than 0.5%
b. Stearoylethanolamide		Not more than 0.2%
c. Myristoylethanolamide		Not more than 0.2%
d. Individual Maximum impurity		Not more than 0.1%
e. Total Impurities		Not more than 1.0%
Heavy Metals (By ICP-MS)	Cadmium (Cd)	NMT 0.5 ppm
	Arsenic (As)	NMT 0.5 ppm
	Mercury (Hg)	NMT 0.1 ppm
	Lead (Pb)	NMT 0.5 ppm
Microbiological Parameters		
Total plate count (TAMC)		Not more than 1000 cfu/g
Total Combined yeasts and moulds counts(TYMC)		Not more than 100 cfu/g
Tests for specified Microorganisms		
Escherichia Coli		Should be absent
Salmonella		Should be absent
Pseudomonas Aeruginosa		Should be absent
Staphylococcus Aureus		Should be absent