

**PLEASE RIP THIS PAGE OFF THE TEST!**

Appropriate formulae from the AQA formulae book

**Binomial series**

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n \quad (n \in \mathbb{N})$$

$$\text{where } \binom{n}{r} = {}^nC_r = \frac{n!}{r!(n-r)!}$$

$$(1 + x)^n = 1 + nx + \frac{n(n-1)}{1.2}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{1.2\dots r}x^r + \dots \quad (|x| < 1, n \in \mathbb{Q})$$

**Differentiation from first principles**

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$



# First Year Maths and Further Maths combined

## Test B11

### AS Mathematics Pure mock

### 2 hours

#### Materials

- You will need the rip off formula sheet on the front of the paper.
- You should have a graphical or scientific calculator that meets the requirements of the specification.

#### Instructions

- Use black ink or a black ball-point pen. Pencil should only be used for drawing.
- Write your name at the top of this page.
- Answer all questions.
- You must answer each question in the space provided for that question. If you need extra space for your answer(s), ask your teacher for extra paper. Write your name on every sheet and write the question number against your answer(s).
- Do **not** write outside the box around each page or on blank pages.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

#### Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 100.

#### Advice

- Unless stated otherwise, you may quote formulae, without proof, from the rip off formula sheet.
- You do not necessarily need to use all the space provided.

LEAVE BLANK. FOR TEACHER USE					
Question	Marks	Question	Marks	Question	Marks
1		8		15	
2		9		16	
3		10		17	
4		11		18	
5		12		19	
6		13		20	
7		14		21	
				Total Marks	

Answer **all** questions in the spaces provided.

- 1** At a point  $P$  on a curve, the gradient of the tangent to the curve is 10

State the gradient of the normal to the curve at  $P$

Circle your answer.

**[1 mark]**

$-10$

$-0.1$

$0.1$

$10$

- 2** Identify the expression below which is equivalent to  $\left(\frac{2x}{5}\right)^{-3}$

Circle your answer.

**[1 mark]**

$\frac{8x^3}{125}$

$\frac{125x^3}{8}$

$\frac{125}{8x^3}$

$\frac{8}{125x^3}$

**3**

Simplify  $\log_2 8^a$

Circle your answer.

**[1 mark]**

$$a^3$$

$$2a$$

$$3a$$

$$8a$$

**4**

It is given that  $\sin \theta = \frac{4}{5}$  and  $90^\circ < \theta < 180^\circ$

Find the value of  $\cos \theta$

Circle your answer.

**[1 mark]**

$$-\frac{3}{4}$$

$$-\frac{3}{5}$$

$$\frac{3}{5}$$

$$\frac{3}{4}$$

**5**

The coefficient of  $x^2$  in the binomial expansion of  $(1 + ax)^6$  is  $\frac{20}{3}$

Find the two possible values of  $a$

**[3 marks]**

---

---

---

---

---

---

---

---

---

---

**Turn over for the next question**

**6 (a)** Find  $\int \left( 2x^3 + \frac{8}{x^2} \right) dx$

**[3 marks]**

---

---

---

---

---

---

---

---

---

**6 (b)** A curve has gradient function  $\frac{dy}{dx} = 2x^3 + \frac{8}{x^2}$

The  $x$ -intercept of the curve is at the point (2, 0)

Find the equation of the curve.

**[2 marks]**

---

---

---

---

---

---

---

---

**[3 marks]**

[illegible]
$$5 \cos^2 \theta - 4 \sin^2 \theta = 0$$

**[2 marks]**

[illegible]



Do not write  
outside the  
box

8

Find the exact solution of the equation  $\ln(x+1) + \ln(x-1) = \ln 15 - 2\ln 7$

Fully justify your answer.

**[5 marks]**

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

**9 (a)** Given that  $y = x\sqrt{x}$ , find  $\frac{dy}{dx}$

**[2 marks]**

---

---

---

---

---

---

**9 (b)** The line,  $L$ , has equation  $6x - 2y + 5 = 0$

$L$  is a tangent to the curve with equation  $y = x\sqrt{x} + k$

Find the value of  $k$

**[5 marks]**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

10

It is given that  $\sin 15^\circ = \frac{\sqrt{6} - \sqrt{2}}{4}$  and  $\cos 15^\circ = \frac{\sqrt{6} + \sqrt{2}}{4}$

Use these two expressions to show that  $\tan 15^\circ = 2 - \sqrt{3}$

Fully justify your answer.

[4 marks]

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

Turn over for the next question

**11 (a)** The curve  $C_1$  has equation  $y = 2x^2 - 20x + 42$

Express the equation of  $C_1$  in the form

$$y = a(x - b)^2 + c$$

where  $a$ ,  $b$  and  $c$  are integers.

**[3 marks]**

---

---

---

---

---

---

---

---

**11 (b)** Write down the coordinates of the minimum point of  $C_1$

**[1 mark]**

---

---

**11 (c)** The curve  $C_1$  is mapped onto the curve  $C_2$  by a stretch in the  $y$ -direction.

The minimum point of  $C_2$  is at  $(5, -4)$

Find the equation of  $C_2$

**[2 marks]**

---

---

---

---

---

---

$$y = 2x^2 + px + 1$$
$$y = 5x - 2$$

Give your answer in exact form.

**[5 marks]**

[illegible]

**13** Points  $P$  and  $Q$  lie on the curve with equation  $y = x^4$

The  $x$ -coordinate of  $P$  is  $x$

The  $x$ -coordinate of  $Q$  is  $x + h$

**13 (a)** Expand  $(x + h)^4$

**[2 marks]**

---

---

---

---

---

---

**13 (b)** Hence, find an expression, in terms of  $x$  and  $h$ , for the gradient of the line  $PQ$

**[1 mark]**

---

---

---

**13 (c)** Explain how to use the answer from part **(b)** to obtain the gradient function of  $y = x^4$

**[2 marks]**

---

---

---


---

---

---

**14** The curve  $C$  has equation  $y = f(x)$

$C$  has a maximum point at  $P$  with coordinates  $(a, 2b)$  as shown in the diagram below.



**14 (a)**  $C$  is mapped by a single transformation onto curve  $C_1$  with equation  $y = f(x + 2)$

State the coordinates of the maximum point on curve  $C_1$

[1 mark]

---



---



---

**14 (b)**  $C$  is mapped by a single transformation onto curve  $C_2$  with equation  $y = 4f(x)$

State the coordinates of the maximum point on curve  $C_2$

[1 mark]

---



---



---

**14 (c)**  $C$  is mapped by a stretch in the  $x$ -direction onto curve  $C_3$  with equation  $y = f(3x)$

State the scale factor of the stretch.

[1 mark]

---



---



---

**15 (a)** Show that

$$\int_1^a \left(6 - \frac{12}{\sqrt{x}}\right) dx = 6a - 24\sqrt{a} + 18$$

**[3 marks]**

---

---

---

---

---

---

---

---


---

---





The shaded region  $R_2$  is bounded by the curve, the line  $x = a$  and the  $x$ -axis.



Find the value of  $a$

Fully justify your answer.

**[4 marks]**

[illegible]

Do not write  
outside the  
box

16

Prove that the sum of the cubes of two consecutive odd numbers is always a multiple of 4.

**[5 marks]**

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

**Turn over for the next question**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**

**Turn over ►**

**17** A continuous curve has equation  $y = f(x)$

The curve passes through the points  $A(2, 1)$ ,  $B(4, 5)$  and  $C(6, 1)$

It is given that  $f'(4) = 0$


Jasmin made two statements about the nature of the curve  $y = f(x)$  at the point  $B$ :

Statement 1: There is a turning point at  $B$

Statement 2: There is a maximum point at  $B$


**17 (a)** Draw a sketch of the curve  $y = f(x)$  such that Statement 1 is correct and Statement 2 is correct.

**[1 mark]**




- 17 (b)** Draw a sketch of the curve  $y = f(x)$  such that Statement 1 is correct and Statement 2 is **not** correct.

[1 mark]



- 17 (c)** Draw a sketch of the curve  $y = f(x)$  such that Statement 1 is **not** correct and Statement 2 is **not** correct.

[1 mark]



**18** Charlie buys a car for £18 000 on 1 January 2016.

The value of the car decreases exponentially.

The car has a value of £12 000 on 1 January 2018.

**18 (a)** Charlie says:

- because the car has lost £6000 after two years, after another two years it will be worth £6000.

Charlie's friend Kaya says:

- because the car has lost one third of its value after two years, after another two years it will be worth £8000.

Explain whose statement is correct, justifying the value they have stated.

**[2 marks]**

---

---

---

---

---

---

---

---

---

---

- 18 (b)** The value of Charlie's car, £ $V$ ,  $t$  years after 1 January 2016 may be modelled by the equation

$$V = Ae^{-kt}$$

where  $A$  and  $k$  are positive constants.

Find the value of  $t$  when the car has a value of £10 000, giving your answer to two significant figures.

**[5 marks]**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

- 10 (c)** Give a reason why the model, in this context, will not be suitable to calculate the value of the car when  $t = 30$

**[1 mark]**

---

---


---

19

A piece of wire of length 66 cm is bent to form the five sides of a pentagon.

The pentagon consists of three sides of a rectangle and two sides of an equilateral triangle.

The sides of the rectangle measure  $x$  cm and  $y$  cm and the sides of the triangle measure  $x$  cm, as shown in the diagram below.



19 (a) (i) You are given that  $\sin 60^\circ = \frac{\sqrt{3}}{2}$

Explain why the area of the triangle is  $\frac{\sqrt{3}}{4}x^2$

[1 mark]

---



---



---

19 (a) (ii) Show that the area enclosed by the wire,  $A \text{ cm}^2$ , can be expressed by the formula

$$A = 33x - \frac{1}{4}(6 - \sqrt{3})x^2$$

[3 marks]

---



---



---



---



---



---



---



---



---



---





**20 (a)** A circle has equation

$$x^2 + y^2 - 10x - 6 = 0$$

Find the centre and the radius of the circle.

**[2 marks]**

---

---


---

---

---

---

**20 (b)** An equilateral triangle has one vertex at the origin, and one side along the line  $x = 8$ , as shown in the diagram below.



**20 (b) (i)** Show that the vertex at the origin lies inside the circle  $x^2 + y^2 - 10x - 6 = 0$

**[1 mark]**

---

---

---

**20 (b) (ii)** Prove that the triangle lies completely within the circle  $x^2 + y^2 - 10x - 6 = 0$  **[4 marks]**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

21


The line  $L_1$  has equation  $x + 7y - 41 = 0$

$L_1$  is a tangent to the circle  $C$  at the point  $P(6, 5)$

The line  $L_2$  has equation  $y = x + 3$

$L_2$  is a tangent to the circle  $C$  at the point  $Q(0, 3)$

The lines  $L_1$  and  $L_2$  and the circle  $C$  are shown in the diagram below.



21 (a)

Show that the equation of the radius of the circle through  $P$  is  $y = 7x - 37$

[3 marks]

---

---

---

---

---

---

---

---

---

---

Do not write  
outside the  
box

**21 (b)**

Find the equation of  $C$

**[4 marks]**

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.