

REVIEW ARTICLE

T-type calcium channels as therapeutic targets in essential tremor and Parkinson's disease

Lillian G. Matthews¹, Corey B. Puryear¹, Susana S. Correia¹, Sharan Srinivasan^{1,2}, Gabriel M. Belfort¹, Ming-Kai Pan^{3,4,5,6,#} & Sheng-Han Kuo^{7,8,#}

¹Praxis Precision Medicines, Boston, Massachusetts, 02110, USA

²Department of Neurology, University of Michigan, Ann Arbor, Michigan, 48109, USA

³Department and Graduate Institute of Pharmacology, National Taiwan University College of Medicine, Taipei, 10051, Taiwan

⁴Neurobiology and Cognitive Science Center, National Taiwan University, Taipei, 10617, Taiwan

⁵Department of Medical Research, National Taiwan University Hospital, Taipei, 10002, Taiwan

⁶Cerebellar Research Center, National Taiwan University Hospital, Yun-Lin Branch, Yun-Lin, 64041, Taiwan

⁷Department of Neurology, Columbia University, New York, New York, 10032, USA

⁸Initiative for Columbia Ataxia and Tremor, Columbia University, New York, New York, 10032, USA

Correspondence

Lillian G. Matthews, Praxis Precision Medicines, 99 High Street, 30th Floor, Boston, MA 02110, USA.
Tel: +61 (0)450 292 682;
E-mail: lmattthews@praxismedicines.com

Received: 11 October 2022; Revised: 20 December 2022; Accepted: 21 December 2022

Annals of Clinical and Translational Neurology 2023; 10(4): 462–483

doi: 10.1002/acn3.51735

#These authors contributed equally to this work.

Abstract

Neuronal action potential firing patterns are key components of healthy brain function. Importantly, restoring dysregulated neuronal firing patterns has the potential to be a promising strategy in the development of novel therapeutics for disorders of the central nervous system. Here, we review the pathophysiology of essential tremor and Parkinson's disease, the two most common movement disorders, with a focus on mechanisms underlying the genesis of abnormal firing patterns in the implicated neural circuits. Aberrant burst firing of neurons in the cerebello-thalamo-cortical and basal ganglia-thalamo-cortical circuits contribute to the clinical symptoms of essential tremor and Parkinson's disease, respectively, and T-type calcium channels play a key role in regulating this activity in both the disorders. Accordingly, modulating T-type calcium channel activity has received attention as a potentially promising therapeutic approach to normalize abnormal burst firing in these diseases. In this review, we explore the evidence supporting the theory that T-type calcium channel blockers can ameliorate the pathophysiologic mechanisms underlying essential tremor and Parkinson's disease, furthering the case for clinical investigation of these compounds. We conclude with key considerations for future investigational efforts, providing a critical framework for the development of much needed agents capable of targeting the dysfunctional circuitry underlying movement disorders such as essential tremor, Parkinson's disease, and beyond.

Introduction

Essential tremor (ET) and Parkinson's disease (PD) are the two most common movement disorders worldwide. In the United States, an estimated 7 million individuals have ET, representing approximately 2% of the total population,¹ which is slightly higher than reported global prevalence estimates of 0.32–1.33%.^{2,3} Prevalence of ET increases with age, with rates reaching 5.8% in patients aged ≥ 65 years and estimated to increase ~ 1.7 -fold with every decade of life; indeed, rates exceeding 20% have been observed in the 9th and 10th decades of life.²

Prevalence estimates for PD indicate there are approximately 1 million affected individuals aged ≥ 45 years in the United States, with prevalence increasing from 0.4% in patients aged 65–74 years, to nearly 2% in those > 80 years.^{4,5} Importantly, with an increasingly aging population, novel approaches for treating neurologic diseases are paramount. A large focus of ET and PD research has centered around the degeneration of specific cell populations,^{6–10} with the objective of developing disease-modifying therapies, although, to date, no such interventions have been approved.^{11,12} More recently, there has been increasing interest in targeting dysregulated neuronal

activity as a novel and promising approach to treating ET and PD.

Aberrant neuronal activity has been implicated in the pathology of several neurologic conditions, including epilepsy, tinnitus, neuropathic pain, dystonia, ET, PD, and major depression.^{13–17} The role of dysfunctional circuitry is particularly well studied in ET and PD.^{18–20} Specifically, the cerebello-thalamo-cortical (CTC) and basal ganglia-thalamo-cortical (BG) circuits play pivotal roles in motor function, with related CTC and BG disturbances widely implicated in ET and PD, respectively.^{21–26} Abnormal burst firing and dysregulated oscillatory neural activity have been identified as important factors mediating tremor and motor symptoms in both conditions.^{18,27–32} Surgical ablation and deep brain stimulation (DBS) provide supportive evidence for firing pattern modulation in ET and PD. Targeted disruption of the thalamic ventral intermediate nucleus (VIM) via DBS can reduce burst firing in patients with ET and improve tremor symptoms.^{33–35} Additionally, surgical thalamic ablation can improve symptoms in both ET and tremor-dominant PD.^{36,37} In patients with PD, DBS of the internal globus pallidus (GPi) or subthalamic nucleus (STN) can improve rigidity, whereas bradykinesia can be improved by targeting the STN.³⁸ Despite success in many patients, DBS is not without risks common to surgical interventions, including perioperative hemorrhage and infection, with reduced efficacy also observed following long-term use.^{39–42} Consequently, treatment options become limited as the disease advances,⁴³ implicating an urgent need for targeted, noninvasive pharmacologic approaches that can modulate the aberrant circuit activity in ET and PD and provide symptomatic relief.

The search for pharmacologic options has led to identification of various approaches to modulating neural circuit activity in the CTC and BG. One of the most promising targets are T-type calcium channels (TTCC), known to be involved in the generation of burst activity in several brain areas.^{44–46} Notably, TTCCs are highly expressed in specific structures of CTC and BG circuits and play an important role in modulating neuronal action potential firing patterns.^{47–53} Furthermore, increasing evidence highlights their role in the aberrant bursting phenomenon observed in ET and PD,^{18,29,54} with TTCC blockade shown to be capable of inhibiting this bursting phenotype and ameliorating symptoms in animal models of both ET and PD.^{29,54–56} Here, we discuss in depth the mechanistic role of TTCCs in the pathophysiology of ET and PD. More specifically, we review the evidence supporting the potential for TTCC blockade as a pharmacologic approach to symptomatic therapy, with the goal of informing future efforts in developing TTCC-targeted therapies for patients with ET and PD.

TTCC Overview and CNS Expression Patterns

TTCCs—named “T-type” due to their transient calcium conductance—are low voltage-activated calcium channels with key roles in membrane excitability.^{57,58} These channels allow calcium entry into cells at relatively hyperpolarized membrane potentials (Ca_v3 ; -70 to -55 mV)^{58,59} compared with high voltage-activated calcium channels, which activate and allow calcium entry at more depolarized membrane potentials (Ca_v1 and Ca_v2 ; > -40 mV).⁵⁸ A brief hyperpolarizing event within a single cell is sufficient to activate TTCCs, facilitating depolarization of the membrane and activation of low-threshold calcium current leading to generation of low voltage-activated calcium spikes, otherwise known as rebound bursting.⁵⁹ This voltage sensitivity allows TTCCs to support a window current that is near the resting membrane potential, in turn regulating neuronal excitability.^{60,61}

$\text{Ca}_v3.1$,⁶² $\text{Ca}_v3.2$,⁶³ and $\text{Ca}_v3.3$ ⁶⁴ constitute the three known TTCC isoforms, each with unique biophysical features and distribution patterns.^{62,63,65–68} In humans, $\text{Ca}_v3.1$, $\text{Ca}_v3.2$, and $\text{Ca}_v3.3$ are encoded by the genes *CACNA1G* (chromosome 17), *CACNA1H* (chromosome 16), and *CACNA1I* (chromosome 22), respectively.⁶⁹ TTCCs are expressed widely throughout the brain and in peripheral tissues, with particularly high expression within cerebellar motor circuits and tremor-related structures (Figures 1 and 2, Figure S1).^{66,70–75} $\text{Ca}_v3.1$ channels are expressed in the soma and dendrites of neurons in several brain regions including the cerebellum, thalamus, hypothalamus, olfactory bulb, basal ganglia, cortex, and hippocampus, as shown by rodent and human data.^{62,65–67,71–75} One of the key functions of $\text{Ca}_v3.1$ is to regulate the oscillatory activity of thalamic neurons, as first described in cat dorsal lateral geniculate nucleus.⁷⁶ While neural oscillation can refer to rhythmic electrical activity of clusters of activated neurons, individual neurons demonstrate rhythmic oscillation, as shown in the work of McCormick and Pape via intracellular recordings within individual cat dorsal lateral geniculate relay neurons.⁷⁶ Notable high expression levels of $\text{Ca}_v3.1$ in mouse and human cerebellum^{71–75} suggest its potential further role in cerebellar circuit function. Although more widely expressed than $\text{Ca}_v3.1$,⁷⁷ $\text{Ca}_v3.2$ channels in mouse are predominantly expressed in the hippocampus (dentate gyrus),^{72,73,78} while in human brain, expression appears greatest in the thalamus and basal ganglia.^{71,74,75} $\text{Ca}_v3.2$ channels have been shown to control NMDA receptor-mediated transmission at central synapses in cultured rat hippocampal cells expressing the human variant.⁷⁹ $\text{Ca}_v3.2$ are also required for acute pain perception⁸⁰ and excitability of spinal cord lamina II neurons, as demonstrated in mice.⁸¹ $\text{Ca}_v3.3$ channels are similarly




Figure 1. Gene expression patterns of TTCCs in mouse brain. Three-dimensional gene expression patterns of *CACNA1G* ($\text{Ca}_v3.1$), *CACNA1H* ($\text{Ca}_v3.2$), and *CACNA1I* ($\text{Ca}_v3.3$) in mouse whole brain. Gene expression patterns for each isoform are shown for all brain regions on the *left* (cortical distribution visible), and for regions of greatest subcortical expression on the *right*—cerebellum, thalamus, and hippocampus. Red and green denote regions of highest and lowest gene expression levels, respectively. The greatest expression of $\text{Ca}_v3.1$ is in the cerebellum and the greatest expression of $\text{Ca}_v3.2$ is in the hippocampus. Images created from the Allen Mouse Brain Atlas using Brain Explorer 2.0.

expressed in various brain regions in both mouse^{72,73} and human brain,^{74,75} and have been shown to facilitate neuronal oscillation in mice, particularly in the thalamus,

where they are key for generating sleep spindles during non-rapid eye movement (NREM) sleep.⁸² Published data to date suggest that $\text{Ca}_v3.3$, and to a lesser extent $\text{Ca}_v3.2$,




Figure 2. Gene expression patterns of TTCCs in human brain. (A) Microarray-based gene expression of *CACNA1G* (Ca_v3.1), *CACNA1H* (Ca_v3.2), and *CACNA1I* (Ca_v3.3) in human brain. Three-dimensional expression patterns from a representative donor brain are shown for each isoform, overlaid on basal ganglia (purple), thalamus (green), and cerebellum (cyan). Data are normalized across the entire set of microarray samples and shown as z-scores. Images created from the Allen Human Brain Atlas using Brain Explorer 2.0. (B) Violin plots depicting in situ hybridization-based gene expression of *CACNA1G* (Ca_v3.1), *CACNA1H* (Ca_v3.2), and *CACNA1I* (Ca_v3.3) in human brain. Gene expression values are shown in transcripts per million, calculated from a gene model with isoforms collapsed to a single gene. No other normalization steps have been applied. Box plots are shown as median and 25th and 75th percentiles. Plots generated and adapted from The Genotype-Tissue Expression (GTEx) Project portal.

plays the dominant role in sleep spindle rhythogenesis, whereas Ca_v3.1 appears to play a minimal role in sleep spindle activity.^{82–85} It should be noted, however, that these reports rely on indirect measurement of sleep spindles (EEG power in the sigma frequency range, 10–15 Hz). As such, a deeper analysis of specific sleep spindle features (e.g., spindle amplitude, frequency, duration, etc.) may highlight more nuanced roles each Ca_v isoform plays in the generation and/or regulation of sleep spindles.

A key function of TTCCs is to regulate neuronal firing patterns (Figure 3).^{86–89} Burst firing and associated sub-threshold membrane potential oscillations, mediated by

TTCC activation, occur in various brain areas and networks, including populations of thalamic, cerebellar, and cortical neurons, and modulate *thalamocortical* oscillations, spike-wave discharges, and neuropathic pain.^{16,44,90–98} Of note, TTCC genetic mutations have been identified in three families diagnosed with ET and have been implicated in spinocerebellar ataxia 42, epilepsy, autism, schizophrenia, developmental disorders, and other neurologic conditions (Table 1).^{87,99–110} Given their key roles in regulating neuronal firing, pharmacologically targeting TTCCs thus has the potential to yield a wealth of therapeutic benefits in various neurologic disorders. Importantly, while various Food




Figure 3. TTCC neuronal firing patterns. (A) Illustrations representing the firing patterns that are dependent on the contribution of TTCC within various brain areas and networks, including populations of thalamic, cerebellar, and cortical neurons. Tonic firing (*left*) is more likely to occur in cells with either low TTCC expression or in neurons with deactivated TTCC due to a depolarized membrane potential. Low threshold burst firing (*right*) is likely to occur in cells that have a higher density of TTCC channels, such as those in the VIM or STN, or in neurons with hyperpolarized membranes that activate TTCC. (B) Membrane hyperpolarization activates TTCC, leading to production of low threshold spikes and subsequent rebound burst firing. Adapted from Park *et al.*⁸⁶ as licensed under CC BY 4.0. (C) Ca_v3.1, Ca_v3.2, and Ca_v3.3 current traces elicited by action potentials in cultured HEK-293 cells. (D) Illustration of the window current (orange shading) of TTCCs which arises due to overlap of steady-state inactivation and activation properties. The window current of TTCCs occurs within the range of the resting membrane potential (blue shading) and has been shown to be enlarged in patients with CACNA1G mutations and inhibited by TTCC blockade.⁸⁷ (C and D) Adapted from Lory *et al.*⁸⁸ as licensed under CC BY 4.0. I_h , HCN channel-mediated currents; I_T , T-type calcium current; Na_v, voltage-gated sodium channel; V_m, membrane potential.

Table 1. TTCC mutations in neurologic disorders.

Isoform	Disease	Protein-level mutation(s)	Effect on TTCC electrophysiological properties
Ca _v 3.1/CACNA1G ⁹⁹	Essential tremor	<ul style="list-style-type: none"> Gly627Arg Arh456Glu Arg1235Gln 	<ul style="list-style-type: none"> Little to no difference versus wild type
Ca _v 3.1/CACNA1G ¹⁰⁰	Spinocerebellar ataxia 42	<ul style="list-style-type: none"> Ala961Thr 	<ul style="list-style-type: none"> Unknown
Ca _v 3.1/CACNA1G ^{101–103}	Spinocerebellar ataxia 42	<ul style="list-style-type: none"> Arg1715His 	<ul style="list-style-type: none"> Shift toward more positive membrane potentials (inactivation and activation)
Ca _v 3.1/CACNA1G ¹⁰⁴	Spinocerebellar ataxia 42	<ul style="list-style-type: none"> Met1574Lys 	<ul style="list-style-type: none"> Unknown
Ca _v 3.1/CACNA1G ¹⁰⁵	Autosomal-dominant cerebellar syndrome	<ul style="list-style-type: none"> Arg1715His 	<ul style="list-style-type: none"> Shift toward more positive membrane potentials (inactivation and activation)
Ca _v 3.1/CACNA1G ¹⁰⁶	Infantile-onset syndromic cerebellar atrophy	<ul style="list-style-type: none"> Ala961Thr Met1531Val 	<ul style="list-style-type: none"> Gain of function Gain of function
Ca _v 3.1/CACNA1G ⁸⁷	Childhood-onset cerebellar atrophy	<ul style="list-style-type: none"> Ala961Thr Met1531Val 	<ul style="list-style-type: none"> Gain of function Gain of function
Ca _v 3.2/CACNA1H ¹⁰⁷	Autism	<ul style="list-style-type: none"> Arg212Cys Arg902Trp GLy2886Cys Ala1874Val/Arg1871Gln 	<ul style="list-style-type: none"> Loss of function Loss of function Loss of function Loss of function
Ca _v 3.3/CACNA1I ¹⁰⁸	Schizophrenia	<ul style="list-style-type: none"> Thr797Met Arg1311His 	<ul style="list-style-type: none"> Unknown Unknown
Ca _v 3.3/CACNA1I ^{109,110}	Schizophrenia	<ul style="list-style-type: none"> Thr797Met Arg1346His 	<ul style="list-style-type: none"> Unknown

and Drug Administration (FDA)-approved agents, including anticonvulsants, calcium channel blockers, and psychotropics, exhibit some degree of TTCC blockade (Table 2),^{111–120} no currently approved agent selectively targets TTCCs as a primary mechanism of action for its indicated use.

TTCCs in Essential Tremor

Essential tremor (ET) is characterized by a 4–12 Hz postural and action tremor, primarily affecting the upper limbs but often accompanied by head and voice tremors.^{121–123} Management of tremor in affected patients is complicated by multiple comorbidities and high treatment discontinuations, all of which contribute to high disease burden.¹²⁴ Several genetic studies of families with ET suggest that an autosomal-dominant inheritance pattern is common.^{125–128} Variants in the *CACNA1G* gene encoding the Ca_v3.1 isoform have been reported in three families based on whole exome (of 37 families) and whole genome (of eight families) sequencing studies of familial ET; however, corresponding functional studies showed little to no change from wild type.^{99,129} Clinical and preclinical data suggest the dysregulation of neuronal activity in

the CTC network in ET.¹⁸ Within the CTC, TTCCs are highly expressed in deep cerebellar nuclei (DCN) and Purkinje cells (PC),^{6,130,131} ventral thalamic nuclei (particularly the VIM),¹³² and motor cortex (Figure 4).⁶⁶ Neuronal burst firing has been recorded in the VIM in ET patients undergoing DBS, and the firing pattern correlates with tremor.¹⁸ Furthermore, bilateral or unilateral VIM-DBS has both been shown to improve tremor in patients with ET.^{33–35} The inferior olive (IO) is a related TTCC-rich⁶⁶ component that modulates CTC activity and may also be implicated in tremor generation.^{133–136} While there are suggestions that broader olivo-cerebello-thalamo-cortical network dysfunction exists in ET,⁷⁰ the traditional olivary model focusing on a central IO disturbance in ET has been increasingly in question, with limited imaging or postmortem evidence to support its primary role in tremor.¹³⁷ On the other hand, the CTC has been at the center of ET pathophysiology, and neurons at each node (cerebellar cortex, DCN, VIM, and motor cortex) are enriched for TTCC expression to modulate the firing patterns in normal and diseased states.

Animal models have provided insights into the mechanisms underlying tremor generation, the most common of which is the harmaline rodent tremor model.¹³⁸

Table 2. FDA-approved agents with TTCC blocking properties.

	Approved clinical uses
<i>Anticonvulsants</i>	
Ethosuximide ¹¹¹	Absence seizures
Zonisamide ¹¹²	Partial seizures, absence seizures, infantile spasms, essential tremor (second line)
Trimethadione ¹¹³	Absence seizures
Valproic acid ¹¹⁴	Partial seizures, absence seizures, bipolar disorder, migraine
<i>Calcium channel blockers</i>	
Verapamil ¹¹⁵	Hypertension, angina
Amlodipine ¹¹⁶	Hypertension, coronary artery disease, angina
Nicardipine ¹¹⁶	Hypertension, angina
Nimodipine ¹¹⁶	Subarachnoid hemorrhage
<i>Psychotropics</i>	
Pimozide ¹¹⁷	Tourette's syndrome
Fluoxetine ^{118,119}	Major depressive disorder, anxiety disorders, obsessive compulsive disorder, eating disorders, premenstrual dysphoric disorder
Trazodone ¹²⁰	Major depressive disorder, insomnia

Harmaline, a central nervous system (CNS) stimulant, induces acute tremor in a variety of animals,^{139–141} with peak tremor frequency varying by species and ranging from 8–10 Hz in monkeys to 10–16 Hz in pigs.^{141–144} Importantly, harmaline induces tremor by augmenting TTCC function.¹⁴⁵ Both knockout and selective knock-down of *CACNA1G* have been shown to eliminate tremor in a harmaline mouse model,¹⁴⁶ and TTCC blockers have shown efficacy in harmaline-induced tremor in mice and rats in a dose-dependent manner.^{55,56,147} Importantly, harmaline studies have contributed critical insights into the role of TTCC-rich CTC nodes in tremor genesis. Subthreshold oscillations (i.e., neuronal oscillations that occur as a result of intrinsic membrane potential fluctuations but which are below the voltage threshold required to cause a neuron to fire) are dependent on TTCC conductance¹⁴⁸ and closely associated with generation of tremor rhythm, and have been shown to be amplified in IO neurons following harmaline treatment.^{146,149,150} In harmaline-induced tremor, hyperactive glutamatergic IO neurons recruit medial regions of cerebellar cortex via climbing fibers to PC dendrites, inducing PC burst firing rhythmically and synchronously.^{151,152} Additionally, harmaline administration results in the expression of c-fos, a marker for neuronal activation, in PC nuclei in rats, further supporting PC hyperexcitability.¹⁵³ Climbing fiber and PC interactions appear to be particularly important in harmaline-induced tremor. Indeed, *pcd* mice with some PC degeneration but intact climbing fibers have an attenuated response to harmaline administration, with reduced tremor frequency and amplitude relative to

wild-type mice.¹⁵⁴ In contrast, *Lurcher* mice which lack PC neurons and climbing fibers do not develop tremor upon harmaline administration.¹⁵⁴ Interestingly, repeated administration of harmaline to rats results in an attenuated tremor response, which may be due to a reduction in PC neuron burst firing and/or degeneration of PC neurons themselves.^{155–157} Within the DCN, dysfunction of parvalbumin-expressing (PV⁺) excitatory neurons results in action tremor in mice, and DCN synaptic transmission block reverses the tremor.¹⁵⁸ Burst firing of DCN neurons travels to the cerebral cortex via the VIM of the thalamus,⁴⁶ with tremor manifesting as a result of rhythmic modulation of primary motor cortex signals that are conveyed to muscles. In line with this, both VIM neuronal bursting^{159,160} and enhanced rhythmic neuronal activity in the primary motor cortex measured with magnetoencephalography have demonstrated coherence with tremor.²¹

The *Grid2*^{dupE3} mouse model is a recently developed tremor model for ET, in which mice have a genetic mutation of the *Grid2* gene (exon 3 duplication) that leads to increased protein degradation and reduced GluRδ2 cerebellar expression levels, resembling the GluRδ2 deficiency observed in patients with ET.²² GluRδ2 is a synaptic organizer for PC, and the reduction of GluRδ2 levels in *Grid2*^{dupE3} mice results in aberrant climbing fibers extending into parallel fiber synaptic territory,²² mimicking observations in the human postmortem ET cerebellum.¹⁶¹ *Grid2*^{dupE3} mice develop 20 Hz cerebellar oscillatory activity (coherent with action tremor), which can be dampened by ethanol and first-line agents used to treat patients with ET, including propranolol and primidone.²² Notably, IO neurons have different firing patterns, including simple spikes and bursting activity that heavily depends on TTCC, and the burst firing of IO neurons has been shown to be phase-coupled with tremor in *Grid2*^{dupE3} mice,²² supporting that TTCC in IO neurons may contribute to tremor. Interestingly, this mouse model may have some translational value since cerebellar oscillatory activity measured by electroencephalography (EEG) has also been described in patients with ET,²² with evidence suggesting that cerebellar oscillations may be a relevant biomarker of dysregulated cerebellar synaptic organization in some patient subpopulations.¹⁶²

Despite the growing body of preclinical evidence, there has been a paucity of studies on the effects of TTCC blockers on human tremor. Moreover, few selective TTCC blockers exist. For example, 1-octanol and its primary metabolite octanoic acid have shown evidence of tremor reduction in adults with ET.^{163,164} Though 1-octanol has been shown to block TTCC currents in vitro,¹⁶⁵ as a long-chain alcohol, the biologic mechanism of action is likely complex and similar to ethanol. The antiepileptic




Figure 4. CTC circuit dysfunction in ET. (A) Key nodes of the CTC circuit (cerebellar cortex, DCN including the dentate nucleus, thalamic VIM, and motor cortex) and the intersecting white matter pathways that connect them. Orange = Guillain Mollaret triangle connecting red nucleus, IO, and dentate nucleus; red = corticospinal tract, blue = dentatothalamic tract, gray = corticopontine tract, black (infratentorial) = pontocerebellar tract, and black (supratentorial) = thalamocortical tract. (B) Tremor generation is mediated by TTCCs in IO neurons which project climbing fibers to PC dendrites within cerebellar cortex, and form synapses on glutamatergic neurons of the DCN, leading to inhibitor responses followed by rebound excitation. TTCCs play a key role in generating subsequent burst firing in the thalamus, with thalamocortical (dashed line) pathways transmitting tremor-related signals to brainstem and spinal cord via corticopontine and corticospinal tracts. DCN, deep cerebellar nuclei; IO, inferior olive; VIM, ventral intermediate nucleus.

drug, zonisamide, was an early agent considered for ET due to its TTCC blocking properties;¹⁶⁶ however, it has also been shown to modulate voltage-gated sodium channels and GABA receptors,^{167,168} and its clinical efficacy has not been consistently demonstrated.^{169–172} Ethosuximide is another antiepileptic agent with nonselective TTCC blocking properties¹⁷³ that has previously shown

promise in mouse models of ET,⁵⁶ but not in patients.¹⁷⁴ The clinical benefit of these compounds thus remains unclear. In recent years, renewed interest in the development of more selective TTCC blockers as a therapeutic approach in ET has led to various clinical trials investigating the safety and efficacy of this mechanism.^{55,175–178} Suvecaltamide (JZP385/CX-8998) has demonstrated some

improvement in tremor and activities of daily living in a Phase 2 proof-of-concept study in ET patients, although the primary efficacy endpoint was not met.^{175,176} Suvacaltamide is currently under evaluation in Phase 2b trials (NCT05122650). Recent reports from a Phase 2 clinical trial of NBI-827104 indicate lack of tremor severity reduction in adult participants with ET, although it will be important to understand the full pharmacokinetic, efficacy, safety, and tolerability profile of this compound and its impact on activities of daily living. Notably, none of these trials included biomarkers of TTCC blockade; as such, it is difficult to determine whether functional TTCC blockade was achieved. PRAX-944 is a selective TTCC blocker previously shown on cryo-electron microscopy to have high binding affinity to the Cav3.1 pore domain.¹⁷⁹ A recent translational study of PRAX-944 included measurement of sleep spindle-related activity (sigma band EEG power) during NREM sleep as a biomarker of functional TTCC blockade, and demonstrated robust sigma band reduction (relative to other spectral frequencies) in both rodents and healthy human participants.⁵⁵ Furthermore, recent PRAX-944 Phase 2 results showed statistically significant and clinically meaningful improvements in both tremor severity and activities of daily living in patients with ET.¹⁷⁷ PRAX-944 is currently under evaluation in a larger Phase 2b trial.¹⁷⁸ These data with more selective TTCC blockers are promising and highlight the importance of biomarkers of TTCC activity in informing selection of doses expected to produce a functional response in relevant circuitry implicated in ET.

TTCCs in Parkinson's Disease


PD is clinically manifested by cardinal symptoms of bradykinesia (slowness of movement), rigidity, or stiffness that impedes normal motor activities, and/or rest tremor.¹⁸⁰ The motor symptoms of PD are frequently disabled and significantly affect daily function.¹⁸¹ Although the primary pathologic mechanisms driving the neurodegenerative processes in PD have not yet been fully identified, the hypokinetic symptoms, bradykinesia, and rigidity

have been linked to the loss of dopamine (DA) neurons within the substantia nigra pars compacta (SNc) and the resulting reduction of striatal DA tone.^{182–184} Importantly, DA is a major regulator of the BG motor circuit (Figure 5), and degeneration of the SNc is correlated with changes in neuronal activity in BG circuits.^{186–189} Thus, the mainstay of PD treatment over the last five decades has focused on levodopa as a DA replacement agent to normalize the proposed underlying BG circuit dysfunction.¹⁹⁰


Although effective, long-term levodopa therapy can eventually lose efficacy or lead to dyskinesia and hallucinations^{191,192} and contribute to impulsivity,¹⁹³ especially when combined with other dopaminergic agents.¹⁹⁴ For PD with tremor or treatment-refractory dyskinesias, DBS is an effective therapy for minimizing motor fluctuations,^{195–197} where it can be placed in one of the three locations depending on indication and surgical preference. As in ET, DBS of the thalamic VIM is commonly used for tremor suppression in PD. However, VIM-DBS does not address other symptoms of PD, and DBS of the GPi and STN is frequently used with added benefit for bradykinesia and rigidity.^{195,198–201} Indeed, the hypothesis that targeting BG circuit dysfunction may improve motor symptoms in patients with PD is supported by the use of GPi- and STN-DBS, with recent studies demonstrating reduction in PD-associated action and rest tremor with both GPi- and STN-DBS, although the former may require more time than STN-DBS before observation of maximum tremor suppression.^{38,199–201} In patients with advanced disease undergoing BG recordings during DBS placement, hyperactivity and increased bursting are consistently observed in the STN and GPi.^{202–204} This is in line with observations in nonhuman primates treated with 1,2,3,6-tetrahydropyridine (MPTP), which produces a DA-deficient state and parkinsonian-like symptoms.^{27,205} Moreover, lesioning or DBS of the STN reduces hypokinetic symptoms in MPTP-treated nonhuman primates.^{20,206} Despite DBS being effective in many patients, the high risk of associated perioperative complications³⁹ provides a strong impetus for developing novel agents capable of pharmacologically recapitulating the effects of DBS in PD.

Figure 5. BG motor circuit dysfunction in PD. (A) Cross-sectional illustration of BG motor circuit components. (B) In the BG motor circuit, sensorimotor cortex neurons project to the putamen. Dopamine (blue) from the SNc activates the direct (green) pathway which promotes movement in the healthy state via excitatory D1 receptors. The direct pathway projects from the putamen to the GPi, continuing to the thalamus before terminating in motor cortex via thalamocortical projections. (C) Dopamine (blue) also activates the indirect (red) pathway which suppresses movement in the healthy state via inhibitory D2 receptors. The indirect pathway projects from the putamen to the GPe, continuing to the STN, GPi, and thalamus before terminating in motor cortex. In the parkinsonian state, the loss of dopamine from the SNc leads to *hypoactivity* of the direct pathway and *hyperactivity* of the indirect pathway, with the imbalance resulting in PD hypokinetic symptoms. (D) The hyperdirect (purple) pathway projects directly from the cortex to the STN. GPe, external globus pallidus; GPi, internal globus pallidus; SNc, substantia nigra pars compacta; SNr, substantia nigra reticulata; STN, subthalamic nucleus. Adapted from Hashemiyoan, et al.¹⁸⁵ as licensed under CC BY 4.0.


(A)




(B)



(C)



(D)



While much of our understanding of the dysfunctional circuitry in PD to date has come from observations of abnormal bursting activity in primate models or in patients with PD, data comparing the brain activity of patients with PD to healthy participants, or to patients with PD before disease onset, are lacking. In one study comparing advanced and early PD, patients with more advanced PD were shown to have neurons that fire at a higher rate while demonstrating higher overall level of STN background activity.²⁰⁷ In the absence of neuronal recordings from healthy brain, studies across different disease populations can provide added insight. A study comparing STN activity in patients with PD versus ET (a disease without known involvement of the BG) showed that the mean firing rate in STN neurons was significantly increased in PD compared to ET, with a larger percentage of neurons in PD exhibiting the burst firing phenotype.²⁰⁸ Additionally, in a comparison between patients with PD and patients with dystonia (a disease also known to involve the BG), mean GPi discharge rate was also higher in patients with PD.²⁰⁴ Thus, despite the lack of available control data, the combined preclinical and clinical evidence to date support the hypothesis of BG circuit hyperactivity in the pathophysiology of PD.

A specific role for TTCCs in modulating the underlying circuit dysfunction in PD can be inferred from preclinical studies, which allude to TTCC involvement in burst firing within BG circuit nodes, notably the STN. The abnormal bursting activity in the BG circuit associated with PD has been causally linked to TTCC activity in preclinical models. A TTCC blocker, ML218, inhibits TTCC current and consequently reduces low threshold spike and rebound burst activity in STN neurons in *ex vivo* thalamic slices, and further reverses haloperidol-induced catalepsy in a rat model of PD.⁵³ However, ML218 has demonstrated no effect on parkinsonian symptoms in MPTP-treated monkeys;^{52,209} in part, due to its sedative effects, which highlights the need for well-tolerated agents. Further evidence for TTCC involvement in parkinsonian symptoms is seen in the 6-hydroxydopamine (6-OHDA) unilateral lesion model, which results in unilateral depletion of striatal DA innervation. In rats and mice, such DA depletion leads to STN neuronal hyperactivity and rebound bursts thought to be mediated by TTCCs.^{29,210–212} To this end, local infusion of TTCC blockers into the STN of 6-OHDA lesioned rats, but not L-type calcium channel blockers, has demonstrated reduced STN bursting activity and improved motor performance.²⁹ Another study found that STN-targeted DBS has differential effects on open field locomotor activity (distance moved, rearing, and movement duration) in 6-OHDA and healthy rats, whereby delivery of constant negative current reduced STN burst activity and eliminated parkinsonian

symptoms in 6-OHDA rats, while delivery of constant positive current increased STN burst activity and induced a parkinsonian state in healthy rats.⁵⁴ Altogether, these studies suggest a strong relationship between STN burst firing and parkinsonian symptoms that can be modulated by both TTCC blockade and DBS.

While the electrical underpinnings of bradykinesia and rigidity seem to relate, at least in part, to disrupted TTCC modulation of BG motor circuits, the mechanisms underlying PD tremor are less clear. Preclinical PD tremor research is particularly scarce due to the lack of a robust rest tremor model.²¹³ Nonetheless, some evidence suggests that aberrant motor circuitry is similarly implicated in tremor genesis in PD. For example, oscillatory activity in the BG has been observed in patients with PD with limb tremor; specifically, neurons with high-frequency oscillatory activity have been found in the STN and denoted “tremor cells” due to their spike rate correlating with limb tremor frequency.²¹⁴ Notably, recordings derived through thalamic DBS exploration in patients with PD have also identified neuronal subpopulations within the ventral nuclear thalamus,²¹⁵ VIM, and ventralis oralis posterior in which neuronal activity correlates with tremor as assessed via electromyography sensors.²¹⁶ These findings allude to a potential alternative mechanism for PD tremor implicating TTCC-rich nodes within the CTC circuit. Indeed, overlapping CTC involvement and oscillatory flow between the cerebellum and sensorimotor cortex have been reported in patients with PD and ET.²¹⁷ Furthermore, longitudinal functional magnetic resonance imaging in patients with PD has shown increased CTC activity with disease progression, namely greater recruitment of cortical motor-associated and cerebellar areas.²¹⁸ In an attempt to reconcile a BG versus CTC hypothesis in PD tremor, a “dimmer-switch” model²¹⁹ has been proposed, which postulates that the BG circuit initiates the tremor episodes, whereas the CTC circuit propagates and produces the tremor. This hypothesis is consistent with findings from a study comparing the functional connectivity of the BG (GPi, external globus pallidus [GPe], putamen, and caudate) and CTC circuits in patients with PD with tremor and those without;²²⁰ wherein the GPi, GPe, and putamen were activated at the start of tremor episodes, while CTC activity was aligned with tremor amplitude.

As with ET, few studies have focused on the effectiveness of TTCC blockers in PD tremor. In the rat tacrine-induced tremulous jaw movement model (used to mimic parkinsonian tremor),²²¹ mibefradil, a non-CNS-penetrant TTCC blocker, has demonstrated largely no effect on tremor suppression, while NNC 55-0396, a CNS-penetrant TTCC blocker analog, has been shown to reduce tremor.²²² In MPTP monkeys with levodopa-responsive




Figure 6. Targets for pharmacologic intervention in ET and PD. (*Middle, top to bottom*) The VIM has a high concentration of TTCCs and is a common target for DBS placement in patients with ET. In patients with PD, targets for DBS placement include the VIM, STN, or GPi, all areas with high expression of TTCCs. VIM-DBS is commonly used for tremor suppression in PD, while the GPi and STN are the preferred targets for managing tremor-dominant PD, with added benefit for bradykinesia and rigidity. Pharmacologic TTCC blockers therefore have the potential to mimic the effects of DBS as an alternative to surgical intervention in ET and PD. (*Left*) TTCC isoforms ($Ca_v3.1$, $Ca_v3.2$, and $Ca_v3.3$) expressed in olivocerebellar (blue shading) and thalamocortical (pink shading) nodes, with key CTC circuit targets for potential pharmacologic intervention in ET highlighted. Inhibitory and excitatory neurons are denoted by red and blue lines, respectively. (*Right*) TTCC isoforms ($Ca_v3.1$, $Ca_v3.2$, and $Ca_v3.3$) expressed in BG (green shading) and thalamocortical nodes (pink shading), with key BG circuit targets for potential pharmacologic intervention in PD highlighted. Inhibitory and excitatory neurons are denoted by red and blue lines. DCN, deep cerebellar nuclei; GPe, external globus pallidus; GPi, internal globus pallidus; nRT, nucleus reticularis of the thalamus; IO, inferior olive; SNc, substantia nigra compacta; SNr, substantia nigra reticulata; STN, subthalamic nucleus. Adapted from Park *et al.*⁸⁶ as licensed under CC BY 4.0.

tremor, ethosuximide, an anticonvulsant with TTCC blocking properties,^{111,223} has demonstrated tremor suppression.²²⁴ Together, these studies lend support to the hypothesis that motor deficits in PD, including both hypokinetic and tremor symptoms, may be reduced via central TTCC blockade. Ongoing and future clinical efforts investigating the safety and efficacy of novel TTCC blockers in patients with PD will be critical to better establish the plausibility of these mechanistic relationships.

Conclusions and Future Directions

The concept of treating ET and PD as diseases of dysfunctional circuitry represents an important paradigm shift in the management of movement disorders. The development of agents with novel mechanisms of action that can pharmacologically mimic the effects of DBS has the potential to provide much needed alternatives to invasive surgical interventions. While identifying suitable targets for pharmacologic intervention has been challenging, the current

mechanistic exploration of ET and PD points to consistent patterns of disrupted circuitry, including cerebellar and thalamic rhythmic oscillations in ET and STN burst firing in PD. Importantly, TTCC activity appears to be implicated in both ET and PD circuit pathophysiology, with TTCC blockade emerging as a promising therapeutic strategy for alleviating the motor symptoms of these movement disorders (Figure 6). Multiple trials have been completed or are now underway to investigate the safety and efficacy of TTCC blockers in ET and PD. Suvcaltamide (JZP385/CX-8998, NCT03101241),¹⁷⁶ PRAX-944 (NCT05021978),¹⁷⁷ and NBI-827104 (NCT04880616) have been assessed in patients with moderate to severe ET. Additional Phase 2 trials in ET are currently underway for suvcaltamide (NCT05122650) and PRAX-944 (NCT05021991).¹⁷⁸ PRAX-944 is also being evaluated in patients with PD with motor fluctuations. Future investigations including those aimed at improved selectivity for certain Ca_v isoforms may lead to the development of treatments targeted to critical neuronal circuit nodes in ET and PD.

We conclude with some considerations for future investigational efforts. To successfully measure the effect of TTCC blockers on motor symptoms, studies need to be designed carefully to (i) optimize CNS target engagement and drug titration, and (ii) monitor/minimize potential adverse events. Firstly, differences in efficacy among TTCC blockers may reflect differences in CNS penetrance, with some agents more able to cross the blood–brain barrier. Assessing target engagement in response to TTCC blockade via CNS pharmacodynamic biomarkers⁵⁵ may therefore be useful for informing dose range finding in clinical trials. Secondly, TTCCs have been linked to hormone secretion, neurotransmitter release, sleep–wake cycles, and feeding behavior, with many other physiologic roles outside the CNS, including vascular and cardiomyocyte functions.^{89,225–229} Thus, attenuation of aberrant neural activity with minimal disruption to normal TTCC activity should be a therapeutic objective when developing novel agents for ET and PD. Relatedly, drug–target affinity determination will be an important consideration for future drug development efforts. Taken together, these recommendations provide a critical framework for the development of much needed agents capable of targeting the dysfunctional circuitry underlying pathologic conditions such as ET, PD, and beyond.

Author Contributions

Conceptualization and literature review: Lillian G. Matthews, Corey B. Puryear, Susana S. Correia, Sharan Srinivasan, Gabriel M. Belfort, Ming-Kai Pan, and Sheng-Han Kuo. Analysis and interpretation: Lillian G. Matthews, Corey B. Puryear, Susana S. Correia, Sharan Srinivasan, Gabriel M. Belfort, Ming-Kai Pan, and Sheng-Han Kuo. Visualization: Lillian G. Matthews, Corey B. Puryear, Sharan Srinivasan, Gabriel M. Belfort, and Sheng-Han Kuo. Writing—original draft: Lillian G. Matthews, Corey B. Puryear, Susana S. Correia, and Sharan Srinivasan. Writing—review and editing: Lillian G. Matthews, Corey B. Puryear, Susana S. Correia, Sharan Srinivasan, Gabriel M. Belfort, Ming-Kai Pan, and Sheng-Han Kuo. All authors read and approved the final article.

Acknowledgments

We thank Sheryl Torr-Brown, Alyssa Theodore, and Lorena Puto of Simpson Healthcare for assistance with preparing the article. The Genotype-Tissue Expression (GTEx) Project was supported by the [Common Fund](#) of the Office of the Director of the National Institutes of Health, and by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. Figure 2 and Figure S1 in this article were obtained from the GTEx Portal on September 9, 2022.

Funding information

Funding for editorial support was provided by Praxis Precision Medicines. SHK is supported by NINDS #R01 NS104423, NINDS #R01 NS118179, and NINDS #R01 NS124854.

Conflict of Interest Statement

LGM serves as a paid consultant to Praxis Precision Medicines and is a Praxis shareholder. CBP, GMB, and SC were employees of Praxis Precision Medicines during article preparation and are Praxis shareholders. MKP and SHK serve as scientific advisors for Praxis Precision Medicines and Sage Therapeutics.

References

1. Louis ED, Ottman R. How many people in the USA have essential tremor? Deriving a population estimate based on epidemiological data. *Tremor Other Hyperkinet Mov* (NY). 2014;4:259.
2. Louis ED, McCreary M. How common is essential tremor? Update on the worldwide prevalence of essential tremor. *Tremor Other Hyperkinet Mov* (NY). 2021;11:28.
3. Song P, Zhang Y, Zha M, et al. The global prevalence of essential tremor, with emphasis on age and sex: a meta-analysis. *J Glob Health*. 2021;11:04028.
4. Marras C, Beck JC, Bower JH, et al. Prevalence of Parkinson's disease across North America. *NPJ Parkinsons Dis*. 2018;4:21. doi:[10.1038/s41531-018-0058-0](#)
5. Pringsheim T, Jette N, Frolkis A, Steeves TD. The prevalence of Parkinson's disease: a systematic review and meta-analysis. *Mov Disord*. 2014;29(13):1583–1590. doi:[10.1002/mds.25945](#)
6. Louis ED, Faust PL, Vonsattel JPG, et al. Neuropathological changes in essential tremor: 33 cases compared with 21 controls. *Brain*. 2007;130(12):3297–3307.
7. Axelrad JE, Louis ED, Honig LS, et al. Reduced Purkinje cell number in essential tremor: a postmortem study. *Arch Neurol*. 2008;65(1):101–107. doi:[10.1001/archneurol.2007.8](#)
8. Greffard S, Verny M, Bonnet AM, et al. Motor score of the unified Parkinson disease rating scale as a good predictor of Lewy body-associated neuronal loss in the substantia nigra. *Arch Neurol*. 2006;63(4):584–588. doi:[10.1001/archneur.63.4.584](#)
9. Louis ED, Lee M, Babij R, et al. Reduced Purkinje cell dendritic arborization and loss of dendritic spines in essential tremor. *Brain*. 2014;137(Pt 12):3142–3148. doi:[10.1093/brain/awu314](#)
10. Choe M, Cortés E, Vonsattel JP, Kuo SH, Faust PL, Louis ED. Purkinje cell loss in essential tremor: random

- sampling quantification and nearest neighbor analysis. *Mov Disord.* 2016;31(3):393-401. doi:[10.1002/mds.26490](https://doi.org/10.1002/mds.26490)
11. Lang AE, Siderowf AD, Macklin EA, et al. Trial of cinpanemab in early Parkinson's disease. *N Engl J Med.* 2022;387(5):408-420. doi:[10.1056/NEJMoa2203395](https://doi.org/10.1056/NEJMoa2203395)
 12. Pagano G, Taylor KI, Anzures-Cabrera J, et al. Trial of prasinezumab in early-stage Parkinson's disease. *N Engl J Med.* 2022;387(5):421-432. doi:[10.1056/NEJMoa2202867](https://doi.org/10.1056/NEJMoa2202867)
 13. Jeanmonod D, Magnin M, Morel A, et al. Thalamocortical dysrhythmia II: clinical and surgical aspects. *Thalamus Related Syst.* 2001;1(3):245-254.
 14. Llinás R, Ribary U, Jeanmonod D, et al. Thalamocortical dysrhythmia I: functional and imaging aspects. *Thalamus Related Syst.* 2001;1(3):237-244.
 15. Llinás RR, Ribary U, Jeanmonod D, Kronberg E, Mitra PP. Thalamocortical dysrhythmia: a neurological and neuropsychiatric syndrome characterized by magnetoencephalography. *Proc Natl Acad Sci U S A.* 1999;96(26):15222-15227.
 16. Lenz FA, Kwan HC, Dostrovsky JO, Tasker RR. Characteristics of the bursting pattern of action potentials that occurs in the thalamus of patients with central pain. *Brain Res.* 1989;496(1-2):357-360.
 17. Fremont R, Tewari A, Khodakhah K. Aberrant Purkinje cell activity is the cause of dystonia in a shRNA-based mouse model of rapid onset dystonia-parkinsonism. *Neurobiol Dis.* 2015;82:200-212. doi:[10.1016/j.nbd.2015.06.004](https://doi.org/10.1016/j.nbd.2015.06.004)
 18. Milosevic L, Kalia SK, Hodaie M, Lozano AM, Popovic MR, Hutchison WD. Physiological mechanisms of thalamic ventral intermediate nucleus stimulation for tremor suppression [published correction appears in *Brain* 2018;141(9):e72]. *Brain.* 2018;141(7):2142-2155.
 19. Mitchell IJ, Clarke CE, Boyce S, et al. Neural mechanisms underlying parkinsonian symptoms based upon regional uptake of 2-deoxyglucose in monkeys exposed to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. *Neuroscience.* 1989;32(1):213-226.
 20. Bergman H, Wichmann T, DeLong MR. Reversal of experimental parkinsonism by lesions of the subthalamic nucleus. *Science.* 1990;249(4975):1436-1438.
 21. Schnitzler A, Müns C, Butz M, Timmermann L, Gross J. Synchronized brain network associated with essential tremor as revealed by magnetoencephalography. *Mov Disord.* 2009;24(11):1629-1635.
 22. Pan MK, Li YS, Wong SB, et al. Cerebellar oscillations driven by synaptic pruning deficits of cerebellar climbing fibers contribute to tremor pathophysiology. *Sci Transl Med.* 2020;12(526):eaay1769.
 23. Connolly AT, Bajwa JA, Johnson MD. Cortical magnetoencephalography of deep brain stimulation for the treatment of postural tremor. *Brain Stimul.* 2012;5(4):616-624.
 24. Louis ED, Vonsattel JP, Honig LS, et al. Essential tremor associated with pathologic changes in the cerebellum. *Arch Neurol.* 2006;63(8):1189-1193.
 25. Reck C, Florin E, Wojtecki L, et al. Characterisation of tremor-associated local field potentials in the subthalamic nucleus in Parkinson's disease. *Eur J Neurosci.* 2009;29(3):599-612.
 26. Hirschmann J, Hartmann CJ, Butz M, et al. direct relationship between oscillatory subthalamic nucleus-cortex coupling and rest tremor in Parkinson's disease. *Brain.* 2013;136(Pt 12):3659-3670.
 27. Bergman H, Wichmann T, Karmon B, DeLong MR. The primate subthalamic nucleus II: neuronal activity in the MPTP model of parkinsonism. *J Neurophysiol.* 1994;72(2):507-520.
 28. Leblois A, Meissner W, Bioulac B, Gross CE, Hansel D, Boraud T. Late emergence of synchronized oscillatory activity in the pallidum during progressive parkinsonism. *Eur J Neurosci.* 2007;26(6):1701-1713.
 29. Tai CH, Yang YC, Pan MK, Huang CS, Kuo CC. Modulation of subthalamic T-type Ca(2+) channels remedies locomotor deficits in a rat model of Parkinson disease. *J Clin Invest.* 2011;121(8):3289-3305.
 30. Pan MK, Tai CH, Liu WC, Pei JC, Lai WS, Kuo CC. Deranged NMDAergic cortico-subthalamic transmission underlies parkinsonian motor deficits. *J Clin Invest.* 2014;124(10):4629-4641.
 31. Sharott A, Gulberti A, Zittel S, et al. Activity parameters of subthalamic nucleus neurons selectively predict motor symptom severity in Parkinson's disease. *J Neurosci.* 2014;34(18):6273-6285.
 32. Pan MK, Kuo SH, Tai CH, et al. Neuronal firing patterns outweigh circuitry oscillations in parkinsonian motor control. *J Clin Invest.* 2016;126(12):4516-4526. doi:[10.1172/JCI88170](https://doi.org/10.1172/JCI88170)
 33. Benabid AL, Pollak P, Gervason C, et al. Long-term suppression of tremor by chronic stimulation of the ventral intermediate thalamic nucleus. *Lancet.* 1991;337(8738):403-406.
 34. Huss DS, Dallapiazza RF, Shah BB, Harrison MB, Diamond J, Elias WJ. Functional assessment and quality of life in essential tremor with bilateral or unilateral DBS and focused ultrasound thalamotomy. *Mov Disord.* 2015;30(14):1937-1943.
 35. Pahwa R, Lyons KE, Wilkinson SB, et al. Long-term evaluation of deep brain stimulation of the thalamus. *J Neurosurg.* 2006;104(4):506-512.
 36. Jankovic J, Cardoso F, Grossman RG, Hamilton WJ. Outcome after stereotactic thalamotomy for parkinsonian, essential, and other types of tremor. *Neurosurgery.* 1995;37(4):680-686.
 37. Goldman MS, Ahlskog JE, Kelly PJ. The symptomatic and functional outcome of stereotactic thalamotomy for

- medically intractable essential tremor. *J Neurosurg.* 1992;76(6):924-928.
38. Krack P, Benazzouz A, Pollak P, et al. Treatment of tremor in Parkinson's disease by subthalamic nucleus stimulation. *Mov Disord.* 1998;13(6):907-914.
 39. Zarzycki MZ, Domitrz I. Stimulation-induced side effects after deep brain stimulation - a systematic review. *Acta Neuropsychiatr.* 2020;32(2):57-64.
 40. Lu G, Luo L, Liu M, et al. Outcomes and adverse effects of deep brain stimulation on the ventral intermediate nucleus in patients with essential tremor. *Neural Plast.* 2020;2020:2486065.
 41. Paschen S, Forstenpointner J, Becktepe J, et al. Long-term efficacy of deep brain stimulation for essential tremor: an observer-blinded study. *Neurology.* 2019;92(12):e1378-e1386.
 42. Enatsu R, Kitagawa M, Morishita T, et al. Effect of cycling thalamosubthalamic stimulation on tremor habituation and rebound in Parkinson disease. *World Neurosurg.* 2020;12(144):64-67.
 43. Louis ED, Rohl B, Rice C. Defining the treatment gap: what essential tremor patients want that they are not getting. *Tremor Other Hyperkinet Mov (NY).* 2015;5:331.
 44. Vickstrom CR, Liu X, Zhang Y, et al. T-type calcium channels contribute to burst firing in a subpopulation of medial habenula neurons. *eNeuro.* 2020;7(4):ENEURO.0201-20.2020.
 45. Deschênes M, Paradis M, Roy JP, Steriade M. Electrophysiology of neurons of lateral thalamic nuclei in cat: resting properties and burst discharges. *J Neurophysiol.* 1984;51(6):1196-1219. doi:[10.1152/jn.1984.51.6.1196](https://doi.org/10.1152/jn.1984.51.6.1196)
 46. Llinás R, Jahnsen H. Electrophysiology of mammalian thalamic neurones in vitro. *Nature.* 1982;297(5865):406-408. doi:[10.1038/297406a0](https://doi.org/10.1038/297406a0)
 47. Molineux ML, McRory JE, McKay BE, et al. Specific T-type calcium channel isoforms are associated with distinct burst phenotypes in deep cerebellar nuclear neurons. *Proc Natl Acad Sci U S A.* 2006;103(14):5555-5560.
 48. Perez-Reyes E. Molecular physiology of low-voltage-activated t-type calcium channels. *Physiol Rev.* 2003;83(1):117-161.
 49. Lauro PM, Lee S, Akbar U, Asaad WF. Subthalamic-cortical network reorganization during Parkinson's tremor. *J Neurosci.* 2021;41(47):9844-9858.
 50. Beurrier C, Congar P, Bioulac B, Hammond C. Subthalamic nucleus neurons switch from single-spike activity to burst-firing mode. *J Neurosci.* 1999;19(2):599-609. doi:[10.1523/JNEUROSCI.19-02-00599.1999](https://doi.org/10.1523/JNEUROSCI.19-02-00599.1999)
 51. Song WJ, Baba Y, Otsuka T, Murakami F. Characterization of Ca(2+) channels in rat subthalamic nucleus neurons. *J Neurophysiol.* 2000;84(5):2630-2637. doi:[10.1152/jn.2000.84.5.263](https://doi.org/10.1152/jn.2000.84.5.263)
 52. Devergnas A, Chen E, Ma Y, et al. Anatomical localization of Cav3.1 calcium channels and electrophysiological effects of T-type calcium channel blockade in the motor thalamus of MPTP-treated monkeys. *J Neurophysiol.* 2016;115(1):470-485.
 53. Xiang Z, Thompson AD, Brogan JT, et al. The discovery and characterization of ML218: a novel, centrally active T-type calcium channel inhibitor with robust effects in STN neurons and in a rodent model of Parkinson's disease. *ACS Chem Neurosci.* 2011;2(12):730-742. doi:[10.1021/cn200090z](https://doi.org/10.1021/cn200090z)
 54. Tai CH, Pan MK, Lin JJ, Huang CS, Yang YC, Kuo CC. Subthalamic discharges as a causal determinant of parkinsonian motor deficits. *Ann Neurol.* 2012;72(3):464-476.
 55. Scott L, Puryear CB, Belfort GM, et al. Translational pharmacology of PRAX-944, a novel T-type calcium channel blocker in development for the treatment of essential tremor. *Mov Disord.* 2022;37(6):1193-1201.
 56. Handforth A, Homanics GE, Covey DF, et al. T-type calcium channel antagonists suppress tremor in two mouse models of essential tremor. *Neuropharmacology.* 2010;59(6):380-387.
 57. Catterall WA, Perez-Reyes E, Snutch TP, Striessnig J. International Union of Pharmacology. XLVIII. Nomenclature and structure-function relationships of voltage-gated calcium channels. *Pharmacol Rev.* 2005;57(4):411-425.
 58. Powell KL, Cain SM, Snutch TP, O'Brien TJ. Low threshold T-type calcium channels as targets for novel epilepsy treatments. *Br J Clin Pharmacol.* 2014;77(5):729-739.
 59. McCormick DA, Huguenard JR. A model of the electrophysiological properties of thalamocortical relay neurons. *J Neurophysiol.* 1992;68:1384-1400.
 60. Chevalier M, Lory P, Mironneau C, Macrez N, Quignard JF. T-type CaV3.3 calcium channels produce spontaneous low-threshold action potentials and intracellular calcium oscillations. *Eur J Neurosci.* 2006;23:2321-2329.
 61. Dreyfus FM, Tscherter A, Errington AC, et al. Selective T-type calcium channel block in thalamic neurons reveals channel redundancy and physiological impact of I(T) window. *J Neurosci.* 2010;30(1):99-109. doi:[10.1523/JNEUROSCI.4305-09.2010](https://doi.org/10.1523/JNEUROSCI.4305-09.2010)
 62. Perez-Reyes E, Cribbs LL, Daud A, et al. Molecular characterization of a neuronal low-voltage-activated T-type calcium channel. *Nature.* 1998;391(6670):896-900.
 63. Cribbs LL, Lee JH, Yang J, et al. Cloning and characterization of alpha1H from human heart, a member of the T-type Ca²⁺ channel gene family. *Circ Res.* 1998;83(1):103-109.
 64. Lee JH, Daud AN, Cribbs LL, et al. Cloning and expression of a novel member of the low voltage-

- activated T-type calcium channel family. *J Neurosci.* 1999;19(6):1912-1921.
65. Monteil A, Chemin J, Bourinet E, Mennessier G, Lory P, Nargeot J. Molecular and functional properties of the human $\alpha 1G$ subunit that forms T-type calcium channels. *J Biol Chem.* 2000;275(9):6090-6100.
 66. Talley EM, Cribbs LL, Lee JH, Daud A, Perez-Reyes E, Bayliss DA. Differential distribution of three members of a gene family encoding low voltage-activated (T-type) calcium channels. *J Neurosci.* 1999;19(6):1895-1911.
 67. Craig PJ, Beattie RE, Folly EA, et al. Distribution of the voltage-dependent calcium channel $\alpha 1G$ subunit mRNA and protein throughout the mature rat brain. *Eur J Neurosci.* 1999;11(8):2949-2964.
 68. Klöckner U, Lee JH, Cribbs LL, et al. Comparison of the Ca^{2+} currents induced by expression of three cloned $\alpha 1$ subunits, $\alpha 1G$, $\alpha 1H$ and $\alpha 1I$, of low-voltage-activated T-type Ca^{2+} channels. *Eur J Neurosci.* 1999;11(12):4171-4178.
 69. Weiss N, Zamponi GW. Genetic T-type calcium channelopathies. *J Med Genet.* 2020;57(1):1-10.
 70. Shukla AW. Reduction of neuronal hyperexcitability with modulation of T-type calcium channel or SK channel in essential tremor. *Int Rev Neurobiol.* 2022;163:335-355. doi:10.1016/bs.irn.2022.02.008
 71. The Genotype-Tissue Expression (GTEx) Project. GTEx Portal. V8. Accessed September 9, 2022. <https://gtexportal.org/home/>.
 72. Dataset: Allen Institute for Brain Science (2004). Allen Mouse Brain Atlas [dataset]. mouse.brain-map.org. RRID:SCR_002978; Primary publication: Lein ES, Hawrylycz MJ, Ao N, et al. Genome-wide atlas of gene expression in the adult mouse brain. *Nature.* 2007;445(7124):168-176. doi:10.1038/nature05453.
 73. Allen Reference Atlas – Mouse Brain [brain atlas]. atlas.brain-map.org.
 74. Dataset: Allen Institute for Brain Science (2010). Allen Human Brain Atlas: Microarray [dataset]. human.brain-map.org. RRID:SCR_007416; Primary publication: Hawrylycz MJ, Lein ES, Guillozet-Bongaarts AL, et al. An anatomically comprehensive atlas of the adult human transcriptome. *Nature.* 2012;489(7416), 391-399. doi:10.1038/nature11405
 75. Allen Reference Atlas – Human Brain [brain atlas]. atlas.brain-map.org.
 76. McCormick DA, Pape HC. Properties of a hyperpolarization-activated cation current and its role in rhythmic oscillation in thalamic relay neurones. *J Physiol.* 1990;431:291-318. doi:10.1113/jphysiol.1990.sp018331
 77. Aguado C, García-Madróna S, Gil-Minguez M, Luján R. Ontogenic changes and differential localization of T-type Ca^{2+} channel subunits Cav3.1 and Cav3.2 in mouse hippocampus and cerebellum. *Front Neuroanat.* 2016;10:83.
 78. Bernal Sierra YA, Haseleu J, Kozlenkov A, Bégay V, Lewin GR. Genetic tracing of $Ca_v3.2$ T-type calcium channel expression in the peripheral nervous system. *Front Mol Neurosci.* 2017;10:70.
 79. Wang G, Bochorishvili G, Chen Y, et al. Cav3.2 calcium channels control NMDA receptor-mediated transmission: a new mechanism for absence epilepsy. *Genes Dev.* 2015;29(14):1535-1551. doi:10.1101/gad.260869.115
 80. Choi S, Na HS, Kim J, et al. Attenuated pain responses in mice lacking $Ca(V)3.2$ T-type channels. *Genes Brain Behav.* 2007;6(5):425-431. doi:10.1111/j.1601-183X.2006.00268.x
 81. Candelas M, Reynders A, Arango-Lievano M, et al. Cav3.2 T-type calcium channels shape electrical firing in mouse lamina II neurons. *Sci Rep.* 2019;9(1):3112. doi:10.1038/s41598-019-39703-3
 82. Astori S, Wimmer RD, Prosser HM, et al. The $Ca(V)3.3$ calcium channel is the major sleep spindle pacemaker in thalamus. *Proc Natl Acad Sci U S A.* 2011;108(33):13823-13828.
 83. Pellegrini C, Lecci S, Lüthi A, Astori S. Suppression of sleep spindle rhythmogenesis in mice with deletion of Cav3.2 and Cav3.3 T-type Ca^{2+} channels. *Sleep.* 2016;39(4):875-885. doi:10.5665/sleep.5646
 84. Anderson MP, Mochizuki T, Xie J, et al. Thalamic Cav3.1 T-type Ca^{2+} channel plays a crucial role in stabilizing sleep. *Proc Natl Acad Sci U S A.* 2005;102(5):1743-1748. doi:10.1073/pnas.0409644102
 85. Lee J, Kim D, Shin HS. Lack of delta waves and sleep disturbances during non-rapid eye movement sleep in mice lacking $\alpha 1G$ -subunit of T-type calcium channels. *Proc Natl Acad Sci U S A.* 2004;101(52):18195-18199. doi:10.1073/pnas.0408089101
 86. Park YG, Kim J, Kim D. The potential roles of T-type Ca^{2+} channels in motor coordination. *Front Neural Circuits.* 2013;7:172.
 87. Chemin J, Siquier-Pernet K, Nicoulet M, et al. De novo mutation screening in childhood-onset cerebellar atrophy identifies gain-of-function mutations in the CACNA1G calcium channel gene. *Brain.* 2018;141:1998-2013.
 88. Lory P, Nicole S, Monteil A. Neuronal Cav3 channelopathies: recent progress and perspectives. *Pflugers Arch.* 2020;472(7):831-844.
 89. Zamponi GW, Striessnig J, Koschak A, Dolphin AC. The physiology, pathology, and pharmacology of voltage-gated calcium channels and their future therapeutic potential. *Pharmacol Rev.* 2015;67(4):821-870.
 90. Gerke MB, Duggan AW, Xu L, Siddall PJ. Thalamic neuronal activity in rats with mechanical allodynia following contusive spinal cord injury. *Neuroscience.* 2003;117(3):715-722.
 91. Iwata M, LeBlanc BW, Kadasi LM, et al. High-frequency stimulation in the ventral posterolateral thalamus reverses electrophysiologic changes and hyperalgesia in a rat

- model of peripheral neuropathic pain. *Pain*. 2011;152(11):2505-2513.
92. Walton KD, Llinás RR. Central pain as a thalamocortical dysrhythmia: a thalamic efference disconnection? In: Kruger L, Light AR, eds. *Translational Pain Research: from Mouse to Man*. CRC Press/Taylor & Francis; 2010.
 93. Sarnthein J, Jeanmonod D. High thalamocortical theta coherence in patients with neurogenic pain. *Neuroimage*. 2008;39(4):1910-1917. doi:[10.1016/j.neuroimage.2007.10.019](https://doi.org/10.1016/j.neuroimage.2007.10.019)
 94. Sarnthein J, Stern J, Aufenberg C, Rousson V, Jeanmonod D. Increased EEG power and slowed dominant frequency in patients with neurogenic pain. *Brain*. 2006;129(Pt 1):55-64. doi:[10.1093/brain/awh631](https://doi.org/10.1093/brain/awh631)
 95. Amir R, Michaelis M, Devor M. Burst discharge in primary sensory neurons: triggered by subthreshold oscillations, maintained by depolarizing afterpotentials. *J Neurosci*. 2002;22(3):1187-1198.
 96. Pouille F, Cavalier P, Desplantez T, et al. Dendro-somatic distribution of calcium-mediated electrogenesis in purkinje cells from rat cerebellar slice cultures. *J Physiol*. 2000;527(Pt 2):265-282. doi:[10.1111/j.1469-7793.2000.00265.x](https://doi.org/10.1111/j.1469-7793.2000.00265.x)
 97. Swensen AM, Bean BP. Ionic mechanisms of burst firing in dissociated Purkinje neurons. *J Neurosci*. 2003;23(29):9650-9663. doi:[10.1523/JNEUROSCI.23-29-09650.2003](https://doi.org/10.1523/JNEUROSCI.23-29-09650.2003)
 98. Womack MD, Khodakhah K. Dendritic control of spontaneous bursting in cerebellar Purkinje cells. *J Neurosci*. 2004;24(14):3511-3521. doi:[10.1523/JNEUROSCI.0290-04.2004](https://doi.org/10.1523/JNEUROSCI.0290-04.2004)
 99. Odgerel Z, Sonti S, Hernandez N, et al. Whole genome sequencing and rare variant analysis in essential tremor families. *PLoS One*. 2019;14(8):e0220512.
 100. Casas-Alba D, López-Sala L, Pérez-Ordóñez M, et al. Early-onset severe spinocerebellar ataxia 42 with neurodevelopmental deficits (SCA42ND): case report, pharmacological trial, and literature review. *Am J Med Genet A*. 2021;185(1):256-260. doi:[10.1002/ajmg.a.61939](https://doi.org/10.1002/ajmg.a.61939)
 101. Kimura M, Yabe I, Hama Y, et al. SCA42 mutation analysis in a case series of Japanese patients with spinocerebellar ataxia. *J Hum Genet*. 2017;62(9):857-859.
 102. Hashiguchi S, Doi H, Kunii M, et al. Ataxic phenotype with altered Cav3.1 channel property in a mouse model for spinocerebellar ataxia 42. *Neurobiol Dis*. 2019;130:104516. doi:[10.1016/j.nbd.2019.104516](https://doi.org/10.1016/j.nbd.2019.104516)
 103. Morino H, Matsuda Y, Muguruma K, et al. A mutation in the low voltage-gated calcium channel CACNA1G alters the physiological properties of the channel, causing spinocerebellar ataxia. *Mol Brain*. 2015;8:89.
 104. Li X, Zhou C, Cui L, et al. A case of a novel CACNA1G mutation from a Chinese family with SCA42: a case report and literature review. *Medicine (Baltimore)*. 2018;97(36):e12148.
 105. Coutelier M, Blesneac I, Monteil A, et al. A recurrent mutation in CACNA1G alters Cav3.1 T-type calcium-channel conduction and causes autosomal-dominant cerebellar ataxia. *Am J Hum Genet*. 2015;97(5):726-737.
 106. Barresi S, Dentici ML, Manzoni F, et al. Infantile-onset syndromic cerebellar ataxia and CACNA1G mutations. *Pediatr Neurol*. 2020;104:40-45.
 107. Splawski I, Yoo DS, Stotz SC, Cherry A, Clapham DE, Keating MT. CACNA1H mutations in autism spectrum disorders. *J Biol Chem*. 2006;281(31):22085-22091.
 108. Gulsuner S, Walsh T, Watts AC, et al. Consortium on the genetics of schizophrenia (COGS), PAARTNERS study group. Spatial and temporal mapping of de novo mutations in schizophrenia to a fetal prefrontal cortical network. *Cell*. 2013;154:518-529. doi:[10.1016/j.cell.2013.06.049](https://doi.org/10.1016/j.cell.2013.06.049)
 109. Schizophrenia Working Group of the Psychiatric Genomics Consortium. Biological insights from 108 schizophrenia-associated genetic loci. *Nature*. 2014;511(7510):421-427. doi:[10.1038/nature13595](https://doi.org/10.1038/nature13595)
 110. Andrade A, Hope J, Allen A, Yorgan V, Lipscombe D, Pan JQ. A rare schizophrenia risk variant of CACNA1I disrupts Cav3.3 channel activity. *Sci Rep*. 2016;6:6. doi:[10.1038/srep34233](https://doi.org/10.1038/srep34233)
 111. Gomora JC, Daud AN, Weiergraber M, Perez-Reyes E. Block of cloned human T-type calcium channels by succinimide antiepileptic drugs. *Mol Pharmacol*. 2001;60(5):1121-1132.
 112. Matar N, Jin W, Wrubel H, Hescheler J, Schneider T, Weiergraber M. Zonisamide block of cloned human T-type voltage-gated calcium channels. *Epilepsy Res*. 2009;83(2-3):224-234. doi:[10.1016/j.epilepsyres.2008.11.010](https://doi.org/10.1016/j.epilepsyres.2008.11.010)
 113. Zhang YF, Gibbs JW 3rd, Coulter DA. Anticonvulsant drug effects on spontaneous thalamocortical rhythms in vitro: ethosuximide, trimethadione, and dimethadione. *Epilepsy Res*. 1996;23(1):15-36. doi:[10.1016/0920-1211\(95\)00079-8](https://doi.org/10.1016/0920-1211(95)00079-8)
 114. Mattson RH, Cramer JA, Williamson PD, Novelly RA. Valproic acid in epilepsy: clinical and pharmacological effects. *Ann Neurol*. 1978;3:20-25. doi:[10.1002/ana.410030105](https://doi.org/10.1002/ana.410030105)
 115. Bergson P, Lipkind G, Lee SP, Duban ME, Hanck DA. Verapamil block of T-type calcium channels. *Mol Pharmacol*. 2011;79(3):411-419. doi:[10.1124/mol.110.069492](https://doi.org/10.1124/mol.110.069492)
 116. Kopecky BJ, Liang R, Bao J. T-type calcium channel blockers as neuroprotective agents. *Pflugers Arch*. 2014;466(4):757-765. doi:[10.1007/s00424-014-1454-x](https://doi.org/10.1007/s00424-014-1454-x)
 117. Bancila M, Copin JC, Daali Y, Schatlo B, Gasche Y, Bijlenga P. Two structurally different T-type Ca²⁺ channel inhibitors, mibefradil and pimozone, protect CA1 neurons from delayed death after global ischemia in rats. *Fundam Clin Pharmacol*. 2011;25(4):469-478. doi:[10.1111/j.1472-8206.2010.00879.x](https://doi.org/10.1111/j.1472-8206.2010.00879.x)

118. Traboulsie A, Chemin J, Kupfer E, Nargeot J, Lory P. T-type calcium channels are inhibited by fluoxetine and its metabolite norfluoxetine. *Mol Pharmacol.* 2006;69(6):1963-1968. doi:[10.1124/mol.105.020842](https://doi.org/10.1124/mol.105.020842)
119. Deák F, Lasztóczy B, Pacher P, Petheö GL, Kecskeméti V, Spät A. Inhibition of voltage-gated calcium channels by fluoxetine in rat hippocampal pyramidal cells. *Neuropharmacology.* 2000;39(6):1029-1036. doi:[10.1016/s0028-3908\(99\)00206-3](https://doi.org/10.1016/s0028-3908(99)00206-3)
120. Kraus RL, Li Y, Jovanovska A, Renger JJ. Trazodone inhibits T-type calcium channels. *Neuropharmacology.* 2007;53(2):308-317. doi:[10.1016/j.neuropharm.2007.05.011](https://doi.org/10.1016/j.neuropharm.2007.05.011)
121. Bhatia KP, Bain P, Bajaj N, et al. Consensus statement on the classification of tremors. From the task force on tremor of the International Parkinson and Movement Disorder Society. *Mov Disord.* 2018;33(1):75-87.
122. Brennan KC, Jurewicz EC, Ford B, Pullman SL, Louis ED. Is essential tremor predominantly a kinetic or a postural tremor? A clinical and electrophysiological study. *Mov Disord.* 2002;17(2):313-316.
123. Louis ED. Clinical practice. Essential tremor. *N Engl J Med.* 2001;345(12):887-891. doi:[10.1056/NEJMcp010928](https://doi.org/10.1056/NEJMcp010928)
124. Vetterick C, Lyons KE, Matthews LG, Pandal R, Ravina B. The hidden burden of disease and treatment experiences of patients with essential tremor: A retrospective claims data analysis. *Adv Ther.* 2022;39(12):5546-5567.
125. Gulcher JR, Jónsson P, Kong A, et al. Mapping of a familial essential tremor gene, FET1, to chromosome 3q13. *Nat Genet.* 1997;17(1):84-87.
126. Higgins JJ, Pho LT, Nee LE. A gene (ETM) for essential tremor maps to chromosome 2p22-p25. *Mov Disord.* 1997;12(6):859-864.
127. Kovach MJ, Ruiz J, Kimonis K, et al. Genetic heterogeneity in autosomal dominant essential tremor. *Genet Med.* 2001;3(3):197-199.
128. Shatunov A, Sambuughin N, Jankovic J, et al. Genomewide scans in north American families reveal genetic linkage of essential tremor to a region on chromosome 6p23. *Brain.* 2006;129(Pt 9):2318-2331.
129. Liu X, Hernandez N, Kisselev S, et al. Identification of candidate genes for familial early-onset essential tremor. *Eur J Hum Genet.* 2016;24(7):1009-1015.
130. Ly R, Bouvier G, Schonewille M, et al. T-type channel blockade impairs long-term potentiation at the parallel fiber-Purkinje cell synapse and cerebellar learning. *Proc Natl Acad Sci U S A.* 2013;110(50):20302-20307. doi:[10.1073/pnas.1311686110](https://doi.org/10.1073/pnas.1311686110)
131. Cavelier P, Lohof AMB, Lonchamp E, et al. Participation of low-threshold Ca²⁺ spike in the Purkinje cells complex spike. *Neuroreport.* 2008;19(3):299-303. doi:[10.1097/WNR.0b013e3282f4cb96](https://doi.org/10.1097/WNR.0b013e3282f4cb96)
132. Pedrosa DJ, Reck C, Florin E, et al. Essential tremor and tremor in Parkinson's disease are associated with distinct 'tremor clusters' in the ventral thalamus. *Exp Neurol.* 2012;237(2):435-443.
133. Beitz AJ, Saxon D. Harmaline-induced climbing fiber activation causes amino acid and peptide release in the rodent cerebellar cortex and a unique temporal pattern of Fos expression in the olivo-cerebellar pathway. *J Neurocytol.* 2004;33:49-74.
134. Hallett M, Dubinsky RM. Glucose metabolism in the brain of patients with essential tremor. *J Neurol Sci.* 1993;114(1):45-48.
135. Boecker H, Wills AJ, Ceballos-Baumann A, et al. The effect of ethanol on alcohol-responsive essential tremor: a positron emission tomography study. *Ann Neurol.* 1996;39(5):650-658.
136. Simantov R, Snyder SH, Oster-Granite ML. Harmaline-induced tremor in the rat: abolition by 3-acetylpyridine destruction of cerebellar climbing fibers. *Brain Res.* 1976;114(1):144-151.
137. Louis ED, Lenka A. The olivary hypothesis of essential tremor: time to lay this model to rest? *Tremor Other Hyperkinet Mov (N Y).* 2017;7:473. doi:[10.7916/D8FF40RX](https://doi.org/10.7916/D8FF40RX)
138. Martin FC, Le AT, Handforth A. Harmaline-induced tremor as a potential preclinical screening method for essential tremor medications. *Mov Disord.* 2005;20:298-305.
139. De Montigny C, Lamarre Y. Effects produced by local applications of harmaline in the inferior olive. *Can J Physiol Pharmacol.* 1975;53(5):845-849.
140. Wilms H, Sievers J, Deuschl G. Animal models of tremor. *Mov Disord.* 1999;14(4):557-571.
141. Lee J, Kim I, Lee J, et al. Development of harmaline-induced tremor in a swine model. *Tremor Other Hyperkinet Mov (N Y).* 2018;8:532.
142. Miwa H. Rodent models of tremor. *Cerebellum.* 2007;6(1):66-72. doi:[10.1080/14734220601016080](https://doi.org/10.1080/14734220601016080)
143. Lamarre Y, Mercier LA. Neurophysiological studies of harmaline-induced tremor in the cat. *Can J Physiol Pharmacol.* 1971;49(12):1049-1058.
144. Kosmowska B, Wardas J. The pathophysiology and treatment of essential tremor: the role of adenosine and dopamine receptors in animal models. *Biomolecules.* 2021;11(12):1813. doi:[10.3390/biom11121813](https://doi.org/10.3390/biom11121813)
145. Handforth A. Harmaline tremor: underlying mechanisms in a potential animal model of essential tremor. *Tremor Other Hyperkinet Mov (NY).* 2012;2:02-92-769-1. doi:[10.7916/D8TD9W2P](https://doi.org/10.7916/D8TD9W2P)
146. Park YG, Park HY, Lee CJ, et al. Ca(V)3.1 is a tremor rhythm pacemaker in the inferior olive. *Proc Natl Acad Sci U S A.* 2010;107(23):10731-10736.
147. Sinton CM, Krosser BI, Walton KD, Llinás RR. The effectiveness of different isomers of octanol as blockers of harmaline-induced tremor. *Pflugers Arch.* 1989;414(1):31-36.

148. Choi S, Yu E, Kim D, et al. Subthreshold membrane potential oscillations in inferior olive neurons are dynamically regulated by P/Q- and T-type calcium channels: a study in mutant mice. *J Physiol*. 2010;588(16):3031-3043.
149. Llinás R, Yarom Y. Oscillatory properties of Guinea-pig inferior olivary neurones and their pharmacological modulation: an in vitro study. *J Physiol*. 1986;376:163-182.
150. Crunelli V, Lightowler S, Pollard CE. A T-type Ca^{2+} current underlies low-threshold Ca^{2+} potentials in cells of the cat and rat lateral geniculate nucleus. *J Physiol*. 1989;413:543-561.
151. Lang EJ. Organization of olivocerebellar activity in the absence of excitatory glutamatergic input. *J Neurosci*. 2001;21(5):1663-1675.
152. Brown AM, White JJ, van der Heijden ME, Zhou J, Lin T, Sillitoe RV. Purkinje cell misfiring generates high-amplitude action tremors that are corrected by cerebellar deep brain stimulation. *Elife*. 2020;9:e51928.
153. Tian JB, Bishop GA. Stimulus-dependent activation of c-Fos in neurons and glia in the rat cerebellum. *J Chem Neuroanat*. 2002;23(3):157-170.
154. Milner TE, Cadoret G, Lessard L, Smith AM. EMG analysis of harmaline-induced tremor in normal and three strains of mutant mice with Purkinje cell degeneration and the role of the inferior olive. *J Neurophysiol*. 1995;73(6):2568-2577.
155. Lorden JF, Stratton SE, Mays LE, Oltmans GA. Purkinje cell activity in rats following chronic treatment with harmaline. *Neuroscience*. 1988;27:465-472.
156. Bernard JF, Buisseret-Delmas C, Compoint C, Laplante S. Harmaline induced tremor III a combined simple units, horseradish peroxidase, and 2-deoxyglucose study of the olivocerebellar system in the rat. *Exp Brain Res*. 1984;57:128-137.
157. O'Hearn E, Molliver ME. Degeneration of Purkinje cells in parasagittal zones of the cerebellar vermis after treatment with ibogaine or harmaline. *Neuroscience*. 1993;55:303-310. doi:10.1016/0306-4522(93)90500-F
158. Zhou M, Melin MD, Xu W, Südhof TC. Dysfunction of parvalbumin neurons in the cerebellar nuclei produces an action tremor. *J Clin Invest*. 2020;130(10):5142-5156. doi:10.1172/JCI135802
159. Hua SE, Lenz FA, Zirh TA, Reich SG, Dougherty PM. Thalamic neuronal activity correlated with essential tremor. *J Neurol Neurosurg Psychiatry*. 1998;64(2):273-276. doi:10.1136/jnnp.64.2.273
160. Hua SE, Lenz FA. Posture-related oscillations in human cerebellar thalamus in essential tremor are enabled by voluntary motor circuits. *J Neurophysiol*. 2005;93(1):117-127. doi:10.1152/jn.00527.2004
161. Lin H, Magrane J, Clark EM, et al. Early VGLUT1-specific parallel fiber synaptic deficits and dysregulated cerebellar circuit in the KIKO mouse model of Friedreich ataxia. *Dis Model Mech*. 2017;10(12):1529-1538.
162. Wong SB, Wang YM, Lin CC, et al. Cerebellar oscillations in familial and sporadic essential tremor. *Cerebellum*. 2022;21(3):425-431. doi:10.1007/s12311-021-01309-9
163. Voller B, Lines E, McCrossin G, et al. Dose-escalation study of octanoic acid in patients with essential tremor. *J Clin Invest*. 2016;126(4):1451-1457. doi:10.1172/JCI83621
164. Haubenberger D, McCrossin G, Lungu C, et al. Octanoic acid in alcohol-responsive essential tremor: a randomized controlled study. *Neurology*. 2013;80(10):933-940. doi:10.1212/WNL.0b013e3182840c4f
165. Joksovic PM, Choe WJ, Nelson MT, Orestes P, Brimelow BC, Todorovic SM. Mechanisms of inhibition of T-type calcium current in the reticular thalamic neurons by 1-octanol: implication of the protein kinase C pathway. *Mol Pharmacol*. 2010;77(1):87-94. doi:10.1124/mol.109.059931
166. Kito M, Maehara M, Watanabe K. Mechanisms of T-type calcium channel blockade by zonisamide. *Seizure*. 1996;5(2):115-119.
167. Leppik IE. Zonisamide: chemistry, mechanism of action, and pharmacokinetics. *Seizure*. 2004;13:S5-S9.
168. Mimaki T, Suzuki Y, Tagawa T, Karasawa T, Yabuuchi H. Interaction of zonisamide with benzodiazepine and GABA receptors in rat brain. *Med J Osaka Univ*. 1990;39(1-4):13-17.
169. Ondo WG. Zonisamide for essential tremor. *Clin Neuropharmacol*. 2007;30(6):345-349.
170. Zesiewicz TA, Ward CL, Hauser RA, Sanchez-Ramos J, Staffetti JF, Sullivan KL. A double-blind placebo-controlled trial of zonisamide (zonegran) in the treatment of essential tremor. *Mov Disord*. 2007;22(2):279-282.
171. Morita S, Miwa H, Kondo T. Effect of zonisamide on essential tremor: a pilot crossover study in comparison with arotinolol. *Parkinsonism Relat Disord*. 2005;11(2):101-103.
172. Handforth A, Martin FC, Kang GA, Vanek Z. Zonisamide for essential tremor: an evaluator-blinded study. *Mov Disord*. 2009;24(3):437-440.
173. Gören MZ, Onat F. Ethosuximide: from bench to bedside. *CNS Drug Rev*. 2007;13:224-239. doi:10.1111/j.1527-3458.2007.00009.x
174. Gironell A, Marin-Lahoz J. Ethosuximide for essential tremor: an open-label trial. *Tremor Other Hyperkinet Mov (NY)*. 2016;6:378.
175. Papapetropoulos S, Lee MS, Boyer S, Newbold EJ. A phase 2, randomized, double-blind, placebo-controlled trial of CX-8998, a selective modulator of the T-type calcium channel in inadequately treated moderate to severe essential tremor: T-CALM study design and methodology for efficacy endpoint and digital biomarker

- selection. *Front Neurol.* 2019;10:597. doi:[10.3389/fneur.2019.00597](https://doi.org/10.3389/fneur.2019.00597)
176. Papapetropoulos S, Lee MS, Versavel S, et al. A phase 2 proof-of-concept, randomized, placebo-controlled trial of CX-8998 in essential tremor. *Mov Disord.* 2021;36(8):1944-1949. doi:[10.1002/mds.28584](https://doi.org/10.1002/mds.28584)
 177. Olhaye O, Belfort G, Raines S, et al. A phase 2 clinical trial evaluating the efficacy, safety, tolerability, and pharmacokinetics of PRAX-944 in adults with essential tremor [abstract]. *Mov Disord.* 2022;37(suppl 1). Accessed September 21, 2022. <https://www.mdsabstracts.org/abstract/a-phase-2-clinical-trial-evaluating-the-efficacy-safety-tolerability-and-pharmacokinetics-of-prax-944-in-adults-with-essential-tremor/>
 178. Olhaye O, Belfort G, Raines S, et al. A phase 2 randomized, double-blind, placebo-controlled trial of PRAX-944 for the treatment of essential tremor [abstract]. *Mov Disord.* 2022;37(suppl 1). Accessed September 21, 2022. <https://www.mdsabstracts.org/abstract/a-phase-2-randomized-double-blind-placebo-controlled-trial-of-prax-944-for-the-treatment-of-essential-tremor/>
 179. Zhao Y, Huang G, Wu Q, et al. Cryo-EM structures of apo and antagonist-bound human Cav3.1. *Nature.* 2019;576:492-497. doi:[10.1038/s41586-019-1801-3](https://doi.org/10.1038/s41586-019-1801-3)
 180. Váradi C. Clinical features of Parkinson's disease: the evolution of critical symptoms. *Biology (Basel).* 2020;9(5):103. doi:[10.3390/biology9050103](https://doi.org/10.3390/biology9050103)
 181. Bock MA, Brown EG, Zhang L, Tanner C. Association of motor and nonmotor symptoms with health-related quality of life in a large online cohort of people with Parkinson disease. *Neurology.* 2022;98(22):e2194-e2203. doi:[10.1212/WNL.000000000000200113](https://doi.org/10.1212/WNL.000000000000200113)
 182. Braak H, Del Tredici K, Rüb U, et al. Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol Aging.* 2003;24(2):197-211.
 183. Gröger A, Kolb R, Schäfer R, Klose U. Dopamine reduction in the substantia nigra of Parkinson's disease patients confirmed by in vivo magnetic resonance spectroscopic imaging. *PLoS One.* 2014;9(1):e84081.
 184. Gerlach M, Gsell W, Kornhuber J, et al. A post mortem study on neurochemical markers of dopaminergic, GABA-ergic and glutamatergic neurons in basal ganglia-thalamocortical circuits in Parkinson syndrome. *Brain Res.* 1996;741(1-2):142-152.
 185. Hashemiyooun R, Kuhn J, Visser-Vandewalle V. Putting the pieces together in Gilles de la Tourette syndrome: exploring the link between clinical observations and the biological basis of dysfunction. *Brain Topogr.* 2017;30(1):3-29.
 186. Wang L, Zhou C, Cheng W, et al. Dopamine depletion and subcortical dysfunction disrupt cortical synchronization and metastability affecting cognitive function in Parkinson's disease. *Hum Brain Mapp.* 2022;43(5):1598-1610.
 187. Albin RL, Young AB, Penney JB. The functional anatomy of basal ganglia disorders. *Trends Neurosci.* 1989;12(10):366-375. doi:[10.1016/0166-2236\(89\)90074-x](https://doi.org/10.1016/0166-2236(89)90074-x)
 188. Alexander GE, Crutcher MD. Functional architecture of basal ganglia circuits: neural substrates of parallel processing. *Trends Neurosci.* 1990;13(7):266-271. doi:[10.1016/0166-2236\(90\)90107-1](https://doi.org/10.1016/0166-2236(90)90107-1)
 189. DeLong MR. Primate models of movement disorders of basal ganglia origin. *Trends Neurosci.* 1990;13(7):281-285. doi:[10.1016/0166-2236\(90\)90110-v](https://doi.org/10.1016/0166-2236(90)90110-v)
 190. Gao LL, Zhang JR, Chan P, Wu T. Levodopa effect on basal ganglia motor circuit in Parkinson's disease. *CNS Neurosci Ther.* 2017;23(1):76-86.
 191. Thanvi BR, Lo TC. Long term motor complications of levodopa: clinical features, mechanisms, and management strategies. *Postgrad Med J.* 2004;80(946):452-458. doi:[10.1136/pgmj.2003.013912](https://doi.org/10.1136/pgmj.2003.013912)
 192. Banerjee AK, Falkai PG, Savidge M. Visual hallucinations in the elderly associated with the use of levodopa. *Postgrad Med J.* 1989;65(764):358-361. doi:[10.1136/pgmj.65.764.358](https://doi.org/10.1136/pgmj.65.764.358)
 193. Molina JA, Sáinz-Artiga MJ, Fraile A, et al. Pathologic gambling in Parkinson's disease: a behavioral manifestation of pharmacologic treatment? *Mov Disord.* 2000;15(5):869-872. doi:[10.1002/1531-8257\(200009\)15:5<869::aid-mds1016>3.0.co;2-i](https://doi.org/10.1002/1531-8257(200009)15:5<869::aid-mds1016>3.0.co;2-i)
 194. Weiss HD, Marsh L. Impulse control disorders and compulsive behaviors associated with dopaminergic therapies in Parkinson disease. *Neurol Clin Pract.* 2012;2(4):267-274. doi:[10.1212/CPJ.0b013e318278be9b](https://doi.org/10.1212/CPJ.0b013e318278be9b)
 195. Wong JK, Viswanathan VT, Nozile-Firth KS, et al. STN versus GPi deep brain stimulation for action and rest tremor in Parkinson's disease. *Front Hum Neurosci.* 2020;14:578615.
 196. Parihar R, Alterman R, Papavassiliou E, Tarsy D, Shih LC. Comparison of VIM and STN DBS for parkinsonian resting and postural/action tremor. *Tremor Other Hyperkinet Mov (NY).* 2015;5:321.
 197. Katlowitz K, Ko M, Mogilner AY, Pourfar M. Effect of deep brain stimulation on arm, leg, and chin tremor in Parkinson disease. *J Neurosurg.* 2018;131:1-6. doi:[10.3171/2018.7.JNS18784](https://doi.org/10.3171/2018.7.JNS18784)
 198. Hacker ML, DeLong MR, Turchan M, et al. Effects of deep brain stimulation on rest tremor progression in early stage Parkinson disease. *Neurology.* 2018;91(5):e463-e471.
 199. Lee DJ, Dallapiazza RF, De Vloo P, Lozano AM. Current surgical treatments for Parkinson's disease and potential therapeutic targets. *Neural Regen Res.* 2018;13(8):1342-1345.
 200. Kumar R, Lozano AM, Kim YJ, et al. Double-blind evaluation of subthalamic nucleus deep brain stimulation in advanced Parkinson's disease. *Neurology.* 1998;51(3):850-855.

201. Dallapiazza RF, De Vloo P, Fomenko A, et al. Considerations for patient and target selection in deep brain stimulation surgery for Parkinson's disease. In: Stoker TB, Greenland JC, eds. *Parkinson's Disease: Pathogenesis and Clinical Aspects*. Codon Publications; 2018:8.
202. Hutchison WD, Allan RJ, Opitz H, et al. Neurophysiological identification of the subthalamic nucleus in surgery for Parkinson's disease. *Ann Neurol*. 1998;44(4):622-628.
203. Benazzouz A, Breit S, Koudsie A, Pollak P, Krack P, Benabid AL. Intraoperative microrecordings of the subthalamic nucleus in Parkinson's disease. *Mov Disord*. 2002;17(Suppl 3):S145-S149.
204. Starr PA, Rau GM, Davis V, et al. Spontaneous pallidal neuronal activity in human dystonia: comparison with Parkinson's disease and normal macaque. *J Neurophysiol*. 2005;93:3165-3176.
205. Wichmann T, Bergman H, DeLong MR. The primate subthalamic nucleus. III. Changes in motor behavior and neuronal activity in the internal pallidum induced by subthalamic inactivation in the MPTP model of parkinsonism. *J Neurophysiol*. 1994;72:521-530.
206. Benazzouz A, Gross C, Féger J, Boraud T, Bioulac B. Reversal of rigidity and improvement in motor performance by subthalamic high-frequency stimulation in MPTP-treated monkeys. *Eur J Neurosci*. 1993;5(4):382-389.
207. Remple MS, Bradenham CH, Kao CC, Charles PD, Neimat JS, Konrad PE. Subthalamic nucleus neuronal firing rate increases with Parkinson's disease progression. *Mov Disord*. 2011;26(9):1657-1662.
208. Steigerwald F, Pötter M, Herzog J, et al. Neuronal activity of the human subthalamic nucleus in the parkinsonian and nonparkinsonian state. *J Neurophysiol*. 2008;100(5):2515-2524. doi:[10.1152/jn.00858.2015](https://doi.org/10.1152/jn.00858.2015)
209. Galvan A, Devergnas A, Pittard D, et al. Lack of antiparkinsonian effects of systemic injections of the specific T-type calcium channel blocker ML218 in MPTP-treated monkeys. *ACS Chem Neurosci*. 2016;7(11):1543-1551. doi:[10.1021/acschemneuro.6b00186](https://doi.org/10.1021/acschemneuro.6b00186)
210. Bichler EK, Cavarretta F, Jaeger D. Changes in excitability properties of ventromedial motor thalamic neurons in 6-OHDA lesioned mice. *eNeuro*. 2021;8(1):ENEURO.0436-20.2021.
211. Walters JR, Hu D, Itoga CA, Parr-Brownlie LC, Bergstrom DA. Phase relationships support a role for coordinated activity in the indirect pathway in organizing slow oscillations in basal ganglia output after loss of dopamine. *Neuroscience*. 2007;144(2):762-776.
212. Magill PJ, Bolam JP, Bevan MD. Dopamine regulates the impact of the cerebral cortex on the subthalamic nucleus-globus pallidus network. *Neuroscience*. 2001;106(2):313-330.
213. Potashkin JA, Blume SR, Runkle NK. Limitations of animal models of Parkinson's disease. *Parkinsons Dis*. 2010;2011:658083.
214. Rodriguez-Oroz MC, Rodriguez M, Guridi J, et al. The subthalamic nucleus in Parkinson's disease: somatotopic organization and physiological characteristics. *Brain*. 2001;124(9):1777-1790. doi:[10.1093/brain/124.9.1777](https://doi.org/10.1093/brain/124.9.1777)
215. Lenz FA, Tasker RR, Kwan HC, et al. Single unit analysis of the human ventral thalamic nuclear group: correlation of thalamic "tremor cells" with the 3-6 Hz component of parkinsonian tremor. *J Neurosci*. 1988;8(3):754-764.
216. Zirh TA, Lenz FA, Reich SG, Dougherty PM. Patterns of bursting occurring in thalamic cells during parkinsonian tremor. *Neuroscience*. 1998;83(1):107-121.
217. Muthuraman M, Raethjen J, Koirala N, et al. Cerebello-cortical network fingerprints differ between essential, Parkinson's and mimicked tremors. *Brain*. 2018;141(6):1770-1781.
218. Sen S, Kawaguchi A, Truong Y, Lewis MM, Huang X. Dynamic changes in cerebello-thalamo-cortical motor circuitry during progression of Parkinson's disease. *Neuroscience*. 2010;166(2):712-719.
219. Helmich RC, Hallett M, Deuschl G, Toni I, Bloem BR. Cerebral causes and consequences of parkinsonian resting tremor: a tale of two circuits? *Brain*. 2012;135(Pt 11):3206-3226.
220. Helmich RC, Janssen MJ, Oyen WJ, et al. Pallidal dysfunction drives a cerebellothalamic circuit into Parkinson tremor. *Ann Neurol*. 2011;69(2):269-281.
221. Salamone JD, Mayorga AJ, Trevitt JT, Cousins MS, Conlan A, Nawab A. Tremulous jaw movements in rats: a model of parkinsonian tremor. *Prog Neurobiol*. 1998;56(6):591-611. doi:[10.1016/s0301-0082\(98\)00053-7](https://doi.org/10.1016/s0301-0082(98)00053-7)
222. Miwa H, Koh J, Kajimoto Y, Kondo T. Effects of T-type calcium channel blockers on a parkinsonian tremor model in rats. *Pharmacol Biochem Behav*. 2011;97(4):656-659.
223. Coulter DA, Huguenard JR, Prince DA. Characterization of ethosuximide reduction of low-threshold calcium current in thalamic neurons. *Ann Neurol*. 1989;25(6):582-593.
224. Gomez-Mancilla B, Latulippe JF, Boucher R, Bédard PJ. Effect of ethosuximide on rest tremor in the MPTP monkey model. *Mov Disord*. 1992;7(2):137-141.
225. Ge W, Ren J. Combined L-/T-type calcium channel blockers: ready for prime time. *Hypertension*. 2009;53(4):592-594. doi:[10.1161/HYPERTENSIONAHA.108.127548](https://doi.org/10.1161/HYPERTENSIONAHA.108.127548)
226. Hagiwara N, Irisawa H, Kameyama M. Contribution of two types of calcium currents to the pacemaker potentials of rabbit sino-atrial node cells. *J Physiol*. 1988;395:233-253. doi:[10.1113/jphysiol.1988.sp016916](https://doi.org/10.1113/jphysiol.1988.sp016916)
227. Hirano Y, Fozzard HA, January CT. Characteristics of L- and T-type Ca^{2+} currents in canine cardiac Purkinje cells. *Am J Physiol*. 1989;256(5 Pt 2):H1478-H1492. doi:[10.1152/ajpheart.1989.256.5.H1478](https://doi.org/10.1152/ajpheart.1989.256.5.H1478)

228. Tseng GN, Boyden PA. Multiple types of Ca^{2+} currents in single canine Purkinje cells. *Circ Res.* 1989;65(6):1735-1750. doi:[10.1161/01.res.65.6.1735](https://doi.org/10.1161/01.res.65.6.1735)
229. Bohn G, Moosmang S, Conrad H, Ludwig A, Hofmann F, Klugbauer N. Expression of T- and L-type calcium channel mRNA in murine sinoatrial node. *FEBS Lett.* 2000;481(1):73-76. doi:[10.1016/s0014-5793\(00\)01979-7](https://doi.org/10.1016/s0014-5793(00)01979-7)

Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Figure S1