

SUSPECTED TOXICOLOGICAL EFFECTS OF PER- AND POLYFLUOROALKYL SUBSTANCES (PFAS) ON THE ENVIRONMENT AND HUMAN HEALTH

Snejana Dineva

Trakia University of Stara Zagora, Faculty of Technics and Technologies of Yambol,
Department of Food Technology,

Graf Ignatiev str. 38, Yambol 8602, Bulgaria

mobile: +359898999457, e-mail: snezhana.dineva@trakia-uni.bg

Abstract: PFAS stands for per- and polyfluoroalkyl substances, which are a group of chemicals that are commonly used in industrial and commercial production. These chemicals have been found to have significant toxicological effects on both humans and wildlife, meaning they can cause harm to living organisms. The toxicity of PFAS mixtures, which are the combinations of different PFAS chemicals that can be found in the environment, is not yet fully understood. In recent decades, there has been an increasing amount of information discovered about chemical organic pollution, which includes thousands of human-made substances that are widely used in various aspects of daily life and are emitted from many industrial processes. The purpose of the article is to review the toxicity of PFAS chemicals, their behavior and existence in the environment, their tendency to accumulate in living organisms, and the potential negative health effects they can have on the biota, which refers to all living organisms in a particular area or ecosystem.

Keywords: per- and polyfluoroalkyl substances (PFAS), perfluoroalkyl acids (PFAA), environmental pollution, human health.

1. INTRODUCTION

Per- and polyfluoroalkyl substances (PFAS) are organic fluorinated chemicals, highly persistent, synthetic contaminants with tremendous health impacts on human populations such as obesity, various cancers, elevated cholesterol levels, liver damage, decreased immune and liver functionalities, and significant birth defects (Jha et al. 2021; Dickman & Aga 2022). PFASs can be referred also as PFCs (per- and polyfluorinated chemicals, or perfluorocarbons, more than 5000 substances (USEPA 2021), perfluoroalkyl acids (PFAA), perfluoroalkyl carboxylic acids (PFCA), and perfluoroalkane sulfonic acids (PFSA). Limited toxicological data are available for most PFAS yet (McCarthy et al. 2021). PFAAs exert human toxicity, high persistence, and environmental mobility, widespread presence in food of plant and animal origin, and poor removability from contaminated matrices (Sznajder-Katarzyńska 2019; Langenbach & Wilson 2021; Lyu et al. 2022; Harvard News 2023).

The purpose of the article is to review the toxicity of per- and polyfluoroalkyl substances, environmental behavior, bioaccumulation, and health effects.

2. METHODS

The review is made by searching open data sources of Google, Google Scholar, PubMed, Taylor & Francis, and others. The searches were conducted using the title of the article, titles of main sectors included in the paper, and different keywords. The articles were chosen by interest and data of publication, usually looking for the last years of publication. The keywords used in the search were numerous: "PFAS toxicity", "PFAS in soil", etc.

3. RESULTS

3.1. PFAS into the environment

The majority PFASs emissions are released into the aquatic environment mainly from point sources such as manufacturing plants and firefighting activities using PFAS-containing foams (Koch et al. 2019). PFAS are released to air from a variety of sources, travel significant distances in air, depending on the physico-chemical properties, registered in Arctic, which demonstrates their long-range atmospheric transport (USEPA 2021).

Surface deposition of atmospheric emissions is followed by leaching to groundwater (de Silva et al., 2021; Jha et al. 2021). PFAS are identified in all parts of biosphere, on surface of the water and groundwater, in the rain, snow, and drinking water (Houtz et al. 2013; Hu et al. 2016; Guardian et al. 2020; Sun et al. 2016; Gobelius et al. 2019; Abunada et al. 2020; Barzen-Hanson et al. 2017; Kaserzon et al. 2019; Abunada et al. 2020; Xu et al. 2021), manufactured goods like cosmetics (Vestergren et al. 2013; Mousavi et al. 2021), food packaging (Trier et al. 2011; Zabaleta et al. 2016), agricultural foodstuffs (Vestergren et al. 2013; Blaine et al. 2013; 2014; Genualdi et al. 2017).

Once into the environment, terrestrial or aquatic, PFAS persist for long periods due to their resistance, the fate of the PFAS in different environmental conditions provides information for the mechanisms of degradation (fig.1; Dickman & Aga 2022).




Figure 1. Fate, effects, and treatment of per- and polyfluoroalkyl substances (PFAS). Source: Dickman & Aga 2022. Open access, used with permission

PFAS consequences depend on carbon chain lengths, head groups, ether bonds, ionic state, and environmental pH, chain fluorination degree, which determines sorption and bioaccumulation, volatilization, and the potential for transformation (Dickman & Aga 2022). PFAS easily undergo bioaccumulation due to their long persistence times (Kurwadkar et al. 2022), and provoke negative health effects on humans and wildlife (Dickman & Aga 2022). Long chain PFAS are more bioaccumulative (Koch et al. 2019).

PFAS are ubiquitous and have negative health effects, leading to monitoring in drinking water, food products, and the environment, with regulatory focus on drinking water and food products, but they exist everywhere in soils, water, plants, livestock, and dairy products, and human exposure comes from various sources (fig.2; Danish Environmental Protection Agency Report, 2015; Ojo et al., 2021; Fenton et al., 2021; Jha et al. 2021; Yadav et al. 2022).



Figure 2. Sources of per-and poly-fluoroalkyl substances (PFAS; Yadav et al. 2022).

Source: Yadav et al. 2022. Used with permission

3.2. PFAS toxicity

PFAS are present in a variety of mixtures in the environment, and their interactions with other chemicals can affect their toxicity. The most widely studied PFAS are PFOA and PFOS, which have been found to have toxic effects on humans and wildlife (Kang 2019; Dickman & Aga 2022). PFOA and PFOS are persistent in the environment, meaning that they do not break down easily and can accumulate in the food chain (Koch et al. 2019; Kurwadkar et al. 2022). Studies have shown that exposure to PFOA and PFOS can lead to a range of health effects, including developmental and reproductive problems, immune system dysfunction, and cancer (ATSDR 2021; Jha et al. 2021). Despite the widespread use and potential health effects of PFAS, there is still much to learn about their toxicity and how they interact with other chemicals in the environment (Lerch et al. 2023).

The concentration of pollutants, such as PFAAs, in fish is a result of two processes: uptake from the aqueous phase and intake of contaminated food (aquatic organisms). Uptake from the aqueous phase refers to the process by which fish absorb pollutants directly from the water they swim in. This can occur through their gills or skin. Intake of contaminated food refers to the process by which fish consume other organisms that have already accumulated pollutants in their tissues. This can include smaller fish, crustaceans, and other aquatic organisms. The combination of these two processes can result in high concentrations of pollutants in fish, which can then be passed up the food chain to fish-eating birds or mammals (Navarro et al. 2020). It is important to consider the entire food chain when assessing the ecological risk of pollutants in aquatic ecosystems, as pollutants can accumulate and magnify as they move up the chain. The effects of two types of per- and polyfluoroalkyl substances (PFAS), namely PFOA and PFOS, were investigated on a type of fish called Medaka (*Oryzias latipes*). The study found that PFOA and PFOS induced different effects on the fish. PFAS affect the endocrine system of fish, which can have implications for their reproductive health and the health of the ecosystems they inhabit (Kang 2019).

3.3. Human health effects of PFAS

Humans can be exposed to PFAS from air, water, food or food contact materials (FCMs). Inhalation of PFAS can occur when PFAS-containing products are used indoors or when dust containing PFAS settles on surfaces (De la Torre et al. 2019; Zhang et al. 2020). Direct contact with PFAS-containing products are also sources of toxicity, when products such as non-stick cookware or waterproof clothing are used (Trudel et al. 2008), or cosmetics (Mousavi et al.

2021). Ingestion of contaminated water occur when PFAS leach into groundwater or surface water from sources such as firefighting foam or landfills (Domingo & Nadal 2019). PFAS can migrate from food contact materials into food, contributing to human dietary exposure and risk assessment procedures should consider this pathway (Sunderland 2018; Susmann et al. 2019).

Exposure to PFAS can lead to various health problems including birth defects, developmental delays, elevated cholesterol levels, increased risk of heart disease and stroke, various cancers, decreased immune and liver functionalities (fig.3; ATSDR 2021; Beans 2021; Dickman & Aga 2022).




Figure 3. Effects of per- and polyfluoroalkyl substances on human health.
Used with permission from the European Environment Agency (2019)

PFAS exposure increased osteoporosis risk, kidney dysfunction, hormonal disregulation and metabolism, most frequently thyroid hormones (Jha et al. 2021). PFAS route exposure in humans is intake with food and water consumption (Sznajder-Katarzyńska et al. 2019). An FDA Total Diet Survey (TDS) confirmed that at least one type of PFAS was detected in 17 TDS samples, 14 of the samples were seafood, representing 44% (14 of 32) of TDS seafood samples. In a targeted survey of seafood in 2022, PFASs were found in 74% (60 of 81) of samples of shellfish, cod, crab, pollock, salmon, shrimp, tilapia and tuna. Data on PFAS in seafood is still limited, but testing suggests that it is likely at higher risk of environmental contamination with PFAS than other types of food (FDA 2023).

PFAS from food contact materials can contribute to dietary exposure and impact human health, but unlike other persistent organic pollutants, they do not bioaccumulate in adipose tissue and primarily distribute to the liver, blood serum, and kidney (Bischel et al. 2011; Lau 2012, 2015; Kato et al. 2015; Lerch et al. 2023). PFAS are present in 99% of the human population and have been linked to adverse health effects, prompting a national health study by ATSDR (CDC 2022; Langenbach & Wilson 2021; ATSDR 2020).

4. CONCLUSION

Overall, the accumulation of PFAS in the human body can have serious health consequences, and it is important to limit exposure to these chemicals as much as possible. This can be done by avoiding products that contain PFAS, such as non-stick cookware and stain-resistant fabrics, and by using alternative products that are safer for human health and the environment.

5. REFERENCES

- [1] Abunada Z., M.Y.D. Alazaiza, M. J.K. Bashir (2020). An Overview of Per- and Polyfluoroalkyl Substances(PFAS) in the Environment: Source, Fate, Riskand Regulations. *Water*. 2020. 12, 3590; doi: 10.3390/w12123590 www.mdpi.com/journal/water.
- [2] ATSDR (2020). Multi-site Health Study – PFAS Cooperative Agreement. Last Modified July 22, 2020. <https://www.atsdr.cdc.gov/pfas/activities/studies/multi-site.html>.
- [3] ATSDR (2021). Toxicological Profile for Perfluoroalkyls: Agency for Toxic Substances and Disease Registry; 2021.
- [4] Barzen-Hanson K.A., S.C. Roberts, S. Choyke, K. Oetjen, A. McAlees, N. Riddell, R. McCrindle, P.L. Ferguson, C.P. Higgins, J.A. (2017). Field Discovery of 40 classes of per- and polyfluoroalkyl substances in historical aqueous film-forming foams (AFFFs) and afff-impacted groundwater. *Environ. Sci. Technol.*, 51 (4) (2017), pp. 2047-2057.
- [5] Beans C. (2021). How “forever chemicals” might impair the immune system. *Proceedings of the National Academy of Sciences*, 118(15), e2105018118. <https://doi.org/10.1073/pnas.2105018118>.
- [6] Bischel H.N., L.A. Macmanus-Spencer, C. Zhang, R.G. Luthy (2011). Strong associations of short-chain perfluoroalkyl acids with serum albumin and investigation of binding mechanisms. *Environ Toxicol Chem* 30 (11): 2423-30. <https://doi.org/10.1002/etc.647>.
- [7] Blaine A.C. , C.D. Rich, L.S. Hundal, C. Lau, M.A. Mills, K.M. Harris, C.P. Higgins (2013). Uptake of perfluoroalkyl acids into edible crops via land applied biosolids: field and greenhouse studies. *Environ. Sci. Technol.*, 47 (24) (2013), pp. 14062-14069.
- [8] Blaine A.C., C.D. Rich, E.M. Sedlacko, L.S. Hundal, K. Kumar, C. Lau, M.A. Mills, K.M. Harris, C.P. Higgins (2014). Higgins Perfluoroalkyl acid distribution in various plant compartments of edible crops grown in biosolids-amended soils. *Environ. Sci. Technol.*, 48 (14) (2014), pp. 7858-7865.
- [9] CDC (2022). National Health and Nutrition Examination Survey. 2022. <https://www.cdc.gov/nchs/nhanes/index.htm>.
- [10] Christensen Jen (2023). EPA proposes first standards to make drinking water safer from ‘forever chemicals’. <https://edition.cnn.com/2023/03/14/health/epa-pfas-standards-wellness/index.html>. [March 15, 2023].
- [11] Danish Environmental Protection Agency Report (2015): Short-Chain Polyfluoroalkyl Substances (PFAS). A Literature Review of Information on Human Health Effects and Environmental Fate and Effect Aspects of Short-Chain PFAS. (Environmental Project No. 1707, 2015).
- [12] De la Torre A., I. Navarro, P. Sanz, M. de los Á. Martínez (2019). Occurrence and human exposure assessment of perfluorinated substances in house dust from three European countries *Science of the Total Environment*, 685 (2019), pp. 308-314, 10.1016/j.scitotenv.2019.05.463.
- [13] De Silva A., J.M. Armitage, A.A. Bruton, C. Dassuncao, W. Heigr-Bernays, X.C. Hu, A. Kärman, B. Kelly, C. Ng, A. Robuck, M. Sun, T.F. Webster, E.M. Sunderland (2021). PFAS exposure pathways for humans and wildlife: a synthesis of current knowledge and gaps in understanding. *Environmental Toxicology and Chemistry* 40 (3) pp 631-651.

- [14] Dickman R.A., Aga D.S. (2022). A review of recent studies on toxicity, sequestration, and degradation of per- and polyfluoroalkyl substances (PFAS). *Journal of Hazardous Materials*, 436, 129120. <https://doi.org/10.1016/j.jhazmat.2022.129120>.
- [15] Domingo J.L., M. Nadal (2019). Human exposure to per- and polyfluoroalkyl substances (PFAS) through drinking water: A review of the recent scientific literature. *Environmental Research*, 177 (July) (2019), 10.1016/j.envres.2019.108648.
- [16] Ebnesajjad S. (2013). *Introduction to Fluoropolymers Materials, Technology and Applications*. ISBN: 9781455774425. eBook ISBN: 9781455775514. [May 11, 2013].
- [17] FDA (2023). *Authorized Uses of PFAS in Food Contact Applications*. <https://www.fda.gov/food/process-contaminants-food/authorized-uses-pfas-food-contact-applications>. [05/31/2023].
- [18] Fenton S.E., A. Ducatman, A. Boobis, J.C. DeWitt, C. Lau, C. Ng, J.S. Smith, S.M. Roberts (2021). Per- and polyfluoroalkyl substance toxicity and human health review: current state of knowledge and strategies for informing future research. *Environ. Toxicol. Chem.*, 40 (3) (2021), pp. 606-630.
- [19] Genualdi S., N. Jeong, L. deJager, T. Begley (2017). Investigation into perfluoroalkyl substances (PFASs) in a cranberry bog: method development and sampling results. *Food Addit. Contam.: Part A*, 34 (12) (2017), pp. 2181-2189.
- [20] Gobelius L., C. Persson, K. Wiberg, L. Ahrens (2019). Calibration and application of passive sampling for per- and polyfluoroalkyl substances in a drinking water treatment plant. *J. Hazard. Mater.* 362 (2019), pp. 230-237.
- [21] Guardian M.G.E., E.G. Boongaling, V.R.R. Bernardo-Boongaling, J.Gamonchuang, T. Boontongto, R. Burakham, P. Arnnok, D.S. Aga (2020). Prevalence of per- and polyfluoroalkyl substances (PFASs) in drinking and source water from Two Asian Countries *Chemosphere*, 256 (2020), Article 127115.
- [22] Harvard News (2023). Protecting against 'forever chemicals'. <https://www.hsph.harvard.edu/news/hsph-in-the-news/protecting-against-forever-chemicals/>. [16.03.2023].
- [23] Houtz E.F., C.P. Higgins, J.A. Field, D.L. Sedlak (2013). Persistence of perfluoroalkyl acid precursors in aff-impacted groundwater and soil. *Environ. Sci. Technol.*, 47 (15) (2013), pp. 8187-8195.
- [24] Hu X.C., D.Q. Andrews, A.B. Lindstrom, T.A. Bruton, L.A. Schaider, P. Grandjean, R. Lohmann, C.C. Carignan, A. Blum, S.A. Balan, C.P. Higgins, E.M. Sunderland (2016). Detection of poly- and perfluoroalkyl substances (PFASs) in U.S. Drinking water linked to industrial sites, military fire training areas, and wastewater treatment plants. *Environ. Sci. Technol. Lett.*, 3 (10) (2016), pp. 344-350.
- [25] Jha G., Kankarla V., McLennon E., Pal S., Sihi D., Dari B., Diaz D., Nocco M. (2021). Per- and Polyfluoroalkyl Substances (PFAS) in Integrated Crop–Livestock Systems: Environmental Exposure and Human Health Risks. *International Journal of Environmental Research and Public Health*, 18(23). <https://doi.org/10.3390/ijerph182312550>.
- [26] Kang J.S., T.-G. Ahn, J.-W. Park (2019). Perfluorooctanoic acid (PFOA) and perfluooctane sulfonate (PFOS) induce different modes of action in reproduction to Japanese Medaka (*Oryzias latipes*) *J. Hazard. Mater.*, 368 (2019), pp. 97-103.
- [27] Kaserzon S.L., S. Vijayasathay, J. Bräunig, L. Mueller, D.W. Hawker, K.V. Thomas, J.F. Mueller (2019). Calibration and validation of a novel passive sampling device for the time integrative monitoring of per- and polyfluoroalkyl substances (PFASs) and precursors in contaminated groundwater. *J. Hazard. Mater.*, 366 (2019), pp. 423-431.
- [28] Kato H., S. Fujii, M. Takahashi, M. Matsumoto, M. Hirata-Koizumi, A. Ono, A. Hirose. (2015). Repeated dose and reproductive/developmental toxicity of perfluorododecanoic acid in rats. *Environ Toxicol* 30 (11):1244-63. <https://doi.org/10.1002/tox.21996>.
- [29] Koch A., Kärman A., L.W.Y. Yeung, M. Jonsson, L. Ahrens, T. Wang (2019). Point source characterization of per- and polyfluoroalkyl substances (PFASs) and extractable

- organofluorine (EOF) in freshwater and aquatic invertebrates. *Environ. Sci.: Processes Impacts*, 2019, 21, 1887–1898.
- [30] Kurwadkar S., D.J. Kanel, S.R. Nadagouda, M.N. Cawdrey, R.W. Ambade, B. Struckhoff G.C., Wilkin R. (2022). Per- and polyfluoroalkyl substances in water and wastewater: A critical review of their global occurrence and distribution. *Science of the Total Environment*, 809, 151003. <https://doi.org/10.1016/j.scitotenv.2021.151003>.
- [31] Langenbach B., Wilson M. (2021). Per- and Polyfluoroalkyl Substances (PFAS): Significance and Considerations within the Regulatory Framework of the USA. *International Journal of Environmental Research and Public Health*, 18(21), 11142. <https://doi.org/10.3390/ijerph182111142>.
- [32] Lau C. (2012). Perfluorinated Compounds. In *Molecular, Clinical and Environmental Toxicology*, edited by A. Luch, 47-86. Vol. *Experientia Supplementum*, 101. Basel: Springer. ISBN No. 978-3-7643-8340-4.
- [33] Lau C. (2015). Perfluorinated Compounds: An Overview. In *Toxicological Effects of Perfluoroalkyl and Polyfluoroalkyl Substances*, edited by J.D. DeWitt. Humana Press. ISBN No. 978-3-319-15517-3.
- [34] Lerch, M., Fengler, R., Mbog, G., Nguyen, K. H., & Granby, K. (2023). Food simulants and real food – What do we know about the migration of PFAS from paper based food contact materials? *Food Packaging and Shelf Life*, 35, 100992. <https://doi.org/10.1016/j.fpsl.2022.100992>.
- [35] Lyu X., Xiao F., Shen C., Chen J., Park C. M., Sun Y., Flury M., Wang D. (2022). Per- and Polyfluoroalkyl Substances (PFAS) in Subsurface Environments: Occurrence, Fate, Transport, and Research Prospect. *Reviews of Geophysics*, 60(3), e2021RG000765. <https://doi.org/10.1029/2021RG000765>.
- [36] McCarthy, C. J., Roark, S. A., & Middleton, E. T. (2021). Considerations for toxicity experiments and risk assessments with PFAS mixtures. *Integrated Environmental Assessment and Management*, 17(4), 697-704. <https://doi.org/10.1002/ieam.4415>.
- [37] Mousavi S.E., J.M. Delgado-Saborit, L. Godderis (2021). Exposure to per- and polyfluoroalkyl substances and premature skin aging. *J. Hazard. Mater.*, 405 (2021), Article 124256.
- [38] Navarro, I., de la Torre, A., Sanz, P., & Martínez, M. D. L. Á. (2020). Perfluoroalkyl acids (PFAAs): Distribution, trends and aquatic ecological risk assessment in surface water from Tagus River basin (Spain). *Environmental Pollution*, 256, 113511. <https://doi.org/10.1016/j.envpol.2019.113511>.
- [39] Ojo A.F., C. Peng, J.C. Ng (2021). Assessing the human health risks of per- and polyfluoroalkyl substances: a need for greater focus on their interactions as mixtures. *J. Hazard. Mater.*, 407 (2021), Article 124863.
- [40] Silva J.A. K., W.A. Martin, J.L. Johnson, J.E. McCray (2019). Evaluating air-water and NAPL-water interfacial adsorption and retention of Perfluorocarboxylic acids within the Vadose zone. *Journal of Contaminant Hydrology*. 223:103472. <https://doi.org/10.1016/j.jconhyd.2019.03.004>.
- [41] Sun M., E. Arevalo, M. Strynar, A. Lindstrom, M. Richardson, B. Kearns, A. Pickett, C. Smith, D.R.U. Knappe (2016). Legacy and emerging perfluoroalkyl substances are important drinking water contaminants in the cape fear river watershed of North Carolina. *Environ. Sci. Technol. Lett.*, 3 (12) (2016), pp. 415-419.
- [42] Sunderland E.M. (2018). A review of the pathways of human exposure to poly- and perfluoroalkyl substances (PFASs) and present understanding of health effects. *Physiology & Behavior*, 176 (5) (2018), pp. 139-148, 10.1038/s41370-018-0094-1.A.
- [43] Susmann H.P., L.A. Schaidler, K.M. Rodgers, R.A. Rudel (2019). Dietary habits related to food packaging and population exposure to PFASs. *Environmental Health Perspectives*, 127 (10) (2019), 10.1289/EHP4092.

- [44] Sznajder-Katarzyńska K., M. Surma, I. Cieřlik (2019). A Review of Perfluoroalkyl Acids (PFAAs) in terms of Sources, Applications, Human Exposure, Dietary Intake, Toxicity, Legal Regulation, and Methods of Determination", *Journal of Chemistry*, vol. 2019, Article ID 2717528, 20 pages, 2019. <https://doi.org/10.1155/2019/2717528>.
- [45] Trier X., K. Granby, J.H. Christensen 2011. Polyfluorinated Surfactants (Pfs) in paper and board coatings for food packaging. *Environ. Sci. Pollut. Res. Int.*, 18 (7) (2011), pp. 1108-1120.
- [46] Trudel D., Horowitz L., Wormuth M., Scheringer M., Cousins I. T., Hungerbühler K. (2008). Estimating Consumer Exposure to PFOS and PFOA. *Risk Analysis*, 28(2), 251-269. <https://doi.org/10.1111/j.1539-6924.2008.01017.x>.
- [47] Uptake of perfluoroalkyl acids into edible crops via land applied biosolids: field and greenhouse studies. *Environ. Sci. Technol.*, 47 (24) (2013), pp. 14062-14069.
- [48] USEPA (2021). PFAS Master List of PFAS Substances (Version 2). 2017. Retrieved May 14, 2021, from https://comptox.epa.gov/dashboard/chemical_lists/pfasmaster.
- [49] Vestergren R., F. Orata, U. Berger, I.T. Cousins (2013). Bioaccumulation of perfluoroalkyl acids in dairy cows in a naturally contaminated environment. *Environ. Sci. Pollut. Res.*, 20 (11) (2013), pp. 7959-7969.
- [50] Xu B., S. Liu, J.L. Zhou, C. Zheng, J. Weifeng, B. Chen, T. Zhang, W. Qiu (2021). PFAS and their substitutes in groundwater: occurrence, transformation and remediation. *J. Hazard. Mater.*, 412 (2021), Article 125159.
- [51] Yadav S., Ibrar I., Al-Juboori R. A., Singh L., Ganbat N., Kazwini T., Karbassiyazdi, E., Samal A. K., Subbiah S., Altaee A. (2022). Updated review on emerging technologies for PFAS contaminated water treatment. *Chemical Engineering Research and Design*, 182, 667-700. <https://doi.org/10.1016/j.cherd.2022.04.009>.
- [52] Zabaleta I., E. Bizkarguenaga, D. Bilbao, N. Etxebarria, A. Prieto, O. Zuloaga (2016). Fast and simple determination of perfluorinated compounds and their potential precursors in different packaging materials. *Talanta*, 152 (Supplement C) (2016), pp. 353-363.
- [53] Zhang Bo, Yuan He, Yingyan Huang, Danhong Hong, Yiming Yao, Lei Wang, Wenwen Sun, Baoqin Yang, Xiongfei Huang, Shiming Song, Xueyuan Bai, Yuankai Guo, Tao Zhang, Hongwen Sun (2020). Novel and legacy poly- and perfluoroalkyl substances (PFASs) in indoor dust from urban, industrial, and e-waste dismantling areas: The emergence of PFAS alternatives in China *Environmental Pollution*, 263 (2020), Article 114461, [10.1016/j.envpol.2020.114461](https://doi.org/10.1016/j.envpol.2020.114461).