

[Lecture-7]

Q. Unsupervised Learning

→ A type of learning where the model works with the unlabeled data to discover patterns.

Example: Clustering (e.g. k-means)

Dimensionality (e.g. PCA)

Clustering: It is a way of grouping the data points into different clusters, consisting of a similar data points.

Q1.

House size (x_i)	Rent (\$1000) ($y_i - \hat{y}_i$)
0.0	1.5
0.4	2.0
0.7	2.5
1.0	3.0

Either error tolerance or no. of steps will be given in the question.

Ans:

We know that,

$$w = w - \alpha \left[\frac{2}{n} \sum_{i=1}^n x_i (y_i - \hat{y}_i) \right]$$

$$b = b - \alpha \left[\frac{2}{n} \sum_{i=1}^n (y_i - \hat{y}_i) \right]$$

$$\text{Cost function MSE} = \left[\frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2 \right]$$

Line for linear regression, $\hat{y} = wx + b$

[Lecture-3] Supervised Learning

Q. Linear Regression!

→ Linear Regression is a fundamental statistical method for predicting continuous values. It models the relationships between the dependent (labels) and independent variables (features) using a straight line. A model with dependent variable is called multivariate linear regression. A model with independent variable is called multiple linear regression.

Q. Linear Regression Algorithm:

- i. Initialize weight (w), bias (b) and learning rate (α)
- ii. Compute the predicted value, \hat{y} .
- iii. Compute the loss using MSE, J .
- iv. Update the weight and bias using Gradient Descent.
$$w = w - \alpha \frac{\partial J}{\partial w}, \quad b = b - \alpha \frac{\partial J}{\partial b}$$
- v. Repeat until convergence.

Q. Polynomial Regression

→ Describes the relationship between the independent and dependent variable (x, y) using an n^{th} -degree polynomial in x .

Types: Linear (degree[1]), Quadratic (degree[2]), Cubic (degree[3+])

[Lecture-4]

Q. Gradient:

→ Gradient of a scalar function is a vector that points in the direction of the greatest rate of increase of the function.

Algorithm:

- i. Pick an initial point x_0
- ii. Gradient Descent Update Rule
$$x_{t+1} = x_t - \alpha \nabla f(x_t)$$
- iii. Iterate until convergence
- iv. $-\nabla f(x_t)$ moves in the negative direction of the gradient to minimize $f(x)$.

where,

x_t = current position at iteration t

α = step size

$\nabla f(x_t)$ = gradient (first derivative) of the function $f(x)$

1st iteration!

here, $w = 0$, $b = 0$, $\alpha = 0.2$

i	$\hat{y}_i = wx + b$	$y_i - \hat{y}_i$	$(y_i - \hat{y}_i)^2$	$x_i(y_i - \hat{y}_i)$
1	$0 \times 0.0 + 0 = 0$	1.5	2.25	$0.00 \times 1.5 = 0.00$
2	0	2.0	4.00	$0.4 \times 2.0 = 0.80$
3	0	2.5	6.25	$0.7 \times 2.5 = 1.75$
4	0	3.0	9.00	$1.0 \times 3.0 = 3.00$
$n=4$	—	$\sum y_i - \hat{y}_i = 9$	$\sum (y_i - \hat{y}_i)^2 = 21.5$	$\sum x_i(y_i - \hat{y}_i) = 5.55$

$$\begin{aligned} \text{So, } w &= w - \alpha \left[-\frac{2}{n} \sum x_i(y_i - \hat{y}_i) \right] \\ &= 0 - 0.2 \left[-\frac{2}{4} \times 5.55 \right] \\ &= 0.555 \end{aligned}$$

$$\begin{aligned} b &= b - \alpha \left[-\frac{2}{n} \sum (y_i - \hat{y}_i) \right] \\ &= 0 - 0.2 \left(-\frac{2}{4} \times 9 \right) \\ &= 0.90 \end{aligned}$$

$$\begin{aligned} \text{MSE} &= \frac{1}{n} \sum (y_i - \hat{y}_i)^2 \\ &= \frac{1}{4} \times 21.5 \\ &= 5.375 \end{aligned}$$

2nd iteration:

Now $W = 0.555$ and $b = 0.90$ and $\alpha = 0.2$

i	$\hat{y} = Wx + b$	$(y_i - \hat{y}_i)$	$(y_i - \hat{y}_i)^2$	$x_i(y_i - \hat{y}_i)^2$
1	$0.555 \times 0.04 + 0.9 = 0.900$	$1.5 - 0.9 = 0.6$	$(0.6)^2 = 0.36$	0
2	$0.555 \times 0.9 + 0.9 = 1.122$	$2.0 - 1.122 = 0.878$	$(0.878)^2 = 0.771$	0.3512
3	$0.555 \times 0.7 + 0.9 = 1.288$	$2.5 - 1.288 = 1.212$	$(1.212)^2 = 1.468$	0.848
4	$0.555 \times 1.0 + 0.9 = 1.455$	$3.0 - 1.455 = 1.545$	2.387	1.545
		4.235	4.986	2.744

$$\therefore W = 0.555 - 0.2 \times \left[-\frac{2}{4} \times 2.744 \right] = 0.8294$$

$$\therefore b = 0.90 - 0.2 \times \left[-\frac{2}{4} \times 4.235 \right] = 1.3235$$

$$\therefore \text{MSE} = \frac{1}{4} \times 4.986 = 1.2465$$

$$\therefore \hat{y} = Wx + b$$

$$= 0.8294x + 1.3235$$

(Ans.)

Q2.

$$f(x) = x^2 \Rightarrow 2x$$

$$\therefore \nabla f(x) = 2x$$

initial, $x_0 = -4$

step size, $\alpha = 0.8$

Ans!

we know that,

$$x_{t+1} = x_t - \alpha \nabla f(x_t)$$

$$\therefore x_1 = -4 - 0.8 \times (2 \times -4) = 2.4$$

$$y = x^2 = 2.4 \times 2.4 = 5.76 \quad \therefore (x_1, y_1) = (2.4, 5.76)$$

$$\therefore x_2 = 2.4 - 0.8 \times (2 \times 2.4) = -1.44$$

$$y = 2.07 \quad \therefore (x_2, y_2) = (-1.44, 2.07)$$

$$\therefore x_3 = -1.44 - \{0.8 \times 2 \times (-1.44)\} = 0.864$$

$$y = (0.864)^2 = 0.746 \quad \therefore (x_3, y_3) = (0.864, 0.746)$$

Q.

Linear	Polynomial	Logistic
1. Predicts continuous outcomes, assumes a straight relationship between the independent and dependent variables	1. Extends linear regression to model non-linear relationships using polynomial terms of the independent variables	1. Predicts probabilities for binary outcomes, using the logistic function to squeeze the output between 0 and 1. It is used for classification rather than regression

Q. K means clustering

→ It is a method for grouping n observations into K clusters.

Algorithm:

- i. Choose the number of clusters K .
- ii. Select K random points from the data as centroids.
- iii. Assign all the points to the closest cluster centroids.
- iv. Recompute the centroids of newly formed clusters.
- v. Repeat steps 3 and 4.

Q. point (x, y) A(1,2) B(2,3) C(3,3) D(6,7) E(7,8) F(8,8)

Ans:

Let, $K = 2$ [means divide the dataset into two clusters]

Centroid 1 (C_1) = A(1,2)

Centroid 2 (C_2) = E(7,8)

By using Manhattan distance:

	C_1	C_2	Decision
A	$ 1-1 + 2-2 = 0$	$ 7-1 + 8-2 = 12$	C_1
B	$ 1-2 + 2-3 = 2$	$ 7-2 + 8-3 = 10$	C_1
C	$ 1-3 + 2-3 = 3$	$ 7-3 + 8-3 = 9$	C_1
D	$ 1-6 + 2-7 = 9$	$ 7-6 + 8-7 = 2$	C_2
E	$ 1-7 + 2-8 = 12$	$ 7-7 + 8-8 = 0$	C_2
F	$ 1-8 + 2-8 = 13$	$ 7-8 + 8-8 = 1$	C_2

Recompute Centroids:

$$C_1 = \left(\frac{1+2+3}{3}, \frac{2+3+3}{3} \right) = (2, 2.67)$$

$$C_2 = \left(\frac{6+7+8}{3}, \frac{7+8+8}{3} \right) = (7, 7.67)$$

Again by using Manhattan distance

	C1	C2	Decision
A	$ 2-1 + 2.67-2 = 1.666$	$ 7-1 + 7.67-2 = 10.666$	C1
B	$ 2-2 + 2.67-3 = 0.334$	$ 7-2 + 7.67-3 = 9.666$	C1
C	$ 2-3 + 2.67-3 = 1.334$	$ 7-3 + 7.67-3 = 8.666$	C1
D	$ 2-6 + 2.67-7 = 5.334$	$ 7-6 + 7.67-7 = 1.666$	C2
E	$ 2-7 + 2.67-8 = 5.334$	$ 7-7 + 7.67-8 = 0.334$	C2
F	$ 2-8 + 2.67-8 = 6.334$	$ 7-8 + 7.67-8 = 1.334$	C2

Recompute Centroids:

$$C_1 = \left(\frac{1+2+3}{3}, \frac{2+3+3}{3} \right) = (2, 2.666)$$

$$C_2 = \left(\frac{6+7+8}{3}, \frac{7+8+8}{3} \right) = (7, 7.67)$$

The C_1 and C_2 remain unchanged ; so the solution is ,

$$C_1 = A, B, C$$

$$C_2 = D, E, F$$