

Output tables for 1xN statistical comparisons.

March 10, 2025

1 Average rankings of Friedman test

Average ranks obtained by each method in the Friedman test.

Algorithm	Ranking
Random	6.875
NN	5
Sweep	5.75
SaveSeq	2.625
SaveParall	1
MJ	2.625
CMT	4.125

Table 1: Average Rankings of the algorithms (Friedman)

Friedman statistic (distributed according to chi-square with 6 degrees of freedom): 43.071429.
P-value computed by Friedman Test: 0.

2 Post hoc comparison (Friedman)

P-values obtained in by applying post hoc methods over the results of Friedman procedure.

i	algorithm	$z = (R_0 - R_i)/SE$	p	Holm	Finner	Li
6	Random	5.439193	0	0.008333	0.008512	0.04566
5	Sweep	4.397645	0.000011	0.01	0.016952	0.04566
4	NN	3.70328	0.000213	0.0125	0.025321	0.04566
3	CMT	2.893188	0.003814	0.016667	0.033617	0.04566
2	SaveSeq	1.504458	0.132464	0.025	0.041844	0.04566
1	MJ	1.504458	0.132464	0.05	0.05	0.05

Table 2: Post Hoc comparison Table for $\alpha = 0.05$ (FRIEDMAN)

Holm's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.025 .
Finner's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.041844 .
Li's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.04566 .

3 Adjusted P-Values (Friedman)

Adjusted P-values obtained through the application of the post hoc methods (Friedman).

i	algorithm	unadjusted p	p_{Holm}
1	Random	0	0
2	Sweep	0.000011	0.000055
3	NN	0.000213	0.000851
4	CMT	0.003814	0.011441
5	SaveSeq	0.132464	0.264927
6	MJ	0.132464	0.264927

Table 3: Adjusted p -values (FRIEDMAN) (I)

i	algorithm	unadjusted p	p_{Finner}	p_{Li}
1	Random	0	0	0
2	Sweep	0.000011	0.000033	0.000013
3	NN	0.000213	0.000426	0.000245
4	CMT	0.003814	0.005715	0.004377
5	SaveSeq	0.132464	0.156772	0.132464
6	MJ	0.132464	0.156772	0.132464

Table 4: Adjusted p -values (FRIEDMAN) (II)