

Output tables for 1xN statistical comparisons.

March 10, 2025

1 Average rankings of Friedman test

Average ranks obtained by each method in the Friedman test.

Algorithm	Ranking
Random	6.375
NN	5.25
Sweep	6.375
SaveSeq	2.875
SaveParall	1.875
MJ	2.375
CMT	2.875

Table 1: Average Rankings of the algorithms (Friedman)

Friedman statistic (distributed according to chi-square with 6 degrees of freedom): 38.625.
P-value computed by Friedman Test: 0.000001.

2 Post hoc comparison (Friedman)

P-values obtained in by applying post hoc methods over the results of Friedman procedure.

i	algorithm	$z = (R_0 - R_i)/SE$	p	Holm	Finner	Li
6	Random	4.16619	0.000031	0.008333	0.008512	0.018767
5	Sweep	4.16619	0.000031	0.01	0.016952	0.018767
4	NN	3.124643	0.00178	0.0125	0.025321	0.018767
3	SaveSeq	0.92582	0.354539	0.016667	0.033617	0.018767
2	CMT	0.92582	0.354539	0.025	0.041844	0.018767
1	MJ	0.46291	0.643429	0.05	0.05	0.05

Table 2: Post Hoc comparison Table for $\alpha = 0.05$ (FRIEDMAN)

Holm's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.016667 .
Finner's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.033617 .
Li's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.018767 .

3 Adjusted P-Values (Friedman)

Adjusted P-values obtained through the application of the post hoc methods (Friedman).

i	algorithm	unadjusted p	p_{Holm}
1	Random	0.000031	0.000186
2	Sweep	0.000031	0.000186
3	NN	0.00178	0.007121
4	SaveSeq	0.354539	1.063618
5	CMT	0.354539	1.063618
6	MJ	0.643429	1.063618

Table 3: Adjusted p -values (FRIEDMAN) (I)

i	algorithm	unadjusted p	p_{Finner}	p_{Li}
1	Random	0.000031	0.000186	0.000087
2	Sweep	0.000031	0.000186	0.000087
3	NN	0.00178	0.003557	0.004968
4	SaveSeq	0.354539	0.481433	0.498571
5	CMT	0.354539	0.481433	0.498571
6	MJ	0.643429	0.643429	0.643429

Table 4: Adjusted p -values (FRIEDMAN) (II)