

Output tables for 1xN statistical comparisons.

March 10, 2025

1 Average rankings of Friedman test

Average ranks obtained by each method in the Friedman test.

Algorithm	Ranking
Random	6.875
NN	5.375
Sweep	5.75
SaveSeq	1.5
SaveParall	2.375
MJ	2.125
CMT	4

Table 1: Average Rankings of the algorithms (Friedman)

Friedman statistic (distributed according to chi-square with 6 degrees of freedom): 43.928571.
P-value computed by Friedman Test: 0.

2 Post hoc comparison (Friedman)

P-values obtained in by applying post hoc methods over the results of Friedman procedure.

i	algorithm	$z = (R_0 - R_i)/SE$	p	Holm	Finner	Li
6	Random	4.976283	0.00001	0.008333	0.008512	0.023009
5	Sweep	3.934735	0.000083	0.01	0.016952	0.023009
4	NN	3.587553	0.000334	0.0125	0.025321	0.023009
3	CMT	2.31455	0.020638	0.016667	0.033617	0.023009
2	SaveParall	0.810093	0.417887	0.025	0.041844	0.023009
1	MJ	0.578638	0.562834	0.05	0.05	0.05

Table 2: Post Hoc comparison Table for $\alpha = 0.05$ (FRIEDMAN)

Holm's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.016667 .

Finner's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.041844 .

Li's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.023009 .

3 Adjusted P-Values (Friedman)

Adjusted P-values obtained through the application of the post hoc methods (Friedman).

i	algorithm	unadjusted p	p_{Holm}
1	Random	0.00001	0.000004
2	Sweep	0.000083	0.000416
3	NN	0.000334	0.001335
4	CMT	0.020638	0.061913
5	SaveParall	0.417887	0.835774
6	MJ	0.562834	0.835774

Table 3: Adjusted p -values (FRIEDMAN) (I)

i	algorithm	unadjusted p	p_{Finner}	p_{Li}
1	Random	0.00001	0.000004	0.000001
2	Sweep	0.000083	0.00025	0.00019
3	NN	0.000334	0.000667	0.000763
4	CMT	0.020638	0.030796	0.045079
5	SaveParall	0.417887	0.477593	0.488726
6	MJ	0.562834	0.562834	0.562834

Table 4: Adjusted p -values (FRIEDMAN) (II)