

Output tables for 1xN statistical comparisons.

March 10, 2025

1 Average rankings of Friedman test

Average ranks obtained by each method in the Friedman test.

Algorithm	Ranking
Random	6.875
NN	4.75
Sweep	6.125
SaveSeq	1
SaveParall	3.5
MJ	2.25
CMT	3.5

Table 1: Average Rankings of the algorithms (Friedman)

Friedman statistic (distributed according to chi-square with 6 degrees of freedom): 44.410714.
P-value computed by Friedman Test: 0.

2 Post hoc comparison (Friedman)

P-values obtained in by applying post hoc methods over the results of Friedman procedure.

i	algorithm	$z = (R_0 - R_i)/SE$	p	Holm	Finner	Li
6	Random	5.439193	0	0.008333	0.008512	0.039623
5	Sweep	4.744828	0.000002	0.01	0.016952	0.039623
4	NN	3.471825	0.000517	0.0125	0.025321	0.039623
3	SaveParall	2.31455	0.020638	0.016667	0.033617	0.039623
2	CMT	2.31455	0.020638	0.025	0.041844	0.039623
1	MJ	1.157275	0.24716	0.05	0.05	0.05

Table 2: Post Hoc comparison Table for $\alpha = 0.05$ (FRIEDMAN)

Holm's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.016667 .
Finner's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.05 .
Li's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.039623 .

3 Adjusted P-Values (Friedman)

Adjusted P-values obtained through the application of the post hoc methods (Friedman).

i	algorithm	unadjusted p	p_{Holm}
1	Random	0	0
2	Sweep	0.000002	0.00001
3	NN	0.000517	0.002068
4	SaveParall	0.020638	0.061913
5	CMT	0.020638	0.061913
6	MJ	0.24716	0.24716

Table 3: Adjusted p -values (FRIEDMAN) (I)

i	algorithm	unadjusted p	p_{Finner}	p_{Li}
1	Random	0	0	0
2	Sweep	0.000002	0.000006	0.000003
3	NN	0.000517	0.001034	0.000686
4	SaveParall	0.020638	0.030796	0.026682
5	CMT	0.020638	0.030796	0.026682
6	MJ	0.24716	0.24716	0.24716

Table 4: Adjusted p -values (FRIEDMAN) (II)