

Output tables for 1xN statistical comparisons.

March 10, 2025

1 Average rankings of Friedman test

Average ranks obtained by each method in the Friedman test.

Algorithm	Ranking
Random	6
NN	5
Sweep	7
SaveSeq	2.5
SaveParall	1.125
MJ	2.375
CMT	4

Table 1: Average Rankings of the algorithms (Friedman)

Friedman statistic (distributed according to chi-square with 6 degrees of freedom): 46.553571.
P-value computed by Friedman Test: 0.

2 Post hoc comparison (Friedman)

P-values obtained in by applying post hoc methods over the results of Friedman procedure.

i	algorithm	$z = (R_0 - R_i)/SE$	p	Holm	Finner	Li
6	Sweep	5.439193	0	0.008333	0.008512	0.039623
5	Random	4.513373	0.000006	0.01	0.016952	0.039623
4	NN	3.587553	0.000334	0.0125	0.025321	0.039623
3	CMT	2.661733	0.007774	0.016667	0.033617	0.039623
2	SaveSeq	1.273003	0.203017	0.025	0.041844	0.039623
1	MJ	1.157275	0.24716	0.05	0.05	0.05

Table 2: Post Hoc comparison Table for $\alpha = 0.05$ (FRIEDMAN)

Holm's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.025 .
Finner's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.041844 .
Li's procedure rejects those hypotheses that have an unadjusted p-value ≤ 0.039623 .

3 Adjusted P-Values (Friedman)

Adjusted P-values obtained through the application of the post hoc methods (Friedman).

i	algorithm	unadjusted p	p_{Holm}
1	Sweep	0	0
2	Random	0.000006	0.000032
3	NN	0.000334	0.001335
4	CMT	0.007774	0.023322
5	SaveSeq	0.203017	0.406034
6	MJ	0.24716	0.406034

Table 3: Adjusted p -values (FRIEDMAN) (I)

i	algorithm	unadjusted p	p_{Finner}	p_{Li}
1	Sweep	0	0	0
2	Random	0.000006	0.000019	0.000008
3	NN	0.000334	0.000667	0.000443
4	CMT	0.007774	0.011638	0.010221
5	SaveSeq	0.203017	0.238379	0.212393
6	MJ	0.24716	0.24716	0.24716

Table 4: Adjusted p -values (FRIEDMAN) (II)