Eclipses by Appointment: 921 Eclipsing Systems, Catalogue Corrections and a Measured Null Planet Search from Public TESS and Gaia DR3 Data

November 5, 2026

The author's role and the use of AI systems are set out in Section 6, Contributions and AI use. Per-object analyses, available light curves and supporting measurements are in the Supplement and repository; catalogue corrections, probable blends and other objects are also documented in the earlier working draft (Appendix B).

AAVSO keywords = Eclipsing Binaries; Catalogues, Databases, Surveys; Period Analysis, Period Change; Extrasolar Planets; Variable Stars (individual), Observing Targets: HIP 1060, HD 170829, V1192 Cas, TYC 4590-364-1

ADS keywords = binaries: eclipsing; binaries: spectroscopic; catalogs; planetary systems; stars: variables: general; techniques: photometric

1. Abstract

We retain 921 accepted eclipsing-system claims in public TESS photometry: 808 confirmed and 113 probable. No prior eclipsing-binary identification was found for these claims in the catalogues searched (Section 2.7). The orbit-informed review archive retains 29 plausible signals outside that count, including eight from a later 49-target review. A separate mixed-method harvest, including period-free dips and complementary leads, has 350 completed selected reviews. Among the 64 physically confirmed or probable cases, 13 are known recoveries, 43 retain unknown catalogue novelty, and eight scoped nonmatches passed claim review and contribute one confirmed and seven probable additions (R725–R732). The 43 pending cases remain outside the 921-system accepted claim count. One hundred seven systems come from a method we call eclipses by appointment: a Gaia DR3 spectroscopic orbit with a period of 20–60 d predicts conjunction windows in which TESS can reveal eclipses that a blind period search misses. Among 3,749 stars with G < 11 the method found about 1.7 new eclipsing binaries per hundred stars; a G 11–11.5 subsample contributes 42 confirmed systems. Gaia-orbit period folds contribute 798 systems, the nearby-M-dwarf-selected search contributes four (including neighbouring hosts), near-edge-on Gaia astrometric orbits contribute four, and the mixed-method period-free follow-up contributes eight. The period-fold sample includes fainter spectroscopic binaries and two probable cases from a near-orbital-period pilot. An adaptive exploratory conjunction pilot adds two confirmed systems while retaining a known recovery separately; its selection and stopping do not support an unbiased yield estimate. The fixed remaining-screened magnitude comparison reviews 60 candidates in each of G 11.5-12 and G 12-12.5; its physical confirmation fractions retain known recoveries and are not completeness or causal magnitude estimates. The analyses combine raw-flux measurements, timing and coverage tests, Gaia covariance, alternative reductions and source attribution, with physical classification kept separate from scoped catalogue status. Twenty Gaia DR3 orbit solutions are contradicted under the paper's stated comparison rule; ten heartbeat systems are physically confirmed, one also eclipsing, and one further system is a candidate; the newly reviewed heartbeat retains incomplete supplementary catalogue checks; eleven existing catalogue entries are corrected. Harvard plates from 1890–1990 support the eclipses of 17 of 31 tested systems, ASAS-SN photometry supports 20 of 55, and ASAS-3 supports 10 of 34. A transit search of about 3,100 nearby M dwarfs found no new

planet where about 1.2 detections were expected under the stated assumptions. We recover both TIC 4206066 candidates and identify TOI-5394.01 as an M-dwarf occultation. Per-object evidence, ephemerides where constrained, status limitations and follow-up tests accompany the sample.

2. Introduction

Gaia DR3 published about 180,000 spectroscopic orbits (Gaia Collaboration, Arenou et al. 2023), each giving a period, an eccentricity, an argument of periastron and a time of periastron for a binary resolved only in velocity. TESS has photometered most of those stars in its full-frame images over the same years. Where the two overlap, an orbit seen within a degree or two of edge-on must eclipse, and TESS records the eclipse whenever a conjunction falls in an observed sector. For periods below about 20 d a period search at Gaia's period finds such eclipses easily. Between 20 and 60 d it does not: a 27-day sector holds at most one eclipse of each kind, and multi-sector baselines of years can be phased only when the period is already known to minutes or the single events are deep enough to be found one by one. Catalogued eclipsing binaries are correspondingly scarce in that period range, and the Gaia orbits that would gain most from an eclipse, which fixes the inclination and with it the companion mass, are the ones TESS searches least well.

Gaia DR3 contains about 800,000 non-single-star solutions for astrometric, spectroscopic and eclipsing binaries (Gaia Collaboration, Arenou et al. 2023). The spectroscopic processing, restricted to single-lined F–K stars brighter than magnitude 12, delivered 181,327 SB1 orbits after filtering (Gosset et al. 2025), and the astrometric processing about 165,500 orbits (Halbwachs et al. 2023a). The SB1 orbits are mostly reliable: of 827 binaries shared with the SB9 catalogue, 655 agree to within 10% in period and eccentricity (Merle et al. 2026), and a comparison with LAMOST and GALAH velocities removed the physically unlikely short-period, high-eccentricity solutions (Bashi et al. 2022). Double-lined solutions fare worse, matching the photometry in period and eccentricity for about half of SB2 eclipsing binaries (Rowan et al. 2023) and 20% of SB2 heartbeat stars, against 85% of SB1 ones (Callahan et al. 2025).

TESS eclipsing binaries are catalogued by Prša et al. (2022; 4,584 systems from sectors 1–26), IJspeert et al. (2021, 2024) and Kostov et al. (2025; 10,001 vetted systems from the full-frame images and about 900,000 unvetted machine-learning detections), and Gaia DR3 adds 2,184,477 candidates from its own photometry (Mowlavi et al. 2023). All favour short periods: the 7,936 new systems of Kostov et al. (2025) have a median period of 3.5 d and a 95th percentile of about 11.6 d. Prša et al. (2022) attribute this to the sector length, which suppresses detections from periods as short as 13 d, and to the box least-squares search, which cannot find single eclipses. Planet searches meet the same limit with single-event methods, estimating the period from one transit's shape (Osborn et al. 2016) or catching a second event from the ground, which also finds eclipsing binaries such as a 61.8-d M dwarf eclipsing a Sun-like star (Gill et al. 2020); "duotransits", one transit in each of two TESS years, typically have periods above 20 d (Hawthorn et al. 2024). In each case the event is found first and the period afterwards.

Heartbeat stars are eccentric binaries that brighten near periastron through tidal distortion, irradiation and Doppler beaming (Kumar et al. 1995; Welsh et al. 2011; Thompson et al. 2012); some show tidally excited oscillations (Fuller and Lai 2012; Fuller 2017). Kepler supplied the first samples (Kirk et al. 2016; Shporer et al. 2016), and the TESS searches have been at short periods: 23 systems at $P = 2.7–20$ d (Li et al. 2024a), 180 at $P \lesssim 10$ d (Solanki et al. 2025) and 112 from Gaia spectroscopic orbits, with the SB1 orbits cut at $P < 13$ d (Callahan et al. 2025). A third body shows in eclipse times through the light-travel-time effect (Irwin 1952); eclipse timing found third bodies in 222 of about 2,600 Kepler binaries (Rappaport et al. 2013; Borkovits et al. 2016), and 403 known eclipsing binaries have a Gaia DR3 solution at a period much longer than the eclipse period (Czavalinga et al. 2023a). Over longer baselines, the Harvard plates, about 500,000 taken from 1885 to 1992 and digitised by DASCH with

photometry good to about 0.1–0.15 mag (Laycock et al. 2010; Grindlay et al. 2012; Tang et al. 2013; Williams 2025), have been combined with TESS eclipse times to follow a binary's period over a century (Zhang et al. 2026).

New variables are registered in the AAVSO International Variable Star Index (VSX; Watson et al. 2006); times of minimum from AAVSO observers appear regularly in this journal (e.g., Samolyk 2022), and eclipsing binaries found in survey and archival data have been reported here, entered in VSX and characterised with the authors' own photometry (Romanov 2021; De Elías 2025). Guinan et al. (2012) set out the follow-up that eclipsing binaries from space missions need from observers.

Orbits and eclipses have been combined in both directions. The EBLM programme measured orbits for 118 binaries first seen as shallow WASP eclipses (Triaud et al. 2017), and joint fits of this kind supply the masses and radii in DEBCat (Southworth 2015). Starting from Gaia, Rowan et al. (2023) measured masses and radii of 122 stars from SB2 orbits and ASAS-SN light curves, Schmidt et al. (2023) matched SB1 orbits to TESS Objects of Interest, and Halbwachs et al. (2023b) selected 41 bright astrometric binaries that could eclipse. The conjunction prediction itself is not new: Tarrants and Mendes (2023) applied it to Gaia DR3 SB1 elements to vet known TESS and K2 planet candidates and called it their least reliable test, and Yamaguchi et al. (2024) applied it to one 358-d orbit in search of self-lensing pulses. Eclipses in Gaia orbits have been found by other routes: Callahan et al. (2025) found 18 among their 112 heartbeat candidates with a periodogram, El-Badry et al. (2026) found 25 among the orbits of compact-object candidates, 14 of them hierarchical triples with an eclipse period much shorter than Gaia's, and Beck et al. (2024) searched the six oscillating red giants whose Gaia astrometric inclinations are within 3° of edge-on, a sample that includes our A60, which they did not flag. The search itself, examining TESS only inside the windows that Gaia DR3 spectroscopic orbits predict, was taken two weeks before this work began. Chopard-Vilhem (2026a) deposited on Zenodo on 2026 September 27 a survey of 9,389 Gaia DR3 SB1 orbits whose conjunctions are predictable to within a day, in up to three TESS sectors each, reporting 93 eclipsing systems at periods of 13–119 d, 90 of them not previously known to eclipse, with masses and radii for 54. Its object list is embargoed until the accompanying paper is accepted, so the overlap with the systems reported here is not yet known; his period range contains ours, and some overlap is likely. A second record (Chopard-Vilhem 2026b) applies the same test to 1,726 Gaia astrometric orbits within 1° of edge-on. Its latest open version (1.2.0) retains seven surviving detections; its ten dossier entries include withdrawn or known systems and are not ten new discoveries. A comparison against all 235 rows of our object ledger, using positions within 20″ in the parent sample and exact TIC identifiers in the dossier, shares one claimed detection (HD 170829, our A20, found by both searches with the same period). A73, A74 and the withdrawn TYC 3707-251-1 are present in the parent sample; their previously noted pilot-window or event correspondences are not shared claimed detections. All 74 physically confirmed R-labelled and returned additions have exact Gaia DR3 identifiers available; none matches the open parent list or dossier, and none matches a dossier TIC identifier. The separate 93-system spectroscopic list remains embargoed, so its overlap is unknown. Neither record was found by the earlier web and arXiv searches, which did not include Zenodo. The dated comparison and its matching limits are retained in `reviews/novelty-reconciliation/README.md`. This paper claims no priority for the method; its contributions include the objects, archival recoveries and contradicted Gaia orbits.

The work reported here began from a planet question. A public report of two transiting-planet candidates around the K dwarf TIC 4206066 (Rabtsevich 2026) led to a transit search of about 3,100 nearby M dwarfs in TESS full-frame-image light curves (Section 2.3). That search found no new planet (Section 4.6), recovered both reported candidates without using the reported periods or epochs, and turned up four uncatalogued eclipsing systems (A1–A4), two of which had Gaia DR3 evidence of a stellar companion that had never been combined with the TESS photometry. We therefore inverted the

search: starting from Gaia's spectroscopic orbits, we tested each for eclipses in TESS. For orbits shorter than 20 d this is a period search at Gaia's period (Section 2.4), and it found 42 uncatalogued eclipsing systems.

For longer periods a Gaia orbit still fixes the conjunction times: the period, the time of periastron, the eccentricity and the argument of periastron together determine when each conjunction falls, to about 2 d at TESS epochs for the median orbit in our sample. Eclipses can only happen at conjunction, so the search can be confined to the predicted windows and use a single-event statistic instead of a period search over years of data. The number of trials falls from a four-year period grid to the windows, which hold about half of the usable data for the median star, and an eclipse that lands on the predicted conjunction is attributed to the star with the Gaia orbit, which is the hardest step in any blended-field TESS search. We call this eclipses by appointment (Section 2.5). It accounts for 107 of the 913 accepted systems in this paper.

The contributions of this work are (i) the measured appointments-search yield, about 1.7 new eclipsing binaries per hundred Gaia orbits searched at G < 11; (ii) 913 accepted, catalogue-screened systems with physical eclipse evidence, with ephemerides where constrained (Table A1 and Appendix C); (iii) eleven catalogue corrections and twenty Gaia DR3 orbit solutions that the eclipses contradict; and (iv) a null planet result with its expected yield estimated under stated assumptions, and the reason it was null. "Uncatalogued" means absent from the seven catalogue groups specified in Section 2.7; literature searches are a separate, limited check, not unseen by any pipeline: 72 of the first 104 carry an automated threshold-crossing event of the SPOC or TESS-SPOC pipelines (Jenkins et al. 2016; Caldwell et al. 2020) that was never vetted or promoted to a TESS Object of Interest, and a full-text search of NASA ADS found a prior mention of eclipses for only one (A7; Bryant et al. 2023). Physical classification and catalogue novelty are separate. The accepted claim sample contains 807 confirmed and 106 probable systems. The orbit-informed review archive has 29 plausible signals outside that count; the separate period-free selection adds 48 plausible outcomes without changing the claim total. Stable archival keys and the distinct period-free register preserve individual evidence and current verdicts (Appendix C). The object labels (A1–A204, B1–B11, C1–C3) were assigned in this work and are used as identifiers throughout; the A-labelled per-object entries with their light curves are in the Supplement, while the individual physical reviews are in the repository, and the data and code are in the deposit described in Appendix B.

3. Data and methods

3.1 TESS photometry

All photometry is from TESS (Ricker et al. 2015), retrieved from MAST. The primary source was the Quick-Look Pipeline (QLP; Huang et al. 2020; Kunimoto et al. 2021) full-frame-image light curves, at 30-min (sectors 1–26), 10-min (27–55) and 200-s (56 onward) cadence. For the long-period work we used the raw aperture flux SAP_FLUX (Section 2.8). Independent reductions of the same pixels served as cross-checks: SPOC 2-min (Jenkins et al. 2016), TESS-SPOC (Caldwell et al. 2020), TGLC (Han and Brandt 2023), eleanor (Feinstein et al. 2019), GSFC-ELEANOR-LITE (Powell et al. 2022), TASOC (Handberg et al. 2021), CDIPS (Bouma et al. 2019), TARS (Boyle et al. 2026), T16 (Hartman et al. 2025) and TEQUILA (Ogunwale et al. 2025). Times are BTJD = $\mathrm{BJD}_{\mathrm{TDB}}$ − 2457000.

3.2 Gaia DR3 orbits

Orbits are from the Gaia DR3 (Gaia Collaboration, Vallenari et al. 2023) table nss_two_body_orbit (Gaia Collaboration, Arenou et al. 2023; Halbwachs et al. 2023a; Gosset et al. 2025): the spectroscopic types SB1, SB2, SB1C, SB2C and AstroSpectroSB1, and the astrometric Orbital type (Section 2.6), matched to the TESS Input Catalog (TIC 8.2; Stassun et al. 2019; Paegert et al. 2021) by Gaia source identifier.

Gaia gives the time of periastron T_p relative to J2016.0 = BTJD 389.0. Conjunction A is at true anomaly $\nu = 90° - \omega$ and conjunction B at $\nu = 270° - \omega$, with ω the argument of periastron of the star Gaia tracks, which is behind its companion at A. From ν follow the eccentric anomaly E, the mean anomaly $M = E - e \sin E$, and

$$t_c = T_p + 389.0 + M\,P / 2\pi \ \ (BTJD),\ \ plus\ integer\ multiples\ of\ P.$$

Uncertainties were propagated by Monte Carlo through Gaia's errors on P, T_p, e and ω; the conjunction tests used Gaia's full covariance matrix. For circular solutions $\omega = 0$ was assumed. “Primary” denotes the deeper eclipse; a transit is the smaller star crossing the larger, an occultation the smaller star hidden.

3.3 Transit searches of nearby M dwarfs

The parent sample was the TIC 8.2 dwarfs with T_{eff} 3200–4300 K, Tmag 8.5–11 and distance < 40 pc, less every star with a TESS Object of Interest (TOI) or community TOI and every star with a TIC contamination ratio of 0.05 or more; 1,299 of these had 3–8 QLP sectors. Further searches covered fainter and smaller stars, crowded fields, single transits, periods of 0.17–0.5 d and 78 known planet hosts. The searches used box least-squares (BLS; Kovács et al. 2002), with completeness measured from injected transits. Surviving peaks faced odd/even tests, a cyclic-roll null that keeps red noise and destroys transits, difference-image centroids in two field orientations, and archive queries.

3.4 Period folds of Gaia orbits with P < 20 d

The first sample was every Gaia DR3 spectroscopic orbit with P < 20 d and parallax > 10 mas: 736 science orbits, plus 14 Gaia eclipsing binaries as controls, all recovered. A coherent BLS over the flattened QLP flux searched $\pm\max(5\sigma_P, 0.1\%)$ around P_{Gaia} and $P_{Gaia}/2$, scoring each star with a z value against 300 random periods on the same star. Triage removed VSX matches, rotators and ellipsoidal variables, and required an eccentric system's eclipse to land on Gaia's predicted conjunction.

The same code then ran on orbits with parallax ≤ 10 mas or missing. At G < 9, an independent re-check kept seven (A8–A12, B6, B7). At 9 ≤ G < 10, 2,559 science targets among 3,022 orbits were searched, and 199 of 218 controls (91%) reached z ≥ 5, which measures the search's completeness. After triage and catalogue searches, 38 folds showed eclipse-shaped dips and 37 were re-analysed to the full protocol of Section 2.7. About 175 eclipse-like stars with a VSX entry within 30″ and nine marginal dips were not pursued, so this yield is a lower bound.

The fold was then extended to 10 ≤ G < 11.5 in batch (8,492 and 7,470 orbits in the two subsamples; 6,919 and 6,870 searched; 479 of 549 controls, 87%, at z ≥ 5 in the first). Here the per-object re-analysis of Section 2.7 was replaced by the same tests run in bulk: triage and the catalogue searches as above, a visual read of every fold, a search of the vetted eclipsing-binary tables, the conjunction test on Gaia's orbit (offset within 0.03 in phase), per-sector depth consistency, and a difference image of the eclipse deficit from TESScut relative to the target and every Gaia neighbour. A review with separately written code then queried ExoFOP, VSX and SIMBAD for every survivor and re-folded 60 of them, drawn at random, from freshly downloaded QLP flux with a blind BLS over 0.5–25 d. The initial survivors were

listed as candidates until they had had the per-object re-analysis of Section 2.7; the eight that the difference imaging had flagged and the 60 shallowest of the others (to 28 ppt) have had it, and the 66 it confirmed are counted (A119–A179 and A200–A204, Section 3.2). Two limits of the batch tests showed in that pass: the conjunction test cannot tell an eclipse from the tidal minimum of a heartbeat star when the conjunction falls near periastron (two of the 35 were heartbeat stars, Section 3.3; in four more the conjunction fell near periastron and the eclipse model won outright, and in two a real tidal signal and a real eclipse coexist), and the difference-image offsets select crowded fields and pixel boundaries rather than off-target signals (seventeen of seventeen examined were artefacts: of crowding, of a target sitting on a pixel boundary, of an uncalibrated pixel scale with the measuring box centred on the rounded pixel, of a half-period stack that mixed two unequal eclipses, or of out-of-eclipse windows that fell inside a long eclipse).

The spectroscopic-orbit period folds also extend to the fainter Gaia SB1 population at G 12.75–13 and Gaia P < 20 d. The selection stratum contains 2,346 orbit rows; the reviewed subset is not a complete census of that stratum. The original 35 included systems have Gaia G = 12.753–12.993 and selection-orbit periods 1.867–19.643 d. These Gaia values describe selection, not newly measured eclipse ephemerides or proof that the eclipses belong to the Gaia orbit. Individual raw-flux, timing, alternative-reduction, attribution and catalogue checks use the same criteria as the brighter period-fold populations. C57 was subsequently promoted from plausible to probable, bringing the active fainter-SB1 claim subtotal to 36 (28 confirmed, eight probable); its measurement limits remain in the reconciled report.

3.5 Eclipses by appointment

Sample and windows. Gaia DR3 holds 11,446 spectroscopic orbits with $20 \le P < 60$ d and $G < 12$. We searched three subsamples, G < 10 (1,422 stars, including 40 Gaia eclipsing binaries as controls), G 10–11 (2,327 stars) and, during the analysis, G 11–11.5 (2,608 stars): 6,357 stars in all. For each orbit the conjunction times and their uncertainty σ_T were computed at every TESS cycle (Section 2.2). Each window had a half-width of $\max(3\sigma_T, 0.25\text{ d})$, capped at 7 d.

Statistic and cuts. In each window the QLP SAP_FLUX (15-min bins) was searched with a single-event box of 1–14 h against the mean of two side windows 0.75–2.5 durations out. Two shape tests followed: the far ratio (the depth against a baseline 3–6 durations out, divided by the near depth, about 1 for an eclipse) and a deficit in both halves of the box. The null was the same statistic in 40 random windows on the same star outside every predicted window, giving $z = (\text{SNR} - \text{median})/\text{MAD}$, a rank against the star's own null rather than a Gaussian significance. A candidate needed SNR ≥ 8, a far ratio in [0.6, 1.6], the event within $3\sigma_T$ of a conjunction, no null window reaching it, and $z \ge 8$ where a null exists. Of the 40 controls, 32 showed an eclipse at the predicted conjunction.

Coverage. Over the 3,708 science stars of the first two subsamples the windows hold a median 51% of the usable cadences, and every cadence for 165 stars (4.5%), where no null exists; for 591 more (16%), z is undefined because all null windows land on the same feature (MAD = 0). Seven of the 54 reported systems therefore never faced the z test (A37, A40 and A42 with no null; A27, A51, A54 and A64 with MAD = 0) and passed on the remaining cuts. One, A18, failed a cut (far ratio 2.19); it was taken from the raw SNR ranking before the cuts were written and rests on six eclipses and its re-analysis.

Classification. Candidates passing the cuts were matched against VSX, ASAS-SN, Prša et al. (2022), IJspeert et al. (2024), Kostov et al. (2025) and SIMBAD; a VSX entry of a non-eclipsing type did not remove a star. The remaining 161 (G < 10) and 280 (G 10–11) stars were classified visually. Every star classified as an eclipser, and every unusual non-eclipse signal, was re-analysed independently (Section 2.7), with three exceptions (Section 3.1): 16 at G < 10 and 54 at G 10–11. At G 11–11.5, 408 stars passed

the cuts, 29 of them catalogued eclipsing binaries recovered blind, and the 379 others were classified visually: 47 eclipsers and three possible ones. Forty-five of these 50 candidates have individual reviews: the original twenty counted systems (A180–A199), followed by 25 historical reviews. The latter comprise one known recovery, one retained plausible signal and the 23 systems reconciled here using the bounded checks of Section 2.7. This reconciliation supports 22 confirmed additions and one plausible signal. Five candidates remain unreviewed and are excluded.

3.6 Edge-on Gaia astrometric orbits

Of the 6,652 Gaia DR3 astrometric orbits with P < 300 d and G < 13, we kept the 191 whose inclination, recovered from the Thiele–Innes coefficients, is within 2° of edge-on; the inclination served only to rank. For the 177 with QLP light curves, raw SAP flux was searched around Gaia's period and its half, and separately with a per-sector box search over 0.3–13 d and an inspection of the deepest single events.

Underexplored selection and observational gaps. A frozen queue of 40 remaining screened period-fold candidates was reviewed in increasing Gaia G, covering G = 11.5033-11.7834 within the G 11.5-12 selection stratum. The same bounded raw-flux, measured timing, source-attribution, alternative-reduction and Gaia-geometry checks support 39 confirmed and one probable system; the probable TIC 253126207 retains unresolved attribution after a completed review. Periods come from measured eclipse midpoints and same-kind cycle counts where depths alternate. Predicted conjunction centers are search guides, not measured timing points. Timing errors retain their individual meanings, including scatter inflation where specified. Operational processing groups are provenance only.

An independent exploratory pilot examined underexplored windows using existing methods. The appointments branch was adaptively extended from 48 to 96 selected science targets: 90 had TIC matches, 89 had usable QLP sectors, six lacked a TIC match and one matched target lacked usable QLP data. It produced two confirmed new systems and one known IJspeert et al. (2024) recovery, retained as a control. The near-TESS-orbital-period branch examined 72 targets (including the initial 24), retaining two probable eclipsing systems because capable close neighbours prevent definitive attribution. Sixteen long-duration targets gave no promoted case; availability queries for eight prior single-event targets found no new sectors. Failed and unavailable attempts remain in the denominators. Adaptive selection and stopping make these pilot counts unsuitable as unbiased occurrence, completeness or yield estimates. The target selections, attempted observations and outcomes are retained separately from the frozen 40-target queue.

Fixed magnitude comparison. Sixty remaining screened Gaia spectroscopic-orbit period-fold candidates were reviewed: 30 at G = 11.5-12 and 30 at G = 12-12.5, the latter split equally between historical chunks 6 and 7. The initial 120-target queue was seeded (2026103101) before review. Four inventory-magnitude boundary substitutions (seed 2026103102; two active and two reserved) occurred while the first processing group was being reviewed, solely by the magnitude rule. Original queues are retained. Historical inventory magnitudes differ slightly from Gaia DR3 values: later crosschecks showed the removed objects remained inside the DR3 intervals. All final 120 satisfy both definitions. This is a sample from boundary-valid historical screening strata, not a pristine random sample of all eligible DR3 stars. Both the first 60 and the reserved second 60 have now completed the bounded review, giving 60 selected targets in each magnitude interval. No reserved target was replaced after review. Scientific summaries use Gaia DR3 G, with inventory G preserved separately.

Completion of the bright screened stratum. The remaining 73 screened Gaia spectroscopic-orbit period-fold candidates at actual Gaia DR3 G = 11.5-12 (range 11.6578-11.9967) completed the same bounded review: 69 confirmed, 3 probable and 1 plausible; 69 remain accepted unique additions (66

confirmed and three probable) after three prior-known recoveries. This follow-up completes that remaining bright stratum and is separate from the fixed balanced 120-target magnitude comparison, whose sample and descriptive results remain unchanged. Historical inventory-boundary exclusions that meet the actual DR3 interval retain their original substitution provenance. No targets were substituted in this follow-up. The screened inventory has now completed individual review. Five weaker conjunction candidates, 58 lower-score candidates and 48 unexecuted exploratory appointment targets are separate populations. These outcomes do not measure blind discovery yield or completeness.

Brightest remaining screened period folds. We reviewed the next 130 candidates in increasing actual Gaia DR3 G (12.0036-12.3371), using the same physical, attribution and catalogue checks defined in Section 2.7. Of these, 103 are physically confirmed, 24 probable and three plausible. Six systems are now known recoveries, so 121 accepted unique additions remain: 99 confirmed and 22 probable; plausible signals remain excluded. These candidate reviews extend the prior G 11.5-12 completion and are separate from the fixed balanced 120-target magnitude comparison. The final 136 candidates in this screened inventory have now completed individual review. Five weaker conjunction appointments, 58 lower-score survivors and 48 unexecuted exploratory pilot targets belong to separate selections. This selected follow-up does not estimate blind discovery yield or completeness.

Completion of the screened period-fold inventory. The final 136 screened candidates, reviewed in increasing actual Gaia DR3 G (12.337704-12.742949), comprise 101 physically confirmed systems, 34 probable systems and one plausible signal. Three systems, including TIC 295771108, are now known recoveries and excluded from additions. The 132 accepted unique additions comprise 100 confirmed and 32 probable systems. TIC 290749976 remains plausible because eclipse morphology and source attribution are insecure. This completes this screened inventory, while five weaker conjunction appointments, 58 lower-score survivors and 48 unexecuted exploratory pilot targets remain separate selections. The fixed balanced 120-target comparison is unchanged. These selected outcomes do not estimate blind discovery yield or completeness.

3.7 Independent re-analysis and catalogue checks

Original all-event re-analysis. In the original analyses, every candidate from the appointments search, the G 9–10 fold and the edge-on search was re-analysed independently, with separately written code and with the aim of refuting the claim (Section 6). Each original re-analysis re-measured every eclipse from raw SAP flux, including flagged cadences; fitted a linear ephemeris to all events, so that the period comes from the eclipses themselves; located the flux deficit in difference images relative to the target and every Gaia neighbour bright enough to produce it; re-measured the eclipses in at least one independent pipeline; compared the eclipse geometry with Gaia's orbit and mass function; and predicted the next eclipses.

Conjunction-selected evidence reconciliation. The 23 later-reviewed G 11–11.5 systems were assessed from the deciding raw-event, timing and coverage, pixel attribution, alternative-reduction and Gaia geometry evidence. Existing validated evidence was reused with hashes; missing material checks were completed where practical, including a repeated eclipse in newer TESS coverage for TIC 12864953. Alternative reductions of the same photons test extraction and processing, not independent observing occasions. This bounded reconciliation is not a new exhaustive all-sector search. A completed review can retain a plausible physical outcome. Reported timing errors retain their source-specific meaning: the KELT+TESS period error for TIC 277542587 is heuristic, and the conservative allowance for TIC 12864953 is not calibrated coverage. Reference epochs are named by eclipse kind rather than assumed to identify the deeper primary. The October 31 catalogue checks cover the catalogue sources defined below; the cached Chopard v1.2.0 exact-identifier comparison found no overlap among these 23, without asserting a newer online release was searched or resolving the 93 embargoed identities.

Catalogues. Each object was cone-searched in VSX (Watson et al. 2006), SIMBAD (Wenger et al. 2000), the Gaia DR3 eclipsing-binary (Mowlavi et al. 2023), classifier (Rimoldini et al. 2023; Eyer et al. 2023) and non-single-star tables, Prša et al. (2022), IJspeert et al. (2024), Kostov et al. (2025) and ASAS-SN (Shappee et al. 2014; Kochanek et al. 2017; Jayasinghe et al. 2018), through VizieR (Ochsenbein et al. 2000), with a known eclipsing binary as positive control. A repeat on 2026 October 13, adding the GCVS and NSV (Samus' et al. 2017), ASAS-3 (Pojmański 2002), ATLAS (Heinze et al. 2018), ZTF (Chen et al. 2020) and Catalina (Drake et al. 2014), found no eclipsing-type entry for any counted A-object. A full-text query of the NASA ADS index for 248 identifiers of the 74 A-objects (2026 October 12) returned one prior report of eclipses (A7; Bryant et al. 2023); a second (October 13) covered 242 identifiers of the 35 G 9–10 systems.

Catalogue scope and status words. **Uncatalogued** is shorthand for no prior eclipsing-binary identification found in the searched sources: (1) VSX; (2) SIMBAD primary classifications; (3) Gaia DR3 `vari_eclipsing_binary`; (4) Prša et al. (2022); (5) IJspeert et al. (2024); (6) the vetted tables 3 and 4 of Kostov et al. (2025); and (7) ASAS-SN. These are seven catalogue sources or source groups, not seven physical classes of binaries. An entry only in Kostov et al.'s unvetted machine-learning table 2, or a generic non-eclipsing variability classification, does not constitute a prior eclipsing-binary identification. Matches to nearby stars are assessed against the measured signal and source attribution rather than automatically assigned to the target. The dated evidence records each query or exact-identifier table comparison, its scope and result; checks span the original October analyses and subsequent extension reviews. Unavailable or invalid responses remain unknown, and an object missing a required catalogue check is excluded from new claims until that check is resolved. Identified prior literature reports also exclude a system from new claims. Successful nonmatches are bounded by the sources, identities, search radii and dates used; they do not establish exhaustive discovery priority. **Confirmed**: the system passed every check above that its data allow (repeated events in raw flux, a period from the eclipses, a difference image on the target, a second pipeline, a geometry consistent with Gaia's orbit). **Probable**: the eclipses are real and repeat, but one check is missing or failed. **Plausible**: a single event, or two with the period untested. **By-catch**: a real variable on the target whose signal does not follow the Gaia orbit that selected it.

3.8 Known systematics

QLP's detrended flux suppresses long eclipses in post-sector-56 files: it removed the 20-h primary of A16. QLP's sector-80 light curves run 2.7 min late against SPOC and TESS-SPOC in the two systems where this was measured (A11, A98); whether the offset applies to the whole sector has not been tested. At 30-min cadence QLP mid-times can be off by minutes: in A92, sectors 14 and 15 are about 6 min late against three independent pipelines. Depths are those measured in the named pipeline and are diluted by any neighbour it does not correct for. Because the QLP aperture changes between sectors, a neighbour can make one sector's eclipse 9–27% shallower than the others (for example A85), so depths are taken from the sectors that agree with difference images or a second pipeline.

Individual review uses the same five deciding checks on representative available events, rather than requiring all-event or all-sector extraction. Review of the G 10–11.5 period-fold candidates includes raw-flux events and uncertainties, timing and coverage, each source's own Gaia covariance, alternative reductions, neighbours and pixel attribution, and physical plausibility. Tested analysis primitives and valid existing evidence are reused where appropriate; these reviews are not all independent-code replications. Alternative reductions, including custom TESScut aperture extractions, share the same TESS photons and test reduction robustness rather than independent astrophysical detections. Original cycle counts and timing baselines are retained in each report; no uniform precision period errors are inferred from point estimates. Physical verdict, catalogue novelty, review limitations and next tests are

recorded separately. All included systems have completed the catalogue checks defined in Section 2.7 without a prior eclipsing-binary identification; known recoveries are excluded from unique additions. The unified scientific review register links physical outcomes and their evidence (Appendix C). Successful shared checks are retained, and failed requests are not negative matches. The audits and row-level coverage are in `reviews/novelty-reconciliation/README.md` and `reviews/2026-10-30-next70/`. Literature searches do not establish exhaustive priority.

3.9 Period-free dip screening and complementary-lead review

We also searched archived TESS QLP light curves for local flux deficits without using a Gaia period or conjunction window, while retaining complementary brightening, fold-only and near-threshold leads. The dip detector used quality-zero, finite positive SAP flux, ten-minute bins where required, and six trial durations from 0.06 to 1.6 d. A supported window required fractional depth at least 0.001 and contrast at least five, where contrast was the depth divided by the larger of sideband residual RMS and robust adjacent-difference noise. Maximizing over windows and durations makes this an ordering statistic, not a calibrated eclipse significance.

The combined inclusive harvest held 2,984 mixed-method leads: dip, brightening, fold-only and near-threshold signals. Removing 80 historical identity overlaps left 2,904 eligible targets. The stratified first 100 comprised 30 cross-method dip, 20 other multi-sector dip, 20 single-event, 10 fold-only, 10 near-threshold fold and 10 brightening leads. Its outcomes informed the order for the remaining 2,804: cross-method dip leads, then other multi-sector dip leads, ordered within each group by maximum dip contrast. The first 250 in that order were reviewed without substitution, yielding 350 distinct reviews and leaving 2,554 mixed-method leads unreviewed. Because the second selection used the first sample's outcomes, the combined fractions are neither a population occurrence rate nor independent validation of that order.

Review inspected representative raw events for recurrence and eclipse morphology against gaps, steps, smooth variability and detector disturbances. Same-kind event timings, integer-cycle alternatives and observed coverage of predicted repeats constrained any clock; missing coverage was not a null. Alternative QLP reductions and TESSCut extractions tested robustness and spatial attribution but reused the same photons. Gaia-orbit association, when proposed, was checked against source-specific conjunction predictions and uncertainty. Physical tier, source association, catalogue status and publication eligibility were recorded separately. A successful scoped catalogue nonmatch is not exhaustive novelty proof; an unavailable query remains unknown. The period-free evidence register retains every selected review and its full report. Among the 64 physically confirmed or probable cases, 13 are known recoveries, 43 retain unknown catalogue novelty, and eight scoped nonmatches passed claim review and contribute one confirmed and seven probable additions (R725–R732). The 43 pending cases remain outside the 921-system accepted claim count. Later direct VSX queries identified five previously pending cases from the earlier 250 as prior eclipsing counterparts. The dated reports remain unchanged; explicit catalogue updates accompany the current register. Direct VSX access subsequently failed its final positive control, and ASAS-SN and vetted Kostov checks remain unavailable for the new queue. Positive identities are retained; failed or unperformed checks are unknown. Source-specific nearby entries were adjudicated separately from exact target matches.

4. Results

4.1 Sample and yield of the appointments search

Table 1 gives the yield of the appointments search stage by stage. In the G < 10 subsample, 181 stars passed the cuts. In the G 10–11 subsample, 447 of 2,327 stars passed, and the 280 that no eclipsing-binary catalogue lists were inspected. Of the 70 stars re-analysed independently, 54 A-labelled eclipsing systems were accepted: 50 confirmed and four probable (A37, A40, A53, A68). Three are new heartbeat stars (Section 3.3). Table 2 lists the rest. Four candidates were refuted; two remain plausible, while TYC 7687-990-1 is now physically probable with catalogue novelty pending. Three on-target stars were already known, and 20 catalogued eclipsing binaries were recovered blind in each subsample. In the G 11–11.5 subsample, 408 of 2,608 stars passed the cuts; 29 were catalogued eclipsing binaries recovered blind, and of the 379 inspected, 47 were classified as eclipsers and three as possible. The original twenty strongest are confirmed eclipsing systems (A180–A199). Of 25 later historical reviews, one known recovery and one plausible signal remain outside the counted sample; the other 23 have now been reconciled under Section 2.7, supporting 22 confirmed additions and one plausible signal (TIC 8853239). Thus this stratum contributes 42 counted confirmed systems. Forty-five of the 50 candidates have individual reviews, and five remain unreviewed.

The automated search missed whole eclipses in two ways. The search first discarded 0.5 d after every data gap, which removed whole eclipses (five of the six of A18); the trim was reduced to 0.1 d and the G < 10 subsample was searched again. And in many systems whole eclipses lie in cadences that QLP flags, mostly for scattered light, although the flux is usable: five of the ten eclipses of A30, for example. These were recovered from flagged cadences, other pipelines or cutout photometry.

4.2 The eclipsing systems




Figure 1. Positions of the A-labelled catalogue-screened systems and comparison objects (equatorial, J2000, Mollweide projection). Marker shape gives the search that found each eclipsing system; the catalogue corrections, heartbeat stars, planet candidates and other objects are shown separately. Counts are in brackets; the R-labelled and returned C-labelled systems of Appendix C are not plotted.




Figure 2. Orbital period and deeper eclipse depth of the A-labelled systems and the catalogue corrections; R-labelled and returned C-labelled systems are not plotted. Depths are as measured in the pipeline aperture, diluted by any blended light; a vertical bar spans a depth quoted as a range. The appointments search (circles) fills the 20–60 d range that the period folds (triangles) do not reach.

The accepted claim sample contains 921 systems (808 confirmed and 113 probable), all meeting the stated seven-group uncatalogued standard. The orbit-informed review archive has 29 plausible signals outside that count, including eight from the later 49-target review. By search method, 107 accepted systems come from predicted spectroscopic conjunctions, 798 from Gaia-orbit period folds, four from the nearby-M-dwarf-selected search (including neighbouring hosts), four from near-edge-on Gaia astrometric orbits, and eight from the mixed-method period-free follow-up. Physical classification, association with a Gaia orbit and catalogue status are separate assessments. The prior 913-claim baseline reconciled 917 earlier public claims, 17 prior-known removals and 13 already accepted private additions; A74 changed from probable to confirmed and C57 from plausible to probable. Eight newly reviewed claims bring that baseline to 921. The removed systems retain physical evidence and archival measurements as known recoveries.

HIP 1060 (A47) illustrates the method (Figure 3). It shows a 34%, 6.9-h total eclipse at conjunction A of its Gaia double-lined orbit and a 36%, 16-h annular transit at B, eleven events over eight years at P = 24.10512 d. Because Gaia measured both velocities, one model gives both masses (1.10 and 1.22 $M_{\odot}$) and radii (1.26 and 1.83 $R_{\odot}$). Three eclipses in sector 106, predicted before those data were examined, fell within 1.5 min of the ephemeris. HD 170829 (A20), at G 6.3 the brightest star here, is a G8 subgiant with an asteroseismic mass, eclipsed every 26.384493 d; VSX lists it as NSV 24480 with type “VAR:” and no period.




Figure 3. HIP 1060 (A47, TIC 70858500). Left: QLP SAP flux, detrended with a 1-d running median with the eclipses masked, folded on P = 24.10512 d; dashed lines mark Gaia's predicted conjunctions A and B. Middle: the six 34%, 6.9-h total eclipses at A, 2018–2026. Right: the five 36%, 16-h annular transits at B.

Appointments. Of the 54 systems, 52 are long-period eclipsing binaries found at Gaia's conjunctions, with periods from 20.29 d (A35) to 57.24 d (A63). Two are by-catch at periods other than Gaia's: A45 (31.15 d, against a Gaia double-lined period of 21.68 d) and A54 (a 3.02-d eclipsing pair whose 39-d Gaia orbit is spurious). A third by-catch object, A46, is a 0.38-d contact or ellipsoidal variable on an unresolved pair and is not counted. A180–A199, from the G 11–11.5 subsample, have periods from 21.68 d (A181) to 56.17 d (A198). Ten have giant or subgiant primaries, and in five of these the deeper eclipse is the occultation of a smaller, hotter companion (A182, A183, A189, A192, A199); three carry VSX rotational classifications that describe the primary's spot modulation and miss the eclipses, so EA is added (A184, A185, A194), and a fourth rotational entry is replaced (A192); A191 is WASP0-TR-06, a single partial event of 2000 rejected as a planet (Kane et al. 2005), which now has its period; two community planet candidates on ExoFOP are the occultations of stellar companions (A197, A199); the Kostov et al. (2025) flag near A196 sits on a neighbour 37″ away; two show one eclipse kind only, as their geometry requires (A194, A199); and the orbital tensions stop short of the rule, the largest being A188's occultation-to-transit spacing, 2.9σ from Gaia's, where the occultation times also wander by about 50 min against the transits in a way that neither apsidal motion nor light-travel time explains, and A196's $e \cos \omega$ at 2.3σ, where APOGEE velocities side with the eclipses.

The later conjunction-selected systems retain individual limitations: TIC 277542587 combines repeated KELT eclipses with TESS at a heuristic working period precision of 0.0002 d; TIC 12864953 has a repeated secondary in sector 107 and a conservative 0.00015-d timing allowance, while its new primary window is quality-flagged and a phase-separation tension of about 3.35σ remains unresolved. Neither has been added to the formal Gaia-orbit contradiction count without the stated comparison-rule analysis. TIC 8853239 retains only one primary and an untested period and remains plausible.

Period folds at P < 20 d. A5–A12 come from the folds at parallax > 10 mas and at G < 9; all eight were known spectroscopic binaries, and the eclipses are new. A5, A6, A7 and A10 are probable, because the fold makes period agreement certain by construction and none has a pixel test or a second pipeline. The fold at G 9–10 produced A75–A109, all 35 confirmed. A93 was already typed as an eclipsing binary in SIMBAD from a SuperWASP flag without a period (Schanche et al. 2019), so 34 are counted. In A95 the eclipse times vary with an amplitude of 3.4 ± 0.3 min on the 541-d period of Gaia's astrometric orbit for the star, consistent with the light-travel time that orbit predicts (chance coincidence ~10%).

Period-fold populations and physical discrimination

The G 10–11.5 period-fold search examined 13,789 Gaia spectroscopic orbits and selected 266 signals. Individual analysis of 242 eclipse candidates supports 230 confirmed targets (including four now-known catalogue recoveries), nine probable targets, two non-eclipsing heartbeat stars and one plausible signal. The screening register retains known planet-candidate signals, catalogue corrections and historical holds separately. The catalogue-screened additions use the same seven-group standard as the conjunction search.

The fainter Gaia SB1 period-fold population now contributes 36 accepted systems: 28 confirmed and eight probable, including the promoted C57. These are detections by the existing Gaia-orbit folding method at periods below 20 d, rather than a separate search technique. Known eclipsing-binary recoveries and a blind recovery of A4 are excluded from additions. Appendix C groups all individual reviews by scientific outcome, with the original identifiers retained solely as evidence keys.

Physical discrimination matters most where a tidal minimum resembles an eclipse or a nearby source contaminates an aperture. Two G 10–11.5 signals, TIC 391918838 and TIC 98149548, are heartbeat stars without eclipses (Section 3.3); A144 combines tidal variation with eclipses. Pixel-level tests resolve apparent centroid offsets caused by crowding, while independent reductions test shallow events and systematic features. Unresolved attribution or incomplete decisive checks retain a probable or plausible classification rather than being absorbed into the confirmed sample. The archived reports preserve fitted depths, event counts, ephemerides, Gaia comparisons and each deciding limitation.

M-dwarf searches. A1–A4 were detected as planet-like signals on nearby M or K dwarfs. A1, A2 and A4 are eclipses on a neighbouring star whose light enters the dwarf's aperture. A3 is on the target and is probable: Gaia shows that its companion is most likely a star, but the companion's orbital period is not known.

Edge-on astrometric orbits. A19 is probable: three eclipses repeat at a period that agrees with Gaia's to 0.7σ, but they fall about 2.5σ from the nearer predicted conjunction, and two exist only in flagged or QLP-only data. A72 eclipses on its 215.0116-d Gaia orbit itself. A74 is now confirmed after the accepted tier review, with two 4.6% TESS eclipses three cycles apart and a ZTF night at the predicted time of the cycle between them. A73 (TYC 4590-364-1) is a 17.54-d eclipsing binary inside a 125-d Gaia orbit. Its eclipses have faded from 33% to 13% in four years and their timings vary by hours (Figure 4). NEOWISE W1 photometry, read during the analysis, has the primary no deeper than 20% in 2016–17 and at least 21% in 2021 August, so the depth rose before it fell. A third body is needed to explain this, though the data do not show that it is Gaia's.




Figure 4. TYC 4590-364-1 (A73), from fits to individual primary (black) and secondary (red) eclipses in QLP sectors of 2019–2024. From top: timing residuals against a linear 17.54-d ephemeris, eclipse depths, total durations, and the phase of the secondary (blue). The eclipses fade from 33% to 13% and narrow, and the timings vary by hours.

The G 11.5-12 spectroscopic-orbit period-fold population has 40 physical reviews (39 confirmed, one probable); 38 are current accepted unique additions after two prior-known recoveries. The near-orbital-period pilot contributes two probable systems. The probable cases retain unresolved target-versus-capable-neighbour attribution. TIC 397586559 retains Gaia drift limitations; TIC 343926224 excludes anomalous windows from timing and uses a scatter-inflated uncertainty; TIC 314041837 has localized eclipse cores atop modulation, and its fit window is not a physical eclipse duration. These limitations do

not convert predicted conjunctions into timing measurements or independent reductions into independent observations. The fresh conjunction pilot contributes TIC 382042754 and TIC 389128842 as confirmed systems; TIC 155303130 is a strong known recovery and adds nothing. These findings demonstrate useful underexplored selection windows, not previously untouched parameter space or an unbiased pilot yield.

Physical confirmation among selected candidates. At G = 11.5-12 (actual DR3 range 11.6764-11.9742), 27/30 are physically confirmed (90.0%; descriptive Wilson 95% interval 74.4-96.5%), 3 probable and 0 plausible; 30 are new counted additions. At G = 12-12.5 (actual DR3 range 12.0029-12.4879), 21/30 are physically confirmed (70.0%; descriptive Wilson 95% interval 52.1-83.3%), 8 probable and 1 plausible; 26 remain accepted unique additions after two further prior-known recoveries. Every selected target remains in its denominator, including any unavailable or execution-incomplete outcome; all 60 completed the bounded review. TIC 377389892 is a confirmed known VSX EA recovery (0.02 arcsec counterpart), included in physical confirmation and excluded from new additions. Probable and plausible outcomes remain separate; plausible cases add nothing. These descriptive confirmation fractions concern remaining screened candidates, not blind discovery yield or completeness. Wilson intervals are binomial model summaries despite finite historical strata. Case mix, prior depletion of the bright stratum below G = 11.663, and selection bookkeeping prevent a causal magnitude interpretation.

Reserved and cumulative magnitude comparison. The second reserved sixty: At G = 11.5-12 (actual DR3 range 11.6648-11.9955), 29/30 are physically confirmed (96.7%; descriptive Wilson 95% interval 83.3-99.4%), 1 probable and 0 plausible; 30 new counted additions. At G = 12-12.5 (actual DR3 range 12.0036-12.4872), 26/30 are physically confirmed (86.7%; descriptive Wilson 95% interval 70.3-94.7%), 2 probable and 2 plausible; 28 new counted additions. Both rounds together: At G = 11.5-12 (actual DR3 range 11.6648-11.9955), 56/60 are physically confirmed (93.3%; descriptive Wilson 95% interval 84.1-97.4%), 4 probable and 0 plausible; 60 new counted additions. At G = 12-12.5 (actual DR3 range 12.0029-12.4879), 47/60 are physically confirmed (78.3%; descriptive Wilson 95% interval 66.4-86.9%), 10 probable and 3 plausible; 54 current accepted unique additions. All 120 selected targets completed the bounded review and remain in their selected denominators. Known recoveries enter physical confirmation but add nothing; probable and plausible outcomes remain separate. The first-round results above are retained unchanged. These are descriptive remaining-screened candidate comparisons, not blind discovery yield, completeness or a causal magnitude effect. Wilson intervals are binomial model summaries despite finite historical strata; case mix, depletion, crowding, coverage and selection bookkeeping limit interpretation.

Completion of the bright screened stratum. The remaining 73 screened Gaia spectroscopic-orbit period-fold candidates at actual Gaia DR3 G = 11.5-12 (range 11.6578-11.9967) completed the same bounded review: 69 confirmed, 3 probable and 1 plausible; 69 remain accepted unique additions (66 confirmed and three probable) after three prior-known recoveries. This follow-up completes that remaining bright stratum and is separate from the fixed balanced 120-target magnitude comparison, whose sample and descriptive results remain unchanged. Historical inventory-boundary exclusions that meet the actual DR3 interval retain their original substitution provenance. No targets were substituted in this follow-up. The screened inventory has now completed individual review. Five weaker conjunction candidates, 58 lower-score candidates and 48 unexecuted exploratory appointment targets are separate populations. These outcomes do not measure blind discovery yield or completeness.

Period, epoch and attribution limits. TIC 16809274 retains a shallower-eclipse reference epoch. TIC 119193784 remains probable because integer-cycle aliases are unresolved; TIC 382631802 remains plausible and excluded because structured variability and broad-model duration tension leave the eclipse interpretation insecure; the pixel evidence favours the target. TIC 343340931 and TIC

363264601 remain probable because capable neighbours prevent definitive attribution. The equal-depth aliases of TIC 76020303 and TIC 168917716 are Gaia-informed interpretations; their eclipse timing uncertainties do not resolve the orbital half/double ambiguity. Full reports preserve measured event centers, quality flags, error meanings and source-specific limitations.

Brightest remaining screened period folds. We reviewed the next 130 candidates in increasing actual Gaia DR3 G (12.0036-12.3371), using the same physical, attribution and catalogue checks defined in Section 2.7. Of these, 103 are physically confirmed, 24 probable and three plausible. Six systems are now known recoveries, so 121 accepted unique additions remain: 99 confirmed and 22 probable; plausible signals remain excluded. These candidate reviews extend the prior G 11.5-12 completion and are separate from the fixed balanced 120-target magnitude comparison. The final 136 candidates in this screened inventory have now completed individual review. Five weaker conjunction appointments, 58 lower-score survivors and 48 unexecuted exploratory pilot targets belong to separate selections. This selected follow-up does not estimate blind discovery yield or completeness.

Measurement and attribution qualifications. The listed periods and BTJD reference epochs are the accepted per-object measurements, not uniform predictions from Gaia. Formal timing errors or local fit errors do not represent fully calibrated future-epoch uncertainty; eclipse kind and candidate cycle aliases are specified in the individual reports. TIC 238406817 and TIC 66072136 are physically confirmed known recoveries excluded from additions. TIC 210418035 and TIC 381076353 remain plausible because eclipse morphology is insecure; TIC 232533575 remains plausible with unresolved cycle aliases. A QLP and TESScut comparison from the same TESS sector shares photons, whereas distinct sectors are separate observing occasions from the same instrument. This distinction is material for TIC 210343372, TIC 1978575696 and TIC 234413494. Source attribution remains limited by the available pixel scale, crowding and reduction systematics where noted in the reports.

Completion of the screened period-fold inventory. The final 136 screened candidates, reviewed in increasing actual Gaia DR3 G (12.337704-12.742949), comprise 101 physically confirmed systems, 34 probable systems and one plausible signal. Three systems, including TIC 295771108, are now known recoveries and excluded from additions. The 132 accepted unique additions comprise 100 confirmed and 32 probable systems. TIC 290749976 remains plausible because eclipse morphology and source attribution are insecure. This completes this screened inventory, while five weaker conjunction appointments, 58 lower-score survivors and 48 unexecuted exploratory pilot targets remain separate selections. The fixed balanced 120-target comparison is unchanged. These selected outcomes do not estimate blind discovery yield or completeness.

Measurement qualifications. Individual reports retain eclipse-kind, alias, timing-error and source-attribution limits. Reference epochs in BTJD are per-object measurements; formal local errors are not fully calibrated future-epoch uncertainties. Same-sector QLP and TESScut reductions share photons; different sectors are separate TESS observing occasions. Current depth values are realized central depths of the fitted event models, rather than triangular amplitudes extrapolated beyond the event center. The correction leaves timings, physical verdicts, secondary-event values and capable-neighbour sets unchanged.

Physical classification and limitations. The accepted claim sample contains 921 systems (808 confirmed and 113 probable). The 29 orbit-informed plausible reviews and 48 separate period-free plausible reviews remain outside that count. Physical classification, association with a Gaia orbit and catalogue status are separate assessments. Repeated discrete eclipses, independent reductions and source attribution distinguish secure eclipsing systems from fold-only or ambiguous signals. A confirmed on-target eclipse does not by itself establish its relationship to a discrepant Gaia orbit.

Report-specific attribution, secondary-localization and reduction limitations remain explicit in Appendix C. The same criteria apply throughout this continuous version 1 analysis; the current reconciliation retains new deciding measurements and changed classifications in the dated evidence.

Period-free and complementary-lead selected follow-up

Eight selected signals passed the target-specific catalogue checks defined in Section 2.7 and individual claim review (R725–R732): TIC 238855958 is confirmed; TIC 219345200, 450652458, 458324895, 236223193, 28059397, 19385783 and 51321183 are probable. These are new accepted claims from a mixed-method follow-up of Gaia spectroscopic-orbit targets, not additions to the Gaia period-fold or predicted-conjunction method totals. Raw repeated events and measured or conditional clocks support the eclipse signals. Probable limits include close capable neighbours, unresolved integer-cycle aliases and photometric clocks that do not establish association with the selecting Gaia orbit. For TIC 51321183 the claim is a repeating signal in the blended aperture, not secure assignment to the TIC star. A nearby VSX eclipser 38.35 arcsec from TIC 236223193 has a different clock and insufficient depth to explain its target-region signal. Current physical and catalogue dispositions for all 350 reviews are given below; no failed request was treated as a nonmatch. Selection was outcome-informed, so these eight additions do not measure survey yield or completeness.

The 350 completed reviews comprise 28 confirmed, 36 probable, 65 plausible, 211 refuted screening signals, 5 non-eclipse variables and 5 unresolved outcomes. Among the 64 physically confirmed or probable cases, 13 are known recoveries, 43 retain unknown catalogue novelty, and eight scoped nonmatches passed claim review and contribute one confirmed and seven probable additions (R725–R732). The 43 pending cases remain outside the 921-system accepted claim count. The latest fixed 100-target extension contributes 73 refuted, 17 plausible, 3 probable, 6 confirmed, 1 non eclipse variable. It adds no new accepted eclipsing claims because known recoveries and incomplete catalogue checks remain separate from physical confidence. A plausible event can merit follow-up while its morphology, source or cycle count remains unsettled; a refutation addresses the selected signal, not every possible binary interpretation. The 2,554 unreviewed leads remain in the frozen ranking. These selected-review fractions do not measure completeness or blind discovery yield.

The 29 plausible signals in the orbit-informed review archive and the 65 plausible results here are distinct review populations. Their counts must not be read as a cumulative claim sample. Appendix C and the dedicated register retain all 350 dispositions, including the eight accepted additions and 43 catalogue-pending physical cases.

Probable blends. Three signals from the crowded-field and known-host searches, A13–A15, are eclipses on faint stars near the targets, which the pixels locate but not to a single star or not with enough sectors. They are listed as probable and are not counted among the 199.

The newly reviewed TIC 311918550 retains robust target-eclipse attribution and conjunction agreement despite a reported 8.56σ period difference from Gaia. This is a newly flagged orbital tension; it has not been added to the twenty contradictions without the same formal comparison-rule analysis.

4.3 Heartbeat stars

Table 3 contains ten physically confirmed heartbeat systems, one also eclipsing, and one candidate. Their characteristic brightening and dimming are locked to periastron. For each we fitted the equilibrium-tide model of Kumar et al. (1995) to the light curve alone and compared the e and ω it returns with Gaia's radial-velocity orbit. Three come from the appointments search. HD 193389 (Gaia P 20.7966 d) recovers Gaia's ω to 0.3° and e to 0.016 (3.2σ by Gaia's error); TYC 8203-301-1 also shows a coherent 842-ppm oscillation at 23.0009 ± 0.0006 times the orbital frequency, which we read as a

tidally excited oscillation. Four more come from the G 9–10 fold. Of nine light curves set aside there for a brightening-and-dip feature, seven put it at Gaia's periastron (about 3 in 10,000 by chance); three of the seven are published heartbeat stars, recovered blind, and four are in no catalogue or heartbeat search. BD+61 2067, BD+37 4854 and CD-66 2078 are confirmed; TYC 9045-989-1 is a candidate, its ω 20° (2.4σ) from Gaia's. Two more come from the batch candidates of Section 3.2, where the conjunction test had read their tidal minimum as an eclipse. In TYC 3347-1439-1 (TIC 391918838, *P* 8.92 d, Gaia *e* 0.44) the 2.2 ppt dip is about 20 h wide where the geometry allows 6 h of eclipse, and the Kumar et al. model with Gaia's *e* and ω held fixed beats any trapezoid by $\Delta\chi^2 = 766$; in HD 339939 (TIC 98149548, *P* 2.4026 d, a near-twin A-star SB2) the tidal curve reproduces the whole 9.2 ppt dip with no eclipse, the limit at the dip being 0.4 ppt (2σ) with injected eclipses recovered down to 0.5 ppt. Both are uncatalogued. The same blind spot has a constructive side. In TYC 3624-860-1 (A144, *P* 7.3057 d) an 18 ppt peak-to-peak tidal curve with periastron at phase 0.25 carries a 12 ppt M-dwarf transit at conjunction A and a 2.1 ppt occultation at phase 0.3625, 0.26σ from the separation Gaia's *e* and ω predict; the tide alone loses to tide plus eclipses by $\Delta\chi^2 > 11000$, and the batch's eye-read of “messy oscillation and dip” was this. In CD−35 10394 (A150) a tidal dip of about 1 ppt sits under a 15 ppt eclipse 18° past periastron. In four more of the 35 the conjunction fell near periastron and the heartbeat model lost to the eclipse by $\Delta\chi^2$ of 5000 or more (A138, A141, A143, A147).

TYC 8588-1489-1 (TIC 438089101). A further spectroscopic-orbit fold yields a confirmed, non-eclipsing heartbeat. An independent re-analysis recovers P = 4.800079 ± 0.000030 d, e = 0.2646 ± 0.0040 and ω = 84.78 ± 2.6° from the light curve alone, consistent with Gaia, with a periastron epoch of BTJD 2998.2452 ± 0.0159. The tidal signal is localized to the target in eight sectors over 2019–2025 and reproduced in an independent aperture reduction of the same exposures. Physically duration-limited eclipse models lose by $\Delta\mathrm{BIC} \geq 3869$; additional eclipse limits at the two conjunctions are 539 and 440 ppm. Two target-localized, phase-coherent oscillations occur at 69.0000 ± 0.0005 and 74.0001 ± 0.0005 times the orbital frequency, with amplitudes 246 and 260 ppm. These detections reverse an earlier null caused by an amplitude-normalization error and an insufficient frequency range. A 4.471 d^{-1} mode instead belongs to a nearby star. The fitted beaming term is aperture dependent and is withdrawn as a radial-velocity check. The tide amplitude remains about 4.4 times the expectation using the TIC radius, and the Gaia robust RV amplitude exceeds twice its fitted semi-amplitude; these tensions remain unresolved. No prior heartbeat identification was found in the searched literature or successful direct catalogue checks, but current ASAS-SN, SB9 and WDS checks remain unavailable. The full review, numerical receipts and figure preserve the measurement uncertainties, corrections and catalogue scope. This non-eclipsing system does not increase the eclipsing-system claim count.

4.4 Gaia DR3 orbit solutions contradicted by the eclipses

Eclipses fix the period, and their spacing fixes e cos ω, independently of any velocity measurement. We count a Gaia solution as contradicted when its period is spurious or wrong, or when the period or e cos ω differs from the eclipse value by 3σ or more of Gaia's quoted error, whether Gaia's covariance is propagated in full or its correlations are ignored, and however the ephemeris is carried. Conjunction offsets alone, and e and ω from eclipse durations, are recorded as tensions. Nineteen of the 184 counted systems compared with a non-circular Gaia orbit meet the rule, and so does A113, whose circular Gaia solution carries the wrong period (Table 4; twenty in all): three periods are spurious or wrong (A54, A9, A113, the last the 63-d scan-cadence alias of the eclipse period), ten are 3.4–8.0σ off (A110 by 21 min per orbit, 4.5σ; A117 by 87 min per orbit, 3.5σ; A147 by 4 min per orbit, 4.3σ, which the batch's conjunction offset had extrapolated to +0.019 in phase; A167 by 1.05 min per orbit, 8.0σ, an error understated by a perturbed double-lined solution whose two conjunctions still match the eclipses at Gaia's own epoch; A202 by 11 s per orbit, 3.4σ, whose $e\cos\omega$ is 3.5σ off as well, the occultation at phase 0.49993 ±

0.00045 making the orbit circular where Gaia found e = 0.054 ± 0.015), and seven have e cos ω 3.7σ or more off (for A145 the secondary eclipse at phase 0.49996 ± 0.00018 gives 0.0000 ± 0.0003 against the +0.036 ± 0.008 of Gaia's e = 0.039, 4.5σ, so the orbit is circular and the eccentricity spurious; for A159 the secondary at phase 0.62842 ± 0.00004 gives 0.2005 ± 0.0002 against Gaia's 0.252 ± 0.013, 3.9σ with Gaia's covariance and 3.3σ without, and Gaia's period is 3.2σ long as well). Three conjunction offsets (A89, A104, A87) move between about 2σ and about 8σ with the treatment and are not counted; A109's period difference falls to 1.8–2.7σ once its observed period change is included.

4.5 Catalogue corrections

Eleven catalogued stars have the wrong type, the wrong period or none (B1–B11, Table 5). Nine corrections are established by the data; two are proposed, B1 because its host is not yet separated from an unresolved neighbour 4″ away and B7 because a contamination ratio of 1.77 leaves a blend possible. V1274 Tau (B5) is typed RS at 6.62 d in the GCVS and 6.448 d in VSX, and ASAS-3 lists it as ESD/EC at 1.727 d (Pojmański 2002). TESS shows 4.5- and 2-ppt eclipses at Gaia's 5.62323 d, and the ASAS-3 photometry itself peaks at the 6.450-d rotation period with no power at 1.727 d. Five appointment systems (A20, A25, A31, A32, A52) are VSX type revisions.

4.6 Planet candidates and other objects

The M-dwarf search recovered both transit candidates of TIC 4206066 (Rabtsevich 2026) blind, at SNR 15 and 10 (C1; P = 11.132690 and 3.182785 d). TOI-5394.01 (HD 89277, C2) is the occultation of the M-dwarf companion in its Gaia SB1 orbit (P 15.195017 d), not a planet transit. The search around known hosts recovered the two TOI-6284 signals and a TOI-1467 threshold-crossing event, all in a preprint (C3; Tschudi 2026). Table 6 lists the other objects, for example GJ 4348, a young M dwarf that is plausibly a near-twin M+M pair at 0.223864 d, a case one spectrum at quadrature would settle.

Table 1. Yield of the appointments search.

	***G* < 10 subsample (1,422 stars)**	***G* 10–11 subsample (2,327 stars)**	***G* 11–11.5 subsample (2,608 stars)**
Sent to independent re-analysis	16 stars (two more, HD 201073 and HD 7423, were dismissed on inspection before any check)	54 stars (TYC 5976-120-1, already in VSX as an eclipser, was not sent)	45 of 50 candidates have individual reviews: 20 original systems plus 25 later historical reviews; 23 of the latter reconciled here, with one known recovery and one retained plausible signal outside this queue; 5 remain unreviewed
Eclipsing systems, uncatalogued	14 confirmed (A117 after submission, from the corrected-null re-run)	42 confirmed + 5 probable (A110–A116 and A118 after submission, from the corrected-null re-run) (A37, A40, A53, A68) long-period; by-catch: TYC 447-247-1 (A54, 3.02 d) and TYC 7897-48-1 (A45, a 31.15-d eclipsing binary); BD+30 2277 (A46) is a contact or ellipsoidal variable and is not counted	42 confirmed and counted: 20 original (A180–A199) plus 22 after current reconciliation; catalogue status is separate from planet-candidate annotations
Confirmed heartbeat stars, new	1 (HD 193389)	2 (TYC 8203-301-1, with tidally excited oscillations at 22, 23 and 24 times the orbital frequency; BD+59 2405)	none examined yet

	G < 10 subsample (1,422 stars)	*G* 10–11 subsample (2,327 stars)	*G* 11–11.5 subsample (2,608 stars)
Recovered as already known	20 catalogued eclipsing binaries recovered blind (not sent for re-analysis)	20 catalogued eclipsing binaries recovered blind; 3 on target (TOI-1897 and TOI-5149, planet candidates that are M-dwarf eclipsing binaries; TYC 1194-680-1, a SuperWASP EA with the wrong period); 2 known eclipsing binaries on neighbours 63″ away	29 catalogued eclipsing binaries recovered blind; one known recovery in the later 25 historical reviews (TIC 318209162), excluded from additions
Refuted	1 (systematics)	3 (all neighbour contamination at 36–60″; two of the neighbours are uncatalogued eclipsers themselves)	0 of 45
Uncounted, unresolved	1 plausible single event (BD+16 2639)	2 plausible single events (Wolf 73 and HD 282670, the latter not the Gaia companion's); TYC 7687-990-1 is physically probable from repeated 2025 eclipses but has earlier covered flat sectors and pending catalogue novelty	2 reviewed plausible signals: TIC 47542584 (historical one-sector pair) and TIC 8853239 (one primary, period untested); 5 candidates unreviewed

Table 2. Appointment candidates that are not new confirmed eclipsing binaries.

Object	Outcome	What the check found
Recovered, already known		
TOI-1897 = TIC 420645189	Known: retired planet candidate	TOI-1897.01 and .02 are this system's primary and secondary eclipses, retired by TFOPWG as FP/EB on 2024-09-17; P = 48.533998 ± 0.000022 d. Recovered blind. New: the Gaia SB1 gives f(M) = 0.0088 $M_\odot$, a 0.25 $M_\odot$ M-dwarf companion (K_1 12.4 km/s against about 50 m/s for the putative planet), and a sixth A eclipse in s86
TOI-5149.01 = TIC 286094277 = TYC 3578-1416-1	Known TOI; reinterpreted as an EBLM	A planet candidate since 2022-02-08; the 1.00–1.11%, 9.4–9.8-h transit is the TOI signal, P = 27.37163 ± 0.00006 d. New: a 602 ± 57 ppm secondary eclipse at phase 0.697 (z = 7.1; also in SPOC, TESS-SPOC and TARS); e cos ω = 0.3128 and e sin ω = −0.093 match Gaia; f(m) = 0.0014 gives m_2 = 0.11–0.13 $M_\odot$, R_2 = 0.15 $R_\odot$, T_2 ≈ 3100 K, so the companion is a low-mass star (an EBLM) rather than a planet. Speckle imaging shows a companion at 0.086″ (Δm 1.97)
TYC 1194-680-1 = TIC 435855781 = 1SWASP J004527.74+210427.9	Known eclipser; period and type revision	Listed as EA/EB with P = 1.66 d in Thiemann et al. 2021 (SuperWASP citizen science; not in VSX). TESS gives P = 22.450574 ± 0.000054 d (Gaia SB2 22.44398 ± 0.0033, 2.0σ off); 45% and 39% eclipses; joint SB2 fit: M = 1.144 + 1.161 $M_\odot$, R = 1.93 + 2.13 $R_\odot$, e = 0.1826, ω = 57.4°. This is a revision of the period and type, not a discovery
HD 181068 and seven catalogued EBs	Known; recovered blind	HD 181068, the Kepler triply eclipsing triple, recovered blind, and seven catalogued eclipsing binaries among the first 711 stars triaged: LW Pup, TIC 188625601, four IJspeert et al. 2024 OBAF eclipsers and one Prša et al. 2022 TESS EB. The full rerun of the G < 10 subsample recovered 20 known eclipsing binaries in all
Refuted		
TIC 71350905 = TYC 4361-615-1	Refuted: systematics	The claimed long periastron eclipse was orbit-edge scattered-light ramps: sector 40's first orbit, with no conjunction within 14 d, shows the same 1% drop. The mass function (0.0025 $M_\odot$) makes the companion a ~0.2 $M_\odot$ M dwarf whose transit could be at most 0.14% deep
TIC 212431508 = TYC 5162-1158-1	Refuted: neighbour, itself an uncatalogued eclipser	Both dips (13% and 4.3%) are on Gaia DR3 4230027841458799744 (G 11.02, 35.9″, same parallax, probably a wide companion), which loses 16.1% and 5.3% while the target loses 0.6% and −0.3%; depth falls 13 → 2.2 → 0.7% as the aperture tightens. The neighbour is itself a Gaia SB1 (P 33.294 d) whose A–B spacing is 2.9σ from its own orbit: a separate lead

Object	Outcome	What the check found
TIC 17238861 = TYC 2508-895-1	Refuted: neighbour, itself an uncatalogued eclipser	The 1.9% dips are 10.5% and 9.6% V-shaped eclipses of TIC 17238865 = TYC 2508-1098-1 (*G* 12.20, 60.4″, same parallax but not bound); the target's core pixels change by no more than 0.1–0.4%. The neighbour's period is any of 732.0556/*N* d (*N* = 1–28); 48.80 d is one of these 28 aliases, the one tied to the target's orbit
TIC 265677843 = TYC 3335-389-1	Refuted: background eclipsing binary	The 7.37% dip is a 32.4% eclipse of Gaia DR3 250601205600534272 (*G* 12.06, 37″, parallax 0.85 mas, a background star); the target loses 0.01 ± 0.04%; TARS 0.83% against QLP 7.37%. Attribution to the target would also require a period 3.9σ from Gaia's
Plausible single events		
BD+16 2639 = TIC 334634917 = HIP 70251	Plausible, unconfirmed	One 0.52%, 2.50-h grazing eclipse at BTJD 2676.6639, 0.4σ from Gaia's A conjunction, identical in five pipelines and on the target's pixels, in the only TESS sector (s50). The geometry ($i \approx 86.3°$) predicts the 2.5–3.0-h duration and no secondary. An unrelated dip has a chance of a few percent of falling inside a ±0.5 d window. Next A: BTJD 4345.49, 4365.60 (±0.17)
Wolf 73 = HIP 6972 = TIC 365156806	Plausible, unconfirmed	One 2.98 ± 0.10%, 3.72-h flat-bottomed dip at BTJD 2481.9196 in SPOC 2-min data, 0.29σ from Gaia's B conjunction, on target; no other conjunction covered. A grazing occultation of a companion of at least 0.50 $M_\odot$. SPOC's own pipeline flagged the same event in 2021 at a spurious 32.139-d period. Next B: BTJD 4327.91 ± 0.47 (Gaia's period only)
HD 282670 = TYC 2392-1644-1 = TIC 366141684	Plausible, unconfirmed	One 2.10 ± 0.01%, 6.8-h eclipse at BTJD 2491.9694 in QLP and TESS-SPOC alike, on target to 1.1″ (3σ) by centroids and a difference image, with 115 other stars in the sector clean at that time; 0.06σ from Gaia's A conjunction, but **not the Gaia companion**: its mass function allows at most a 0.9% transit, and no companion of the 3.5–5.2 $R_\odot$ F primary fits the duration at any period the other sectors allow. Best read as an eclipse in an unresolved bound F/A third star carrying ≥ 6% of the light (Gaia duplicated_source, RUWE 1.22); period unknown, > 35 d if circular. SuperWASP DR1 holds a second eclipse on 2004-09-13 (–1.8% on two cameras over 3.3 h, a comparison star flat; the TESS trapezoid fitted to it gives depth 2.10% and mid-time JD 2453261.681), so the period is 6230.29/N d; the TESS non-repeat condition, SuperWASP's other nights and 2,000 KELT-North points from 2006–2008 leave 32 integer-N values between 36.9 and 188.8 d, none at Gaia's period. ASAS-SN or Gaia DR4 epoch photometry should cut these to a few, and one ground night at a predicted mid-time settles a candidate; one high-resolution spectrum would settle the third star
Unresolved		
TYC 7687-990-1 = TIC 271053559	Probable; repeated 2025 eclipses, catalogue novelty pending	In s89: a 31.33% V-shaped eclipse and two flat 5.19% and 5.51% eclipses, all on the target (difference images, TARS, clean common mode), at Gaia's 24.65-d period (*P* = 24.6548 ± 0.0018 d). However, the same ephemeris falls on covered, flat data in s8, s9 and s62, and a scan of every period from 24.58 to 24.76 d (the s89 value at ±50σ, Gaia's at ±5σ) finds, for each one, at least one of s8, s9, s35 and s62 covering the A conjunction with unflagged data and nothing deeper than 0.63% there, so the absence before 2025 is not an artefact of the one-sector ephemeris. Gaia also has a 300.8-d astrometric orbit on the star; a compact triple and nodal precession are unproved possible explanations; the earlier covered flat sectors remain a failed stability check. The current review retains a probable physical verdict and unresolved catalogue novelty. Earlier conditional A predictions: BTJD 4327.29, 4351.95 (±0.05; an unconfirmed third body could alter timing)

Table 3. Heartbeat systems: ten physically confirmed (one also eclipsing) and one candidate. The newly reviewed TIC 438089101 retains the catalogue limitations stated below. The equilibrium-tide model of Kumar et al. (1995), fitted to the light curve alone, is compared with Gaia's independent orbit.

Name	TIC	*T* (mag)	*P* (d)	*e*	Origin	Status	Note
HD 193389	51223058	7.063	20.7974 ± 0.0015	0.36	App	new	F5.5 bright giant; the model recovers Gaia's ω to 0.3° and *e* to 0.016 (3.2σ)

Name	TIC	*T* (mag)	*P* (d)	*e*	Origin	Status	Note
BD+59 2405	405590521	9.886	25.90	0.59	App	new	A2 star in Trumpler 37; *P* is Gaia's
TYC 8203-301-1	147413927	10.134	22.0440 ± 0.0005	0.68	App	new	tidally excited oscillation: 842 ppm at 23.0009 ± 0.0006 times the orbital frequency, weaker peaks at 22 and 24 times
BD+61 2067	365670788	8.653	8.216284 ± 0.000094		Fold	new	the light curve alone returns Gaia's *e* and ω
BD+37 4854	333169278	9.38	5.077463 ± 0.000021		Fold	new	Gaia's ω returned once Doppler beaming is added
CD−66 2078	302055860	9.546	5.040480 ± 0.000030		Fold	new	Gaia SB2, near-twin F stars; returns Gaia's *e* and ω
TYC 9045-989-1	350856983	9.368	4.682234 ± 0.000021	0.12	Fold	candidate	Gaia SB2, near-twin F stars; ω 20° (2.4σ) from Gaia's; largely ellipsoidal
TYC 3347-1439-1	391918838	9.860	8.9222	0.44	Batch	new	the 2.2 ppt batch "eclipse" is a 20-h tidal minimum at periastron, 0.16 d after Gaia's conjunction; Kumar model with Gaia's *e*, ω fixed, $\Delta\chi^2$ 766 over any trapezoid
HD 339939	98149548	10.107	2.402590 ± 0.000010	0.15	Batch	new	Gaia SB2, near-twin A stars; 15 ppt peak-to-peak tidal curve with its minimum at conjunction A; eclipse limit at the dip 0.4 ppt (2σ), injections recovered to 0.5 ppt; *P* 3.6σ from Gaia's 2.40241
TYC 3624-860-1	200398115	10.765	7.305664 ± 0.000049	0.21	Batch	new, eclipsing (A144)	18 ppt peak-to-peak tidal curve with periastron at phase 0.25, carrying a 12 ppt M-dwarf transit at conjunction A and its 2.1 ppt occultation at phase 0.3625; the tide alone loses by $\Delta\chi^2 >$ 11000
TYC 8588-1489-1	438089101	11.86	4.800079 ± 0.000030	0.274	Fold	confirmed; catalogue checks incomplete	Light-curve e = 0.2646 ± 0.0040; target-localized TEOs at orbital harmonics 69 and 74; no eclipse; beaming unusable as a velocity check

T: TESS magnitude from the TIC. *e*: Gaia's eccentricity where the text quotes it. HD 193389: Gaia SB1 *P* 20.7966 d. Origin as in Table A1: the first three were found by the appointments search (Section 2.5), the other four set aside by the *G* 9–10 Gaia-orbit fold (Section 2.4); Batch, the three found among the shallow candidates of the *G* 10–11.5 fold by the per-object pass (Section 3.2). None of the first nine eclipses at the fitted inclination; TYC 3624-860-1 does, and is counted as A144 in Table A1. TIC 438089101 comes from the G 12–12.25 spectroscopic-orbit fold and is also non-eclipsing. Its physical confirmation is separate from novelty: direct VSX, Gaia and SIMBAD checks and nine heartbeat-paper searches found no prior identification, but ASAS-SN, SB9 and WDS checks remain unavailable. These are bounded searches, not exhaustive priority evidence.

Table 4. Gaia DR3 orbit solutions contradicted by the eclipses: 19 of the 184 counted systems compared with a non-circular Gaia spectroscopic orbit, and A113, whose circular Gaia solution carries the wrong period. A solution is counted as contradicted when its period is spurious or wrong, or when its period or *e* cos ω differs from the eclipse value by 3σ or more of Gaia's quoted error under every treatment of its covariance.

Label	Name	Gaia P (d)	Eclipse P (d)	Fails in	Discrepancy
A9	HD 7601	13.42 (SB2)	27.0822013 ± 0.0000180	period (wrong)	$2P_{\mathrm{Gaia}}$ differs by 100σ; $(M_1 + M_2)\sin^3 i$ = 0.49 $M_\odot$, impossible for this pair
A12	HD 217492		14.0343984 ± 0.0000110	period	4.2σ (conjunction phase 5.5σ)
A21	TYC 4244-623-1		30.763077 ± 0.000013	$e\cos\omega$	3.7σ (RUWE 35)
A28	CD−32 4901	51.311 ± 0.014	51.37333 ± 0.00020	$e\cos\omega$; period	5.7–6.1σ; period 4.5σ
A29	BD−01 648	29.5888 ± 0.0053 (SB2)	29.60666 ± 0.00002	period	3.4σ
A30	BD+29 2845		27.94676 ± 0.00001	$e\cos\omega$	5.3–5.6σ
A52	TYC 5406-1905-1	40.543 ± 0.015 (SB2)	40.46953 ± 0.00005	period	4.9σ
A54	TYC 447-247-1	39 (SB2)	3.02348 ± 0.00001	period (spurious)	not an alias; would need 14 + 13.5 $M_\odot$
A63	TYC 3801-1499-1		57.2400 ± 0.0003	$e\cos\omega$	3.8σ (4.1–4.3σ from the eclipse spacing)
A65	BD+27 4098		26.768321 ± 0.000009	$e\cos\omega$	4.0–5.0σ in the eclipse spacing
A86	BD−01 2238		15.798442 ± 0.000003	period	4.1σ short (eclipse spacing 2.7σ)
A102	HD 194269		1.9385707 ± 0.0000005	period	5.0σ short
A110	TYC 7212-2039-1	22.054690 (SB2)	22.039984 ± 0.000046	period	4.5σ long, 21 min per orbit; the secondary at phase 0.506 also differs from the 0.526 that Gaia's e, ω predict
A113	TYC 4251-1144-1	37.6646 (SB2, circular)	23.63714 ± 0.00001	period	Gaia's period is the 63-d scan-cadence alias of the eclipse period (1/23.637 − 1/37.665 = 1/63.5 d); two primaries 23.637 d apart within one sector exclude it by themselves, and $e\cos\omega$ = 0.280 excludes the circular solution
A117	TYC 629-1389-1	36.6007 ± 0.0172	36.54015 ± 0.00003	period	3.5σ long, 87 min per orbit; Gaia's e and ω hold
A145	TYC 3344-341-1	4.746792 ± 0.000199	4.746808 ± 0.000003	$e\cos\omega$	the secondary eclipse at phase 0.49996 ± 0.00018 gives $e\cos\omega$ = 0.0000 ± 0.0003; Gaia's +0.036 ± 0.008 (e = 0.039) predicts 0.5231 ± 0.0051, 4.5σ away; the orbit is circular and Gaia's eccentricity spurious
A147	TYC 2136-2756-1	14.801518 ± 0.000644 (SB2)	14.804287 ± 0.000006	period	4.3σ short, 4 min per orbit; at Gaia's own epoch the eclipse is 1.06 h (2.4σ) after the predicted conjunction B, and the batch's +0.019 phase offset at TESS epochs was this period error extrapolated

Label	Name	Gaia *P* (d)	Eclipse *P* (d)	Fails in	Discrepancy
A159	TYC 6635-619-1	11.874084 ± 0.001091	11.870552 ± 0.000001	$e \cos \omega$	the secondary eclipse at phase 0.62842 ± 0.00004 gives $e \cos \omega$ = 0.2005 ± 0.0002; Gaia's +0.252 ± 0.013 (e = 0.299, ω = 32.6°) predicts 0.6603 ± 0.0081, 3.9σ away with Gaia's covariance and 3.3σ without; the period is 3.2σ long as well; Gaia's goodness of fit of 2.6 suggests underestimated errors, and the K dwarf's 15–25% of the light can bias a single-lined solution
A167	TYC 8994-1423-1	4.173881 ± 0.000092 (SB2)	4.1731519 ± 0.0000014	period	8.0σ long, 1.05 min per orbit; at Gaia's own epoch the two eclipses fall 2.9 h and 1.2 h after the predicted conjunctions A and B (0.01–0.04 in phase), so the orbit is the right one and the error is understated: a perturbed double-lined solution (goodness of fit 12.2, RUWE 2.70) on a star with a ±3 min eclipse-timing variation common to both eclipses, which suggests a third body; the secondary at phase 0.49985 ± 0.00012 also gives $e \cos \omega$ = −0.0002 ± 0.0002 against Gaia's +0.0256 ± 0.0097 (2.7σ)
A202	TYC 5995-1398-1	2.5540664 ± 0.0000374	2.553938 ± 0.000005	$e \cos \omega$; period	3.5σ in $e \cos \omega$: the occultation at phase 0.49993 ± 0.00045 gives −0.0001 ± 0.0007 against the +0.051 ± 0.015 of Gaia's e = 0.054 ± 0.015, so the orbit is circular and the eccentricity spurious; and the period is 3.4σ long, 11 s per orbit, with Gaia's conjunction A drifting from +0.1σ to +2.4σ across the TESS sectors; conditioned on a circular orbit, both conjunctions match the eclipses at Gaia's own epoch (−2.0σ and −1.6σ), so the attribution holds; the primary is a spotted RS CVn-like subgiant whose eclipse timings jitter by ±6 min

Discrepancies are in units of Gaia's quoted error. A blank Gaia period is not quoted in the text (for A30 and A65 it agrees with the eclipse period); the Supplement entries give it. Eclipse periods are those of Table A1, except for A29, where the value used in the comparison is given (Table A1: 29.606659 ± 0.000010 d). Three conjunction offsets are in tension without being counted, because they move with the treatment of Gaia's covariance: A87, 4.3σ at Gaia's epoch and 1.7σ at the TESS epoch carried with Gaia's period; A89, 6.7σ with the full covariance of a poorly fitting double-lined solution and 2.2σ with the correlations ignored; A104, 8.2σ with Gaia's epoch carried with the TESS period and 0.2σ with Gaia's own period (its eclipse spacing gives $e \cos \omega$ 2.9σ from Gaia's).

Table 5. Catalogue corrections: the eleven numbered corrections B1–B11, the further corrections from the candidate checks, and the VSX type revisions among the new eclipsing systems.

Label	TIC	Name	Catalogue entry	TESS	Note
B1	48875187	Gaia DR3 5451130921097245056	VSX RS (Gaia classifier), no period	detached twin EB, *P* 0.862264 d, equal 9.1% eclipses	held: host not separated from two possible neighbours

Label	TIC	Name	Catalogue entry	TESS	Note
B2	382523691	ASASSN-V J005242.03-513706.0	VSX ROT 5.3637 d	EB at Gaia's SB1 period, *P* 4.915053 d, 24 + 12 ppt	
B3	397233697	Gaia DR3 5223177460339438088	VSX ROT (Gaia); Gao ROT 8.40 d	EB, *P* 17.13557 d, 100 + 5 ppt	secondary at φ 0.3
B4	333792991	HAT-177-0002605	VSX VAR 9.69 d; ASAS-SN ROT:	EB, *P* 6.21467 d, 45 + 13 ppt	
B5	385557214	V1274 Tau	GCVS RS 6.62 d; VSX RS 6.448 d; ASAS-3 ESD/EC 1.727 d	EB at Gaia's *P* 5.62323 d, 4.5 + 2 ppt	ASAS-3 photometry shows only the 6.45-d rotation
B6	464315748	HD 171459	VSX DSCT|GDOR 3.19721 d	EB, *P* 6.3955 d, 12.7/13.4 ppt	3.19721 d is the eclipse spacing
B7	189269584	CD−28 7251	VSX VAR 0.0146 d	twin EB, *P* 6.9610 d, 20.7/23.9%	proposed: contamination ratio 1.77 leaves a blend possible
B8	99028319	LP 852-5	TESS-EB (Prša et al. 2022) P 9.4045943 ± 0.00045 d; not in VSX	EB, *P* 9.400436 d, 32.9/18.5 ppt	
B9	80158129	CzeV3749	VSX EA, no period	*P* 10.898518 d, no secondary	blend, 9″ from the peak
B10	120660110	WISE J105540.2-521127	VSX EW/EA 0.8152227 d	*P* 0.817052 d	blend, 11″
B11	306658481	NSV 3445	VSX suspected variable, no type or period	eclipsing, 1.66 d	1.5″; depth sector-dependent
Further corrections from the candidate checks					
	435855781	TYC 1194-680-1	SuperWASP EA/EB 1.66 d (Thiemann et al. 2021); not in VSX	EB, *P* 22.450574 ± 0.000054 d, 45% and 39%	Gaia SB2 22.44398 d, 2.0σ short
		Gaia DR3 3327090097954378752	VSX E 3.23146 d	EB, *P* 5.725588 ± 0.000018 d, circular	G 14.8, 63″ from TIC 206952704
		Gaia DR3 5956319690145167336	VSX E 12.3202 d	EB, *P* 1.5432 d	G 13.9, 53″ from TIC 34344890
VSX type revisions among the new eclipsing systems (Table A1)					
A20	235666667	HD 170829 = NSV 24480	VSX VAR:, no type or period	EB, *P* 26.384493 d	
A25	258780392	BD+12 121	VSX MISC 91.5 d	EB, *P* 43.8230 d	91.5 d is spot modulation of the giant
A31	2048935928	V1192 Cas	VSX SRA 37.4 d	EB, *P* 37.4133 d	37.4 d is the orbital period
A32	399758284	TYC 97-228-1	VSX ROT 39.81 d	EB, *P* 40.51032 d	39.81 d is the giant's near-synchronous rotation
A52	307170556	TYC 5406-1905-1	VSX MISC 79.3 d	EB, *P* 40.46953 d	

Depths are as measured in TESS (B1 in TGLC photometry). B1 and B7 are proposed and await the further step in their notes; the other nine are established by the data. For B9 and B10 the TIC star is the aperture in which the blended signal was found. The two Gaia DR3 stars of the second block are neighbours of conjunction-window targets, detected through their contamination of the targets' apertures. TYC 1194-680-1 is a confirmed eclipsing binary but not new: the revision is of its period and type. A21's only prior record is an ExoFOP CTOI that is the star's own secondary eclipse.

Table 6. Other objects: the planet candidates C1–C3, variables that are not eclipsing binaries, instructive false positives and rejected candidates, one line each; the Supplement has the details.

Object	TIC	What	Status
Planet candidates			
C1: TIC 4206066 b, c	4206066	two transiting-planet candidates (Rabtsevich 2026), *P* 3.182785 and 11.132690 d, recovered blind by the M-dwarf search	candidates; independently confirmed in public data; not our discovery
C2: TOI-5394.01 = HD 89277	61109252	the TOI signal is the occultation of an M-dwarf companion; *P* 15.195017 d is Gaia's SB1 period	explained
C3: TOI-6284, TOI-1467	286763141, 240968774	three transit signals around known hosts: 7.349277 and 5.244544 d (TOI-6284), 3.154127 d (TOI-1467)	independently recovered; in a preprint
Other objects			
GJ 4348	206327797	0.223864-d double wave; a detached near-twin M+M pair plausibly on top of a young star	open; one spectrum at quadrature settles it
TYC 3763-577-1	322347036	δ Scuti star, 15.32, 20.53 and 27.14 d^{-1}, ~1 ppt (diluted)	uncatalogued
TIC 234495456	234495456	BY Draconis spot rotator at ~0.7623 d	uncatalogued
TIC 189882802	189882802	a 3,444-ppm, 4-h single event made by two flares	dead
TIC 96202086	96202086	0.63048 d, 307 ± 32 ppm; ~40 such peaks expected over 1,299 stars	not a detection
HD 151770	337393383	real 16.3 + 11.2 ppt eclipses; $e \cos \omega$ ~7σ from Gaia's SB2 and both conjunctions miss	open
HD 34757, HD 44342, HD 196402	309416862, 38022998, 100614055	demoted fold candidates of 0.9–1.6 ppt	dead or unresolvable
EBLM J0021–16 = BD–16 52	406352710	published EB recovered blind by the *G* 9–10 fold; *P* 5.9672641 ± 0.0000012 d	known; a blind recovery
BD+30 9	22503559	ellipsoidal variable on a Gaia SB1 orbit, *P* 1.5786673 ± 0.0000010 d, 16.1 ppt peak to peak; no eclipse	new ellipsoidal variable
TYC 3707-251-1	316045876	4%, ~3-h dips every 5.854343 d; withdrawn as an eclipser on a 197-d Gaia astrometric orbit	new 5.854-d EB; triple probable, host not pixel-verified
TIC 271520202	271520202	two eclipses (18.5% and 17.3%) 4 × 192.391 d apart; Gaia 193.38 ± 0.81 d	plausible; one more primary would confirm
TIC 4382254	4382254	a lone 7,328-ppm, 0.72-h dip in sector 70	not a candidate
Marginal folds		eclipse-like folds in no catalogue	recorded, not chased

C3: all three signals are in the preprint of Tschudi (arXiv:2607.23781). TIC 316045876 and TIC 271520202 come from the edge-on astrometric sample (Section 2.6). The TIC numbers of the marginal folds and the Gaia binary suspects among bright late-M dwarfs are in the Supplement.

5. Discussion

5.1 Why the appointments method works, and its yield

Gaia's elements fix the conjunction times in advance: a single dip either lands in a window fixed beforehand or it does not. The search found 14 new eclipsing systems among 1,422 stars at G < 10 (1.0 per hundred) and 49, five of them probable, among 2,327 at G 10–11 (2.1 per hundred): 1.7 per hundred overall. At G 11–11.5 the first twenty of fifty candidates re-analysed were all eclipsing systems, 0.8 per hundred so far among 2,608 stars, with thirty candidates outstanding. At G 11–11.5 the first twenty of fifty candidates re-analysed were all eclipsing systems, 0.8 per hundred so far among 2,608 stars, with thirty candidates outstanding. At G 11–11.5 the first twenty of fifty candidates re-analysed were all

eclipsing systems, 0.8 per hundred so far among 2,608 stars, with thirty candidates outstanding. The period fold of Gaia orbits at P < 20 d gave 0.4 per hundred at G < 9 and 1.3 at G 9–10; at G 10–11.5 the batch fold gave 1.8 candidates per hundred, and 58 of the 60 shallowest candidates, and all eight of the difference-image-flagged ones, survived the per-object re-analysis. The searches were built differently and the rates are not directly comparable.

5.2 What the eclipses say about the Gaia orbits

Eclipses give the inclination and the relative radii that a spectroscopic orbit leaves free. Most systems with a Gaia orbit at the eclipse period agree with it within about 2σ. Nineteen of the 184 counted systems compared with a non-circular Gaia orbit do not, by 3σ or more in period or $e \cos \omega$ under every treatment of Gaia's covariance, and A113's circular solution carries the wrong period (Section 3.4). Among the G 9–10 systems, 13 of 31 non-circular orbits have a largest offset of 2.3σ or more where about two are expected, and Gaia's quoted errors for these orbits appear to be underestimated by a factor of about 1.5. Below $e \approx 0.05$ the eclipses also test Gaia's $e \sin \omega$, through the durations and the depth of the secondary eclipse, and the result is mixed: six orbits want a circular orbit where Gaia found a small eccentricity (A140, A145, A161, A167, A202, A203; the direction of the bias of Lucy and Sweeney 1971), two want more eccentricity than Gaia found (A137, A171; A204 leans the same way at 1.9σ), and one shows Gaia's 2.3σ eccentricity to be real at 19σ (A176). The period, the conjunction at Gaia's epoch and $e \cos \omega$ therefore carry the attribution, and $e \sin \omega$ is treated as a soft prior.

In A54 a 3.02-d eclipsing pair sits under a spurious 39-d Gaia SB2 solution. El-Badry et al. (2026) found 14 hierarchical triples of the same kind, with the eclipse period much shorter than the Gaia period, and report about half of their Gaia SB1 solutions for compact-object candidates to be spurious. The eclipsers here are a different population, but the contradicted solutions show that a Gaia orbit requires independent confirmation.

5.3 The eclipses on the Harvard plates and in ASAS-SN

The ephemerides can also be run backwards. The Harvard plates, digitised by DASCH (Grindlay et al. 2012; data release 7), cover most of the sky from 1890 to 1990 with a scatter of about 0.13 mag per plate at G 10–11, enough to show a 10–40% eclipse if enough plates fall inside the window. The 31 systems among A1–A74 and B1–B11 with an eclipse of 8% or more and G ≤ 12.5 were folded back 600–12,000 cycles, with the cuts, the statistic and a permutation null fixed before the first fold. Seventeen of the 31 show the eclipse at our ephemeris, 15 of them at the TESS period unchanged; A30's 42% eclipse is a 0.6-mag dip on 27 plates (z = 19). Two systems are untestable, six are too shallow for their plates, and six are below threshold with depths consistent with TESS. The last of these, A63, was first read as an absence: nine plates in its window give 0.066 ± 0.036 mag against the 0.21 mag peak depth of its 18% eclipse, and injected eclipses were recovered 10 times in 10. But the injections were boxes at the peak depth and the eclipse is V-shaped, so the window mean it predicts is 0.105 mag, 1.1σ from the measurement; fitted with the TESS eclipse shape, the plates allow 0.62 ± 0.29 of the TESS depth, where nine of the detected systems fitted the same way give a median of 0.95 with a scatter of 0.25 between stars. A63 is consistent with its TESS depth, not absent. The same fold, applied to the five systems among A75–A109 that meet the cuts, adds A108 (TYC 2747-436-1), whose eclipse appears on 258 in-window plates from 1899 to 1990 (z = 19, the third-strongest detection); A83 is suggestive (FAP 0.005) at a period 4 eP below the TESS value; A82 and A86 are not seen, the plates allowing 0.04 ± 0.31 and 0.28 ± 0.23 of their TESS depths; A98 is untestable. Where TESS covers one or two sectors the century of baseline can refine the period: the plates detect A37 at 22.6131 ± 0.0002 d, twenty times tighter than the TESS value, which Table A1 keeps. The century of plates also tests whether the eclipses have changed: for the 19 systems the plates detect, the TESS eclipse profile was fitted for its amplitude in 20-

year bins from 1890 to 1990, and no system changed depth at 3σ (three are at 2–2.9σ, where one is expected by chance, and each resolves into a single bad era or a single plate series), so A73 remains the only system whose eclipses have changed. Three systems (A38, A28, A16) are 1.4–2.0 times deeper on the blue-sensitive plates than in the TESS band, a colour term that constrains the temperatures of their components.

ASAS-SN V-band photometry (2014–2018; Shappee et al. 2014; Kochanek et al. 2017) gives a second archival test, independent of the plates and reaching shallower eclipses. Of the 118 objects with an ephemeris, 55 have an ASAS-SN light curve with a measured depth, a duration and an in-eclipse epoch. Folded on the ephemerides of Table A1, a backward extrapolation of 15 to 630 cycles into data that had no part in deriving them, and tested against a null that slides the eclipse window to 10,000 random phases, 20 of the 55 are fainter in eclipse at $p < 0.05$, where 2.75 are expected by chance, and six at $p < 0.01$ (A6, A33, A35, A38, A39, A47), where 0.55 are expected. Seven of the 20 are also on the plates (A4, A6, A33, A38, A47, A55, A56); ASAS-SN recovers 13 that the plate fold could not, four of which (A22, A50, A65, A70) the plates tested and did not detect. For the 11 systems detected under a shuffle-permutation null, measured and predicted depths agree (median ratio 1.04, Pearson $r = 0.785$, permutation $p = 0.003$).

ASAS-3 V photometry (2000–2009; Pojmański 2002) fills the gap between the plates and ASAS-SN for the 49 objects south of declination +28°, 34 of them with usable light curves. Folded on the same ephemerides with the same sliding-window null, nine are fainter in eclipse at $p < 0.01$ (A6, A10, A22, A28, A45, A47, A57, A58, A66), and A54 is a tenth once its window is narrowed to its 2-h eclipse; A47 measures 0.456 mag against 0.451 predicted, 193 cycles back. Five of the ten (A10, A45, A57, A58, A66) have no other non-TESS recovery, and A57's excess depth may be the giant's own variability. Among A75–A109, 13 of the 19 systems with usable curves are detected, out to 4,707 cycles (A106). Calibrating the sliding-window null on the light curves themselves, with 440,000 fake ephemerides placed away from the eclipses, gives a false-positive rate of 0.006 at $p < 0.01$ and 0.042 at $p < 0.05$, so these counts and the ASAS-SN counts are, if anything, conservative.

5.4 Limits

The completeness of the appointments search was measured by injection: trapezoid primary eclipses at every conjunction of an ephemeris drawn from Gaia's own errors, into the light curves of 300 of the searched stars, run through the search and the pre-set cuts unchanged (7,200 injections). Where TESS covers an eclipse (94% of injections), the cuts pass 79% of eclipses of 5–10% depth, 67–74% at 2%, 50–69% at 1% and 33–56% at 0.5%, the longer durations recovered better; at every depth down to 0.3% the eclipse is in its window at SNR ≥ 8 in 97–99% of covered injections, so the losses are in the cuts. Most of the loss is one defect: the random windows of the null test were allowed to overlap the star's other predicted windows, so a multi-sector eclipser could be scored against its own eclipses and rejected. The reported systems do not depend on it (54 of the 55 pass the cuts with the null corrected, the other was found by eye), but re-running the search with the null corrected passes 343 further stars that the original cuts rejected, which are being examined. The first of them, TYC 7212-2039-1 (TIC 158544874, a Gaia SB2 at 22.05 d), is confirmed and counted as A110: 13–30% eclipses of both kinds in seven sectors, on the star by difference imaging, with an eclipse period of 22.039984 ± 0.000046 d, 21 min per orbit shorter than Gaia's (4.5σ). Eclipses longer than the 14-h maximum box and the false-alarm rate are not measured. Seven of the 54 reported systems were admitted on the remaining pre-set cuts without the z test. The conjunction test inherits Gaia's period: A9 passed it at Gaia's 13.42 d, and its true period is 27.08 d. The literature check (Section 2.7) read the titles of the documents the NASA ADS full-text

index returned, so a mention in the body or a table of an unrelated paper, or under a name not queried, could be missed, and it did not include Zenodo, which is how the records of Section 1 were missed. No spectroscopy was used. The sample stops at G 11.

5.5 Predicted eclipses and follow-up

Appendix A gives the next eclipses, in geocentric UTC, for the original 199 A-labelled catalogue-screened systems except A3. Reports for the Appendix C systems give conditional predictions where their timing baseline permits; these should not all be assumed to have minute-level precision. Most predictions are good to a few minutes. Most of the stars are G 9–11, with eclipses 1–40% deep lasting hours, so many are within reach of small telescopes. The TESS pipelines reduce the same pixels; an eclipse seen from the ground at the predicted time is the independent observation. A missed eclipse shows an ephemeris error, and an early or late one measures eclipse-timing variations (hinted at, for example, in A47). Timings of both eclipses measure $e \cos \omega$ and so test the contradicted Gaia orbits; depths of A73 would show whether its fade continues; timings of A95 would test the light-travel time. HIP 1060 (A47), with an annular eclipse on 2026 November 2 at 06:37 UTC, has been submitted to the AAVSO VSX and is pending.

The catalogue is bright (median T = 9.6). Every object saturates ZTF, the 27 with no ASAS-SN source are 0.81 mag brighter in the median than those with one, and ASAS-SN's scatter rises from 0.02 mag at V 11–12 to 0.08–0.09 mag brighter than V 9.5. For about a quarter of these systems a small telescope is the only instrument left that can follow them.

5.6 The null planet search

The transit searches of about 3,100 nearby M dwarfs (Section 2.3) found no new planet. Under the estimator's assumptions, the repeated-transit searches were expected to yield about 1.2 detections, with a probability of none of 0.30 and a range of about 1 to 8 allowed by those assumptions; the single-transit search, which the TOI list does not cover, expected about 3 more and is the only one with a real shortfall (2.5σ). A naive occurrence-rate calculation (Dressing and Charbonneau 2015) gives 75–95. The difference lies in the sample. Every search but the one around known hosts excluded stars with a TOI or CTOI; the cut removed 2.2% of the stars but all 41 known hosts of a detectable 0.5–15 d planet in this population (95% upper bound on the surviving fraction 7.3%). The searches recovered both TIC 4206066 candidates blind and showed TOI-5394.01 to be the occultation of an M-dwarf companion (Supplement, S3.7).

6. Conclusions

1. We retain 921 accepted eclipsing-system claims in public TESS photometry: 808 confirmed and 113 probable. No prior eclipsing-binary identification was found for these claims in the catalogues searched (Section 2.7); 29 orbit-informed plausible cases remain excluded. The prior 913-claim baseline removed 17 prior-known systems from 917 earlier public claims, added 13 already accepted private systems, and recorded the A74 and C57 tier updates; eight newly reviewed claims now raise it to 921. Scoped catalogue non-matches do not establish exhaustive literature priority.
2. One hundred seven came from eclipses by appointment. Among 3,749 stars at G < 11 it found about 1.7 new eclipsing systems per hundred stars, and a third subsample of 2,608 stars at G 11–11.5 contributes 42 confirmed systems, with five candidates unreviewed. The fresh exploratory appointments pilot adds two confirmed systems; across these populations, 102 systems are confirmed and five probable.

3. Gaia-orbit period folds contribute 798 systems, including the fainter spectroscopic-binary population; four systems come from the nearby-M-dwarf-selected search (including neighbouring hosts), four from near-edge-on Gaia astrometric orbits, and eight from the mixed-method period-free follow-up. The same physical and catalogue criteria apply across these populations. Known recoveries, plausible signals, non-eclipsing tidal variability and contaminating sources remain visible in the evidence register and are excluded from unique additions.
4. The eclipses contradict twenty Gaia DR3 orbit solutions in period or $e \cos \omega$.
5. Ten heartbeat systems are physically confirmed, one also eclipsing, and one further system is a candidate. TIC 438089101 adds target-localized tidal oscillations at harmonics 69 and 74, with its beaming interpretation withdrawn and supplementary catalogue checks still incomplete. We also report three probable blends and eleven catalogue corrections.
6. For 17 of 31 deep systems, Harvard plates from 1890–1990 show the eclipse at our ephemeris, and ASAS-SN photometry from 2014–2018 shows it for 20 of 55 systems tested, with the depths matching the predictions in the 11 examined.
7. A transit search of about 3,100 nearby M dwarfs found no new planet where about 1.2 detections were expected under the estimator's assumptions (probability of zero 0.30), because removing known planet candidates removed most stars where a detectable planet is known. Both TIC 4206066 candidates were recovered, and TOI-5394.01 is the occultation of an M-dwarf companion.
8. Physical confirmation among selected candidates. At G = 11.5-12 (actual DR3 range 11.6764-11.9742), 27/30 are physically confirmed (90.0%; descriptive Wilson 95% interval 74.4-96.5%), 3 probable and 0 plausible; 30 remain accepted unique additions. At G = 12-12.5 (actual DR3 range 12.0029-12.4879), 21/30 are physically confirmed (70.0%; descriptive Wilson 95% interval 52.1-83.3%), 8 probable and 1 plausible; 26 remain accepted unique additions after two further prior-known recoveries. Every selected target remains in its denominator, including any unavailable or execution-incomplete outcome; all 60 completed the bounded review. TIC 377389892 is a confirmed known VSX EA recovery (0.02 arcsec counterpart), included in physical confirmation and excluded from new additions. Probable and plausible outcomes remain separate; plausible cases add nothing. These descriptive confirmation fractions concern remaining screened candidates, not blind discovery yield or completeness. Wilson intervals are binomial model summaries despite finite historical strata. Case mix, prior depletion of the bright stratum below G = 11.663, and selection bookkeeping prevent a causal magnitude interpretation.
9. Reserved and cumulative magnitude comparison. The second reserved sixty: At G = 11.5-12 (actual DR3 range 11.6648-11.9955), 29/30 are physically confirmed (96.7%; descriptive Wilson 95% interval 83.3-99.4%), 1 probable and 0 plausible; 30 current accepted unique additions. At G = 12-12.5 (actual DR3 range 12.0036-12.4872), 26/30 are physically confirmed (86.7%; descriptive Wilson 95% interval 70.3-94.7%), 2 probable and 2 plausible; 28 current accepted unique additions. Both rounds together: At G = 11.5-12 (actual DR3 range 11.6648-11.9955), 56/60 are physically confirmed (93.3%; descriptive Wilson 95% interval 84.1-97.4%), 4 probable and 0 plausible; 60 current accepted unique additions. At G = 12-12.5 (actual DR3 range 12.0029-12.4879), 47/60 are physically confirmed (78.3%; descriptive Wilson 95% interval 66.4-86.9%), 10 probable and 3 plausible; 54 current accepted unique additions. All 120 selected targets completed the bounded review and remain in their selected denominators. Known recoveries enter physical confirmation but add nothing; probable and plausible outcomes remain separate. The first-round results above are retained unchanged. These are descriptive remaining-screened candidate comparisons, not blind discovery yield, completeness or a causal magnitude effect. Wilson intervals are binomial model summaries despite finite historical strata; case mix, depletion, crowding, coverage and selection bookkeeping limit interpretation.

10. Completion of the bright screened stratum. The remaining 73 screened Gaia spectroscopic-orbit period-fold candidates at actual Gaia DR3 G = 11.5-12 (range 11.6578-11.9967) completed the same bounded review: 69 confirmed, 3 probable and 1 plausible; 69 remain accepted unique additions (66 confirmed and three probable) after three prior-known recoveries. This follow-up completes that remaining bright stratum and is separate from the fixed balanced 120-target magnitude comparison, whose sample and descriptive results remain unchanged. Historical inventory-boundary exclusions that meet the actual DR3 interval retain their original substitution provenance. No targets were substituted in this follow-up. The screened inventory has now completed individual review. Five weaker conjunction candidates, 58 lower-score candidates and 48 unexecuted exploratory appointment targets are separate populations. These outcomes do not measure blind discovery yield or completeness.
11. Brightest remaining screened period folds. We reviewed the next 130 candidates in increasing actual Gaia DR3 G (12.0036-12.3371), using the same physical, attribution and catalogue checks defined in Section 2.7. Of these, 103 are physically confirmed, 24 probable and three plausible. Six systems are now known recoveries, so 121 accepted unique additions remain: 99 confirmed and 22 probable; plausible signals remain excluded. These candidate reviews extend the prior G 11.5-12 completion and are separate from the fixed balanced 120-target magnitude comparison. The final 136 candidates in this screened inventory have now completed individual review. Five weaker conjunction appointments, 58 lower-score survivors and 48 unexecuted exploratory pilot targets belong to separate selections. This selected follow-up does not estimate blind discovery yield or completeness.
12. Completion of the screened period-fold inventory. The final 136 screened candidates, reviewed in increasing actual Gaia DR3 G (12.337704-12.742949), comprise 101 physically confirmed systems, 34 probable systems and one plausible signal. Three systems, including TIC 295771108, are now known recoveries and excluded from additions. The 132 accepted unique additions comprise 100 confirmed and 32 probable systems. TIC 290749976 remains plausible because eclipse morphology and source attribution are insecure. This completes this screened inventory, while five weaker conjunction appointments, 58 lower-score survivors and 48 unexecuted exploratory pilot targets remain separate selections. The fixed balanced 120-target comparison is unchanged. These selected outcomes do not estimate blind discovery yield or completeness.
13. A separate mixed-method harvest includes period-free dips and complementary leads. The 350 completed reviews comprise 28 confirmed, 36 probable, 65 plausible, 211 refuted screening signals, 5 non-eclipse variables and 5 unresolved outcomes. Among the 64 physically confirmed or probable cases, 13 are known recoveries, 43 retain unknown catalogue novelty, and eight scoped nonmatches passed claim review and contribute one confirmed and seven probable additions (R725–R732). The 43 pending cases remain outside the 921-system accepted claim count. Outcome-informed selection does not estimate occurrence or completeness.

HIP 1060 (A47) has been submitted to the AAVSO VSX and is pending; submissions for the others are in preparation.

7. Contributions and AI use

The author posed the question, directed the work, commissioned and read an internal adversarial review of the draft, decided what is claimed and at what status, and takes responsibility for the content of this paper. The searches, measurements, checks and text were carried out by AI language-model agents using local and cloud execution: Claude Opus 5.5, Claude Opus 5, Claude Fable 5 and Claude Fable 5.1 (Anthropic 2026), and GPT-5.6, GPT-6 and GPT-6.1 (OpenAI 2026). In outline, Claude Opus 5.5 designed and ran the M-dwarf transit searches and the first Gaia-orbit fold, explained TOI-5394.01,

and carried out the independent re-analysis of every candidate of the appointments search, the G 9–10 fold and the edge-on search, and the review of the G 10–11.5 candidates; Claude Opus 5 ran the late-M searches, the expected-yield analysis of Section 4.6, the live catalogue check and the ASAS-SN archival fold of Section 4.3; Claude Fable 5 ran the scaled Gaia-orbit fold and its continuation to G 11.5; Claude Fable 5.1 devised and ran the appointments method, folded the Harvard plates and assembled the paper; GPT-5.6 ran the edge-on astrometric-orbit search and the look-elsewhere test for TIC 4206066; GPT-6 (Astra) supervised the 174-object follow-up campaign, reviewed exceptional cases, reconciled the evidence and integrated the results; fresh GPT-6.1 Sol leads and GPT-6.1 Terra helpers carried out the detailed photometry, covariance, catalogue and attribution checks. Fable 5.1 led detailed period-fold reviews with Opus 5.5 helpers, preserving report-specific caveats and separating physical evidence from catalogue novelty. These candidates were originally found by Fable 5's scaled Gaia-orbit search. The project coordinator thanks all contributors who assisted with screening, evidence collection, analysis, review and reconciliation. Per-object credits are recorded in the Supplement and repository review register (Appendices B and C).

The adversarial review was run by a separate group of agents with its own code and no access to the agents writing the paper. It re-derived about 2,500 of the paper's numbers from the analysis files and re-reduced eight systems from the raw light curves; a second review of the same kind read the 35 systems added afterwards and re-measured eight of them. Both were internal quality control with independently written code, not journal peer review. Only public data were used; no telescope time, spectroscopy or proprietary data were involved. The use of AI systems is disclosed here and in the Acknowledgements, as the Journal's editorial policy asks.

The completed fainter SB1 review used Sol/Terra measurements under Astra supervision, following original Fable discovery and screening. The project coordinator coordinated the work; agent measurements are attributed to their performing systems.

8. Acknowledgements

This paper includes data collected by the TESS mission, which are publicly available from the Mikulski Archive for Space Telescopes (MAST); funding for the TESS mission is provided by NASA's Science Mission Directorate. We used the TESS light curves and full-frame-image products of the MIT Quick-Look Pipeline (QLP), the Science Processing Operations Center (SPOC) at NASA Ames Research Center and its TESS-SPOC full-frame-image products, TGLC, eleanor and GSFC-ELEANOR-LITE, the TESS Asteroseismic Science Operations Center (TASOC), CDIPS, TARS, T16 and TEQUILA, and the TESScut service; we thank the teams that built and maintain them. This work has made use of data from the European Space Agency (ESA) mission Gaia, processed by the Gaia Data Processing and Analysis Consortium (DPAC); funding for the DPAC has been provided by national institutions, in particular the institutions participating in the Gaia Multilateral Agreement. This research has made use of the VizieR catalogue access tool and the SIMBAD database, operated at CDS, Strasbourg, France; of the Exoplanet Follow-up Observation Program (ExoFOP) website and the NASA Exoplanet Archive, operated by the California Institute of Technology under contract with NASA; of the International Variable Star Index (VSX) database, operated at the AAVSO, Cambridge, Massachusetts, USA; and of the DASCH digitized Harvard plate archive. The analysis used NumPy (Harris et al. 2020), Astropy (Astropy Collaboration et al. 2022) and Matplotlib (Hunter 2007).

The analysis, the checks and the text of this paper were produced by AI language-model agents (Anthropic 2026; OpenAI 2026) directed by the author, as set out in Section 6. We thank P. Rabtsevich, whose public report of the TIC 4206066 candidates prompted the checks of Section 3.6.

The fainter spectroscopic-binary review used Sol/Terra measurements under Astra supervision, following original Fable discovery and screening. The project coordinator coordinated the work; agent measurements are attributed to their performing systems.

We thank all contributors who assisted with screening, evidence collection, analysis, review and reconciliation.

References

Anthropic 2026, Claude Opus 5, Claude Opus 5.5, Claude Fable 5 and Claude Fable 5.1, language models used through Claude Code, https://www.anthropic.com/claude, accessed July–October 2026.

Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, Astrophys. J., 935, 167 (DOI: 10.3847/1538-4357/ac7c74).

Bashi, D., Shahaf, S., Mazeh, T., et al. 2022, Mon. Not. Roy. Astron. Soc., 517, 3888 (DOI: 10.1093/mnras/stac2928).

Beck, P. G., Grossmann, D. H., Steinwender, L., et al. 2024, Astron. Astrophys., 682A, 7 (DOI: 10.1051/0004-6361/202346810).

Borkovits, T., Hajdu, T., Sztakovics, J., et al. 2016, Mon. Not. Roy. Astron. Soc., 455, 4136 (DOI: 10.1093/mnras/stv2530).

Bouma, L. G., Hartman, J. D., Bhatti, W., et al. 2019, Astrophys. J., Suppl. Ser., 245, 13 (DOI: 10.3847/1538-4365/ab4a7e) (CDIPS HLSP, DOI: 10.17909/t9-ayd0-k727).

Boyle, A. W., Bouma, L. G., and Mann, A. W. 2026, Astrophys. J., Suppl. Ser., 284, 75 (DOI: 10.3847/1538-4365/ae6657) (TARS HLSP, DOI: 10.17909/2sff-fn29).

Bryant, E. M., Bayliss, D., and Van Eylen, V. 2023, Mon. Not. Roy. Astron. Soc., 521, 3663 (DOI: 10.1093/mnras/stad626).

Caldwell, D. A., Tenenbaum, P., Twicken, J. D., et al. 2020, Res. Notes AAS, 4, 201 (DOI: 10.3847/2515-5172/abc9b3) (TESS-SPOC HLSP, DOI: 10.17909/t9-wpz1-8s54).

Callahan, J., Rowan, D. M., Kochanek, C. S., et al. 2025, Open J. Astrophys., 8, arXiv:2506.14869 (DOI: 10.33232/001c.142226).

Chen, X., Wang, S., Deng, L., et al. 2020, Astrophys. J., Suppl. Ser., 249, 18 (DOI: 10.3847/1538-4365/ab9cae).

Chopard-Vilhem, T. 2026a, Eclipsing spectroscopic binaries from Gaia DR3 orbits and TESS: survey data products (93 systems at periods of 13–119 days), Zenodo, deposited 2026 September 27 (DOI: 10.5281/zenodo.22998328).

Chopard-Vilhem, T. 2026b, Eclipsing binaries at the conjunctions predicted by near-edge-on Gaia DR3 astrometric orbits: data, predictions and code, Zenodo, version 1.2.0 (DOI: 10.5281/zenodo.23405354).

Czavalinga, D. R., Mitnyan, T., Rappaport, S. A., et al. 2023a, Astron. Astrophys., 670A, 75 (DOI: 10.1051/0004-6361/202245300).

De Elías, J. 2025, J. Amer. Assoc. Var. Star Obs., 53, 3.

Drake, A. J., Graham, M. J., Djorgovski, S. G., et al. 2014, Astrophys. J., Suppl. Ser., 213, 9 (DOI: 10.1088/0067-0049/213/1/9).

Dressing, C. D., and Charbonneau, D. 2015, Astrophys. J., 807, 45 (DOI: 10.1088/0004-637X/807/1/45).

El-Badry, K., Latham, D. W., Rix, H.-W., et al. 2026, Open J. Astrophys., 9, arXiv:2608.06453 (DOI: 10.33232/001c.172274).

Eyer, L., Audard, M., Holl, B., et al. 2023, Astron. Astrophys., 674A, 13 (DOI: 10.1051/0004-6361/202244242).

Feinstein, A. D., Montet, B. T., Foreman-Mackey, D., et al. 2019, Publ. Astron. Soc. Pacific, 131, 094502 (DOI: 10.1088/1538-3873/ab291c) (eleanor HLSP, DOI: 10.17909/t9-ncv5-bb52).

Fuller, J. 2017, Mon. Not. Roy. Astron. Soc., 472, 1538 (DOI: 10.1093/mnras/stx2135).

Fuller, J., and Lai, D. 2012, Mon. Not. Roy. Astron. Soc., 420, 3126 (DOI: 10.1111/j.1365-2966.2011.20237.x).

Gaia Collaboration, Arenou, F., Babusiaux, C., et al. 2023, Astron. Astrophys., 674A, 34 (DOI: 10.1051/0004-6361/202243782).

Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2023, Astron. Astrophys., 674A, 1 (DOI: 10.1051/0004-6361/202243940).

Gill, S., Cooke, B. F., Bayliss, D., et al. 2020, Mon. Not. Roy. Astron. Soc., 495, 2713 (DOI: 10.1093/mnras/staa1248).

Gosset, E., Damerdji, Y., Morel, T., et al. 2025, Astron. Astrophys., 693A, 124 (DOI: 10.1051/0004-6361/202450600).

Grindlay, J., Tang, S., Los, E., and Servillat, M. 2012, in New Horizons in Time-Domain Astronomy, IAU Symp. 285, 29 (DOI: 10.1017/S1743921312000166) (DASCH, https://dasch.cfa.harvard.edu, data release 7).

Guinan, E. F., Engle, S. G., and Devinney, E. J. 2012, J. Amer. Assoc. Var. Star Obs., 40, 467.

Halbwachs, J.-L., Pourbaix, D., Arenou, F., et al. 2023a, Astron. Astrophys., 674A, 9 (DOI: 10.1051/0004-6361/202243969).

Halbwachs, J.-L., Kiefer, F., and Lebreton, Y. 2023b, in SF2A-2023: Proc. Annual Meeting of the French Society of Astronomy and Astrophysics, 191.

Han, T., and Brandt, T. D. 2023, Astron. J., 165, 71 (DOI: 10.3847/1538-3881/acaaa7) (TGLC HLSP, DOI: 10.17909/610m-9474).

Handberg, R., Lund, M. N., White, T. R., et al. 2021, Astron. J., 162, 170 (DOI: 10.3847/1538-3881/ac09f1) (TASOC HLSP, DOI: 10.17909/t9-4smn-dx89).

Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, Nature, 585, 357 (DOI: 10.1038/s41586-020-2649-2).

Hartman, J. D., Bakos, G. Á., Bouma, L. G., et al. 2025, Publ. Astron. Soc. Pacific, 137, 024501 (DOI: 10.1088/1538-3873/adad42) (T16 HLSP, DOI: 10.17909/8nxx-tw70).

Hawthorn, F., Gill, S., Bayliss, D., et al. 2024, Mon. Not. Roy. Astron. Soc., 528, 1841 (DOI: 10.1093/mnras/stad3783).

Heinze, A. N., Tonry, J. L., Denneau, L., et al. 2018, Astron. J., 156, 241 (DOI: 10.3847/1538-3881/aae47f).

Huang, C. X., Vanderburg, A., Pál, A., et al. 2020, Res. Notes AAS, 4, 204 (DOI: 10.3847/2515-5172/abca2e) (QLP HLSP, DOI: 10.17909/t9-r086-e880).

Hunter, J. D. 2007, Comput. Sci. Eng., 9, 90 (DOI: 10.1109/MCSE.2007.55).

IJspeert, L. W., Tkachenko, A., Johnston, C., et al. 2021, Astron. Astrophys., 652A, 120 (DOI: 10.1051/0004-6361/202141489).

IJspeert, L. W., Tkachenko, A., Johnston, C., and Aerts, C. 2024, Astron. Astrophys., 691A, 242 (DOI: 10.1051/0004-6361/202450507).

Irwin, J. B. 1952, Astrophys. J., 116, 211 (DOI: 10.1086/145604).

Jayasinghe, T., Kochanek, C. S., Stanek, K. Z., et al. 2018, Mon. Not. Roy. Astron. Soc., 477, 3145 (DOI: 10.1093/mnras/sty838).

Jenkins, J. M., Twicken, J. D., McCauliff, S., et al. 2016, Proc. SPIE, 9913, 99133E (DOI: 10.1117/12.2233418).

Kane, S. R., Collier Cameron, A., Horne, K., et al. 2005, Mon. Not. Roy. Astron. Soc.,364, 1091 (DOI: 10.1111/j.1365-2966.2005.09656.x).

Kirk, B., Conroy, K., Prša, A., et al. 2016, Astron. J., 151, 68 (DOI: 10.3847/0004-6256/151/3/68).

Kochanek, C. S., Shappee, B. J., Stanek, K. Z., et al. 2017, Publ. Astron. Soc. Pacific, 129, 104502 (DOI: 10.1088/1538-3873/aa80d9).

Kostov, V. B., Powell, B. P., Fornear, A. U., et al. 2025, Astrophys. J., Suppl. Ser., 279, 50 (DOI: 10.3847/1538-4365/ade2d8).

Kovács, G., Zucker, S., and Mazeh, T. 2002, Astron. Astrophys., 391, 369 (DOI: 10.1051/0004-6361:20020802).

Kumar, P., Ao, C. O., and Quataert, E. J. 1995, Astrophys. J., 449, 294 (DOI: 10.1086/176055).

Kunimoto, M., Huang, C., Tey, E., et al. 2021, Res. Notes AAS, 5, 234 (DOI: 10.3847/2515-5172/ac2ef0).

Laycock, S., Tang, S., Grindlay, J., et al. 2010, Astron. J., 140, 1062 (DOI: 10.1088/0004-6256/140/4/1062).

Li, M.-Y., Qian, S.-B., Zhou, A.-Y., et al. 2024a, Mon. Not. Roy. Astron. Soc., 534, 281 (DOI: 10.1093/mnras/stae2057).

Lucy, L. B., and Sweeney, M. A. 1971, Astron. J., 76, 544 (DOI: 10.1086/111159).

Merle, T., Jorissen, A., Alexandre, S., et al. 2026, Mon. Not. Roy. Astron. Soc., 547, stag351 (DOI: 10.1093/mnras/stag351).

Mowlavi, N., Holl, B., Lecoeur-Taïbi, I., et al. 2023, Astron. Astrophys., 674A, 16 (DOI: 10.1051/0004-6361/202245330).

Ochsenbein, F., Bauer, P., and Marcout, J. 2000, Astron. Astrophys., Suppl. Ser., 143, 23 (DOI: 10.1051/aas:2000169).

Ogunwale, B. B., Zaguri, Y., Perdelwitz, V., et al. 2025, Astron. Astrophys., 704A, 317 (DOI: 10.1051/0004-6361/202555521) (TEQUILA HLSP, DOI: 10.17909/pj9v-7t21).

OpenAI 2026, GPT-5.6 and GPT-6, language models used through Codex, https://openai.com/, accessed July–October 2026.

Osborn, H. P., Armstrong, D. J., Brown, D. J. A., et al. 2016, Mon. Not. Roy. Astron. Soc., 457, 2273 (DOI: 10.1093/mnras/stw137).

Paegert, M., Stassun, K. G., Collins, K. A., et al. 2021, arXiv e-prints, arXiv:2108.04778 (https://arxiv.org/abs/2108.04778).

Pojmański, G. 2002, Acta Astron., 52, 397.

Powell, B. P., Kruse, E., Montet, B. T., et al. 2022, Res. Notes AAS, 6, 111 (DOI: 10.3847/2515-5172/ac74c4) (GSFC-ELEANOR-LITE HLSP, DOI: 10.17909/j2yt-t417).

Prša, A., Kochoska, A., Conroy, K. E., et al. 2022, Astrophys. J., Suppl. Ser., 258, 16 (DOI: 10.3847/1538-4365/ac324a).

Rabtsevich, P. 2026, report of two transiting-planet candidates around TIC 4206066: thread on X (https://x.com/p_rabtsevich/status/2107803696058822742) and write-up (https://tic4206066.pages.dev), both accessed 9 October 2026.

Rappaport, S., Deck, K., Levine, A., et al. 2013, Astrophys. J., 768, 33 (DOI: 10.1088/0004-637X/768/1/33).

Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, J. Astron. Telesc. Instrum. Syst., 1, 014003 (DOI: 10.1117/1.JATIS.1.1.014003).

Rimoldini, L., Holl, B., Gavras, P., et al. 2023, Astron. Astrophys., 674A, 14 (DOI: 10.1051/0004-6361/202245591).

Romanov, F. D. 2021, J. Amer. Assoc. Var. Star Obs., 49, 130 (https://arxiv.org/abs/2111.11086).

Rowan, D. M., Jayasinghe, T., Stanek, K. Z., et al. 2023, Mon. Not. Roy. Astron. Soc., 523, 2641 (DOI: 10.1093/mnras/stad1560).

Samolyk, G. 2022, J. Amer. Assoc. Var. Star Obs., 50, 255.

Samus', N. N., Kazarovets, E. V., Durlevich, O. V., et al. 2017, Astron. Rep., 61, 80 (DOI: 10.1134/S1063772917010085).

Schanche, N., Collier Cameron, A., Almenara, J. M., et al. 2019, Mon. Not. Roy. Astron. Soc., 488, 4905 (DOI: 10.1093/mnras/stz2064).

Schmidt, S. P., Schlaufman, K. C., Ding, K., et al. 2023, Astron. J., 166, 225 (DOI: 10.3847/1538-3881/ad0135).

Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, Astrophys. J., 788, 48 (DOI: 10.1088/0004-637X/788/1/48).

Shporer, A., Fuller, J., Isaacson, H., et al. 2016, Astrophys. J., 829, 34 (DOI: 10.3847/0004-637X/829/1/34).

Solanki, S., Cieplak, A. M., Schnittman, J., et al. 2025, Astrophys. J., Suppl. Ser., 276, 17 (DOI: 10.3847/1538-4365/ad8a62).

Southworth, J. 2015, in Living Together: Planets, Host Stars and Binaries, ASP Conf. Ser. 496, Astron. Soc. Pacific, San Francisco, 164 (https://arxiv.org/abs/1411.1219).

Stassun, K. G., Oelkers, R. J., Paegert, M., et al. 2019, Astron. J., 158, 138 (DOI: 10.3847/1538-3881/ab3467).

Tang, S., Grindlay, J., Los, E., and Servillat, M. 2013, Publ. Astron. Soc. Pacific, 125, 857 (DOI: 10.1086/671760).

Tarrants, T., and Mendes, E. 2023, arXiv e-prints, arXiv:2310.12380 (https://arxiv.org/abs/2310.12380).

Thiemann, H. B., Norton, A. J., Dickinson, H. J., et al. 2021, Mon. Not. Roy. Astron. Soc., 502, 1299 (DOI: 10.1093/mnras/stab140).

Thompson, S. E., Everett, M., Mullally, F., et al. 2012, Astrophys. J., 753, 86 (DOI: 10.1088/0004-637X/753/1/86).

Triaud, A. H. M. J., Martin, D. V., Ségransan, D., et al. 2017, Astron. Astrophys., 608A, 129 (DOI: 10.1051/0004-6361/201730993).

Tschudi, Y. 2026, arXiv e-prints, arXiv:2607.23781 (https://arxiv.org/abs/2607.23781).

Watson, C. L., Henden, A. A., and Price, A. 2006, The Society for Astronomical Sciences 25th Annual Symposium on Telescope Science, The Society for Astronomical Sciences, Rancho Cucamonga, CA, 47.

Welsh, W. F., Orosz, J. A., Aerts, C., et al. 2011, Astrophys. J., Suppl. Ser., 197, 4 (DOI: 10.1088/0067-0049/197/1/4).

Wenger, M., Ochsenbein, F., Egret, D., et al. 2000, Astron. Astrophys., Suppl. Ser., 143, 9 (DOI: 10.1051/aas:2000332).

Williams, P. K. G. 2025, arXiv e-prints, arXiv:2501.12977 (https://arxiv.org/abs/2501.12977) (DASCH Data Release 7).

Yamaguchi, N., El-Badry, K., and Sorabella, N. 2024, Publ. Astron. Soc. Pacific, 136, 124202 (DOI: 10.1088/1538-3873/ad9955).

Zhang, H.-T., Qian, S.-B., Liao, W.-P., et al. 2026, Astrophys. J., 1008, 32 (DOI: 10.3847/1538-4357/ae8e79).

Appendix A. Ephemerides and next eclipses

Table A1 gives, for the A-labelled confirmed and probable systems, the period and epoch from the eclipses, the eclipse depths and the primary duration where measured, and the next predicted primary eclipse after the date of this paper (none for A3, which has no timing fit), in geocentric UTC (light-travel time to the barycentre removed). The R-labelled, returned C-labelled and period-free reviews are registered in Appendix C; their reported ephemerides and limitations are in the review registers and reports. The next secondary eclipses for the A-labelled systems are in the Supplement. Epochs are BJD_{TDB}. The per-object entries in the Supplement give the measurements behind each row. Table A1 preserves measured historical ephemerides; the reconciled claim register supersedes its earlier count flags. A74 is now confirmed; A103, A125 and A157 remain measured physical systems but are prior-known recoveries excluded from the active A-claim count. Fourteen other prior-known removals occur in R-labelled reviews. C57 is now probable. The current claim flags are in the reconciled register and ledger.

Table A1. Catalogue of the 199 eclipsing systems (187 confirmed, 12 probable) in label order, with the by-catch variable A46 and the previously typed A93, which are not counted, and the three probable eclipsing binaries with unresolved hosts (A13–A15) in a separate block: ephemerides, depths and the next primary eclipse after 2026 October 14 (after 2026 October 19 for A119–A159 after 2026 October 21 for A160–A179 and after 2026 October 22 for A180–A204).

The catalogue is printed in four panels, joined by object label. All columns and notes from Table A1 are retained.

Table A1a. Identification and classification.

Label	TIC	Name	*T* (mag)	Origin	Status
A1	311958594	UCAC4 083-054218	9.822	MD	confirmed
A2	183494856	UCAC4 070-000968	14.810	MD	confirmed
A3	85379511	HIP 83929 = G 181-29	9.656	MD	probable
A4	70665829	UCAC4 537-050325	12.294	MD	confirmed
A5	22388285	HD 106770	8.982	Fold	probable
A6	348773952	BD−13 1046	9.596	Fold	probable
A7	271321097	UCAC4 130-138342	10.930	Fold	probable
A8	402800288	HD 131614	8.470	Fold	confirmed
A9	399009076	HD 7601	8.557	Fold	confirmed
A10	300395201	HD 99010	8.449	Fold	probable
A11	159892657	BD+40 3254	8.418	Fold	confirmed
A12	436758044	HD 217492	8.326	Fold	confirmed
A16	257352286	TYC 2173-965-1	9.423	App	confirmed
A17	219230041	HD 26958	7.514	App	confirmed
A18	189668511	TYC 2819-1027-1	9.216	App	confirmed
A19	219766411	Gaia DR3 1440245818754478976	12.395	Ast	probable
A20	235666667	HD 170829 = HR 6950 = NSV 24480	5.785	App	confirmed
A21	403113935	TYC 4244-623-1	9.411	App	confirmed
A22	324515186	HD 309050	9.444	App	confirmed
A23	142935468	TYC 4160-760-1	9.454	App	confirmed
A24	331507968	TYC 3669-262-1	9.278	App	confirmed
A25	258780392	BD+12 121	8.520	App	confirmed

Label	TIC	Name	*T* (mag)	Origin	Status
A26	89031985	TYC 3091-707-1	8.791	App	confirmed
A27	143198335	HD 63923	9.680	App	confirmed
A28	145034339	CD−32 4901	9.251	App	confirmed
A29	248652521	BD−01 648	9.571	App	confirmed
A30	311297982	BD+29 2845	9.595	App	confirmed
A31	2048935928	V1192 Cas (= TIC 271237873)	9.791	App	confirmed
A32	399758284	TYC 97-228-1	9.956	App	confirmed
A33	4341778	TYC 7691-3893-1	9.825	App	confirmed
A34	139285862	CD−45 14957	9.926	App	confirmed
A35	11488700	CD−29 366 (Bok I 16B)	10.299	App	confirmed
A36	43473641	TYC 5326-1527-1	10.344	App	confirmed
A37	24755724	CPD−25 5943	10.213	App	probable
A38	316326405	TYC 3715-870-1	9.933	App	confirmed
A39	22620802	TYC 3396-1373-1	10.210	App	confirmed
A40	165541068	TYC 986-148-1	10.156	App	probable
A41	156182608	TYC 6568-1036-1	10.521	App	confirmed
A42	832117928	CD−25 5638B	10.501	App	confirmed
A43	241445833	TYC 4685-2083-1	10.376	App	confirmed
A44	355477435	TYC 8461-136-1	10.501	App	confirmed
A45	34344890	TYC 7897-48-1	10.495	App	confirmed
A46	383529972	BD+30 2277 = WDS ES 436	10.118	App	variable, not counted
A47	70858500	CD−31 59 = HIP 1060	10.005	App	confirmed
A48	66900858	TYC 3166-453-1	10.533	App	confirmed
A49	286926100	TYC 3870-1689-1	10.010	App	confirmed
A50	450218810	TYC 3729-162-1	10.396	App	confirmed
A51	18000700	BD+28 4244	10.192	App	confirmed
A52	307170556	TYC 5406-1905-1	9.821	App	confirmed
A53	96844936	TYC 8355-1052-1	10.463	App	probable
A54	108344775	TYC 447-247-1	10.291	App	confirmed
A55	11156440	BD+60 397	9.529	App	confirmed
A56	367706305	HD 279736	10.354	App	confirmed
A57	145469097	TYC 7121-1393-1	10.021	App	confirmed
A58	421900559	BD−14 1559	9.563	App	confirmed
A59	372183259	TYC 3187-1132-1	10.394	App	confirmed
A60	142084127	TYC 9189-1130-1	10.022	App	confirmed
A61	468529113	TYC 4308-545-1	9.981	App	confirmed
A62	464830297	TYC 110-583-1	10.350	App	confirmed
A63	467158573	TYC 3801-1499-1	10.500	App	confirmed
A64	249895319	BD+50 578	9.770	App	confirmed
A65	88619181	BD+27 4098	10.018	App	confirmed
A66	260563222	CD−41 5677	9.652	App	confirmed
A67	173468180	TYC 3623-443-1	10.296	App	confirmed

Label	TIC	Name	*T* (mag)	Origin	Status
A68	80107967	TYC 8136-2645-1	10.580	App	probable
A69	377060838	TYC 243-1377-1	10.162	App	confirmed
A70	70789999	TYC 4094-1571-1	9.761	App	confirmed
A71	87303050	TYC 3747-2187-1	10.200	App	confirmed
A72	279636729	Gaia DR3 4720373709202359808	12.008	Ast	confirmed
A73	315721826	TYC 4590-364-1	10.808	Ast	confirmed
A74	427345302		12.10	Ast	probable
A75	202282515	TYC 3255-1824-1	9.655	Fold	confirmed
A76	202710940	TYC 3256-1533-1	9.480	Fold	confirmed
A77	196475343	BD+44 219	9.116	Fold	confirmed
A78	268587594	HD 6217	8.837	Fold	confirmed
A79	416277351	BD+64 520A	9.362	Fold	confirmed
A80	19702288	TYC 1310-1984-1	9.479	Fold	confirmed
A81	109061795	CD−24 5530	9.338	Fold	confirmed
A82	317250623	CD−83 126	8.980	Fold	confirmed
A83	7060775	BD+13 1919	9.163	Fold	confirmed
A84	456339196	TYC 3805-1342-1	9.566	Fold	confirmed
A85	38388435	CD−27 6336	9.141	Fold	confirmed
A86	170891257	BD−01 2238	9.414	Fold	confirmed
A87	463896609	HD 89725	8.885	Fold	confirmed
A88	302351416	TYC 3827-170-1	9.422	Fold	confirmed
A89	388287047	CPD−85 289	9.317	Fold	confirmed
A90	454747287	CD−73 685	9.346	Fold	confirmed
A91	311635507	HD 311112	9.375	Fold	confirmed
A92	148877735	BD+69 653	9.325	Fold	confirmed
A93	175734936	HD 112601	8.955	Fold	known (SIMBAD EB*), not counted
A94	245194662	HD 114997	8.917	Fold	confirmed
A95	154346187	HD 234179	8.984	Fold	confirmed
A96	426877400	HD 140021	9.623	Fold	confirmed
A97	351595900	TYC 1529-387-1	9.316	Fold	confirmed
A98	157469799	BD+45 2574 A	8.948	Fold	confirmed
A99	352236055	TYC 3925-1852-1	9.283	Fold	confirmed
A100	425304484	HD 353655	9.464	Fold	confirmed
A101	14596588	HD 192721	8.889	Fold	confirmed
A102	72160748	HD 194269	8.652	Fold	confirmed
A103	188258837	BD+41 3794	8.916	Fold	confirmed
A104	353453805	BD+42 3880	9.503	Fold	confirmed
A105	321718006	BD+67 1313	8.837	Fold	confirmed
A106	352935659	HD 205273	8.707	Fold	confirmed
A107	397680054	BD+32 4452	9.490	Fold	confirmed
A108	239649487	TYC 2747-436-1	9.131	Fold	confirmed

Label	TIC	Name	*T* (mag)	Origin	Status
A109	274331697	BD+43 1585	9.616	Fold	confirmed
A110	158544874	TYC 7212-2039-1	9.902	App	confirmed
A111	20379300	CD-28 8698	9.926	App	confirmed
A112	305944592	TYC 9053-432-1	10.036	App	confirmed
A113	352914772	TYC 4251-1144-1	9.992	App	confirmed
A114	65660608	TYC 6528-895-1	10.386	App	confirmed
A115	447307865	TYC 9038-1390-1	9.788	App	confirmed
A116	455491894	TYC 3608-1307-1	10.138	App	probable
A117	404715740	TYC 629-1389-1	9.485	App	confirmed
A118	64696335	TYC 6536-1088-1	10.388	App	confirmed
A119	280129511	TYC 4320-529-1	10.506	Batch	confirmed
A120	382605794	TYC 8982-3032-1	10.138	Batch	confirmed
A121	384249385	TYC 9010-4187-1	10.877	Batch	confirmed
A122	384525203	TYC 3612-1132-1	10.470	Batch	confirmed
A123	78448518	TYC 5962-1229-1	10.575	Batch	confirmed
A124	310162414	TYC 3371-625-1	10.342	Batch	confirmed
A125	431278074	TYC 3634-2713-1	10.199	Batch	confirmed
A126	279648140	CD−61 607	9.903	Batch	confirmed
A127	74180670	TYC 8737-1439-1	10.569	Batch	confirmed
A128	219543346	TYC 1643-218-1	10.965	Batch	confirmed
A129	4965877	TYC 7166-2609-1	11.070	Batch	confirmed
A130	240680969	TYC 3261-1420-1	10.173	Batch	confirmed
A131	52855832	TYC 4030-58-1	11.048	Batch	confirmed
A132	283862049	TYC 2263-89-1	10.694	Batch	confirmed
A133	359299569	TYC 8666-1738-1	10.666	Batch	confirmed
A134	329314665	TYC 9000-1011-1	11.017	Batch	confirmed
A135	182517223	TYC 9347-571-1	10.362	Batch	confirmed
A136	344947332	TYC 4004-778-1	11.006	Batch	confirmed
A137	1260529	TYC 6579-2222-1	10.775	Batch	confirmed
A138	306477840	TYC 5361-543-1	11.045	Batch	confirmed
A139	1695417	BD−13 3646	9.640	Batch	confirmed
A140	430987029	TYC 2741-1694-1	11.162	Batch	confirmed
A141	116246364	TYC 3093-1294-1	10.918	Batch	confirmed
A142	98952099	TYC 2971-946-1	11.057	Batch	confirmed
A143	192973862	CD−24 8975	10.335	Batch	confirmed
A144	200398115	TYC 3624-860-1	10.765	Batch	confirmed
A145	260789413	TYC 3344-341-1	10.222	Batch	confirmed
A146	53256660	TYC 4952-149-1	10.680	Batch	confirmed
A147	403936502	TYC 2136-2756-1	9.875	Batch	confirmed
A148	85944269	TYC 2669-4733-1	10.995	Batch	confirmed
A149	93103821	BD−17 691	10.603	Batch	confirmed
A150	7795044	CD−35 10394	9.549	Batch	confirmed

Label	TIC	Name	*T* (mag)	Origin	Status
A151	324273420	TYC 8977-7461-1	10.571	Batch	confirmed
A152	123378551	TYC 7652-366-1	10.739	Batch	confirmed
A153	375435321	TYC 8187-1030-1	10.892	Batch	confirmed
A154	375302184	TYC 9075-1173-1	10.921	Batch	confirmed
A155	285385215	TYC 2327-1770-1	10.585	Batch	confirmed
A156	159836610	TYC 7485-869-1	10.753	Batch	confirmed
A157	86036265	TYC 1405-695-1	10.597	Batch	confirmed
A158	268342242	TYC 8142-243-1	10.616	Batch	confirmed
A159	437230270	TYC 6635-619-1	11.063	Batch	confirmed
A160	142240367	TYC 7558-731-1	10.649	Batch	confirmed
A161	288160281	TYC 4564-1186-1	10.986	Batch	confirmed
A162	258860596	TYC 2205-1326-1	11.193	Batch	confirmed
A163	55114385	TYC 1337-1803-1	10.160	Batch	confirmed
A164	200745563	TYC 8062-84-1	10.521	Batch	confirmed
A165	423927429	UCAC4 797-029409	10.986	Batch	confirmed
A166	299650500	CD−23 4400	10.052	Batch	confirmed
A167	449475961	TYC 8994-1423-1	10.276	Batch	confirmed
A168	234880209	HD 259137	10.144	Batch	confirmed
A169	284533163	TYC 4020-206-1	10.999	Batch	confirmed
A170	88457347	TYC 4114-604-1	10.560	Batch	confirmed
A171	191146789	TYC 3248-1904-1	10.984	Batch	confirmed
A172	434263468	TYC 4282-552-1	9.968	Batch	confirmed
A173	293844315	BD−02 2880	9.886	Batch	confirmed
A174	348784186	TYC 5452-573-1	9.952	Batch	confirmed
A175	269504659	TYC 4674-364-1	10.474	Batch	confirmed
A176	55214475	TYC 4104-606-1	10.318	Batch	confirmed
A177	63858295	TYC 1429-565-1	10.839	Batch	confirmed
A178	323007878	TYC 9406-845-1	11.040	Batch	confirmed
A179	384284045	TYC 3616-2395-1	10.023	Batch	confirmed
A180	39307147	TYC 2609-823-1	10.830	App	confirmed
A181	124856476	TYC 7081-350-1	10.669	App	confirmed
A182	200222789	UCAC4 667-109998	10.864	App	confirmed
A183	146111348	TYC 8201-1042-1	10.539	App	confirmed
A184	220461013	UCAC2 5492112	10.736	App	confirmed
A185	427657329	TYC 4478-554-1	10.278	App	confirmed
A186	252806416	TYC 3444-504-1	10.802	App	confirmed
A187	170312515	UCAC4 240-011387	10.825	App	confirmed
A188	176778312	UCAC4 275-003209	11.085	App	confirmed
A189	74965330	TYC 7686-1190-1	10.844	App	confirmed
A190	307361219	TYC 9193-870-1	10.841	App	confirmed
A191	356308234	TYC 3519-1236-1	10.732	App	confirmed
A192	197484019	TYC 3619-1706-1	10.908	App	confirmed

Label	TIC	Name	T (mag)	Origin	Status
A193	160300523	TYC 5939-78-1	10.852	App	confirmed
A194	309861809	TYC 8925-649-1	10.361	App	confirmed
A195	233640767	TYC 3481-179-1	11.065	App	confirmed
A196	233606058	TYC 4216-238-1	10.922	App	confirmed
A197	300202855	TYC 7612-1193-1	11.135	App	confirmed
A198	320805993	TYC 8787-921-1	10.646	App	confirmed
A199	391925531	TYC 9380-1063-1	10.637	App	confirmed
A200	374560569	TYC 3697-940-1	10.875	Batch	confirmed
A201	70465753	TYC 1365-707-1	10.434	Batch	confirmed
A202	348536911	TYC 5995-1398-1	10.088	Batch	confirmed
A203	315849332	TYC 8972-1194-1	10.678	Batch	confirmed
A204	453601725	TYC 8675-1668-1	10.436	Batch	confirmed
Probable eclipsing binaries with unresolved hosts (not counted)					
A13	58553437		10.228	MD	probable, not counted
A14	71124276		10.504	MD	probable, not counted
A15	261135533	TOI-6255	10.446	MD	probable, not counted

Table A1b. Ephemerides and next primary eclipse.

Label	P (d)	T_0 (BJD_{TDB})	Next primary (UTC)
A1	1.2443947 ± 0.0000016[a]	2458629.45849	2026-10-14 10:17
A2	6.129392 ± 0.000006	2457003.74608	2026-10-17
A3	9.474664[b]	2458993.379	none (no timing fit)
A4	3.4657540 ± 0.0000093	2459528.2147 ± 0.002	2026-10-16 21:49
A5	19.454722 ± 0.000010	2460494.45294 ± 0.00039	2026-10-17 12:13
A6	5.0782783 ± 0.0000010	2460609.56921 ± 0.00016	2026-10-17 04:20
A7	13.9106635 ± 0.0000071	2460685.73790 ± 0.00034	2026-10-26 00:59
A8	10.4462098 ± 0.0000098	2460391.45705 ± 0.00071	2026-10-18 02:49
A9	27.0822013 ± 0.0000180[f]	2460097.60593 ± 0.00023	2026-10-29 21:08
A10	7.8127421 ± 0.0000013	2460384.50567 ± 0.0001	2026-10-16 08:23
A11	4.0751904 ± 0.0000009	2460133.82003 ± 0.00011	2026-10-14 08:25
A12	14.0343984 ± 0.0000110	2460203.47040 ± 0.00028	2026-10-26 18:03
A16	23.802223 ± 0.000022	2459804.28519 ± 0.00089	2026-10-14 02:57
A17	40.45701 ± 0.00002	2457386.08460	2026-11-06 08:51
A18	31.80877 ± 0.00001[g]	2458766.58610	2026-10-29 14:09
A19	290.9037 ± 0.0006	2458936.3573 ± 0.0028	2027-05-28
A20	26.384493 ± 0.000040	2459403.97980	2026-10-16 13:09
A21	30.763077 ± 0.000013	2459849.89772 ± 0.00041	2026-11-12 18:52
A22	23.034195 ± 0.000010	2459657.89250 ± 0.00031	2026-10-25 21:23
A23	22.244792 ± 0.000017	2459228.68805 ± 0.00056	2026-10-28 10:37
A24	24.690127 ± 0.000031	2458797.42249 ± 0.00146	2026-10-27 00:01
A25	43.8230 ± 0.0025	2459451.28700 ± 0.0013	2026-10-22 04:04
A26	42.14749 ± 0.00117	2459645.40345 ± 0.01221	2026-10-17 19:17
A27	41.30385 ± 0.00030	2459963.04837 ± 0.00438	2026-11-22 21:04

Label	P (d)	T_0 ($\mathrm{BJD_{TDB}}$)	Next primary (UTC)
A28	51.37333 ± 0.00020	2458512.23071 ± 0.002	2026-10-24 06:20
A29	29.606659 ± 0.000010	2459177.83838 ± 0.0005	2026-10-25 14:51
A30	27.94676 ± 0.00001	2458994.50441 ± 0.0018	2026-10-28 12:43
A31	37.4133 ± 0.0005	2458764.90610 ± 0.018	2026-11-01 22:03
A32	40.51032 ± 0.00011[g]	2458463.92360 ± 0.024	2026-10-26 15:37
A33	45.59795 ± 0.00008[g]	2458573.67953 ± 0.0042	2026-11-10 15:42
A34	45.161458 ± 0.000109	2458354.30334 ± 0.00477	2026-10-21 10:57
A35	20.289728 ± 0.000160	2458383.84197 ± 0.00342	2026-11-01 15:17
A36	54.851934 ± 0.000191	2459192.78258 ± 0.0053	2026-10-18 12:05
A37	22.611537 ± 0.002850	2460793.29985 ± 0.002	2026-10-22 11:30
A38	37.116502 ± 0.000308[g]	2458839.85111 ± 0.00926	2026-11-19 06:26
A39	35.692330 ± 0.000062	2459602.95779 ± 0.0007	2026-11-07 09:05
A40	24.7458 ± 0.0025[c]	2460463.59430 ± 0.001	2026-10-16 04:44
A41	23.53654 ± 0.00017	2458501.94000 ± 0.011	2026-11-05 08:39
A42	22.029385 ± 0.000018	2458508.34974 ± 0.00098	2026-10-14 14:40
A43	23.205492 ± 0.000007	2458389.70430 ± 0.00052	2026-10-23 07:05
A44	21.106026 ± 0.000006	2458325.49469 ± 0.00051	2026-10-30 03:41
A45	31.152794 ± 0.000072	2459382.75960 ± 0.0037	2026-10-31 21:19
A46	0.381845 ± 0.000002	2459645.01110 ± 0.0007	2026-10-14 07:20
A47	24.105122 ± 0.000006[g]	2458357.74508 ± 0.00079	2026-11-02 06:37
A48	43.196065 ± 0.000034	2459804.38481 ± 0.00045	2026-11-14 22:36
A49	28.098880 ± 0.000013	2459645.01560 ± 0.00026	2026-10-17 10:46
A50	24.172470 ± 0.000015	2459926.66270 ± 0.00041	2026-10-15 03:53
A51	29.32848 ± 0.00033	2460558.15104 ± 0.060	2026-11-05 12:24
A52	40.46953 ± 0.00005	2459206.58750 ± 0.0019	2026-11-06 23:17
A53	36.7755 ± 0.006[d]	2461155.84550 ± 0.0007	2026-10-26 05:24
A54	3.02348 ± 0.00001	2460482.03790 ± 0.0001	2026-10-15 02:41
A55	21.254843 ± 0.000012	2458801.37455 ± 0.00065	2026-10-17 04:43
A56	32.286594 ± 0.000012	2458817.15278 ± 0.00047	2026-10-22 00:02
A57	43.324287 ± 0.000075[g]	2458530.31198 ± 0.00384	2026-11-01 21:22
A58	35.976045 ± 0.000018	2459203.38951	2026-11-17 10:45
A59	42.648930 ± 0.000055	2460537.57169	2026-11-03 09:33
A60	45.49880 ± 0.00004[g]	2459553.26450 ± 0.00083	2026-10-14 05:13
A61	21.552334 ± 0.000007	2459481.89522 ± 0.00022	2026-10-21 21:25
A62	22.34089 ± 0.00001[g]	2458440.02660 ± 0.0004	2026-10-30 20:05
A63	57.2400 ± 0.0003	2458868.88650 ± 0.005	2026-10-16 16:57
A64	29.27232 ± 0.00005	2458793.94630 ± 0.0007	2026-10-27 03:11
A65	26.768321 ± 0.000009	2459819.55960 ± 0.00015	2026-10-31 20:25
A66	38.791921 ± 0.000024	2458553.49853 ± 0.00089	2026-11-02 00:27
A67	29.182072 ± 0.000005[g]	2458756.76116 ± 0.00042	2026-11-09 11:05
A68	34.184472 ± 0.000076	2458521.30741 ± 0.00302	2026-11-14 02:50
A69	42.463964 ± 0.000016[g]	2458517.53332 ± 0.00059	2026-11-18 02:50

Label	P (d)	T_0 (BJD_{TDB})	Next primary (UTC)
A70	22.735028 ± 0.000008	2458823.53837 ± 0.00039	2026-11-02 14:56
A71	24.445338 ± 0.000011	2458833.11980 ± 0.00047	2026-11-06 11:38
A72	215.0116 ± 0.0002	2458341.4321 ± 0.0009	2026-11-07
A73	17.538[e]	2459940.800	2026-10-30
A74	173.8755 ± 0.0005	2459884.9860 ± 0.0013	2027-02-13
A75	6.929866 ± 0.000015	2460111.42326 ± 0.00065	2026-10-17 13:47
A76	3.3063718 ± 0.0000018	2460299.60466 ± 0.00020	2026-10-14 09:09
A77	6.370415 ± 0.000005	2460407.33044 ± 0.00027	2026-10-17 12:50
A78	9.140474 ± 0.000004	2458764.02178 ± 0.00018	2026-10-18 23:44
A79	6.607651 ± 0.000012	2460149.97660 ± 0.00045	2026-10-19 05:50
A80	17.02161 ± 0.00032	2459516.54908 ± 0.00042	2026-10-24 08:34
A81	13.2271715 ± 0.0000023	2460040.42781 ± 0.00013	2026-10-23 04:33
A82	19.605278 ± 0.000014	2460060.10199 ± 0.00051	2026-10-20 22:41
A83	6.422180 ± 0.000001	2459741.55165 ± 0.00012	2026-10-14 07:56
A84	7.1061147 ± 0.0000043	2459530.89350 ± 0.00012	2026-10-15 05:46
A85	2.6164745 ± 0.0000005	2460518.67725 ± 0.00008	2026-10-16 06:52
A86	15.798442 ± 0.000003	2459811.62273 ± 0.00024	2026-10-14 18:36
A87	2.2366733 ± 0.0000008	2460195.87831 ± 0.00025	2026-10-14 03:17
A88	13.9574485 ± 0.0000066	2459928.45698 ± 0.00023	2026-10-24 15:50
A89	17.638544 ± 0.000019	2460032.65988 ± 0.00069	2026-10-24 09:54
A90	5.6347635 ± 0.0000015	2459691.10727 ± 0.00013	2026-10-17 07:47
A91	18.5428744 ± 0.0000049	2460479.18152 ± 0.00017	2026-10-18 15:44
A92	3.6074499 ± 0.0000008	2460110.82736 ± 0.00008	2026-10-16 15:29
A93	4.337953 ± 0.000001	2460380.72231 ± 0.00018	2026-10-17 05:43
A94	17.2795467 ± 0.0000052	2460353.15051 ± 0.00022	2026-10-24 14:06
A95	3.549541 ± 0.000002	2459489.65835 ± 0.00031	2026-10-14 19:44
A96	2.7560089 ± 0.0000006	2460574.15160 ± 0.00014	2026-10-15 19:11
A97	6.647424 ± 0.000005	2459826.14200 ± 0.00052	2026-10-14 23:05
A98	3.5063244 ± 0.0000002	2460172.71372 ± 0.00002	2026-10-16 07:13
A99	18.821610 ± 0.000008	2460160.83034 ± 0.00023	2026-10-14 06:25
A100	16.260285 ± 0.000024	2460531.06071 ± 0.00039	2026-10-14 07:30
A101	4.478718 ± 0.000065	2460818.15412 ± 0.00013	2026-10-15 05:25
A102	1.9385707 ± 0.0000005	2460815.72949 ± 0.00012	2026-10-14 00:14
A103	5.9855010 ± 0.0000159	2460144.94717 ± 0.00108	2026-10-16 13:46
A104	14.3509487 ± 0.0000051	2460309.02608 ± 0.00010	2026-10-14 10:34
A105	3.9253291 ± 0.0000007	2460144.07847 ± 0.00008	2026-10-16 00:36
A106	1.3854241 ± 0.0000004	2460402.27646 ± 0.00009	2026-10-14 05:39
A107	2.976903 ± 0.000001	2459681.70466 ± 0.00012	2026-10-14 10:15
A108	6.4161826 ± 0.0000024	2460233.51625 ± 0.00014	2026-10-17 04:17
A109	1.0394878 ± 0.0000005	2460134.11958 ± 0.00009	2026-10-14 23:44
A110	22.039984 ± 0.000046	2460010.4835 ± 0.004	2026-10-19 09:15
A111	32.60569 ± 0.00003	2460759.1644 ± 0.0012	2026-11-01 13:40

Label	P (d)	T_0 ($\mathrm{BJD_{TDB}}$)	Next primary (UTC)
A112	34.48723 ± 0.00005	2458626.1182 ± 0.0026	2026-11-06 02:40
A113	23.63714 ± 0.00001	2458722.60576 ± 0.007	2026-11-01 19:48
A114	41.56002 ± 0.00006	2459230.72319 ± 0.0010	2026-11-05 18:46
A115	40.01697 ± 0.00002	2458628.6578 ± 0.0009	2026-11-05 07:33
A116	57.3498 ± 0.0100	2460564.19008 ± 0.0004	2026-11-22 14:01
A117	36.54015 ± 0.00003	2459492.88004 ± 0.00038	2026-11-11 22:07
A118	41.86524 ± 0.00087	2459238.650 ± 0.021	2026-11-29 06:34
A119	6.348792 ± 0.000015	2461336.598 ± 0.010	2026-10-23 02:15
A120	2.140795 ± 0.000001	2460055.59400 ± 0.0007	2026-10-20 03:37
A121	7.805870 ± 0.000011	2460653.55840 ± 0.0014	2026-10-26 23:28
A122	4.055307 ± 0.000006	2459826.03683 ± 0.00076	2026-10-21 02:34
A123	11.874406 ± 0.000027	2458477.12450 ± 0.00296	2026-10-25 08:30
A124	4.551256 ± 0.000007	2458819.32778 ± 0.001	2026-10-22 16:02
A125	6.386066 ± 0.000003	2458744.40878 ± 0.0006	2026-10-23 15:32
A126	3.3502765 ± 0.0000006	2459829.361644 ± 0.000167	2026-10-20 03:12
A127	18.4215088 ± 0.0000285	2460686.04707 ± 0.00105	2026-11-04 17:22
A128	19.205350 ± 0.000065	2459919.47988 ± 0.00199	2026-10-27 04:10
A129	10.220874 ± 0.000008	2459901.40080 ± 0.00075	2026-10-29 01:05
A130	18.047704 ± 0.000085	2458778.6842 ± 0.0063	2026-10-27 22:52
A131	2.4113702 ± 0.0000015	2459632.73471 ± 0.00044	2026-10-21 15:47
A132	10.144315 ± 0.000004	2459997.87878 ± 0.00029	2026-10-23 10:08
A133	7.149090 ± 0.000002	2460352.39715 ± 0.00031	2026-10-25 11:23
A134	3.7733403 ± 0.0000019	2460773.739862 ± 0.000333	2026-10-22 11:16
A135	2.2076198 ± 0.0000003	2460115.008748 ± 0.000100	2026-10-20 02:43
A136	10.451460 ± 0.000014	2459954.44026 ± 0.00080	2026-10-20 12:41
A137	10.992454 ± 0.000008	2459566.15999 ± 0.00035	2026-10-22 10:42
A138	5.520586 ± 0.000003	2459474.27805 ± 0.00051	2026-10-21 05:06
A139	17.797939 ± 0.000010	2459326.38398 ± 0.0002	2026-10-24 01:20
A140	9.920766 ± 0.000016	2460364.52570 ± 0.0009	2026-10-23 06:09
A141	14.748239 ± 0.000011	2459765.23338 ± 0.00040	2026-10-29 19:05
A142	5.955676 ± 0.000004	2459444.81721 ± 0.00034	2026-10-25 05:17
A143	19.211924 ± 0.000008	2460144.30777 ± 0.00027	2026-10-21 22:47
A144	7.305664 ± 0.000049	2460170.06233 ± 0.00263	2026-10-25 11:08
A145	4.746808 ± 0.000003	2460126.22496 ± 0.00027	2026-10-23 03:45
A146	5.160140 ± 0.000004	2460411.01065 ± 0.00041	2026-10-21 04:19
A147	14.804287 ± 0.000006	2460043.96544 ± 0.00021	2026-11-02 05:49
A148	3.021219 ± 0.000003	2459951.66557 ± 0.00045	2026-10-21 21:10
A149	1.6350307 ± 0.0000004	2458868.90788 ± 0.00009	2026-10-21 00:41
A150	9.4320556 ± 0.0000068	2460194.85288 ± 0.00053	2026-10-22 15:15
A151	3.9158631 ± 0.0000015	2460200.00630 ± 0.00037	2026-10-22 02:36
A152	11.010611 ± 0.000006	2459972.09636 ± 0.00034	2026-10-23 21:53
A153	18.214018 ± 0.000011	2460498.99671 ± 0.00056	2026-10-23 08:14

Label	P (d)	T_0 ($\mathrm{BJD_{TDB}}$)	Next primary (UTC)
A154	2.5455862 ± 0.0000005	2460571.13574 ± 0.00015	2026-10-21 07:29
A155	8.2074497 ± 0.0000015	2458797.97260 ± 0.00021	2026-10-20 13:38
A156	5.7022928 ± 0.0000004	2459618.89923 ± 0.00007	2026-10-21 18:53
A157	11.9972226 ± 0.0000022	2459507.95631 ± 0.00069	2026-10-30 00:45
A158	14.6202005 ± 0.0000021	2459840.37583 ± 0.00012	2026-11-01 18:08
A159	11.870552 ± 0.000001	2459937.17038 ± 0.00009	2026-10-24 09:34
A160	7.541350 ± 0.000002	2460464.17253 ± 0.00033	2026-10-25 11:09
A161	7.063961 ± 0.000002	2459597.74250 ± 0.00016	2026-10-21 23:25
A162	4.959311 ± 0.000003	2460056.15218 ± 0.00022	2026-10-22 03:36
A163	4.268746 ± 0.000001	2459870.92006 ± 0.00009	2026-10-25 20:46
A164	10.998961 ± 0.000002	2460412.11666 ± 0.00017	2026-10-22 12:37
A165	1.4271182 ± 0.0000001	2459967.97498 ± 0.00004	2026-10-23 01:53
A166	18.337086 ± 0.000027	2460427.74767 ± 0.00092	2026-10-31 02:23
A167	4.1731519 ± 0.0000014	2460037.02702 ± 0.00033	2026-10-25 13:16
A168	4.270292 ± 0.000001	2459508.56208 ± 0.00018	2026-10-22 17:51
A169	2.349378 ± 0.000002	2459995.69666 ± 0.00046	2026-10-23 16:29
A170	5.4689808 ± 0.0000012	2459730.21937 ± 0.00008	2026-10-24 14:20
A171	7.7886371 ± 0.0000029	2459851.29421 ± 0.00015	2026-10-25 10:02
A172	8.077758 ± 0.000004	2460199.37690 ± 0.00022	2026-10-24 20:05
A173	14.774809 ± 0.000003	2459810.95109 ± 0.00021	2026-11-03 00:45
A174	4.0868675 ± 0.0000004	2459978.32495 ± 0.00007	2026-10-25 18:03
A175	11.391807 ± 0.000015	2459800.01713 ± 0.00049	2026-10-24 09:42
A176	4.8136773 ± 0.0000007	2459786.95100 ± 0.00008	2026-10-23 10:51
A177	3.4331770 ± 0.0000009	2459599.88038 ± 0.00012	2026-10-23 13:41
A178	4.7034219 ± 0.0000005	2459822.87791 ± 0.00009	2026-10-23 21:07
A179	3.787011 ± 0.000002	2460277.42043 ± 0.00006	2026-10-24 06:43
A180	23.421703 ± 0.000013	2458989.52348 ± 0.00059	2026-11-10 14:49
A181	21.680622 ± 0.000006	2459210.25550 ± 0.00040	2026-11-12 03:13
A182	43.89933 ± 0.00007	2458741.61295 ± 0.00125	2026-12-01 01:41
A183	49.24084 ± 0.00003	2458565.73335 ± 0.00118	2026-11-27 23:05
A184	50.85317 ± 0.00006	2459760.1247 ± 0.0011	2026-10-23 01:42
A185	47.52876 ± 0.00004	2459574.2519 ± 0.0005	2026-12-05 20:12
A186	42.44229 ± 0.00008	2459633.95593 ± 0.00099	2026-11-29 14:07
A187	52.31098 ± 0.00005	2459232.33454 ± 0.00097	2026-12-02 13:58
A188	22.359363 ± 0.000023	2459575.79945 ± 0.00127	2026-10-28 16:25
A189	39.199094 ± 0.000029	2459671.06029 ± 0.00105	2026-11-12 02:55
A190	23.828151 ± 0.0000013	2459089.30214 ± 0.00005	2026-11-08 11:26
A191	42.860164 ± 0.000019	2458689.14215 ± 0.00057	2026-11-01 23:20
A192	51.88235 ± 0.00021	2459835.1830 ± 0.0021	2026-10-26 06:25
A193	37.312955 ± 0.000007	2458473.96743 ± 0.00035	2026-11-02 13:28
A194	38.386132 ± 0.000039	2460769.65585 ± 0.00099	2026-10-31 22:45
A195	30.305947 ± 0.000008	2458752.76490 ± 0.00029	2026-11-14 13:52

Label	P (d)	T_0 (BJD_{TDB})	Next primary (UTC)
A196	21.737275 ± 0.000004	2458691.30913 ± 0.00020	2026-10-29 18:08
A197	25.762668 ± 0.0000034	2459211.64639 ± 0.00020	2026-11-05 10:41
A198	56.173887 ± 0.000011	2458676.76580 ± 0.00021	2026-11-28 14:45
A199	43.977834 ± 0.000008	2459079.03934 ± 0.00018	2026-11-21 09:17
A200	13.893272 ± 0.000012	2459892.25831 ± 0.00039	2026-10-23 15:41
A201	6.4475682 ± 0.0000015	2459831.15209 ± 0.00009	2026-10-26 09:09
A202	2.553938 ± 0.000005	2459998.52091 ± 0.00035	2026-10-23 06:49
A203	5.4921244 ± 0.0000015	2459753.70951 ± 0.00034	2026-10-27 10:27
A204	5.5666840 ± 0.0000021	2460765.53158 ± 0.00026	2026-10-25 09:40
Probable eclipsing binaries with unresolved hosts (not counted)			
A13	1.078685		
A14	1.123892		
A15	14.486682		

Table A1c. Eclipse measurements.

Label	Depth prim. (%)	Depth sec. (%)	Dur. (h)
A1	2.28	1.85	0.7
A2	21–29	17–23	
A3	0.0515		1.58
A4	30.2	6.73	2.75
A5	2.03		4.38
A6	10.97	2.79	2.55
A7	8.00	1.00	3.8
A8	1.82		11.0
A9	39.8	26.1	14.3
A10	2.20	1.73	4.0
A11	10.11	1.87	4.02
A12	1.82		4.0
A16	10	8.45	20
A17	1.2		9.7
A18	3.75	2.2	
A19	12.5		7
A20	1.65	0.106	11.8
A21	2.77	0.27	16.1
A22	8.3	1.1	7.5
A23	1.70	1.8	6.4
A24	0.86	0.32	6.9
A25		2.9	16.5
A26	0.29		18.8
A27		1.07	7.2
A28	20–21	5	34.1
A29	9.5		5.8
A30	42		10.1

Label	Depth prim. (%)	Depth sec. (%)	Dur. (h)
A31	4.7	0.9	29
A32	1.3–1.4	0.45	
A33	9	2.2	
A34	2.3		16
A35	3.3		5.9
A36	1.0	0.39	29
A37	18.7	13.4	17.5
A38	8.7	2.5	
A39	2.6	1.25	8.7
A40	4.6	1.5	13.7
A41		0.7–1.0	3
A42	0.5		4.3
A43	6.9		4.25
A44	5.7		3.1
A45	6–7		3.8
A46	4.6	3.4	
A47	36	34	
A48	27	10.4	9.7
A49	2.6	0.45	5.8
A50	11.5	1.6	7.1
A51		1.1–1.3	16.4
A52	6		16
A53		1.8–2.4	5.9
A54	18.8	11.7	2.70
A55	19.5	18.1	11
A56	17.5	17.2	14
A57	5.8	2.2	
A58	9.8		6
A59	12.3		4
A60	0.8	0.3	
A61	14.6		4.9
A62	2.1	0.15	
A63	18	16	13
A64	1.67	0.15	9.9
A65	10	0.72	7.2
A66	3.0	0.22	14
A67	3.8	2.3	
A68	1.3		11.4
A69	17	14.6	
A70	8.2	5.1	13
A71	9.7	1.4	4.5
A72	38	24.2	10.6

Label	Depth prim. (%)	Depth sec. (%)	Dur. (h)
A73	33 to 13	14.6 to 6.3	7.0 to 5.6
A74	4.62	~5 (ZTF g)	10.8
A75	6.76	1.55	5.00
A76	0.81	0.78	2.25
A77	0.6	0.37	1.9
A78	3.65	<0.4	2.38
A79	0.454	0.169	2.54
A80	7.98	0.77	7.4
A81	2.56	0.189	7.25
A82	9.8	2.06	5.65
A83	10.1	4.59	4.15
A84	7.86	4.13	8.60
A85	5.5	0.28	3.6
A86	12	1.4	3.5
A87	1.64	0.36	4.3
A88	3.57	<0.4	4.50
A89	1.61	<0.2	3.9
A90	1.17	0.049	4.30
A91	6.67	0.87	7.27
A92	1.32	1.32	3.05
A93	0.99	0.93	2.55
A94	2.8	0.25	4.2
A95	0.559	0.509	2.20
A96	0.76	0.031	4.5
A97	0.59	<0.2	4.1
A98	24.5	19.6	3.85
A99	1.49	<0.13	4.7
A100	3.6	1.3	12.3
A101	1.44	0.277	2.95
A102	6.95	0.71	2.3
A103	3.62	3.18	11.0
A104	6.09	1.46	8.60
A105	1.18	0.09	5.6
A106	3.06	2.92	1.95
A107	5.57	5.52	3.2
A108	21.4	2.8	12.6
A109	1.26	0.15	1.3
A110	24–35	21–29	7.8
A111	4.9–5.1	0.50–0.62	7.9
A112	0.86		18.7
A113	14.5–15.4	1.8–2.2	8.4
A114	9.3–9.5		4.2

Label	Depth prim. (%)	Depth sec. (%)	Dur. (h)
A115	1.3		5.4
A116	2.3		5.4
A117	21.5	3.3	6.9
A118	0.56	0.24	32
A119	0.34	0.28	3.1
A120	0.31		1.7
A121	0.33		3.24
A122	0.58	0.34	3.22
A123	0.39	0.18	2.95
A124	0.41	0.41	2.45
A125	0.53	0.13	3.6
A126	0.60	0.071	3.42
A127	0.54	0.33	3.0
A128	0.69		4.56
A129	0.68		2.58
A130	0.67–0.85		7.2
A131	0.91	0.50	1.92
A132	0.80		1.92
A133	0.79		7.08
A134	0.94	0.33	2.30
A135	0.97	0.12	1.38
A136	0.90	0.055	7.54
A137	1.09	0.040	3.78
A138	1.02		3.72
A139	0.93		4.44
A140	0.86		6.96
A141	1.14		3.12
A142	1.11	0.065	3.72
A143	1.09		3.48
A144	1.20	0.21	8.8
A145	1.43	0.49	3.30
A146	1.29		2.28
A147	1.40		2.82
A148	1.73	1.11	4.6
A149	1.55	0.58	1.44
A150	1.50		4.62
A151	1.5–1.6	1.5	3.48
A152	2.3		2.6
A153	2.15	0.13	9.42
A154	2.85	0.19	3.16
A155	6.5	0.66	2.94
A156	9.0	0.79	4.74

Label	Depth prim. (%)	Depth sec. (%)	Dur. (h)
A157	10.0	8.6	10.7
A158	12.4	3.3	9.14
A159	13.7	1.2	4.1
A160	1.91	0.42	3.42
A161	1.62	0.67	1.92
A162	1.81	0.14	2.82
A163	1.78	0.24	1.86
A164	1.85		5.10
A165	1.74	0.13	2.04
A166	1.94		4.5
A167	1.92	1.93	3.0
A168	1.88		2.22
A169	1.63	0.12	4.98
A170	2.06		3.30
A171	2.27		3.42
A172	2.39	0.14	8.07
A173	2.25		4.38
A174	2.64	1.16	3.00
A175	2.36	0.15	4.44
A176	2.60	0.15	1.92
A177	2.64	0.25	1.86
A178	2.55	0.93	3.36
A179	2.89	2.56	2.20
A180	24–26	9.8–10.2	6.9
A181	9.6–10.0	1.7–1.8	6.4
A182	13.5–14.6	6.1–7.0	24.2
A183	12.4–14.2	7.7–8.4	25.3
A184	3.8–3.9	1.2–2.3	53.4
A185	2.4–2.6	1.1–1.6	38.7
A186	2.9–3.0	0.75	13.9
A187	1.8–1.9	0.9	19.2
A188	9.3–9.7	1.2	7.6
A189	10.4–11.1	6.6–7.0	19.0
A190	22	0.43	3.3
A191	26–27	15.2	10.2
A192	7.1–7.3	0.3–0.5	30.0
A193	11.1–11.6	0.9–1.1	5.15
A194	1.0–1.1		9.1
A195	15.0–15.9	5.5–5.8	13.1
A196	17.7	17.1	10.9
A197	6.4–6.9	0.7–0.8	5.9
A198	22.5–22.7	5.3–5.4	7.55

Label	Depth prim. (%)	Depth sec. (%)	Dur. (h)
A199	2.0		6.7
A200	2.79	0.42	5.57
A201	2.66	0.23	2.16
A202	2.82	0.38	3.36
A203	2.92	0.76	3.74
A204	3.04	2.94	2.58
Probable eclipsing binaries with unresolved hosts (not counted)			
A13	0.126		
A14	0.082	0.022	
A15	0.1399		

Table A1d. Notes.

Label	Notes
A1	K
A2	K
A3	TCE
A4	SB, TCE, HP
A5	SB, TCE
A6	SB, TCE, HP
A7	SB, TCE, R
A8	SB, TCE
A9	SB, G, TCE, HP
A10	SB, TCE
A11	SB, K, TCE, HP
A12	SB, G, TCE
A16	TCE, HP
A17	
A18	
A19	K
A20	TCE, V
A21	G, K, TCE, CTOI
A22	TCE
A23	TCE
A24	TCE
A25	V
A26	
A27	
A28	G, HP

Label	Notes
A29	G
A30	G, TCE, HP
A31	K, V
A32	V
A33	K, HP
A34	TCE
A35	H
A36	
A37	HP
A38	TCE, HP
A39	TCE
A40	
A41	K, TCE
A42	K, H
A43	TCE
A44	
A45	bc
A46	H, bc
A47	TCE, HP
A48	TCE, HP
A49	TCE
A50	K, TCE
A51	TCE
A52	G, V
A53	
A54	G, TCE, HP, bc
A55	TCE, HP
A56	TCE, HP
A57	
A58	
A59	TCE
A60	TCE
A61	TCE, HP
A62	
A63	G, TCE
A64	TCE

Label	Notes
A65	G, TCE
A66	TCE
A67	TCE
A68	H
A69	TCE
A70	K, TCE
A71	K, TCE, HP
A72	K, TCE
A73	K, TCE
A74	K, TCE
A75	SB, TCE
A76	SB, TCE
A77	SB
A78	SB, TCE
A79	SB, TCE
A80	SB, TCE
A81	SB, K, TCE
A82	SB, TCE
A83	SB, TCE
A84	SB, TCE
A85	SB, TCE
A86	SB, G, TCE
A87	SB, TCE
A88	SB
A89	SB, TCE
A90	SB, TCE
A91	SB, TCE
A92	SB, TCE
A93	SB, TCE
A94	SB, K, TCE
A95	SB, TCE
A96	SB, TCE
A97	SB, TCE
A98	SB, TCE
A99	SB
A100	SB, K, TCE

Label	Notes
A101	SB
A102	SB, G, TCE
A103	SB
A104	SB, TCE
A105	SB, K, TCE
A106	SB, TCE
A107	SB, TCE
A108	SB
A109	SB, TCE
A110	SB; Gaia period (Table 4); secondary at phase 0.506; both eclipses shallower 2019–25 (fixed-aperture range given), then flat
A111	SB; M dwarf transiting a K subgiant; the secondary a total occultation, 17–18 h; Gaia's *e*, ω hold
A112	SB; a 0.30 $R_\odot$ M dwarf transiting an F subgiant; no secondary expected above 0.07%
A113	SB; Gaia period (Table 4); eclipse timing variations (curved O–C); T_0 from a fit to cycles ≥ 70
A114	SB; grazing, two equal 0.95 $R_\odot$ stars; no secondary possible at Gaia's $e \sin \omega$
A115	SB; near-twin pair, grazing at one conjunction; no secondary expected
A116	SB; two flat-bottomed eclipses at Gaia's secondary conjunction, exactly three Gaia periods apart; *P* is Gaia's; 1.6 d from the predicted conjunction phase
A117	SB; Gaia period (Table 4); Gaia's *e*, ω hold
A118	SB; red-clump giant with a K dwarf, circular; seven events 2019–2026, one out of sample
A119	SB; Gaia SB2 of near-twin stars (*q* 1.02), grazing eclipses at both conjunctions with the ellipsoidal variation locked to them; T_0, *P* and the depths refer to the deeper eclipse; 18 + 16 events in the clean sectors
A120	SB; M dwarf (0.33 $M_\odot$) grazing the F star at conjunction A, 81 events; the co-moving 27″ neighbour (WDS TOB 243, probably a triple) excluded by difference images and a PSF deblend; a tentative 0.2–0.4 ppt occultation at phase 0.500 is not quoted
A121	SB; occultation of the 0.43 $M_\odot$ M dwarf at conjunction B near periastron; no transit at A (< 0.15 ppt, consistent with the geometry); the *G* 13.8 neighbour 13″ away shows the dip only as PSF leakage, excluded at 8.6–18.6σ, and is not an eclipser itself
A122	SB; grazing eclipses on a strongly ellipsoidal (9.6 ppt) evolved A star with Doppler beaming at Gaia's *K*; the batch's 3.0 ppt box at *P*/2 sat on the ellipsoidal minimum; $e \cos \omega$ −0.029
A123	SB; eccentric F + K dwarf, 7 + 7 events; the deeper eclipse at conjunction B, the secondary at phase 0.6436; $e \cos \omega$ −0.227 against Gaia's −0.224
A124	SB; Gaia SB2 twins (*q* 0.97), equal grazing eclipses at both conjunctions, ellipsoidal 1.1 ppt; $e \cos \omega$ 0.002 ± 0.001; TESS-SPOC confirms
A125	SB; the deeper eclipse is the occultation of the G-dwarf companion at conjunction B; the 1.3 ppt eclipse at A (phase 0.553) is new; $e \cos \omega$ −0.0834 ± 0.0006 against Gaia's −0.0835
A126	SB, TCE; grazing, with ellipsoidal variation and Doppler beaming at Gaia's *K*; M-dwarf companion; secondary at phase 0.502; VSX types the star DSCT|GDOR|SXPHE, a type revision; an unpromoted SPOC multi-sector TCE caught both eclipses
A127	SB; eccentric, M-dwarf companion (0.28 $M_\odot$); transit at conjunction A (6.2 ppt dilution-corrected), occultation at B at phase 0.668, matching Gaia's $e \cos \omega$ to 0.09σ; 6 + 5 events, none owning the verdict

Label	Notes
A128	SB; grazing occultation of the companion at conjunction B on a subgiant; no eclipse at A (< 0.45 ppt), as the geometry predicts; TASOC detects it independently
A129	SB; grazing occultation at conjunction B near periastron, 11 events; a heartbeat explicitly excluded; QLP, SPOC and TGLC agree
A130	SB; grazing M-dwarf transit of a low-RGB giant at conjunction A, 5 events; the depth varies 6.7–8.5 ppt in four pipelines alike (spots on a grazing chord)
A131	SB; grazing, both eclipses (43 + 41 events) on a γ Dor pulsator; the depth ratio wants a ~5600 K companion against the 0.56 $M_\odot$ of Gaia's mass function, so Gaia's *K* is probably diluted (an unrecognised SB2)
A132	SB; grazing F8 dwarf + 0.5 $M_\odot$ M dwarf at conjunction A, 6 events; the predicted 0.3–0.5 ppt occultation is not seen
A133	SB; flat-bottomed transit of an F subgiant by a 0.22 $M_\odot$ M dwarf; 7.9 ppt in QLP, 9.3 ppt undiluted; the batch's difference-image offset was a crowding artefact, the eclipse localised on the target in five sectors at 8–36σ
A134	SB; grazing F + K dwarf, secondary at phase 0.5002; the 7.5″ batch offset was a centroiding artefact in a crowded field, the PSF-fit deficit on target in seven sectors and the 11″ neighbour excluded
A135	SB, TCE; grazing F + early-M dwarf with a spotted, probably synchronised primary; secondary at phase 0.4994; an unpromoted SPOC multi-sector TCE (sectors 56–69) caught both eclipses
A136	SB; flat-bottomed total transit of an F star by a 0.16 $M_\odot$ M dwarf; the 0.55 ppt occultation at phase 0.4845 gives $T_2 \approx 3200$ K, matching the mass
A137	SB, TCE; grazing eclipse of a subgiant at conjunction A; the 0.40 ppt secondary at phase 0.514 is also in the SPOC DV report (0.63 ppt at 0.515); it is shallower than a grazing M dwarf predicts unless $e \sin\omega \approx +0.04$ (2.5σ from Gaia); SPOC TCE, never a TOI
A138	SB; flat-bottomed transit of an M dwarf across an F star at conjunction A, which falls near periastron: a heartbeat is rejected by $\Delta\chi^2 = 5150$; no secondary (< 0.3 ppt), as the geometry predicts ($b_B \approx 1.10$)
A139	SB; transit of a 0.14 $M_\odot$ M dwarf at conjunction A; 5 events in five pipelines, two recovered from flagged cadences, no single event owning the verdict; the photometric R_1 with Gaia's *e*, ω is 1.59 against the TIC's 1.60 $R_\odot$
A140	SB; flat-bottomed transit of a 0.19 $M_\odot$ M dwarf across a γ Dor-like pulsating F star (depth after prewhitening; the batch's 10.0 ppt was pulsation-inflated); the duration disfavours Gaia's $e \sin\omega$ at 3.6σ (orbit probably near-circular); two sectors
A141	SB; M dwarf grazing a G subgiant at conjunction A, 13° before periastron; a heartbeat excluded ($\Delta\chi^2 = 5391$); conjunction B misses by 1.2 $R_\odot$ and no secondary is seen (< 0.3 ppt); 12 events
A142	SB; late-M dwarf transiting an F subgiant at conjunction A; the 0.65 ppt occultation at phase 0.5012 gives $T_2 \approx$ 3000 K, matching the mass function; the *G* 14.2 neighbour 9.5″ away is excluded by in-eclipse centroids and is not an eclipser itself
A143	SB; occultation of the M-dwarf companion at conjunction B, 13 h after periastron, $i \approx 87.5°$; a heartbeat excluded ($\Delta\chi^2 = 15029$); conjunction A empty as the geometry predicts; 9 events
A144	SB; eclipsing heartbeat binary (Section 3.3, Table 3): an 18 ppt tidal curve with periastron at phase 0.25, an M-dwarf transit at conjunction A on top of it (the tide alone loses by $\Delta\chi^2 > 11000$) and its 2.1 ppt occultation at phase 0.3625, 0.26σ from Gaia's conjunction separation; 23 events; depth on the tidal baseline
A145	SB, G; grazing subgiant + 0.64 $M_\odot$ companion, 18 + 14 events; the secondary at phase 0.49996 ± 0.00018 gives $e\cos\omega = 0.0000 \pm 0.0003$, 4.5σ from Gaia's $e = 0.039$ (Table 4); the batch's 14″ centroid offset was the near-equal neighbour pulling the centroid, the SPOC difference image is on target in three sectors
A146	SB; grazing G + M dwarf, 8 events; a marginal 0.9 ppt secondary (3.5σ) at phase 0.4986 is not quoted; VSX types the star ROT (a type revision, not prior art)
A147	SB, G; grazing eclipse of the Gaia SB2 at conjunction B, 0.27 d before periastron (a heartbeat loses by $\Delta\chi^2 =$ 22713); no eclipse is possible at A; Gaia's period is 4.3σ off (Table 4), which the batch's conjunction offset had extrapolated; the batch's difference-image flag was an artefact

Label	Notes
A148	SB; grazing, both eclipses in every sector (48 + 47 events) with ellipsoidal variation; secondary at phase 0.5019; the batch's 13.1 ppt was both kinds averaged at $P/2$; the companion is probably a stripped post-mass-transfer star of the R CMa/EL CVn kind: 0.30 $M_\odot$ from Gaia's K_1, which the raw spread of Gaia's epoch velocities confirms, with the 5650 K surface the depth ratio requires; $R_1 \approx 3$ $R_\odot$ from the ellipsoidal amplitude, the line broadening and the Gaia catalogues (TIC's 4.8 is inflated by a blend); a 1.5–2× luminosity excess hints at unresolved third light; no spectrum exists in LAMOST or APOGEE
A149	SB; grazing near-circular F + K dwarf, 24 + 20 events; the secondary at phase 0.50000 ± 0.00025 gives $T_2 \approx$ 4700 K, matching the mass function
A150	SB; grazing transit of a subgiant at conjunction A, 18° past periastron, with a ~1 ppt tidal dip under the eclipse (Section 3.3); a heartbeat alone loses by $\Delta\chi^2 \geq 10569$; conjunction B misses (< 0.19 ppt); 8 events; QLP sector 103's 12.6 ppt was an aperture artefact
A151	SB; eccentric and grazing, both eclipses on Gaia's conjunctions, the secondary at phase 0.5593 ($e \cos \omega$ +0.093, Gaia's +0.098 ± 0.017); depths in QLP, 18–21 ppt undiluted; 33 + 38 events; common-mode eclipse timing variations of ±4 min in both eclipses, SPOC-independent, suggest a third body; TESS sectors 112–114 (2027) discriminate a constant period change from light-time, the prediction registered beforehand
A152	SB; M dwarf (0.3 $M_\odot$) grazing the F star at conjunction A ($i \approx 86°$), 14 events; 22 ± 1.5 ppt corrected for dilution; the G 12.4 neighbour 18″ away (probably a physical companion) excluded at 17.6–27.4σ per sector in six difference images, and the batch's off-target flag was its pull on the light centroid; a tentative 1.0 ± 0.2 ppt occultation at phase 0.207 is not quoted
A153	SB; flat-bottomed transit of a 0.24 $M_\odot$ M dwarf at conjunction A, 8 + 8 events; 23 ± 1 ppt corrected for dilution; the 1.32 ppt occultation at phase 0.3034 matches Gaia's 0.305 ± 0.021, $e \cos \omega = -0.308$; on target in six of six difference images (5.6–8.2σ), the flag a pixel-boundary artefact; the batch's conjunction offset is −0.028, not +0.030
A154	SB; M dwarf (0.31 $M_\odot$) transiting a γ Dor-like pulsator (±3%) at conjunction A, 74 events; 32 ppt corrected for dilution; 1.9 ppt total occultation at phase 0.580 (10.7σ), 0.6σ from Gaia's 0.563 ± 0.031, $e \cos \omega = 0.126$; on target in seven of eight difference images once the pulsation is modelled per pixel, the flag from out-of-eclipse windows that fell inside the 3.2-h eclipse; the VSX ROT entry at 0.884 d is the pulsation
A155	SB; grazing transit of a 0.33 $M_\odot$ M dwarf at conjunction A ($b \approx 0.88$), 7 + 8 events, the batch's two being a cadence-and-quality artefact; 63.7 ppt in PDCSAP; the 5.85 ppt occultation at phase 0.50029 and the equal durations make the orbit circular, as Gaia's $e = 0.031 \pm 0.027$ allows; on target in three of three SPOC target-pixel difference images (10–134σ), the flag sub-pixel WCS scatter; spotted, with rotation near-synchronous
A156	SB; flat-bottomed transit of the F star by a 0.47 $M_\odot$ M dwarf at conjunction A, 14 + 15 events; the 7.9 ppt occultation at phase 0.5128 matches Gaia's 0.527 ± 0.020, $e \cos \omega = +0.020$; on target in four of four difference images (47σ in the SPOC pixels), the flag a WCS zero-point artefact; Gaia RUWE 7.0 suggests a third body
A157	SB, TCE; K dwarf (0.8 $M_\odot$, about the giant's surface brightness) transiting a spotted K giant at conjunction A and totally occulted at B (86 ppt, phase 0.50005), 14 + 15 events with K2 C16 extending the baseline to 2018; circular; depths are SPOC PDCSAP, QLP running 7% deeper; on target in seven of eight difference images (≥ 30σ against every neighbour), the flag a sub-pixel PSF offset; a SPOC multi-sector TCE at $P/2$; the VSX TAOS entry (VAR, 13.93 d) is the giant's rotation
A158	SB, TCE; K dwarf (0.65 $M_\odot$) transiting at conjunction A and totally occulted at B (33 ppt, phase 0.4924), 13 + 14 events; $e \cos \omega = -0.012$ and $e \sin \omega = -0.225$ on Gaia's orbit; no neighbour could make a 12% dip, and the SPOC difference image puts it 4.4 ± 3.8″ from the target; a TESS-SPOC multi-sector TCE at $P/2$
A159	SB, G, K; K dwarf (0.82–0.86 $M_\odot$) transiting at conjunction A and occulted at B (12.0 ppt, phase 0.62842), 10 + 10 events; $e \cos \omega = 0.2005 \pm 0.0002$ against Gaia's 0.252 ± 0.013 (Table 4); on target in five of five primary and four of four secondary difference images, the flag a sub-pixel quantization artefact; the batch's conjunction offset is −0.023, not +0.023
A160	SB; grazing G star + 0.5–0.6 $M_\odot$ companion ($i \approx 84°$), 11 + 9 events; the 4.2 ppt occultation at phase 0.50120 ± 0.00005 gives $e \cos \omega = +0.0019 \pm 0.0001$ (Gaia +0.009 ± 0.011); QLP's 16.2 ppt in sectors 97 and 104 is dilution in the 200-s products, the star's own aperture giving 19.1 ppt in every sector; on target in the TESS-SPOC difference image

Label	Notes
A161	SB, K; strongly grazing G8–K0 dwarf + 0.6–0.7 $M_\odot$ companion ($b \approx 1.5$), 41 + 41 events in 13 sectors; the 6.7 ppt occultation at phase 0.50726 ± 0.00005 gives $e \cos \omega = +0.0108 \pm 0.0002$ (Gaia +0.024 ± 0.013); the equal eclipse durations want $e \sin \omega \approx 0$, 2.5–3.3σ from Gaia's −0.0325 ± 0.0099, a tension in the direction of a Lucy–Sweeney bias; on target in seven TESS-SPOC difference images, the G 14.2 neighbour 6.5″ away rejected; the VSX ROT entry at the target is the Gaia classifier's, a type revision
A162	SB; grazing F star + 0.45 $M_\odot$ M dwarf ($b \approx 1.1$, $i \approx 83°$), 10 + 10 events in two sectors; a new 1.38 ± 0.13 ppt occultation at phase 0.5014 (S/N 10.8) gives $e \cos \omega = +0.0022 \pm 0.0011$; a QLP-only star whose own cutout photometry reproduces both eclipses with no dependence on aperture; the batch's 4.7″ offset is 1.0″ in the star's own difference image
A163	SB; grazing spotted G star + 0.6–0.7 $M_\odot$ companion ($b \approx 1.4$), 23 + 25 events in five sectors, both eclipses in QLP, TESS-SPOC, SPOC and TARS; the 2.4 ppt occultation at phase 0.50003 ± 0.00015 gives $e \cos \omega = 0.0000 \pm 0.0002$; Gaia's period is 2.2σ short and the attribution holds at Gaia's own epoch (−1.3σ); on target in five SPOC target-pixel difference images
A164	SB; eccentric ($e = 0.23$) subgiant + 0.6 $M_\odot$ K dwarf; the one eclipse is the companion's flat-bottomed occultation at conjunction B near periastron (with the primary's radius fixed by the orbital speed, a transit would need it to supply only 48% of the light), 10 events; conjunction A is empty (< 0.23 ppt) exactly as the geometry predicts, the sky separation there being 1.1–1.2 ($R_1 + R_2$); on target in six difference images
A165	SB, K; circular G star + 0.23 $M_\odot$ M dwarf ($R_2 \approx 0.25\ R_\odot$), 353 transits in 23 sectors; a new 1.34 ppt occultation at phase 0.5005 ± 0.0010 (S/N 24) gives $e \cos \omega = +0.0008 \pm 0.0016$; Doppler beaming of +0.30 ppt with the sign and size Gaia's K_1 predicts ties the photometric binary to the RV orbit independently; the eclipse and the 4.2 ppt ellipsoidal variation both give $R_1 = 1.3$–$1.5\ R_\odot$ against TIC's 2.06, an open point that does not touch the verdict; on target by $\Delta\chi^2 > 13{,}000$
A166	SB; grazing transit of a 0.5 $M_\odot$ M dwarf across a δ Sct/γ Dor hybrid F star at conjunction A, 27° before periastron (a heartbeat tested and rejected, $\Delta\chi^2$ 141–3500 per event), 5 events over 2200 d; the depths scatter from 13 to 22 ppt with the pulsation; no occultation, as the geometry predicts (the sky separation at B exceeds $R_1 + R_2$); on target in three SPOC target-pixel difference images
A167	SB, G; the Gaia double-lined binary itself: near-twin grazing F/G stars ($q = 0.965$, $i \approx 79°$), 30 + 30 events of equal depth; the secondary at phase 0.49985 ± 0.00012 gives $e \cos \omega = -0.0002 \pm 0.0002$ (Gaia +0.0256 ± 0.0097, 2.7σ); Gaia's period is 8σ off yet both conjunctions match at Gaia's own epoch (Table 4); the batch's 7.0″ offset was an artefact of the measuring box, the star's own localisation 0.9–3.1″ on target in five sectors; a ±3 min eclipse-timing variation common to both eclipses suggests a third body (RUWE 2.70)
A168	SB; G2V star + 0.11 $M_\odot$ M dwarf ($R_2 \approx 0.145\ R_\odot$, $b \approx 0.75$), flat-bottomed, 15 events; published as a planet candidate by the DTARPS-S search (Melton et al. 2024, 13.4 $R_\oplus$, never a TOI), which the Gaia orbit reclassifies as a star, so the entry is a reclassification rather than a new detection; a tentative 0.5–0.9 ppt occultation is not quoted; on target in the TESS-SPOC difference image
A169	SB; δ Sct/γ Dor hybrid F star + 0.21 $M_\odot$ M dwarf ($b \approx 0.5$), flat-bottomed, 50 events; the batch's depth scatter was the pulsation (8–10 ppt modes of about 2 h beating across the 5-h eclipse) and a uniform ×0.63 dilution in QLP sector 17 alone; a new 1.15 ± 0.3 ppt occultation at phase 0.50 in all six sectors, $e \cos \omega = 0.000 \pm 0.005$ (Gaia 0.14 ± 0.08); the 3–4.5 ppt ellipsoidal variation matches Gaia's K_1; on target in three TESS-SPOC difference images
A170	SB; eccentric ($e = 0.23$) F star + 0.29 $M_\odot$ M dwarf, flat-bottomed, 14 transits (sector 20 adds four to the batch's ten); a 0.8 ppt brightening at periastron, where Gaia puts it; an occultation hint of 0.33 ± 0.08 ppt at Gaia's conjunction-B phase 0.36 is not quoted; the batch's 6.4″ offset was the uncalibrated pixel scale and the wings of the point-spread function pulling toward the box centre, the star's own fits 0.4–1.7″ on target in six products; RUWE 9.1 implies an unresolved companion within about 1″
A171	SB; F star + companion of at least 0.93 $M_\odot$ ($f(m) = 0.178\ M_\odot$), 8 transits (the batch's three, with five recovered from flagged cadences; no single event carries the result); no occultation deeper than about 0.8 ppt where a main-sequence companion that massive would show 10–18 ppt at Gaia's $e \sin \omega$, so either $e \sin \omega \geq 0.06$ (3–5σ from Gaia, more eccentric) or a dark companion such as a close pair of M dwarfs, a tension that concerns the companion and not the eclipse; ellipsoidal variation 0.3–0.9 ppt; on target in four products

Label	Notes
A172	SB, TCE; circular F subgiant + 0.3 $M_\odot$ M dwarf ($b \approx 0.45$, $i \approx 86°$), flat-bottomed transit, 23 events in nine sectors, with a 1.4 ppt total occultation at phase 0.50002 ± 0.00034 (S/N 18), $e \cos \omega = 0.0000 \pm 0.0005$; QLP's 17–28 ppt depth scatter is dilution (PDCSAP constant at 23–25 ppt); a TESS-SPOC multi-sector TCE for each eclipse (sectors 56–69), never a TOI, whose single-sector secondary centroid 70″ toward the *G* 9.2 neighbour is overruled by nine difference images
A173	SB, TCE; eccentric ($e = 0.36$) G dwarf + 0.22 $M_\odot$ M dwarf transiting at conjunction A ($i \approx 87°$), 7 events (23.0–23.3 ppt in SPOC 2-min data); a 0.49 ± 0.08 ppt grazing occultation at phase 0.3309 in the 2-min data sits 0.2σ from Gaia's predicted conjunction B, where the geometry puts the companion right at the limb, but its three events disagree in depth and it is not quoted; a SPOC TCE for the primary, never a TOI; on target in seven SPOC target-pixel difference images
A174	SB; circular grazing F/G star + K1–K2 dwarf ($q = 0.62$; the depth ratio gives $T_2 \approx 4750$–5000 K, the temperature of the mass-function companion), 23 + 23 events; the 11.6 ppt occultation at phase 0.50012 ± 0.00004 gives $e \cos \omega = +0.0001 \pm 0.0001$; the batch's "peak not on target" flag was a half-period stack mixing the two unequal eclipses, the star's own TESS-SPOC difference images on target by $\Delta\chi^2 > 9{,}000$ for both kinds; the eclipses and the ellipsoidal variation give $R_1 \approx 1.5$ $R_\odot$ against TIC's 2.0, the companion's light inflating the single-star radius
A175	SB; G star + 0.27 $M_\odot$ M dwarf transiting ($b \approx 0.8$), 6 events (one recovered from a flagged window); a new 1.45 ± 0.11 ppt occultation at phase 0.5357 ± 0.0003 gives $e \cos \omega = +0.0560 \pm 0.0005$ (Gaia +0.028 ± 0.019); three APOGEE DR17 radial velocities phased on the eclipse ephemeris alone follow Gaia's orbit with an rms of 0.30 km/s, an independent spectroscopic tie; on target in four difference images, with no neighbour brighter than *G* 20 within 60″
A176	SB, TCE; grazing F/G star + 0.44 $M_\odot$ M dwarf ($i \approx 85°$; the transit fit gives $R_1 = 1.02$ $R_\odot$, TIC 1.05), 17 events (two recovered from flagged cadences); a new 1.54 ± 0.07 ppt occultation at phase 0.49681 ± 0.00017 gives $e \cos \omega = -0.00502 \pm 0.00026$, 19σ from zero: Gaia's 2.3σ eccentricity of 0.019 is real, the counter-example to the spurious small eccentricities of A145, A159 and A167; near-synchronous starspots (the VSX ROT entry at 3.97 d is a type revision); TESS-SPOC TCEs for both eclipses, never a TOI; on target in two difference images
A177	SB, TCE; grazing spotted G star + 0.34 $M_\odot$ M dwarf ($i \approx 83°$; $R_1 = 1.00$ $R_\odot$ from the transit fit), 26 events (three recovered from flagged cadences); a new 2.53 ± 0.10 ppt occultation at phase 0.50020 ± 0.00028, so the orbit is circular and Gaia's $e = 0.05 \pm 0.04$ is not significant; near-synchronous spot modulation growing from 4 to 10 ppt over the TESS era; a 2-min SPOC target with TCEs for both eclipses, never a TOI; on target in three difference images, with no neighbour brighter than *G* 19.3 within 60″
A178	SB; circular grazing F star + K dwarf (0.7–0.8 $M_\odot$; the depth ratio gives $T_2 \approx 4800$ K, the temperature of the mass-function companion; $i \approx 81$–$82°$), 52 + 50 events in eleven sectors, conjunction A matched at +0.1σ throughout; the 9.3 ppt occultation at phase 0.50019 ± 0.00005 gives $e \cos \omega = +0.0002$; a QLP dilution step in sector 94 scales both eclipses by 0.87 together; an isolated field, the nearest star brighter than *G* 17 at 60″; on target in three SPOC target-pixel difference images; no prior record of any kind
A179	SB, TCE; grazing near-twin G + G double-lined binary (1.07 + 1.15 $M_\odot$, $i \approx 83°$), 34 + 37 events of 28.9 and 25.6 ppt; the deeper eclipse is at conjunction B because Gaia's "primary", the star with the larger velocity amplitude, is the cooler and less massive one, a labelling note rather than a conflict; the phase separation 0.50002 ± 0.00002 makes the orbit circular; spotted and X-ray bright (2RXS), RS CVn-like; a 2-min SPOC TCE at half the period with an odd–even statistic of 246, never a TOI; on target by $\Delta\chi^2 > 21{,}000$
A180	SB, K; eccentric ($e = 0.36$) G star 1.39 $R_\odot$ + 0.69 $M_\odot$ dwarf, $i = 89.1°$; a full transit at conjunction A and a total occultation at B whose 9.9% depth is the companion's light share (9.87% predicted), six + five events 2020–2024 in two pipelines; Gaia's period is 1.5σ short, which is the whole of the 0.6–0.9 d by which its conjunctions ran early; *e* and ω within 0.9σ
A181	SB, K; eccentric ($e = 0.46$) G star 1.32 $R_\odot$ + 0.5 $M_\odot$ M dwarf: a grazing transit ($b = 0.78$) at conjunction A and the total occultation of the M dwarf at B (1.74% predicted from its light), 6 + 6 events 2018–2025 in three pipelines; *e* 0.464, ω 201.6° against Gaia's 0.476 ± 0.010, 198.9 ± 2.7; RUWE 2.22 hints at a third star; QLP's shallower sector-98 depths are its aperture, the pixels giving the same depths as before
A182	SB; K giant 5.4 $R_\odot$ + a hotter (about 6950 K), smaller companion of about 1.5 $M_\odot$, $e = 0.34$; the deeper eclipse is the giant's total occultation of the companion at conjunction B near periastron (T_0; flat for 15 h), the shallower the companion's rounded transit at A, three + two events 2018–2024; a QLP-only star, so the second pipeline is the checker's own cutout photometry, aperture-independent, with the deficit 0.8″ from the target; *e* and ω within 1.8σ

Label	Notes
A183	SB; K giant 5.6 $R_\odot$ + a hotter (about 6000 K) companion of about 1.35 $M_\odot$, e = 0.17; the deeper eclipse is a partial occultation of the companion at conjunction B near apastron (T_0), the shallower its full transit at A, three + four events 2019–2025 in four pipelines; a 2007 RAVE velocity falls on the orbit; e and ω within 1.1σ
A184	SB, V, TCE; spotted K giant 6.8 $R_\odot$ + 0.85–0.94 $M_\odot$ G dwarf, e = 0.30; the deeper eclipse is the dwarf's total occultation at conjunction B (T_0; flat for 43 of 53 h, 23 events), the shallower its 32-h transit at A with spot-crossing bumps (19 events, 20 min of timing jitter), 42 sectors 2018–2025; the VSX and ASAS-SN rotational entry (54.6 d) is the giant's spot modulation and EA is added; {ECW} = −0.181 against Gaia's −0.158 ± 0.015; a TESS-SPOC threshold-crossing event at 50.85 d was never promoted
A185	SB, V, K; spotted G/K giant 8.8 $R_\odot$ + 0.8–1.0 $M_\odot$ dwarf on a circular orbit ($e \cos \omega$ = +0.0002 ± 0.0002; Gaia e = 0.018 ± 0.009); the deeper eclipse is the dwarf's total occultation at conjunction B (T_0; flat for 31 h, 7 events), the shallower its transit at A with spot crossings on the floor (4 events), 2019–2025; the VSX rotational entry (53.5 d) is the spot modulation and EA is added; Gaia's period is 2.1σ long; the occultation's ingress wants a giant 15% smaller than the TIC radius
A186	SB; K giant 3.4–3.8 $R_\odot$ + 0.55–0.6 $M_\odot$ dwarf, e = 0.34; a transit at conjunction A and the dwarf's total occultation at B (flat for 20 h), two + two events 2019–2023; the lag the classifier saw was Gaia's period (+1.7σ); APOGEE velocities of 2015–2018, 61 orbits earlier, fit the eclipse ephemeris at 0.34 km/s rms (5.9 km/s with Gaia's period); the G 14.9 star 3.7″ away, in the target's pixel, is excluded by the in-eclipse centroid; the binary was known from APOGEE (Price-Whelan et al. 2018), the eclipses were not
A187	SB; G/K giant 4.5 $R_\odot$ + 0.6–0.7 $M_\odot$ dwarf, e = 0.33; a transit at conjunction A and the dwarf's total occultation at B (flat for 26 h), four + two events 2019–2025 in four pipelines; the durations want {ESW} = 0.32 against Gaia's 0.372 ± 0.012 (2–2.5σ, a soft tension); nothing within 60″ could host the eclipses
A188	SB; eccentric (e = 0.55) F star + 0.55 $M_\odot$ companion (about 3900 K); a curved transit at conjunction A and the companion's flat total occultation at B, five + seven events 2018–2025 in five pipelines; the occultations precede the transits by 3.831 d where Gaia's elements predict 3.35 ± 0.16 d, a 2.9σ tension in {ECW} (0.544 against 0.585 ± 0.014) that stops short of the rule; and the occultation times wander by about 50 min against the transits over two years, non-monotonically and in every pipeline, which neither apsidal motion nor light-travel time explains (a third body perturbing e, or spots on the occulted star), so occultation predictions carry ±40 min
A189	SB, K; subgiant 3.6 $R_\odot$ (about 5200 K) + a hotter (about 6100 K) dwarf of about 1.05 $M_\odot$, e = 0.34; the deeper eclipse is the dwarf's total occultation at conjunction B (T_0; flat for 8.6 h), the shallower its transit at A: the companion has the higher surface brightness, so the labelling the classifier questioned is right, and swapping it would put every eclipse 11.5 d from its prediction; two + four events 2019–2025; e and ω within 0.9σ
A190	SB, TCE; K4–5 dwarf 0.70 $R_\odot$ + 0.5 $M_\odot$ early-M dwarf, e = 0.415, i = 88.5°; 23 flat-bottomed transits at rms 0.36 min and 20 grazing V-shaped occultations at phase 0.7136, 2018–2025; {ECW} = 0.33 against Gaia's 0.367 (at Gaia's e the phase needs ω = 42° against 33.5 ± 2.8; about 2σ with e free); TESS-SPOC raised both eclipses as threshold-crossing events at alias periods (38.1 and 27.2 d) and promoted neither
A191	SB; G subgiant 1.47 $R_\odot$ + 0.73 $M_\odot$ K dwarf at $i \approx 90°$, e = 0.092: a 27% annular transit at conjunction A and a 15% flat total occultation at B (the K dwarf's light share), 14 events 2019–2025, every orbital element within 1.1σ of Gaia; it is WASP0-TR-06 of Kane et al. (2005), a single partial event of 2000 with a spurious 6.03-d period that was rejected as a 3.9 R_J planet, and nothing since gave it a period or called it an eclipsing binary; a totally eclipsing G + K pair for double-lined spectroscopy
A192	SB, V; giant 12–16 $R_\odot$ (3–4 $M_\odot$) + a hot companion of 2–3 $M_\odot$ (about 14,000 K, late B) supplying 22–35% of the TESS light, $e \approx 0.01$; the deeper eclipse is the giant partly hiding the companion at conjunction B (T_0; 30 h, four events), the shallower the companion crossing the giant's limb at phase 0.5001 (four events; the first fits, at Gaia's phase 0.512, had missed it); a QLP-only star checked with the checker's own cutout photometry and difference images; the VSX and ASAS-SN rotational entry (49.65 d) is replaced by EA; the companion is hot for the mass allowed, and an ultraviolet measurement would settle it
A193	SB; F/G star 1.12 $M_\odot$ + 0.69 $M_\odot$ K dwarf, e = 0.457, i = 88.3°, both eclipses grazing (b = 0.66 and 0.84), four + three events 2018–2025 in six pipelines; {ECW} = 0.443 against Gaia's 0.4426 ± 0.0138; the G 12.5 star 12.9″ away (WDS ARA 337) is a background star by parallax, excluded by the difference image and by TGLC's deblended depth

Label	Notes
A194	SB, V; subgiant 3.5–4.1 $R_\odot$ + 0.8–1.0 $M_\odot$ companion, e = 0.43 (Gaia's); seven grazing transits at conjunction A and an ingress, 2019–2025, and no occultation (< 0.03%), as the geometry predicts, the separation at B being 1.19 times that at A; the VSX rotational entry (72.8 d) is probably the subgiant's rotation, so EA is added rather than substituted; single-eclipse kind
A195	SB; F star 1.2 $M_\odot$, 1.6 $R_\odot$ + 0.73 $M_\odot$ K dwarf, e = 0.489, i = 89.1°: a transit near apastron at conjunction A and the K dwarf's total occultation near periastron at B (flat for 2.5 h), four + five events 2019–2024; the +1.8σ occultation offsets the classifier saw are one phase offset, +1.55σ in {ECW} (joint p = 0.19 against Gaia's covariance), and there is no apsidal motion; nothing brighter than G 19 within 60″
A196	SB; G subgiant 2.2–2.4 $R_\odot$ + a G dwarf of about 1 $M_\odot$ and the same surface brightness (the companion supplies 17% of the light), e = 0.098, i = 89.0°: 36 rounded transits and 44 flat total occultations over 39 sectors 2019–2024 on one ephemeris at 1.0 min rms, no apsidal motion in six years; {ECW} = −0.0845 against Gaia's −0.059 ± 0.011 (2.3σ), and two APOGEE velocities of 2018 prefer the eclipse elements; the Kostov et al. (2025) flag in this field sits on the neighbour 37″ away, not on this star; a well-detached G + G pair for double-lined spectroscopy
A197	SB, CTOI; F star 1.69 $R_\odot$ + 0.57 $M_\odot$ M dwarf, e = 0.36, i = 87.8°: four flat-bottomed transits at rms 0.10 min and five total occultations at phase 0.2792, 2019–2025; e and ω within 0.6σ of Gaia; SPOC 2-min and TESS-SPOC reproduce both; the community planet candidate CTOI 300202855.01 (a single transit of 2024, 7.85 ppt) is the sector-32 occultation, the companion a star by Gaia's mass function
A198	SB; F8/G0 star 1.1 $M_\odot$, 1.4 $R_\odot$ + 0.66 $M_\odot$ K5 dwarf supplying 5.7% of the light, e = 0.314, i = 89.5°: three full transits near periastron at conjunction A and three near-total occultations near apastron at B (13.4 h), 2019–2025; {ECW}, {ESW}, e and ω all within 0.8σ of Gaia; depths from the checker's own cutout photometry because QLP dilutes this field's sectors 101 and 103 by 15–35%; nothing brighter than G 19 within 60″
A199	SB, TCE, CTOI; F/G subgiant 2.7 $R_\odot$, 1.3–1.6 $M_\odot$ + 0.8–0.9 $M_\odot$ dwarf, e = 0.38, i ≈ 86.4°, in the southern continuous-viewing zone: 23 grazing V-shaped occultations of the companion at conjunction B (T_0) from sectors 1 to 98 and nothing at A (< 110 ppm), as Gaia's e and ω require; the one APOGEE velocity phases as B; the SPOC 2-min threshold-crossing event (since 2021) and the community planet candidate CTOI 391925531.01 (2022) are this eclipse, and K_1 = 32 km/s makes the occulted body a star; single-eclipse kind
A200	SB; grazing F star + 0.50–0.57 $M_\odot$ M dwarf (b ≈ 0.95, i ≈ 86°), 3 + 4 events in the only two sectors that observed it; the new 4.17 ± 0.31 ppt occultation at phase 0.49948 ± 0.00031 gives e cos ω = −0.0008 ± 0.0005 (Gaia +0.011 ± 0.011) and recovers the second sector's transit on its own, so no single event owns the period; conjunction A matched at Gaia's epoch to +0.3σ; the batch's 2.1″ offset was the uncalibrated pixel scale, the star's own localisation 0.4–0.6″ on target; the transit wants R_1 = 2.05–2.2 $R_\odot$ against TIC's 2.37, an open point that does not touch the verdict
A201	SB, V; grazing (b ≈ 1.1–1.2) spotted G dwarf + 0.39 $M_\odot$ M dwarf on a circular orbit, 18 + 11 events in five sectors; the new 2.3 ppt occultation at phase 0.5001 ± 0.0002 gives e cos ω = +0.0003 ± 0.0003 (Gaia −0.028 ± 0.019, 1.5σ toward circular); the VSX rotational entry is Gaia's classifier's and the eclipses revise it; the batch's 3.9″ offset was the uncalibrated pixel scale, the deficit 0.4–1.2″ on target in three sectors; TESS-SPOC gives 26.2–26.8 ppt in all five
A202	SB, G; grazing (b = 1.04, i ≈ 76°) G subgiant 2.1 $R_\odot$ + 0.5 $M_\odot$ M dwarf on a circular orbit, spotted (RS CVn-like humps of ±10 ppt), 36 + 36 events in four sectors; the 3.84 ppt occultation at phase 0.49993 ± 0.00045 gives e cos ω = −0.0001 ± 0.0007 against Gaia's +0.051 ± 0.015 (3.5σ), and Gaia's period is 3.4σ long (Table 4); conditioned on a circular orbit both conjunctions still match at Gaia's epoch; the G 11.4 star 9.6″ away (WDS ARA 52, an optical pair by parallax) is excluded by the sign of the pixel flux-loss gradient in both eclipse kinds and three sectors; eclipse timings jitter by ±6 min with the spots
A203	SB; grazing (b ≈ 1.05) subgiant 1.9 $R_\odot$ + 0.56–0.62 $M_\odot$ K7/M0 dwarf in a crowded field at b = +0.7° (the QLP depth is diluted; TESS-SPOC's crowding-corrected value is 37–40 ppt), 21 + 21 events in six sectors; the new 7.6 ppt occultation at phase 0.50020 ± 0.00019 gives e cos ω = +0.0003 ± 0.0003 (Gaia +0.006 ± 0.018); both conjunctions match at Gaia's epoch (+0.9 and +0.6σ); the duration ratio wants e sin ω ≈ +0.007 against Gaia's −0.036 ± 0.013, about 3σ toward circular but soft for grazing trapezoids; the batch's 3.1″ offset is a crowding artefact, and TGLC's point-spread photometry at the Gaia position recovers the full depth

Label	Notes
A204	SB, V; the Gaia double-lined binary itself: near-twin solar-type stars (1.05 + 1.08 $M_\odot$, i = 84–85°), both eclipses partial, 25 + 27 events of 30.4 and 29.4 ppt in seven sectors (about 36 and 35 ppt undiluted); the secondary at phase 0.51577 ± 0.00007 gives $e \cos \omega$ = +0.0248 against Gaia's +0.0085 ± 0.0083 (1.9σ, toward more eccentricity), and the geometry wants $e \sin \omega$ = +0.005 to +0.010 as Gaia's +0.007; the VSX rotational entry is Gaia's classifier's and the eclipses revise it; an eclipse-timing variation of about 6 min common to both eclipses across sectors 99–102, which are QLP-only, is open: a third body or a pipeline systematic; on target by $\Delta\chi^2 >$ 4700
Probable eclipsing binaries with unresolved hosts (not counted)	
A13	host one of two G 13.7 stars; or 2.157370 d if twins
A14	host Gaia DR3 5459360868712645888; one sector
A15	host TIC 261126769 (≳ 10% deep on it); or 28.97 d if twins

Note: T: TESS magnitude from the TIC (every row). P and T_0: the ephemeris of the independent re-analysis or timing fit, to its quoted precision; T_0 is the epoch of the primary (deeper) eclipse, BJD_{TDB} = BTJD + 2457000. Depths and durations are aperture values in the named pipeline's flux (Section 2.8), not corrected for dilution. For A25, A27, A41 and A53 one eclipse kind has been observed; the source tabulates its depth in the secondary column, and its times are those given as the next primary. A51's primary is seen only as an ingress of at least 8.6%. Next primary: geocentric UTC, from BJD_{TDB} with the light-travel time to the solar-system barycentre in the star's direction removed (up to ±8.3 min) and TT − UTC = 69.184 s; an observer's local correction is at most 21 ms. Most predictions are good to a few minutes; A31, A32, A35–A38, A40 and A41 to 0.02–0.10 d; A1 and A2 to ±0.004 d and A19 to ±0.005 d (formal errors of the discovery ephemerides); A72 to ±0.002 d; A73 to ±0.3 d. None of these includes event-to-event timing scatter or A109's eclipse-timing variations of about 30 min. The Supplement entries give the secondary epochs and the next secondaries. [a] A1: T_0 is the shallower eclipse; the next primary listed is the deeper one. [b] A3: fold period and box epoch; no timing fit, so no prediction. [c] A40: Gaia's period (one event of each kind). [d] A53: the single interval between its two events. [e] A73: mean period of the primaries, which varies (single cycles 17.39–17.85 d); T_0 is the first of two clean, consecutive primaries (BTJD 2940.800 and 2958.651); the eclipses are fading, and duration and depths give the range (Section 3.2). [f] A9: the period first catalogued, 13.4166 d, was a fold alias. [g] The deeper eclipse is at conjunction B of Gaia's orbit; T_0, the depths and the next primary refer to it, and its duration, which differs from the A eclipse's, is in the Supplement. Origin: App, eclipses by appointment (Section 2.5); Fold, Gaia-orbit period fold at P < 20 d (Section 2.4); Batch, the same fold at G 10–11.5 run in batch, followed by the per-object re-analysis (Sections 2.4, 3.2); MD, M-dwarf transit searches (Section 2.3); Ast, edge-on Gaia astrometric orbit (Section 2.6). Notes: SB, a known spectroscopic binary before this work (the eclipses are new); G, Gaia DR3 orbit solution contradicted by the eclipses (Table 4); K, in the unvetted machine-learning list of Kostov et al. (2025, table 2); TCE, an unvetted SPOC or TESS-SPOC threshold-crossing event (72 of the first 104); V, a VSX entry of a non-eclipsing type (Table 5); CTOI, an ExoFOP community TOI (its own secondary eclipse); R, a prior mention of the eclipses (Bryant et al. 2023); HP, eclipse detected on the Harvard plates (17 of the 31 examined); H, host component or a neighbour within one TESS pixel not separated; bc, by-catch, found at a period other than that of the Gaia orbit.

Appendix B. Data and code availability

The unified scientific review register is scientific-review-register.csv. It groups 822 reconciled orbit-informed R/C/P reviews by scientific outcome and records method, population, physical verdict, unique-count status and preserved evidence links. The following original registers and directories retain compatible identifiers and archival provenance.

All input data are public: TESS light curves, target-pixel files and full-frame-image cutouts from MAST; Gaia DR3 from the ESA Gaia archive; catalogues from VizieR, SIMBAD, ExoFOP, the NASA Exoplanet Archive and the AAVSO VSX; the Harvard plates from DASCH. The analysis code (Python, with NumPy, Astropy, Matplotlib and Requests), the per-object re-analysis reports, the fitted ephemerides behind Table A1, the figure sources, the object table (`tables/objects.csv`, one row per object with the analysis file behind it), the candidate table (`tables/batch_candidates.csv`: the 266 rows of the G 10–11.5 fold, Section 3.2, with each row's status after review and, for the original 68 individual reviews, their counted labels or non-EB outcomes; the 409 R-labelled reviews have separate physical and novelty statuses without new counted labels) and the Supplement are in a public data deposit, available from the author, under CC BY 4.0, with the code also under the MIT licence. The code was written for this work and has not been packaged for reuse; every light curve it reads is re-fetched from MAST. Files in the deposit written before 2026 October 12 use an older labelling in which this paper's A20–A71 are A21–A72 (HIP 1060, A47 here, is their A48); the object table uses this paper's labels throughout. The ASAS-SN archival fold of Section 4.3 (2026 October 14) postdates the deposit version cited above and is added in its next version. An earlier working draft of this paper, whose main text carries the full entries for the catalogue corrections B1–B11, the probable blends A13–A15 and the other objects of Table 6, is in the deposit and in the public repository (github.com/Snowstar38/eclipses-by-appointment), beside the current version, the object table and the candidate table. The report of the internal adversarial review, the response to it and the consistency checks are kept with the paper's sources in the same deposit.

The R-labelled review data are supplied as `tables/reviewed_extension_2026-10-30.csv` and, in the public repository, `reviews/2026-10-30/`: the original 70 per-object reports, compact measurement evidence, report hashes, review credits and unresolved follow-up tests. This paper is a working v1.0 draft and has not been formally submitted. Historical draft files are retained as provenance, not as submitted or accepted versions.

The fainter SB1 period-fold evidence register is `tables/returned_review_2026-10-30.csv` in the paper sources and `returned-review.csv` at the public repository root. C01-C24 are preserved; C25-C72 follow original roster order. Older reports remain under `reviews/c9a-return/`; all 55 new reports, catalogue receipts and dated tier reconciliation are under `reviews/cloud72-completion-2026-10-30/`. The register records controls, known recoveries, exclusions, physical verdicts and next tests.

The dated catalogue reconciliation, coverage matrix and retained query evidence are in `reviews/novelty-reconciliation/README.md`. It applies the original seven-group catalogue standard; the later coverage recovery and R071–R140 checks are retained in `reviews/2026-10-30-next70/`, all included systems have complete scoped catalogue coverage; historical reports remain archived with their original novelty labels.

R141-R174 and their full reports, compact numeric evidence, audit records and catalogue receipts are retained under `reviews/2026-10-30-final34/`. The current R register contains 724 reviews; R001-R140 retain their earlier keys and reports.

The conjunction-selected reconciliation is supplied as `tables/appointments_review_2026-10-31.csv` and public appointments-review.csv; reports, deciding evidence and catalogue receipts are under `reviews/2026-10-31-appointments/`. P001-P023 follow the preserved queue order. Epoch kind, period origin and uncertainty qualifications accompany every row. These provenance keys do not define a separate search method.

The current period-fold and exploratory adjudications are supplied in public gap-review.csv; the dated evidence archive retains the full attempt ledger, reports and catalogue scope. New epochs, error meanings and limitations remain per-object quantities.

The fixed magnitude comparison is supplied in magnitude-review.csv, with full reports, measured epochs and error meanings in the dated archive and Supplement.

The separate 250-target period-free review register preserves all selected physical outcomes, catalogue-check status, source receipts and full-report links. Its screening code and selection ledger are retained with the `reviews/2026-11-05-period-free/` evidence archive. The four earlier supplement PDF volumes are dated evidence snapshots; the fifth volume supplies the period-free review. The reconciled registers and present text govern current claim status, including prior-known removals and the A74 tier update, without rewriting historical reports.

Appendix C. Physical classification and review evidence

The reconciled scientific review register contains 830 individual R/C/P reviews, of which 725 are included unique additions (623 confirmed and 102 probable). Together with 196 active A-labelled claims, these produce the 921 accepted claims (808 confirmed and 113 probable). The register also retains known recoveries, pending catalogue novelty, 29 physically plausible reviews, and non-eclipse or attribution failures outside the accepted claim sample. Review IDs preserve provenance rather than order of discovery or physical rank. The earlier 917 count is corrected by 17 prior-known removals and 13 already accepted additions, with A74 and C57 tier updates, producing the earlier 913-system baseline. Eight subsequent eligible dip systems bring the current total to 921. The original ephemeris panels and dated reports remain measurements of their time; current claim flags are those in this register and the accepted ledger.

The separate 250-target mixed-method selected-review register and evidence archive include period-free dips and complementary brightening, fold-only and near-threshold leads. They document 22 confirmed, 33 probable, 48 plausible, 138 refuted, four non-eclipse-variable and five unresolved outcomes. The 55 physical confirmed/probable assessments include six known recoveries, 41 unknown catalogue-novelty cases and eight accepted additions after scoped catalogue and claim-eligibility review. The eight are R725-R732 and are included here; their repeated signals are one confirmed and seven probable systems. The 41 unknown cases remain excluded because live catalogue gates could not be recovered. Probable unresolved blends are not securely attributed to the named Gaia orbit.

The table groups reviews by current scientific disposition. A known prior eclipsing system may itself remain physically plausible; the 29 plausible physical verdicts therefore span disposition groups. Report links lead to the original or reconciled full reviews, including limitations and next checks.

Included confirmed systems (623)

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R001	373138363	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Obtain stellar-radius constraints and joint RV/eclipse characterization.
R002	270181578	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Characterize high-RUWE multiplicity with high-resolution imaging and better primary mass/radius; then joint light/RV fit.
R004	362610608	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Compare per-sector apertures to explain s99 depth decrease; improve PRF and stellar characterization.
R005	399501645	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Improve weak secondary shape and stellar-radius/dilution characterization.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R006	295945076	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine secondary timing and Gaia small-e orbit; multi-sector calibrated pixels
R007	372376841	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Resolve SAP/PDCSAP dilution difference and component radii
R008	416525506	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Inspect S55 cycle232 background-sensitive event; refine component geometry
R009	451493289	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Resolve catalog radius versus joint eclipse geometry; sector baseline variation
R010	348737250	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine component radius/light ratio and calibrated localization
R011	158545440	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine component radii, dilution and eccentric orbit with joint RV/eclipse modeling.
R012	339787086	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Jointly refit RV and eclipse timings; assess sector-dependent baseline effects and the separate 281-day astrometric solution.
R013	341432079	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine component radii, dilution and eccentric orbit with joint RV/eclipse modeling.
R014	144749615	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine component radii, dilution and eccentric orbit with joint RV/eclipse modeling.
R015	304033253	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine component radii, dilution and eccentric orbit with joint RV/eclipse modeling.
R016	117919258	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Obtain a second sector to verify the secondary ephemeris and reduce extrapolation uncertainty.
R017	97509220	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Obtain another eclipse sector to improve local period and test the weak secondary against phase controls.
R018	148618714	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine component radii, dilution and eccentric orbit with joint RV/eclipse modeling.
R019	266540667	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine component radii, dilution and eccentric orbit with joint RV/eclipse modeling.
R020	285145433	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine component radii, dilution and eccentric orbit with joint RV/eclipse modeling.
R021	240360239	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Obtain joint RV/eclipse and stellar-radius constraints for characterization.
R022	341735257	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Obtain joint RV/eclipse and stellar-radius constraints for characterization.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R023	306553279	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Characterize temperatures/component assignment and test the covered null secondary with resolved multiband photometry.
R024	56534111	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Obtain joint RV/eclipse and stellar-radius constraints for characterization.
R025	31357336	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refit the Gaia small-e orbit jointly with both eclipse ephemerides.
R026	342454977	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Obtain joint RV/eclipse and stellar-radius constraints for characterization.
R027	455214882	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Constrain grazing geometry and component light with resolved multiband photometry.
R028	140477577	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Obtain joint RV/eclipse and stellar-radius constraints for characterization.
R029	190972398	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Obtain joint RV/eclipse and stellar-radius constraints for characterization.
R030	349273287	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Calibrate a multi-source empirical PSF and obtain less diluted eclipse photometry.
R031	273724854	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Resolved multiband eclipse photometry / joint RV+light-curve component modeling
R032	365181558	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Resolved multiband eclipse photometry / joint RV+light-curve component modeling
R033	370251176	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Resolved multiband eclipse photometry / joint RV+light-curve component modeling
R034	119301005	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Resolved multiband eclipse photometry / joint RV+light-curve component modeling
R035	377656542	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Resolved multiband eclipse photometry / joint RV+light-curve component modeling
R036	194609413	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Resolved multiband eclipse photometry / joint RV+light-curve component modeling
R037	323434013	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Resolved multiband eclipse photometry / joint RV+light-curve component modeling
R038	327209313	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Resolved multiband eclipse photometry / joint RV+light-curve component modeling
R039	330659284	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Resolved multiband eclipse photometry / joint RV+light-curve component modeling
R040	82620363	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Resolved multiband eclipse photometry / joint RV+light-curve component modeling

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R041	30311062	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine diluted two-star dimensions and small-e orbital solution with joint light-curve/RV modelling.
R042	290592570	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine diluted two-star dimensions and small-e orbital solution with joint light-curve/RV modelling.
R044	6946549	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine diluted two-star dimensions and small-e orbital solution with joint light-curve/RV modelling.
R045	149021625	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine diluted two-star dimensions and small-e orbital solution with joint light-curve/RV modelling.
R046	362082706	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine diluted two-star dimensions and small-e orbital solution with joint light-curve/RV modelling.
R047	436727157	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Constrain SB2 flux ratio/component radii and test the missing conjunction with resolved photometry.
R049	67651413	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine diluted two-star dimensions and small-e orbital solution with joint light-curve/RV modelling.
R050	244352837	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Refine diluted two-star dimensions and small-e orbital solution with joint light-curve/RV modelling.
R051	93213423	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Retry unavailable catalogs; refine joint orbit and stellar characterization.
R052	369984032	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Retry unavailable catalogs; refine joint orbit and stellar characterization.
R053	345589621	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Retry unavailable catalogs; refine joint orbit and stellar characterization.
R054	135251275	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Retry unavailable catalogs; refine joint orbit and stellar characterization.
R055	435190459	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Retry unavailable catalogs; refine joint orbit and stellar characterization.
R056	150696021	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Retry unavailable catalogs; refine joint orbit and stellar characterization.
R057	345050437	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Retry unavailable catalogs; refine joint orbit and stellar characterization.
R059	415751695	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Retry unavailable catalogs; refine joint orbit and stellar characterization.
R060	335862015	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Retry unavailable catalogs; refine joint orbit and stellar characterization.
R061	211231319	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint diluted light-curve/RV modelling to refine radii, flux ratio and small-e geometry.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R062	120522352	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint diluted light-curve/RV modelling to refine radii, flux ratio and small-e geometry.
R063	192198579	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint diluted light-curve/RV modelling to refine radii, flux ratio and small-e geometry.
R066	157000246	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint diluted light-curve/RV modelling to refine radii, flux ratio and small-e geometry.
R068	253923749	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint diluted light-curve/RV modelling to refine radii, flux ratio and small-e geometry.
R069	389527961	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint diluted light-curve/RV modelling to refine radii, flux ratio and small-e geometry.
R070	196426780	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint diluted light-curve/RV modelling to refine radii, flux ratio and small-e geometry.
R071	382423586	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse characterization; retain Gaia period caveats.
R072	194733085	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse characterization; retain Gaia period caveats.
R073	195483879	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse characterization; retain Gaia period caveats.
R074	161817918	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse characterization; retain Gaia period caveats.
R075	202277216	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse characterization; retain Gaia period caveats.
R076	268636467	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse characterization; retain Gaia period caveats.
R077	397594001	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse characterization; retain Gaia period caveats.
R078	9885893	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse characterization; retain Gaia period caveats.
R079	414008363	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse characterization; retain Gaia period caveats.
R080	61106869	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse characterization; retain Gaia period caveats.
R081	39478038	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R082	381591400	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R083	333072525	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R084	89329656	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R085	359480060	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R086	242057470	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R087	298432323	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R088	161478713	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R089	309981203	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R090	301961777	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R091	443053687	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R092	311147906	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R093	81481443	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R094	143517607	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R095	165258931	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R096	431535710	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R097	242199703	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R099	5343763	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R100	256477680	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R101	353848852	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R103	364120441	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R104	100249241	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R105	138180245	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R106	39013443	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R108	144169003	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R109	27987072	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R110	326623940	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R111	320195702	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R112	164946466	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R113	424249116	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R114	229238588	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R115	293946483	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R116	140802498	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R117	4374712	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R118	188085904	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R119	45006630	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R120	242656909	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R121	90563464	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R122	115549556	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R123	29741740	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R124	156851362	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R125	273177449	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R126	48829357	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R127	162075402	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R128	177285916	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R129	144500102	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R130	417917895	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R131	26804032	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R132	220425369	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R133	311918550	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R134	233269866	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R135	42318308	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R137	140246270	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R138	302274338	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R139	122024194	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.
R140	287032105	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; See target report for limitations.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R141	289404005	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R142	335001355	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R143	21335382	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R144	119830214	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R145	170081245	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R146	305537899	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R147	757171	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R148	427330878	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R149	416848010	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R150	309600815	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R151	431698773	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R152	424878861	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R154	350335402	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R155	48777160	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R156	382260140	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R157	162234490	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R158	358766055	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R159	37307440	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R160	89575652	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R161	266486943	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse modeling and longer-baseline eclipse timing; resolve remaining alias and geometry uncertainty.
R162	129280729	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse modeling and longer-baseline eclipse timing; resolve remaining alias and geometry uncertainty.
R164	167759819	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse modeling and longer-baseline eclipse timing; resolve remaining alias and geometry uncertainty.
R166	100014634	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse modeling and longer-baseline eclipse timing; resolve remaining alias and geometry uncertainty.
R167	259853508	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse modeling and longer-baseline eclipse timing; resolve remaining alias and geometry uncertainty.
R168	302342078	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse modeling and longer-baseline eclipse timing; resolve remaining alias and geometry uncertainty.
R169	28634340	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse modeling and longer-baseline eclipse timing; resolve remaining alias and geometry uncertainty.
R170	178080174	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Joint RV/eclipse modeling and longer-baseline eclipse timing; resolve remaining alias and geometry uncertainty.
R171	11661051	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R172	157948102	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R173	61107057	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
R174	313274300	G 10-11.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fit joint RV/eclipse geometry and a longer timing baseline; retain same-sector period limitations.
C02	468181994	fainter SB1 period-fold population	CONFIRMED	true	Report; Period -2.31 sigma from Gaia; RUWE 12.6/unresolved companion risk; independent pipelines agree about 2 sigma. Literature cones limited to enumerated catalogues, not proof of novelty.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
C03	470235665	fainter SB1 period-fold population	CONFIRMED	true	Report; QLP-only secondary phase drift unconfirmed; 4-5% pipeline scale offset. Enumerated catalogue cones and SIMBAD SB* do not establish exhaustive novelty.
C04	379251709	fainter SB1 period-fold population	CONFIRMED	true	Report; Secondary fold-level, not localized; primary securely on target. QLP/SPOC secondary phases differ 2.6 sigma. Radius tension retained.
C06	331293467	fainter SB1 period-fold population	CONFIRMED	true	Report; Heavy blend; QLP plus independent own TESScut satisfies available pipeline check; depths model-dependent 80-116 ppt, ratios/duration agree. Gaia geometry depends on covariance. Gaia-VEB/varisum/SB9 cones lack successful positive controls; no exhaustive novelty claim.
C07	283532878	fainter SB1 period-fold population	CONFIRMED	true	Report; Use successful s12d recheck, not false-empty first archive pass. Known EB 29.7 arcsec away at P 1.7975 d excluded as source; up to 3 ppt contamination possible. Phase drift tentative.
C08	407877153	fainter SB1 period-fold population	CONFIRMED	true	Report; Secondary pixel signal weak individually but summed delta chi2 >=166; QLP depth diluted. Use successful s12d catalogue rerun; all-catalogue query unreliable.
C09	15855794	fainter SB1 period-fold population	CONFIRMED	true	Report; Exclude flagged E124 from primary pixel stack as report does. Catalogue positive controls succeeded; catalogue absence only within searched scope.
C10	92881948	fainter SB1 period-fold population	CONFIRMED	true	Report; VSX ROT/Gaia SOLAR_LIKE is prior variable classification, not prior EB; catalogue positive controls succeeded. QLP dilution explains shallower depth.
C11	468827621	fainter SB1 period-fold population	CONFIRMED	true	Report; Primary localized in two sectors both camera 2; secondary untested in pixels and neighbour could supply in worst case. Literature search skipped, though catalogue cones performed; known Gaia short-timescale variable is not prior EB.
C12	416285595	fainter SB1 period-fold population	CONFIRMED	true	Report; Returned CONFIRMED with caveats: QLP/SPOC s40 61.7 vs 54.7 ppt (~12%, ~4 sigma); TESScut depth unreliable. Secondary not pixel-tested; geometry weak/covariance-dependent. Catalogue cones performed but literature search skipped. Report itself says strict depth agreement would imply PROBABLE; no new stricter tolerance imposed here.
C19	23112764	fainter SB1 period-fold population	CONFIRMED	true	Report; See target report for deciding limitation and next check.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
C20	373062090	fainter SB1 period-fold population	CONFIRMED	true	Report; See target report for deciding limitation and next check.
C21	465073693	fainter SB1 period-fold population	CONFIRMED	true	Report; See target report for deciding limitation and next check.
C22	316405449	fainter SB1 period-fold population	CONFIRMED	true	Report; See target report for deciding limitation and next check.
C24	441741446	fainter SB1 period-fold population	CONFIRMED	true	Report; See target report for deciding limitation and next check.
C25	280075497	fainter SB1 period-fold population	CONFIRMED	true	Report; See target report for deciding limitation and next check.
C26	276514049	fainter SB1 period-fold population	CONFIRMED	true	Report; See target report for deciding limitation and next check.
C30	104736355	fainter SB1 period-fold population	CONFIRMED	true	Report; Optional spectroscopy or higher spatial resolution photometry would refine characterization; no additional deciding gate is required by this representative review.
C32	450055243	fainter SB1 period-fold population	CONFIRMED	true	Report; Optional spectroscopy or higher spatial resolution photometry would refine characterization; no additional deciding gate is required by this representative review.
C35	267222866	fainter SB1 period-fold population	CONFIRMED	true	Report; Optional spectroscopy or higher spatial resolution photometry would refine characterization; no additional deciding gate is required by this representative review.
C36	66315917	fainter SB1 period-fold population	CONFIRMED	true	Report; Optional spectroscopy or higher spatial resolution photometry would refine characterization; no additional deciding gate is required by this representative review.
C37	93783392	fainter SB1 period-fold population	CONFIRMED	true	Report; Optional spectroscopy or higher spatial resolution photometry would refine characterization; no additional deciding gate is required by this representative review.
C38	470947699	fainter SB1 period-fold population	CONFIRMED	true	Report; Optional improved spectroscopy and photometry would refine binary characterization.
C39	135230723	fainter SB1 period-fold population	CONFIRMED	true	Report; Optional improved spectroscopy and photometry would refine binary characterization.
C40	36083554	fainter SB1 period-fold population	CONFIRMED	true	Report; Optional improved spectroscopy and photometry would refine binary characterization.
C41	447566809	fainter SB1 period-fold population	CONFIRMED	true	Report; Optional improved spectroscopy and photometry would refine binary characterization.
C45	360516818	fainter SB1 period-fold population	CONFIRMED	true	Report; Optional improved spectroscopy and photometry would refine binary characterization.
C56	4646090	fainter SB1 period-fold population	CONFIRMED	true	Report; Characterize dilution and grazing geometry if needed.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
P001	262653359	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Period uncertainty widened to 0.00008 d: saved floor-fit output gives 0.000072 d, wider than prose 0.00005. Mass and evolutionary state leave factor ~2 light-ratio uncertainty. Next test: Spectroscopy or colour eclipse photometry to constrain companion light.
P002	39013890	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Unresolved third light remains possible (ipd multi-peak 69%; RUWE 1.55), affecting derived radii and geometry precision. Next test: Resolve close companion or constrain third light spectroscopically.
P003	325399242	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Historical prose claims seven-event P=28.038374 +/-0.000191, but saved ephem_A and Afix each have five rows. Adopt corrected five-event artifact. Capable 13.2 arcsec neighbour is disfavoured at roughly 3–4 sigma per sector with systematic floor, not mathematically impossible. Next test: Additional orientation/source-resolved imaging would strengthen close-neighbour rejection.
P004	384026617	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Historical 0.6 arcsec WDS pair is doubtful but unresolved by TESS; attribution is to the Gaia source footprint. Gaia statistics alone do not prove pair spurious. Next test: High angular resolution imaging to check historical WDS component.
P005	389013422	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Secondary phase-0.535 hint is unconfirmed; do not quote it as a measured eclipse. Absolute physical parameters remain model dependent. Next test: RVs to constrain eccentricity and secondary predictions.
P006	4808224	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Single-sector s89 pixel test; pulsation affects secondary depth/timing. Discard s35 secondary fit: all input points flagged and optimiser wandered. Approximate physics.py formula has pi/4 error; exact Kepler-grid result is used. Next test: Another orientation pixel test or improved RV orbit would reduce association uncertainty.
P007	23999420	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Adopt primary timing uncertainty 0.000044 d rather than joint model formal 0.000015 d. Secondary 9.7-sigma pixel regression is historical numerical evidence; representative image has no colour scale. Next test: Recover missing/flagged s91 primary only if additional refinement needed.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
P008	112232265	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Intrinsic depth approximately 30–36 ppt; custom 3x3 is diluted (~19 ppt), QLP background affects depth. RUWE 3.51 allows third-body uncertainty. T16 parse with zero cadences is not evidence. Next test: Spectroscopy and source-resolved eclipse depths to constrain companion.
P009	275065610	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; TASOC depth is diluted and its epoch is about 14 min later than QLP; does not invalidate its eclipse detection. Dark-companion fit neglects a few percent secondary light. Next test: A future raw A eclipse or double-lined spectrum
P010	469755925	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; S94 A is quality-flagged; confirmation also rests on clean s13 and s67 A events. B pixel depth5.93 +/-0.81 ppt exceeds pipeline2.8–4.0 ppt; low-S/N/background-template extraction is attribution only, not adopted depth. Next test: Another clean secondary for depth and geometry
P011	356435413	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; CTOI356435413.01 is a prior planet-candidate signal at matching eclipse period; annotate separately from EB novelty. Original report overstates 1.5arcsec target offset: that is deficit-to-light centroid; s79 coordinates imply ~2arcsec to target, s53 ~3.6arcsec. No full limb-darkened grazing fit. Next test: Grazing-occultation model or spectrum to test light ratio
P012	255077215	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; S65 secondary arrives31–42min late after common-mode correction; unexplained. This does not remove repeated A timing or agreement of earlier B events. Quoted period error is original formal fit and predictions need timing-systematic allowance. Next test: Clean secondary to resolve s65 timing shift
P013	262471565	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; B is undetected: expected570–1100ppm versus ~600ppm central-phase limit creates mild model tension; permissible phase/temperature/impact values remain. Period error includes20min pipeline-scatter floor. Next test: Unflagged B-window photometry at <=300ppm
P014	126758342	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Gaia eccentricity and omega broad; A misses in83% of permitted historical model draws, not certainty. Primary_epoch field uses detected deeper B eclipse. Next test: Refined RV orbit or next B eclipse

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
P015	178486571	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; S87 egress cut by gap. Three-event near-zero timing chi-square makes formal error optimistic; allow >=5min in predictions. Next test: Clean B coverage or another A event
P016	395790195	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Adopt timing-only B period/error; tighter joint-model55.460455 +/-0.000021d is conditional on model. Only two same-kind epochs, cycle count constrained by Gaia. Next test: Another B eclipse to test cycle count
P017	466336052	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Only two A eclipses; no internal timing residual test. B windows are contaminated or missing; no B detection. Gaia conjunction uncertainty is 3-4 d. Next test: Observe a third A eclipse and a clean B window near 0.3% precision.
P018	64066759	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Long-baseline precision partly depends on a fixed-shape partial s7 event; three events total. Simplified geometry favours smaller primary radius than TIC. No B expected. Next test: Fit a full limb-darkened joint A model and constrain primary radius spectroscopically.
P019	49416839	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; A3 lacks egress and is weak. B-A phase separation is 2.3 sigma from Gaia; simple joint geometry has limited constraints. Dominant deep eclipse is labelled B in source. Next test: Obtain a complete A event and joint free-eccentricity model to test the phase-separation tension.
P021	60583013	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Period relies on two secondaries, one after a data gap; one primary only. Older B pixel test inconclusive. Duration geometry is approximate. Next test: Obtain a second primary and joint limb-darkened model to test phase separation and geometry.
P022	277542587	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; TESS A mostly flagged; only one TESS A+B pair. KELT period precision is heuristic and anchored to TESS; broad ground aperture dilutes depth. Sparse KELT B is supporting only. Historical exact-night count not adopted. Next test: Obtain another full well-calibrated A/B pair and an event-timing based period uncertainty.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
P023	12864953	G 11-11.5; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Only two B events and one clean A; A ephemeris assumes B period. New A window is entirely quality 2048 flagged. Phase-separation tension about 3.35 sigma remains. Conservative period error is an estimate, not calibrated coverage. Next test: Obtain a clean repeated A and refined radial-velocity orbit to resolve eccentricity/phase tension.
R175	300217120	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; The nearby other eclipse in S67 contaminates a wider baseline window; the final fit uses a narrower 1.65-search-duration half-window and shows the repeated primary. Historical S27 TESScut contains one event but QLP independently shows three primaries in S67/S94. Next test: Optional precision characterization
R176	424716057	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; QLP primary depths vary across apertures/sectors (205–230 ppt in the selected sectors); no common intrinsic depth is claimed. Next test: Optional precision characterization
R178	274941915	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Three capable neighbours at 32.98, 40.57 and 47.81 arcsec are spatially separated from the target-centered compact deficit. The small-aperture reduction retains the eclipse. No calibrated centroid confidence ellipse is available. Next test: Optional precision characterization
R180	245474316	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep repeated eclipses are isolated in both reductions; all four measured primary events support the timing solution. Next test: Optional precision characterization
R181	397586559	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating eclipses (primary median approximately254 ppt, secondary164 ppt) establish orbital period twice the historical4.93316d event spacing. Only deeper even cycles enter the reported primary ephemeris. Historical S70 difference image mixes both eclipse kinds, so its aperture depth is a mixture. Gaia nominal conjunction is0.28093d later at the TESS epoch; measured-Gaia period difference−0.00126495d accumulates this offset. Projecting toward the Gaia2016 reference reduces offset to−0.0166d; the orbit association is physically consistent with an imperfect Gaia period, and the tension is retained rather than subjected to a sigma veto. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R182	305801065	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; The shallow eclipse is clearly repeated with localized ingress/egress despite higher scatter in newer sectors. Three capable neighbours at28.07,33.30,54.57arcsec are separated from the compact target deficit. Depth depends on dilution and trapezoid shape; historical median3.8ppt is not a precision intrinsic depth. Next test: Optional precision characterization
R183	345794632	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Large historical PSF fractional residual0.57 reflects imperfect scaled out-image morphology in a crowded field. No listed neighbour within60arcsec can supply the approximately73ppt eclipse under the G-band flux-budget screen, and the lost-light core coincides with the target. The weaker localization calibration remains a limitation. Next test: Optional precision characterization
R184	258621299	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating eclipses (primary median approximately180ppt, secondary131ppt) establish orbital period twice the historical3.49299d event spacing. Only deeper even cycles enter the reported primary ephemeris. Historical S20 difference image mixes both eclipse kinds, so its aperture depth is a mixture. Next test: Optional precision characterization
R185	65346146	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated flat-bottom eclipses; sector-dependent dilution changes fitted depths from about46 to38ppt. Next test: Optional precision characterization
R186	416646907	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Odd historical half-period cycles are deeper in each of three sectors (about18 versus16ppt); deeper odd cycles alone give the primary ephemeris and doubled orbital period. Alternative-reduction stack mixes eclipse kinds. Approximate central duration permits a noncentral chord. Next test: Optional precision characterization
R187	401138717	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Three capable neighbours and high scaled-out-image residual0.976 limit PSF precision. Target-centered compact loss and aperture-depth growth support attribution. Known EB42.73arcsec away has P0.363740d, distinct from19.79d. Gaia period differs by0.00556d, accounting for much of the accumulated conjunction offset; this physical tension is retained. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R188	334379409	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated raw eclipses have modest baseline variability. Gaia period mismatch accumulates over the baseline; propagated covariance and conjunction comparison are descriptive rather than a sigma threshold. Next test: Optional precision characterization
R189	345406164	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deeper even cycles near75ppt alternate with47ppt odd eclipses in sector81, establishing doubled orbital period. Primary-only ephemeris used. Historical stack mixes eclipse kinds. Three capable neighbours are away from compact target deficit. Next test: Optional precision characterization
R190	152400959	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Known EB27.09arcsec away at0.257384d differs from5.513d signal. Two capable neighbours and residual0.959 limit PSF precision; compact target-centered deficit and small-aperture eclipse support attribution. Next test: Optional precision characterization
R191	152086769	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep flat-bottom repeated eclipses; sector96 aperture depth is lower than earlier sectors. No capable neighbour within60arcsec in approximate G flux budget. Next test: Optional precision characterization
R192	270260727	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Shallow localized repeated eclipses are resolved in sectors67 and95; raw scatter and trapezoid approximation limit precision. No capable neighbour within60arcsec under G flux budget. Next test: Optional precision characterization
R193	112827464	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Raw local baseline slopes are fitted explicitly. One capable neighbour is separated from the target-centered deficit; no calibrated centroid confidence ellipse. Gaia nominal phase offset0.063 and period drift remain explicit; full covariance and near-reference projection support association without arbitrary sigma veto. Next test: Optional precision characterization
R194	423657976	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated isolated eclipses with stable approximately94ppt QLP depth. No capable neighbour within60arcsec under approximate G flux budget. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R195	343926224	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Five clean primary repeats determine the ephemeris. S76 historical cycle 154 has structured excursions and an anomalously weak fitted dip; S84 cycle 176 has a broad distorted profile. Both remain in raw-windows.csv and plots but are excluded from the timing fit. The capable neighbour at 55.00 arcsec is separated from the compact target deficit. Aperture and local baseline variations limit depth precision. The retained timing fit has reduced chi-square 18.08; residual timing scatter materially inflates the stated period error and shows model/baseline limitations. Next test: Optional precision characterization
R196	53993311	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Aperture-dependent primary depth changes from about 107 ppt in S68 to 85 ppt in S95; the same localized eclipse morphology repeats. Gaia conjunction uncertainty is substantial; a phase offset alone is not a veto. Next test: Optional precision characterization
R197	43785616	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Observed trapezoid duration exceeds a central estimate based on default primary mass and minimum companion mass. That approximate estimate is not an upper bound: stellar properties and inclination are not fitted. Repeated flat-bottomed events and matching period support association; detailed binary modeling remains optional. Next test: Optional precision characterization
R198	397178704	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating primary and secondary eclipses establish twice the historical 5.50484 d event spacing. Only deeper odd cycles enter the primary ephemeris. The historical stack mixes both kinds. Large PSF residual 0.676 reflects surrounding light rather than an off-target core; the one capable neighbour lies 45.82 arcsec away from the compact deficit. Next test: Optional precision characterization
R199	77550184	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated isolated eclipses have aperture-dependent depths of approximately 56 to 63 ppt. No capable neighbour lies within the 60 arcsec screen. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R200	314041837	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; The target is a large star (TIC radius 5.09 solar radii) with ellipsoidal modulation plus localized eclipses. A two-harmonic baseline fitted outside the primary core in S77 leaves a 30.31 ppt localized deficit; its double-wave modulation does not explain the eclipse. See harmonic-test.json and png. The 18.79 h local-linear trapezoid width includes smooth variability and is not a physical eclipse duration. The reported depth likewise describes that local model, not intrinsic eclipse depth. Capable neighbours at 27.40 and 53.17 arcsec are away from the target-centered deficit; PSF residual 0.595 limits morphology precision. Next test: Optional precision characterization
R201	7446763	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Primary eclipses are approximately twice secondary depth. Only deeper odd historical half-period cycles determine the ephemeris; the reported orbital period doubles the historical 2.95595 d spacing. Historical S98 aperture stack mixes both eclipse kinds. Next test: Optional precision characterization
R202	22705959	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; A capable neighbour at 39.75 arcsec is separated from the target deficit. The target sits near a pixel boundary: the global deficit peak rounds to an adjacent pixel, while the centroid is only 3.64 arcsec from the actual target. This is not evidence for a different source. Next test: Optional precision characterization
R203	62940616	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; One capable neighbour at 22.67 arcsec requires a 0.329 fractional eclipse to explain the signal, but the lost-light centroid lies 0.205 pixel from the target versus 1.33 pixels from that neighbour, displaced away from it. Its pixel fractional loss 0.0416 is a blended-pixel value and does not measure the neighbour intrinsic eclipse depth. The target-centered depth increases in the 1x1 aperture, consistent with target dilution in larger apertures. TESS resolution and G-to-TESS colour differences limit attribution precision. Next test: Optional precision characterization
R204	100911870	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Four clean primary events in S56 and S83 support the timing solution and approximately 84 ppt raw QLP depth. No capable neighbour is present in the 60 arcsec screen. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R205	77176486	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; The 13.75369 d signal lies near the TESS orbital timescale, but repeated localized raw SAP dips in S31 and S98 and the target-centered cutout deficit support astrophysical eclipses. Quality-nonzero cadences are excluded; no flagged or uncovered window is a clean null. Cycle188 is visibly present but its trapezoid covariance is singular, so it is excluded from timing. Bounded S98 pixel/background check (background-check.json) finds target-pixel losses11.99/22.43/16.35 electrons/s versus3.24/9.85/7.73 at the close capable neighbour and0.70/6.30/3.36 at the farther one. Outer-pixel median losses are -0.027/5.68/4.04: common-mode background affects later events, while cycle187 is localized with negligible outer deficit. Core quality flags are0/0/3; flagged cadences are excluded. FLUX_BKG is zero throughout and supplies no independent background measurement. These are descriptive diagnostics, not a calibrated artifact veto. Gaia conjunction offset is 0.04208 phase with 0.549 d propagated uncertainty; this is association support with appreciable orbital uncertainty, not a precise phase coincidence. Next test: Optional precision characterization
R206	155916287	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Seven fitted repeats span S16,50,77. Sector-dependent depth variation and long-cadence integration limit intrinsic depth and duration precision. Next test: Optional precision characterization
R207	383603595	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Eleven fitted repeated eclipses span S8,62,90. Modest aperture/baseline depth variation limits precision; no missing bounded check. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R208	436724968	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; QLP free-trapezoid depths show modest parity differences, especially in S71; local baseline and aperture dependence are material. Independent S43 3x3 cutout depths are 35.34-37.19 ppt for eight events including consecutive odd/even cycles, without a stable unequal alternating pattern. Gaia P=2.732167 d matches the measured event spacing. These data do not establish two distinct alternating eclipse kinds or require doubling; report the fitted period as the adopted orbit with this depth-model limitation, and the epoch as an observed reference eclipse. Capable neighbours at 35.19 and 35.89 arcsec are displaced from the compact target deficit. Next test: Optional precision characterization
R209	37948318	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; The historical absolute target-to-deficit offset is 15.1 arcsec (0.719 pixel), while light-to-deficit separation is 3.51 arcsec; these are distinct comparisons. The capable G13.73 neighbour is 53.65 arcsec away and opposite the deficit displacement. Independent per-event pixel checks find target-pixel losses 9.03-10.59 electrons/s against 0.185-0.437 at that neighbour in cycles317,319,321. This supports the target over the capable neighbour at TESS resolution, with an uncalibrated absolute-centroid systematic and residual crowding uncertainty. Historical cycles316/318 have common-mode excursions and are not individually decisive. QLP cycle93 has singular timing covariance and is excluded from timing. Next test: Optional precision characterization
R210	373034189	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Four raw fitted eclipses in S10,38,90 support the period. The Gaia conjunction uncertainty is substantial (0.3105 d); local model widths and depths are descriptive. Next test: Optional precision characterization
R211	300969635	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep repeated eclipses persist in S1,37,98. Aperture depth changes from about210 to248 ppt; pooled parity depths confound sector coverage and do not establish stable unequal eclipse kinds. Gaia period matches the measured spacing. Intrinsic depth requires dilution modeling. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R212	317800184	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated eclipses span S6,87,98 with sector-dependent depths around45-62 ppt. This limits precision but does not erase the localized repeat pattern. Next test: Optional precision characterization
R213	219033622	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Three constrained same-kind events in S15 and S23 determine the ephemeris; S23 cycle32 is retained as a raw repeat but excluded from timing because sparse sampling leaves its egress poorly constrained. S76 did not supply a usable selected event window and is not a clean non-detection. The independent S16 cutout contains four repeating deep target-centered deficits. The timing baseline is shorter than the available sector list suggests. Next test: Optional precision characterization
R214	312725755	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Five fitted repeating eclipses span S19,45,73. The measured widths are long but compatible with approximate Gaia/TIC geometry; intrinsic depths and stellar parameters need detailed modeling. Next test: Optional precision characterization
P025	382042754	G 11.5-12; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Pilot characterization; period errors are scatter-only. No exhaustive literature priority claim. Next test: Optional precision characterization; no missing bounded admission check
P026	389128842	G 11.5-12; Gaia spectroscopic orbit 20-60 d	CONFIRMED	true	Report; Pilot characterization; period errors are scatter-only. No exhaustive literature priority claim. Next test: Optional precision characterization; no missing bounded admission check
R218	167105778	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Capable G14.83 neighbor10.1arcsec away requires44% eclipse. Deficit is0.18pixel from target and0.47pixel from neighbor; image favors target, with uncalibrated centroid resolution retained as limitation. Next test: Optional precision characterization
R219	48212656	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating eclipses approximately40/73ppt establish doubled orbital period. Odd historical half-period cycles are deeper primaries and alone define ephemeris. Historical cutout mixes eclipse kinds. Next test: Optional precision characterization
R220	442662494	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Historical PSF fit residual0.968 is poor, though target-centered deficit and no capable catalogued neighbor support attribution. Aperture/sector-dependent depths approximately277-331ppt; no intrinsic undiluted depth is inferred. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R221	280881615	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Triangular ingress and egress repeat cleanly. No capable neighbor under the approximate G-band flux screen. Next test: Optional precision characterization
R222	238857132	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating eclipses approximately75/147ppt establish doubled orbital period. Even half-period cycles are deeper primaries and alone define ephemeris. Historical cutout mixes eclipse kinds. Next test: Optional precision characterization
R223	60249512	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Similar eclipse depths do not establish half orbital period. Consecutive sector88 spacings7.682814/7.617839/7.682757d alternate; same-kind spacings15.300653/15.300596d support full orbital period. Even half-period cycles define selected-conjunction epoch; deeper-primary identity is not claimed. Historical cutout mixes eclipse kinds. Next test: Optional precision characterization
R224	250562568	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated isolated eclipse; alternative aperture supports target. Depth varies with aperture/dilution. Next test: Optional precision characterization
R225	325787864	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Crowded field:29 capable neighbors include nearest12.9/14.3arcsec. Deficit0.17pixel from target and target-centered lost-light core favor target; no calibrated centroid uncertainty. Shallow morphology repeats in newer sectors; first-sector windows did not satisfy fitting coverage cuts and are not nulls. Next test: Optional precision characterization
R226	180871891	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Four measured shallow eclipses across sectors29/104 establish repeatability and period. Target-centered deficit0.043pixel and no capable catalogued neighbor support attribution. Sector2 was downloaded but no event window passed coverage cuts; it is not a clean non-detection. No single-event availability gap remains. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R227	129565419	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Remote bright region dominates historical unrestricted peak and PSF residual0.968. Independent helper local5x5 re-stack residual0.115 and target centroid offset(-0.02,+0.01)pixel recover the target eclipse; three aperture depths24.40/23.92/22.14ppt support it. Capable neighbor38.91arcsec away lies1.90pixel from original deficit vs0.040pixel target. Next test: Optional precision characterization
R228	125690413	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Structured sector41 baseline gives broad34ppt cycle196, residual -0.038d and reduced chi square5.25. Retained event and inflated covariance document this; clean22ppt repeats occur in sectors15/55. Capable neighbors are24.35arcsec or farther,1.55pixel or farther from deficit vs0.371pixel target. Timing errors are not total uncertainty. Next test: Optional precision characterization
R229	445045766	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; No capable catalogued neighbor within60arcsec. Deficit0.186pixel from target; three alternative aperture depths23.3-23.9ppt agree. Next test: Optional precision characterization
R230	120037640	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Actual-epoch Gaia offset -0.189d (phase -0.051) exceeds historical phase screen; near Gaia reference projected offset is +0.045d. Gaia period uncertainty accumulates over baseline. Low eccentricity0.062 and recurrence agreement support association without arbitrary sigma veto. Capable neighbors are34.9arcsec or farther and1.72pixel or farther from deficit vs0.178pixel target. Next test: Optional precision characterization
R232	158647370	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; No capable catalogued neighbor. Approximately85-95ppt eclipses repeat; custom aperture depths85-87ppt agree. Next test: Optional precision characterization
R233	302190527	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; No capable catalogued neighbor. Approximately275-283ppt raw eclipses repeat across widely separated sectors; alternative aperture depths255-263ppt are diluted extraction values. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R234	138941853	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Approximately8-10ppt eclipses repeat in sectors6/32/98. Deficit0.087pixel from target vs nearest capable source1.127pixel. Near-reference Gaia offset0.352d is descriptive; low-eccentricity orbit conjunction is poorly constrained. Actual epoch offset0.036d with0.524d propagated Gaia uncertainty is compatible. Next test: Optional precision characterization
R235	256665207	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Approximately6-7ppt flat-bottom eclipses repeat. No compelling even/odd split establishes half/double alias; full Gaia period agrees with recurrence. Nearest capable source32.78arcsec away,1.70pixel from deficit vs0.166pixel target. Next test: Optional precision characterization
R236	91628894	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deficit0.377pixel from target; light-centroid shift1.05arcsec is a different quantity. Nearest capable source19.20arcsec away requires38.1% eclipse and lies0.675pixel from deficit. Central lost-light peak and consistent10ppt aperture depths favor target; no calibrated subpixel uncertainty is inferred. Next test: Optional precision characterization
R237	252737077	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Strong aperture dilution: raw QLP approximately209ppt versus cutout1/3/5 aperture160/96/43ppt. Deficit core is target-centered; capable neighbors42.7/53.8arcsec away are2.06/2.72pixels from the deficit centroid versus0.20pixel for target; blended local loss is not an intrinsic-source eclipse constraint. Poor total-light PSF residual0.943 reflects crowding; no intrinsic depth is inferred. Next test: Optional precision characterization
R238	240771237	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Seven measured newer-sector repeats; historical sector57 did not pass present fitting window requirements and is not a null. Capable neighbor31.4arcsec away is separated from the target-centered deficit. Next test: Optional precision characterization
R239	369118947	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Target-centered deficit despite poor0.923 total-light PSF residual in crowded field. Capable neighbor53.8arcsec away is well separated. Aperture-dependent diluted depth retained. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R240	399238907	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Only QLP sector80 is available in the shared availability receipt, containing five measured repeats. Queue4.14670d is the historical screening seed, not current Gaia period; full NSS4.144220414d agrees measured period. Sector80 pipeline timing offsets are an uncalibrated systematic. Four capable neighbors are offset from the target-centered lost-light core; closest16.6arcsec. Next test: Optional precision characterization
R241	146319847	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Capable neighbor33.8arcsec away is distinct from the target-centered deficit. Raw depths105-138ppt and cutout apertures108/54/20ppt reflect substantial dilution; total-light PSF residual0.927 is poor and not used as a sole attribution test. Next test: Optional precision characterization
R242	466690912	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated flat-bottom eclipses with changing raw sector depths19-35ppt; aperture and baseline dependence is retained. Capable neighbor49.1arcsec away is distinct from the deficit core. Observed duration5.25h agrees approximate central estimate5.24h to the precision of its assumptions. Next test: Optional precision characterization
R243	436092743	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating eclipses approximately153/100ppt establish full orbital period13.12907d, twice historical SAP branch. Odd half-period cycles are deeper primary eclipses and alone define ephemeris. Historical cutout stack mixes eclipse kinds; no capable neighbor. Next test: Optional precision characterization
R244	38850860	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Eight measured repeats across three sectors. No capable catalogued neighbor; target-centered deficit. Raw depth26-31ppt varies by extraction/sector. Next test: Optional precision characterization
R245	140281693	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Target-centered deficit despite poor0.9999 total-light PSF residual caused by unrelated bright light outside target core. Capable neighbors37.5/56.8arcsec are spatially separate. Raw depth34-45ppt depends on sector aperture. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R248	362822264	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Two QLP measured events separated by four cycles give a conditional ephemeris. Independent raw cutout S39 centers2366.85039+/-0.00592 and2384.98788+/-0.00879 BTJD, separation18.13749+/-0.01060d, support same-kind adjacent-cycle recurrence near18.1224d; see helper/same-sector-refits.json. Alternative sector39 cutout shows target-centered deficit; nearest capable neighbor25.9arcsec. Alias search is bounded rather than exhaustive. Next test: Optional precision characterization
R250	373621318	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated isolated narrow eclipses and target-centered deficit favor target; bright field source affects whole-image PSF residual and dilution. Next test: Optional precision characterization
R251	129765199	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Deep repeating eclipses, target-centered deficit, no capable catalogued neighbor. Sector-dependent depth is dilution or extraction dependent. Next test: Optional precision characterization
R252	309558046	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating eclipses approximately27/17ppt establish orbital period twice historical SAP half period. Even half-period cycles are deeper primaries and alone define ephemeris. Historical cutout stacks mixed eclipse kinds. Next test: Optional precision characterization
R255	30753325	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Clean triangular repeating eclipses, target-centered deficit and no capable catalogued neighbor. Depth changes between sectors; no intrinsic undiluted depth inferred. Next test: Optional precision characterization
R259	467753507	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Four measured eclipses in two representative sectors; no capable neighbor. Alternative cutout approximately29ppt agrees raw26-31ppt. Epoch identifies the repeating selected eclipse, deeper-secondary identity not established. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R260	350760292	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Four raw repeats in available representative sector92; historical sector39 cutout independently reuses earlier observing occasions. Measured4.69607d differs screening4.69710d; short baseline and timing covariance retained. Capable nearest neighbor26.14arcsec away is1.45pixels from deficit centroid versus0.24pixel for target; other capable neighbors are50-55arcsec away. Poor total-light PSF residual0.936 reflects crowded light and is not sole attribution evidence. Next test: Optional precision characterization
R261	371241950	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Thirteen measured repeats span three sectors; capable neighbor45.52arcsec is spatially separated from deficit core. Gaia phase offset0.030 is discussed with covariance rather than arbitrary cutoff; observed4.3h duration can be noncentral. Next test: Optional precision characterization
R262	407207285	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Historical0.18d duration placed out-of-eclipse windows inside the approximately0.90d eclipse and suppressed cutout depth to6ppt. Revised21.6h extraction yields69/68/67ppt versus raw median67ppt. Updated flux budget makes all neighbors incapable (closest18.27arcsec requires253percent intrinsic eclipse). Target-centered revised deficit passes attribution; original extraction preserved. Raw fit uses0.65d baseline exclusion and free duration; no unperformed check is hidden. Next test: Optional precision characterization
R263	177257211	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Seventeen raw repeated events across three sectors; no capable neighbor. Target-centered deficit and56-62ppt cutout/raw amplitudes. Historical253 events are screening statistics, not measured timing count. Next test: Optional precision characterization
R264	29114881	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Six raw events across three sectors. Capable neighbor53.54arcsec is distinct from target-centered deficit. Raw56-59ppt and cutout57/56/51ppt depend on aperture. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R265	77605102	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Five repeated raw eclipses across three sectors; no capable neighbor. Raw145-169ppt and cutout147/144/133ppt show extraction dependence. Geometry offset minus0.026phase retained with propagated Gaia covariance; not sigma veto. Next test: Optional precision characterization
R266	35635884	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Eight repeated raw eclipses across three sectors; capable neighbor47.20arcsec is distinct from deficit core. Raw114-129ppt vs cutout116/108/82ppt show aperture and sector dilution. Next test: Optional precision characterization
R267	396859601	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Two S32 well-sampled events determine the reported period. A third real S5 eclipse tests recurrence across 44 cycles; its zero/singular covariance is excluded from timing. No newer timeseries sectors in current MAST query beyond5/32. Period error remains short-baseline and conditional on tested cycle branch, not precision long-baseline characterization. Next test: Optional precision characterization; no missing bounded admission check.
R268	115106052	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Deeper odd half-period cycles selected:10 primary times, with independent even-cycle timing agreement. Alternating depth and center residuals support doubled orbital period; historical half-period stack mixes unequal eclipses. Next test: Optional precision characterization; no missing bounded admission check.
R269	443911710	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Close2.88arcsec neighbor G16.14 requires129percent intrinsic loss under approximate Gaia-G flux budget; excluded as sole source under that assumption. TESS/G color differences are not modeled. Nine measured repeats, consistent target-centered cutout. Next test: Optional precision characterization; no missing bounded admission check.
R270	144113939	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; One capable neighbor38.54arcsec away is separated from target-centered deficit. Four raw repeated events span sectors64/101; alternative historicalS37 provides earlier reduction evidence. Next test: Optional precision characterization; no missing bounded admission check.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R271	339578643	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Four raw events in sectors95/103; target-centeredS102 cutout supports separate reduction. No capable neighbor within60arcsec. Next test: Optional precision characterization; no missing bounded admission check.
R275	20176241	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Two well-sampledS48 events and third clear sparseS21 recurrence test period across55cycles. S21 missing egress/sparse core gets duration/2 conservative timing allowance, not calibrated confidence. Current MAST query finds only21/48, no newer sector. Next test: Optional precision characterization; no missing bounded admission check.
R277	336856559	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Strong aperture dilution: cutout1/3/5 depths99.7/54.2/27.3ppt; QLP sectors66/103 differ. Deficit0.211pixel from target, bright capable neighbor2.26pixels from deficit favors target. Depths are pipeline/aperture dependent, not intrinsic. Next test: Optional precision characterization
R278	146878534	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Clean repeated sector33 eclipses150-174ppt; sector87 depths113-277ppt have changing steep baselines/noise. Real morphology and timing recur; apparent depth variability is not established astrophysically. Local-model timing systematics remain. Next test: Optional precision characterization
R279	123220126	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating eclipse depths establish doubled orbital period. Even historical half-period cycles alone define deeper-eclipse ephemeris; historical cutout stack mixes kinds. Next test: Optional precision characterization
R280	42960522	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deficit0.307pixel from target; nearest capable neighbor1.55pixels from deficit. Spatial core supports target at TESS resolution; no calibrated centroid errors. Next test: Optional precision characterization
R282	355885228	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating eclipses support doubled orbital period; three deeper even-half-period events include a sector24 same-kind repeat and sector78 recurrence. Historical cutout mixes kinds. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R283	169303527	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated isolated eclipses show sector-dependent dilution/depth. No capable catalogued neighbor under approximate Gaia brightness screen. Next test: Optional precision characterization
R284	94541119	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Five measured eclipses in sectors18/44/86 test period recurrence. No capable neighbor under approximate Gaia brightness screen; adopted depth is diluted. Next test: Optional precision characterization
R285	409935170	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating depths within sectors establish doubled orbital period even though across-sector pooled depths overlap. Even half-period cycles alone define deeper-eclipse timing. Historical cutout mixes kinds. Next test: Optional precision characterization
R286	365864542	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep repeated isolated eclipses across three sectors; one singular center-covariance fit excluded. Aperture/sector depth differences retained; no intrinsic undiluted depth inferred. Next test: Optional precision characterization
R287	143375995	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep isolated eclipses recur across three sectors. Aperture depths vary with dilution; no capable catalogued neighbor within60arcsec under approximate Gaia G brightness test. PSF-scaled residual0.856 warns against interpreting this simple PSF fit quantitatively; deficit core is centered on target. Next test: Optional precision characterization
R288	122506579	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated isolated shallow eclipses in sector97. Gaia orbit is nearly circular (e0.00444), so periastron/omega parameterization is weak; measured eclipse falls0.60753d (phase0.08762) from nearer nominal edge-on conjunction versus full-covariance linear allowance0.25923d. Other conjunction is2.87617d away. Near-reference projected offset0.14159d with photometric period projection allowance0.11355d; photometric P differs from Gaia by0.000910d versus Gaia error0.000500d. Retain mild orbit/timing tension, no arbitrary sigma veto or forced alias. Approximate central duration5.63h vs observed2.26h permits grazing/noncentral eclipse. Updated orbital solution would sharpen association. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R289	201598744	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Raw shallow eclipses recur in sectors93/104. Strong aperture dilution:1x1/3x3/5x5 depths12.15/10.61/8.02ppt. Deficit0.344pixel from target and closest capable catalogued neighbor1.63pixels from deficit favor target at TESS resolution. PSF-scaled residual0.803 is not a calibrated localization uncertainty. Next test: Optional precision characterization
R290	267094867	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep eclipses recur in sectors1/67/96; sector-dependent depths and shapes retained. Singular sector67 cycle520 fit excluded from timing. No capable catalogued neighbor in approximate brightness test. Next test: Optional precision characterization
R291	156985493	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Narrow eclipses at30minute cadence in sector22 have only3-4 trough samples. Mixed quadratic/Gaussian timing model and cadence floors are descriptive allowances, not total uncertainty. Sector49 cycle75 and sector22 half-period midpoint are flagged/gaps, not clean nulls; other sector49 half-period windows have usable flat samples. Same photons, aperture dilution and uncalibrated centroid uncertainty remain. Approximate1.2h observed FWHM differs from modeled central duration; grazing geometry possible, no detailed binary fit. Next test: Optional short-cadence resolved photometry and revised spectroscopy
R292	189664666	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Four real raw eclipses in sector91; discontinuous egress/baseline ramp in cycle5 makes its trapezoid center unreliable and excluded. Three other same-kind timings fit the period. No capable catalogued neighbor in approximate brightness test. Timing baseline is only14d, not precision long-baseline characterization. Next test: Optional precision characterization
R293	309308701	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Four shallow isolated eclipses in sectors73/86, including adjacent-cycle pairs, test recurrence. Target-deficit0.155pixel vs nearest capable neighbor1.50pixels favors target; no calibrated centroid errors. Gaia nominal conjunction offset0.6554d but full-covariance linear allowance0.8420d and near-reference offset0.0239d support compatible geometry. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R294	141811748	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated isolated eclipses across three sectors; sector/aperture-dependent dilution retained. Target-deficit0.187pixel vs closest capable neighbor1.67pixels favors target; no calibrated centroid errors. Sector37 actual adjacent cycles144/145/146 show raw core depths38.52/37.30/37.68ppt (11/11/12 in-event cadences); cycle145 fails fit side-window threshold but tests recurrence. Missing even sector98 cycles have no raw coverage. Sparse sector1 cycles0/1/2 also recur. Thus apparent alternating-sector98 pattern is coverage, not doubled period; see coverage-alias.json. Next test: Optional precision characterization
R295	5781196	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Two precisely fitted deep events in sectors36/99 are supplemented by helper raw sector89 THREE adjacent repeated eclipses near BTJD3719/3730/3741 separated by11.0018d. Those actual raw repeats test long-baseline integer-cycle aliases; they are not predicted centers used as measured timings. Strong aperture/sector dilution differences remain; no capable catalogued neighbor within60arcsec. Next test: Optional precision characterization
R296	370329832	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Five measured shallow eclipses include adjacent repeated pairs in sectors38/90, testing period recurrence. Target-deficit0.071pixel vs closest capable neighbor1.86pixels favors target. Sector63 alternative extraction depths22.30/21.22/17.45ppt show dilution. No calibrated localization error. Next test: Optional precision characterization
R297	379550793	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Crowded field: deficit0.315pixel from target; closest capable neighbor0.856pixel from deficit (G13.91,21.4arcsec). Peak/core favors target, with no calibrated centroid errors; aperture depths12.04/10.48/7.51ppt and sector depths differ through dilution. Intrinsic neighbor eclipse requirements are not directly compared with blended pixel losses. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R298	161823655	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Five measured events over three sectors, including adjacent same-kind repeats. Raw sector85 has changing baselines but clear narrow core; depth variation alone is not established astrophysically. No capable neighbor under approximate Gaia flux-budget screen. Next test: Optional precision characterization
R299	341118777	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep repeated events in sectors13/101. Sector104 cycle371 singular center covariance is excluded from timing; its real partial/noisy morphology is retained as documentation. No capable neighbor under approximate flux-budget screen. Next test: Optional precision characterization
R300	21952826	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Four measured repeats, with consecutive events in sector62. Deficit0.344pixel from target; nearest capable neighbor2.08pixels from deficit. Sector-dependent depth may reflect aperture dilution. Next test: Optional precision characterization
R301	308986181	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Eleven clear measured eclipses in three widely separated sectors. No capable neighbor under approximate Gaia flux-budget screen; target deficit0.176pixel from target. Next test: Optional precision characterization
R302	262082690	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Four real measured events in sectors99/102; adjacent cycles280/281 test recurrence. Historical sector38 adds independently extracted repeated events. Target deficit0.006pixel from target; nearest capable neighbor0.74pixel from deficit. Crowding and aperture dilution remain, but exact target-centered deficit/core favors target without a calibrated localization error. Next test: Optional precision characterization
R303	220568520	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated flat-bottomed raw eclipses over sectors1/30/97. Near-gap asymmetric sector97 cycle141 has singular center covariance and is excluded from timing; sector1 cycle0 and clean sector97 cycle142 are retained. No capable neighbor under approximate flux-budget screen. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R304	276277940	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep raw eclipses recur across three sectors, with sector-dependent depths130-160ppt. Sector83 cycle271 singular center covariance is excluded from timing. Historical cutout1/3/5 depths109.7/109.5/90.1ppt show aperture dilution; no capable catalogued neighbor under approximate brightness screen. Next test: Optional precision characterization
R305	406863699	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Three clear measured eclipses, including consecutive cycles88/89 in sector66 and recurrence in101; historical cutout65 also has two events. No capable neighbor under approximate flux-budget screen. Gaia conjunction offset0.053phase has propagated epoch uncertainty and approximate geometry, not an arbitrary significance veto. Next test: Optional precision characterization
R306	207148517	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Shallow repeat signal is measured in sectors69/97; individual fits vary and broad sector69 cycle236 is baseline-sensitive. Median depth is diluted. Historical PSF-scaled residual0.9997 is poor because bright off-center source dominates wide fit; local deficit core0.130pixel from target versus capable neighbor1.84pixels away supports target. Neighbor intrinsic requirement0.736 is not compared directly with its blended pixel fractional loss. Precision timing/shape is limited by low signal and baseline model. Next test: Optional precision characterization
R307	169657557	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Broad phase modulation surrounds a sharper eclipse core. Quadratic-baseline narrow-core fit replaces the broad linear-baseline timing/depth model; core depth16.34ppt and duration2.41h are model-dependent. Earlier broad-model period agrees within formal errors. Target deficit0.099pixel, closest capable neighbor1.41pixels; all capable sources checked geometrically. No calibrated centroid uncertainty. Next test: Optional precision characterization
R308	363534683	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Ten raw eclipses across three sectors include adjacent repeats. Target deficit0.125pixel versus capable neighbor1.78pixels. Sector depths vary through aperture dilution; no claim of intrinsic constancy. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R309	21295315	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Five raw eclipses include consecutive cycles63/64/65 in sector47, testing recurrence. Target deficit0.022pixel versus capable neighbor2.73pixels. Gaia conjunction linear uncertainty0.73d limits phase inference. Next test: Optional precision characterization
R310	445438807	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Twelve quality-zero measured raw events over three sectors test recurrence. No capable neighbor within60arcsec under approximate Gaia flux budget; target deficit0.188pixel. Finite-cadence trapezoid and aperture dilution limit precision. Next test: Optional precision characterization
R311	394386248	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Two consecutive quality-zero sector81 eclipses measure only a13.58d timing baseline, with formal/cadence period error0.00164d; no long-baseline precision asserted. Custom TESScut sector40 contains two further13.5769d-spaced raw repeats, all cores flag2048; these are flagged supportive recurrence, not clean nulls or precise timing points. Flagged difference image target offset0.15pixel with no capable neighbor under approximate flux screen. Flagged-image systematic and 9x9 residual0.657 limit localization; raw sector81 eclipses are unflagged. Next test: Optional precision characterization
R312	403154475	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Raw dips recur near2.535d, but Gaia orbit is5.06996d. Fit uses even half-period cycles1106,1108,1110,1112,1126,1128 as same-kind events; even median7.954ppt versus odd7.308ppt is not decisive physical-primary identification. Odd-only independent period5.0695421d agrees. Orbital5.0696716d interpretation uses same-kind parity and Gaia geometry; half-period dip recurrence is not silently called orbital period. No capable neighbor under flux-budget screen. Next test: Optional precision characterization
R313	137837801	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Eight deep raw eclipses over sectors28/68/96 include adjacent cycles0/1/2; aperture and sector depth differences reflect dilution/modeling. No capable neighbor under approximate Gaia flux budget. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R314	442606056	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Crowded unresolved field: deficit0.189pixel from target versus closest capable G12.6neighbor0.62pixel. Deficit maximum is target pixel and centroid is displaced away from this neighbor, favoring target. No calibrated localization uncertainty, so resolved photometry remains useful. Intrinsic neighbor requirement is not compared directly to blended pixel loss. Next test: Optional precision characterization
R315	50993470	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Five measured raw eclipses in sectors33/87 include consecutive cycles78/79 and237/238/239, testing recurrence. Sector7 lacks qualifying fitted events and is not claimed a clean null. Target deficit0.322pixel versus nearest capable neighbor1.32pixels; centroid uncertainty uncalibrated. Baseline slope and dilution affect depth. Next test: Optional precision characterization
R316	367362188	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fourteen raw eclipses across three sectors include consecutive cycles285/286 and multiple later repeats. Target deficit0.214pixel versus capable neighbor1.71pixels; depth and timing remain finite-cadence/model-dependent. Next test: Optional precision characterization
R317	388152396	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Reference epoch is the selected shallower eclipse; a deeper eclipse near phase0.34 supports an eccentric binary at the retained period. S54 selected-event depth varies about110–134ppt; no common intrinsic depth is inferred. An unrelated local dip at +0.20..0.45d in S81 cycle67 was removed from baseline fitting, with baseline on both sides retained. The period changes by only0.00000312d relative to the unmasked fit; this is a model-sensitivity comparison, not a total uncertainty. Next test: Optional resolved photometry or spectroscopy for precision characterization.
R318	63577915	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Large scaled-out-image PSF residual0.902 is not alone a failed attribution check: deficit core is on target, and no listed Gaia neighbour within60arcsec can supply the eclipse under the approximate G-band flux budget. No calibrated centroid ellipse is available. Next test: Optional resolved photometry or spectroscopy for precision characterization.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R319	11249710	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Crowded field with four capable neighbours. The deficit is0.306pixel from target and1.283pixels from the closest capable neighbour, so the target is favored at TESS resolution; precise exclusion probabilities are unavailable. Alternative S55 target-centered aperture gives about62ppt in three events, versus QLP about100ppt; dilution and aperture differences prevent an intrinsic-depth equality claim. Second eclipse near phase0.34 supports retained orbital period. Next test: Optional resolved photometry or spectroscopy for precision characterization.
R320	150582131	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated deep events include consecutive S79 cycles84/85. Approximate central-duration estimate is shorter than the fitted width; TIC radii and assumed main-sequence minimum-mass companion are uncertain, so this is not a fitted binary-radius solution or a quantitative duration match. Next test: Optional resolved photometry or spectroscopy for precision characterization.
R321	47653432	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Shallow4–7ppt locally normalized events recur with resolved localized shape; the whole-sector raw fold is dominated by baseline trends. Four capable neighbours are43arcsec or farther away from the compact target deficit. Observed duration about1.54h versus approximate central5.17h can arise from projected chord/radius assumptions; no measured central-duration match is claimed. Next test: Optional resolved photometry or spectroscopy for precision characterization.
R322	94113033	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Large scaled-out-image PSF residual0.947 is not alone a failed attribution check: deficit core is on target and no listed neighbour passes the approximate flux-budget screen. No calibrated centroid uncertainty is available. Next test: Optional resolved photometry or spectroscopy for precision characterization.
R323	314991763	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal phase-zero primary and phase-half secondary establish the retained full orbital period. One capable neighbour at56.26arcsec is separated from the compact target deficit. Next test: Optional resolved photometry or spectroscopy for precision characterization.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R325	115048464	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fitted duration and depth change across cadence/sector apertures; no precision intrinsic depth or common duration is claimed. The capable neighbour at51.03arcsec is separated from the target-core deficit. Next test: Optional resolved photometry or spectroscopy for precision characterization.
R327	240786248	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Seven measured eclipses include consecutive cross-sector cycles102/103 and170/171 after checking all five available QLP sectors17,57,58,84,85. Even/odd depths are confounded with sectors/apertures; they do not independently exclude an orbital double alias. Repeated morphology, actual spacing and Gaia period support the retained association. Depths depend on aperture dilution and local baseline. Pixel centroid uncertainties and timing systematics are not calibrated. Next test: Optional precision characterization
R328	322048799	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Representative same-kind raw eclipses recur within a sector. Photometric period is fitted from measured centers; the historical photometric branch guides extraction. No exhaustive blind alias scan claimed. A distinct secondary occurs near phase.20; it is not mixed into selected-event timings. Depths depend on aperture dilution and local baseline. Pixel centroid uncertainties and timing systematics are not calibrated. Next test: Optional precision characterization
R329	249834973	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Representative same-kind raw eclipses recur within a sector. Photometric period is fitted from measured centers; the historical photometric branch guides extraction. No exhaustive blind alias scan claimed. Depths depend on aperture dilution and local baseline. Pixel centroid uncertainties and timing systematics are not calibrated. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R330	162289987	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Representative same-kind raw eclipses recur within a sector. Photometric period is fitted from measured centers; the historical photometric branch guides extraction. No exhaustive blind alias scan claimed. A distinct secondary occurs near phase.53; it is not mixed into selected-event timings. Depths depend on aperture dilution and local baseline. Pixel centroid uncertainties and timing systematics are not calibrated. Next test: Optional precision characterization
R331	367461405	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Representative same-kind raw eclipses recur within a sector. Photometric period is fitted from measured centers; the historical photometric branch guides extraction. No exhaustive blind alias scan claimed. A distinct secondary occurs near phase.61; it is not mixed into selected-event timings. Depths depend on aperture dilution and local baseline. Pixel centroid uncertainties and timing systematics are not calibrated. Next test: Optional precision characterization
R332	22026010	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Representative same-kind raw eclipses recur within a sector. Photometric period is fitted from measured centers; the historical photometric branch guides extraction. No exhaustive blind alias scan claimed. A distinct secondary occurs near phase.60; it is not mixed into selected-event timings. Depths depend on aperture dilution and local baseline. Pixel centroid uncertainties and timing systematics are not calibrated. Next test: Optional precision characterization
R333	321044210	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Representative same-kind raw eclipses recur within a sector. Photometric period is fitted from measured centers; the historical photometric branch guides extraction. No exhaustive blind alias scan claimed. Depths depend on aperture dilution and local baseline. Pixel centroid uncertainties and timing systematics are not calibrated. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R334	365325928	G 12-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Representative same-kind raw eclipses recur within a sector. Photometric period is fitted from measured centers; the historical photometric branch guides extraction. No exhaustive blind alias scan claimed. A distinct secondary occurs near phase.52; it is not mixed into selected-event timings. Depths depend on aperture dilution and local baseline. Pixel centroid uncertainties and timing systematics are not calibrated. Next test: Optional precision characterization
R337	135362193	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Shallow repeated eclipses across sectors53/80; capable neighbors at18.9/22.9arcsec lie0.91/1.06pixel from deficit versus target0.11pixel. Target-centered core supports attribution at TESS resolution; no calibrated localization errors. Sector80 pipeline offsets remain systematic uncertainty. Next test: Optional precision characterization
R338	346865542	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Four deep isolated eclipses test within-sector recurrence and long-baseline cycle count. Aperture dilution retained; fitted depths are not intrinsic. Next test: Optional precision characterization
R339	131331287	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep repeated events have reduced depths in sector102 relative to37/64; aperture dilution or baseline changes possible, no intrinsic depth variation claimed. Next test: Optional precision characterization
R340	48752905	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating eclipses establish doubled period relative to historical SAP half-period. Deeper even historical cycles alone define epoch and orbital timing; mixed historical cutout stack is not a same-kind depth measurement. Next test: Optional precision characterization
R341	337881909	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Crowded field: eight capable neighbors, nearest17.8arcsec; deficit0.06pixel from target versus0.80pixel from nearest capable source. Target-centered core favors target, but highPSF fractional residual0.83 limits precise localization. No calibrated errors or resolved binary parameters claimed. Next test: Optional precision characterization
R342	274578846	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Same-kind repeat within sector53 and84 secures long-baseline cycle count. Modest aperture/sector depth differences retained. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R343	14291901	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated isolated eclipses at multiple successive cycles; Gaia conjunction offset retained with propagated orbit covariance and radius/model limitations. Next test: Optional precision characterization
R344	180179209	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Crowded field: eleven capable neighbors, nearest16.6arcsec; deficit0.23pixel from target versus1.00pixel from nearest capable source. PSF residual0.82 limits localization precision; target-centered core supports attribution at TESS resolution. Final partial/singular sector101 fit excluded from timing. Next test: Optional precision characterization
R345	351335839	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating eclipse depths establish doubled orbital period; deeper even historical cycles define ephemeris. Initial duration bound broadened to capture deepV morphology. Poor PSF residual0.99 limits detailed image modeling, but target-centered deficit and no capable listed neighbors support source. Mixed historical cutout stack cannot be equated to primary depth. Next test: Optional precision characterization
R346	89553209	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Sector14 cycle-1 lacks ingress coverage and is excluded from timing. Remaining five measured events include successive sector54/81 pairs and a full sector14 event. Historical prediction drift is fitted freely; partial raw event is not a precise timing. Broad eclipse duration and radius-based central estimate agree approximately. Next test: Optional precision characterization
R347	60555220	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Representative raw eclipses recur. Depths remain diluted and sector dependent; timing errors omit uncalibrated extraction and shape systematics. Next test: Optional precision characterization
R348	142932503	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Three separated measured eclipses at cycles0,48,89. Independent integer-count scan1-25d finds unique near18.69d fit; clean sector47 nulls reject identical deep events at divisors2,3,4. This is bounded alias evidence, not exhaustive all-period exclusion. Next test: Optional precision characterization
R349	256260982	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Representative raw eclipses recur. Depths remain diluted and sector dependent; timing errors omit uncalibrated extraction and shape systematics. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R350	11176109	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep repeated eclipses. Target0.142pixel from deficit versus nearest capable neighbor0.722pixel, which would require76percent intrinsic loss. Target favored by concentrated deficit core, without calibrated localization uncertainty. Next test: Optional precision characterization
R351	279823568	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Representative raw eclipses recur. Depths remain diluted and sector dependent; timing errors omit uncalibrated extraction and shape systematics. Next test: Optional precision characterization
R352	179857938	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Representative raw eclipses recur. Depths remain diluted and sector dependent; timing errors omit uncalibrated extraction and shape systematics. Next test: Optional precision characterization
R353	220617848	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Representative raw eclipses recur. Depths remain diluted and sector dependent; timing errors omit uncalibrated extraction and shape systematics. Next test: Optional precision characterization
R354	254723054	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Stable alternating depths establish doubled orbital period. Four deeper even historical half-period events alone define ephemeris; historical cutout stack mixes both kinds. Nearest capable neighbor is0.883pixel from deficit versus target0.100pixel; target favored at image resolution. Next test: Optional precision characterization
R355	257483992	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Representative raw eclipses recur. Depths remain diluted and sector dependent; timing errors omit uncalibrated extraction and shape systematics. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R356	388052602	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Stable alternating depths establish doubled orbital period. Ten deeper odd historical half-period events alone define ephemeris; historical cutout stack mixes both kinds. Gaia conjunction offset0.307d (0.068phase) exceeds the linear propagated0.101d error, but near-reference projected offset is0.06826d with independently propagated Gaia uncertainty0.02888d (about2.36 descriptive sigma) and the measured period differs from Gaia by0.000499d. Independent helper geometry388.py finds about3.03 descriptive sigma at the observed epoch. These are linear model uncertainties, not calibrated total coverage. Period accumulation explains much of the distant-epoch offset; low-eccentricity omega/periastron are strongly correlated. This association tension is retained without a sigma veto; updated spectroscopy would test it. Mass function0.126solar and approximate central duration5.61h versus observed3.48h admit noncentral binary geometry. Next test: Updated spectroscopy to test Gaia ephemeris tension
R357	157850336	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated isolated eclipses and successive raw cycles secure recurrence; target deficit0.066pixel versus capable neighbor2.53pixels. Aperture dilution retained. Next test: Optional precision characterization
R359	48870240	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Six measured raw events in sector91 include four successive cycles5-8 and the cycle0/1 pair, establishing recurrence. Capable neighbor1.39pixels from deficit versus target0.056pixel; target core favors attribution. Next test: Optional precision characterization
R360	178199444	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Three raw events in sector31 include consecutive cycles110/111 and a two-cycle separation108/110. Single selected sector supplies repeated cycles; this is not a single-event case. Next test: Optional precision characterization
R361	375058760	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep repeated V eclipses show sector-dependent morphology and fitted depths; raw baseline/shape systematics remain and fitted depths are diluted. No detailed binary characterization claimed. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R362	167923097	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep repeated V eclipses in sectors9/62/100; aperture differences and0.44pixel deficit offset retained; no capable listed neighbor. Next test: Optional precision characterization
R363	406297889	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated raw events across sectors14/55/82 with stable isolated morphology; no capable listed neighbor. Next test: Optional precision characterization
R364	160258534	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated eclipses across sectors15/56/83; capable neighbors2.51/2.73pixels from deficit versus target0.079pixel. Pixel core supports target at TESS resolution. Next test: Optional precision characterization
R365	16809274	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Persistent unequal alternating eclipses establish doubled orbital period relative to historical SAP half-period7.363d. Even historical cycles are the shallower event; only those five measured events define the ephemeris, explicitly not called deeper primary. Odd deeper events remain in raw evidence. Capable neighbor1.046pixels from deficit versus target0.160pixel; target core favors attribution. Historical cutout stack mixes eclipse kinds. Next test: Optional precision characterization
R366	279065540	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep repeated isolated eclipses across sectors58/85; no capable listed neighbor. Aperture dilution retained. Next test: Optional precision characterization
R367	439808137	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Representative raw eclipses recur. Depths remain diluted and sector dependent; timing errors omit uncalibrated extraction and shape systematics. Next test: Optional precision characterization
R369	239572145	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Stable alternating depths near150/55ppt establish doubled orbital period; six deeper even historical half-period events define ephemeris. Historical pixel stack mixes both eclipse kinds; concentrated target deficit supports localization. Closest Gaia conjunction offset+0.1907d (0.0302phase) versus linear covariance0.0717d is retained as descriptive association tension without arbitrary sigma veto. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R370	90285860	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Three measured eclipses include within-sector S41 recurrence, testing long-baseline cycle count. Closest Gaia conjunction offset-0.4519d (-0.0243phase) versus propagated0.2020d is retained as descriptive association tension, without arbitrary sigma veto; updated spectroscopy would test it. Next test: Optional precision characterization
R371	74717164	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Representative raw eclipses recur. Depths remain diluted and sector dependent; timing errors omit uncalibrated extraction and shape systematics. Next test: Optional precision characterization
R372	134193638	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated sharp raw eclipses coexist with strong oscillatory variability, especially sectors34/88. Free trapezoids can absorb background oscillation; timing reduced chi-square about9 is scatter inflated and uncalibrated shape/baseline systematics remain. Next test: Optional precision characterization
R373	34588199	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Representative raw eclipses recur. Depths remain diluted and sector dependent; timing errors omit uncalibrated extraction and shape systematics. Next test: Optional precision characterization
R374	146463944	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Representative raw eclipses recur. Depths remain diluted and sector dependent; timing errors omit uncalibrated extraction and shape systematics. Next test: Optional precision characterization
R375	113810700	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Representative raw eclipses recur. Depths remain diluted and sector dependent; timing errors omit uncalibrated extraction and shape systematics. Next test: Optional precision characterization
R376	120029059	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Representative raw eclipses recur. Depths remain diluted and sector dependent; timing errors omit uncalibrated extraction and shape systematics. Next test: Optional precision characterization
R377	66787531	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated isolated eclipses; deficit0.233pixel from target versus nearest capable neighbor1.675pixels. Aperture dilution limits intrinsic depth. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R378	382603504	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Persistent unequal alternating eclipses: even historical half-period cycles are deeper; only those17 measured events define the orbital ephemeris. Half-period is rejected by unequal parity depths in both sectors. Target deficit0.089pixel versus capable neighbor1.360pixels. Next test: Optional precision characterization
R379	305651410	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated flat-bottom eclipses; crowded field includes capable neighbor0.550pixel from deficit versus target0.059pixel. Full difference core favors target but subpixel crowding limits precision. Next test: Optional precision characterization
R381	421951253	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated isolated eclipses across widely separated sectors; no capable listed neighbor. Diluted fitted depths vary by sector aperture. Next test: Optional precision characterization
R382	128825695	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating eclipses establish doubled orbital period relative to historical SAP half branch. Even11 deeper events define ephemeris; odd secondary timings excluded. Target deficit0.083pixel versus capable neighbor2.531pixels. Next test: Optional precision characterization
R383	283635479	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Four repeated deep eclipses across sectors17/57/84; sector84 consecutive cycles146/147 test integer recurrence. Target deficit0.206pixel versus capable neighbor1.714pixels. Next test: Optional precision characterization
R384	9262715	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Seven deep repeated eclipses across sectors20/47/60, including sector47 consecutive cycles100-103. No capable listed neighbor. Next test: Optional precision characterization
R385	394620087	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Five deep repeated eclipses across sectors54/81, including consecutive cycles266/267. Target deficit0.123pixel versus capable neighbor1.436pixels. Next test: Optional precision characterization
R386	443900773	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep repeated flat-bottom eclipses across sectors7/46/88; no capable listed neighbor. Fitted depths differ by aperture and sector; no detailed binary characterization. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R387	438188407	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep repeated flat-bottom eclipses in three widely separated sectors. Capable neighbors are more than 2.24 pixels from the deficit, versus target 0.173 pixel. Gaia period drift is acknowledged through propagated covariance; no arbitrary formal-error veto is used. Next test: Optional resolved photometry or spectroscopy for precision characterization
R388	62337815	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Four isolated flat-bottom eclipses in sectors 76/83 with consecutive cycles in each. The 12.25 arcsec capable neighbor is 0.833 pixel from the deficit versus target 0.249 pixel; full image favors target, with residual subpixel crowding limitation. Next test: Optional resolved photometry or spectroscopy for precision characterization
R389	399285183	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Seven repeated localized shallow eclipses, with clean flat surrounding baseline in sector 81 and consecutive measured cycles. Capable neighbors are at least 1.135 pixels from the deficit versus target 0.142 pixel. Next test: Optional resolved photometry or spectroscopy for precision characterization
R391	457003443	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated isolated flat-bottom shallow eclipses across sectors 43/45/72. Full deficit image favors target at 0.218 pixel; capable neighbors are at least 1.708 pixels away. Aperture dilution and baseline variability limit depth interpretation. Next test: Optional resolved photometry or spectroscopy for precision characterization
R392	76020303	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Near-equal alternating eclipses recur every 1.85264 d. Same-parity even events give the reported orbital period 3.705279 d, consistent with Gaia 3.705347 d. Photometry alone retains the half-period interpretation because parity depths are indistinguishable; Gaia supplies the orbital interpretation, not an unannounced alias replacement. Both conjunctions are plausible for the paired eclipses. The target is 0.057 pixel from deficit versus capable neighbors beyond 2.12 pixels. Next test: Optional resolved photometry or spectroscopy for precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R393	251798466	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Three complete deep eclipses in sector 81, consecutive cycles 81-83, measure the period. Earlier sector 54 cycle 0 is a visibly partial raw repeat and supports recurrence but contributes no precision timing. Only the short complete-event baseline sets the stated formal error. Target deficit 0.063 pixel versus capable neighbor 2.789 pixels. Next test: Optional resolved photometry or spectroscopy for precision characterization
R395	399089869	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep repeated isolated eclipses across sectors 23/46/91; sector 46 consecutive cycles 70/71 test recurrence. No capable neighbor in the saved 60 arcsec cone. Aperture-dependent dilution limits intrinsic depth. Next test: Optional resolved photometry or spectroscopy for precision characterization
R396	329559935	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Only sector 80 supplied usable selected QLP data. Unequal alternating depths (odd median about 87.97 ppt; even about 41.28 ppt) establish the doubled orbital period, with five measured odd deeper eclipses spanning cycles -1 to 15. Half-period recurrence mixes two unequal eclipse types. Measured 2.603735 d agrees with Gaia 2.603862 d within the descriptive timing uncertainty; queue 2.60568 d is a historical search value, not adopted. Target deficit 0.081 pixel versus capable neighbors beyond 1.362 pixels. Next test: Optional resolved photometry or spectroscopy for precision characterization
R398	300148401	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated narrow eclipse cores on curved variable baseline; early S6 is sparse at30min cadence, while S33/98 resolve recurrence. Linear baseline and depth precision are limited by curvature; no capable listed neighbor. Next test: Optional precision characterization
R399	167125054	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated broad flat-bottom eclipses in S1/36/99; depth near60ppt. Target deficit0.239pixel versus capable neighbor2.036pixels. Next test: Optional precision characterization
R400	60514418	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated V-shaped eclipses in S43/44/71; consecutive S44 cycles test recurrence. No capable listed neighbor; observed duration3.72h is shorter than illustrative central duration8.77h, compatible with noncentral eclipse. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R401	257047728	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Two fully covered S48 eclipses are consecutive at15.20707d, directly testing recurrence rather than long-baseline integer counting. Historical S40 two-event difference stack supplies separate raw-pixel morphology; no capable listed neighbor. Timing precision is single-sector and not calibrated total uncertainty. Next test: Optional precision characterization
R402	168917716	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Shallow repeating eclipses across S14/74/82 with nearly equal alternating depths. Photometry supports6.701d repetition; selected even cycles give13.40244d consistent with Gaia orbit. Equal depths alone cannot reject half-period orbital identity; retained doubled branch explicitly uses Gaia and eclipse parity, with primary identity unresolved. Deficit0.086pixel versus nearest capable neighbor0.859pixel. Next test: Optional precision characterization
R403	156103314	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated flat-bottom eclipses across S14/41/74 with stable31-32ppt depth. Target deficit0.173pixel versus capable neighbor1.864pixels. Next test: Optional precision characterization
R404	140340353	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Thirteen narrow eclipses across S33/87 repeat coherently. No capable listed neighbor. Gaia conjunction offset-0.05phase (-0.167d) compared with full-covariance0.161d uncertainty: association is physically consistent within orbital-epoch uncertainty, without arbitrary phase veto. Next test: Optional precision characterization
R405	143467555	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Unequal alternating depths persist across S19/59/86: odd deeper events about145/167/162ppt versus even76/82/84ppt. Only13 odd deeper measured events define4.27351856d orbital period; half-period aliases are rejected by persistent parity morphology. Target deficit0.207pixel; mixed-kind historical difference stack still locates the signal. Next test: Optional precision characterization
R406	57358968	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Seven deep V-shaped eclipses across S45/71/72 with multiple consecutive local cycles; target-localized difference core and capable-neighbor geometry support attribution. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R408	451086560	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Deep flat-bottom eclipses repeat in S10/64/100. Partial S64 cycle179 ingress is excluded from timing. Aperture dilution and changing baseline can contribute to depth variation; deficit0.096pixel from target versus nearest capable neighbor1.679pixels. Adjacent S100 cycles directly test8.30d recurrence. A doubled orbital alias with equal eclipse kinds is not ruled out by parity alone; Gaia supports the retained branch. Next test: Optional precision characterization
R409	229308711	G 11.5-12 spectroscopic-orbit population	CONFIRMED	true	Report; Four long flat-bottom eclipses repeat in consecutive pairs in S6 and71. Curved/sloped out-of-eclipse variability and asymmetric troughs limit simple trapezoid midtimes. The timing fit allows scatter inflation when needed (none applied here), but does not include total shape systematics. Historical custom pixel stack confirms target deficit0.118pixel; no capable listed neighbor. Consecutive pairs test recurrence and long-baseline cycle count. Equal eclipse kinds could conceal a doubled orbital alias; Gaia supports the retained branch. Next test: Optional precision characterization
R410	401917014	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Eight deep raw eclipses across S12/93/102; capable G-band neighbor 46.1 arcsec away requires an extreme intrinsic loss, while the deficit core is target-centered. Pixel residual limits subpixel claims. Next test: Optional precision characterization
R411	442634143	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Three deep raw eclipses in S12/65. Observed duration exceeds a rough central radius/minimum-mass estimate; the estimate is model dependent and does not resolve detailed binary geometry. Next test: Optional precision characterization
R412	400057775	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Ten repeated raw eclipses S19/60/86. A capable neighbor 29.4 arcsec away is distinct from the target-centered deficit; high image-model residual limits precision. Next test: Optional precision characterization
R413	241089793	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; S18 clean raw eclipses at successive cycles -1 and 0 plus cycle 2 test the 5.632-day recurrence; measured S85 cycles 324/326 extend the timing baseline. S85 cycle 325 and S18 cycle 1 are flagged, not clean nulls. S18 partial cycle 3 is excluded from timing. The formal period error is not total calibrated uncertainty. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R414	72733549	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Three deep raw eclipses S52/79, including successive S79 cycles. A capable neighbor 48.1 arcsec away is distinct from the target-centered deficit. Next test: Optional precision characterization
R415	9172045	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Five repeated raw eclipses S47/60. A capable neighbor 18.8 arcsec away can in principle produce the blended loss, but the inspected deficit core falls on the target and shows no distinct neighbor peak; attribution is favored at TESS resolution, not proven subpixel. Next test: Optional precision characterization
R416	103809278	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Nine deep raw eclipses across S14/47/75; target-centered difference core and no capable Gaia neighbor under the brightness screen. Next test: Optional precision characterization
R417	131731452	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Five deep raw eclipses S9/63/99. A capable neighbor 32.4 arcsec away is distinct from the target-centered core; high pixel-model residual and changing aperture dilution limit intrinsic depth interpretation. Next test: Optional precision characterization
R418	150969574	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Eight repeated raw eclipses S57/84; target-centered difference core and no capable Gaia neighbor under the brightness screen. Next test: Optional precision characterization
R420	368260545	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; A similarly bright G=12.30 Gaia neighbor 22.87 arcsec away is flux-capable, but the deficit is 0.206 pixel from the target versus 0.92 pixel from that neighbor and peaks on the target pixel. A targeted recomputation of all four stacked windows puts every event deficit closer to the target (0.13–0.47 pixel) than the neighbor (0.75–1.07 pixel). Positional uncertainty remains descriptive rather than calibrated. Next test: Additional pixel orientations or spatially resolved photometry can strengthen attribution and refine the ephemeris.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R421	142427408	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Representative-sector timing and pixel evidence; period uncertainty is formal fit/cadence with scatter inflation where needed, not calibrated total coverage. Alternating half-period events were separated by eclipse kind before orbital timing. Crowded pixel PSF fit is poor, but all flux-capable neighbors are at least 27 arcsec from the target and farther from the deficit. Next test: Additional independent observations can refine the ephemeris and source attribution.
R422	21455449	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Representative-sector timing and pixel evidence; period uncertainty is formal fit/cadence with scatter inflation where needed, not calibrated total coverage. Gaia duration estimate is crude and exceeds the observed narrow event; geometry phase and period remain consistent. Next test: Additional independent observations can refine the ephemeris and source attribution.
R423	37531624	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Representative-sector timing and pixel evidence; period uncertainty is formal fit/cadence with scatter inflation where needed, not calibrated total coverage. Next test: Additional independent observations can refine the ephemeris and source attribution.
R424	39768877	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Representative-sector timing and pixel evidence; period uncertainty is formal fit/cadence with scatter inflation where needed, not calibrated total coverage. Next test: Additional independent observations can refine the ephemeris and source attribution.
R426	288605754	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Representative-sector timing and pixel evidence; period uncertainty is formal fit/cadence with scatter inflation where needed, not calibrated total coverage. Next test: Additional independent observations can refine the ephemeris and source attribution.
R428	159499138	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Representative-sector timing and pixel evidence; period uncertainty is formal fit/cadence with scatter inflation where needed, not calibrated total coverage. Next test: Additional independent observations can refine the ephemeris and source attribution.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R429	247204810	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Representative-sector timing and pixel evidence; period uncertainty is formal fit/cadence with scatter inflation where needed, not calibrated total coverage. Alternating half-period events were separated by eclipse kind before orbital timing. Next test: Additional independent observations can refine the ephemeris and source attribution.
R430	248688713	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Adjacent S32 cycles and S98 repeats establish recurrence. The brighter capable neighbor at 31.6 arcsec is distinct from the target-centered deficit; high image-model residual limits precision. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R433	77571117	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Six raw repeats span three sectors. The deficit is target-centered and no Gaia neighbor is capable under the approximate flux budget. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R435	270214638	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Six deep raw repeats span three sectors. The deficit is target-centered and no Gaia neighbor is capable under the approximate flux budget. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R437	31405122	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Thirteen deep raw repeats include six adjacent-cycle pairs. The deficit is target-centered and no Gaia neighbor is capable under the approximate flux budget. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R438	156976058	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Fourteen deep raw repeats include adjacent cycles in S53 and S80. The only nominally capable neighbor is 55.8 arcsec away, distinct from the target-centered deficit. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R439	335502443	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Five raw repeats include adjacent S60 cycles. A capable neighbor 16.2 arcsec away is distinct from the target-centered deficit core at TESS resolution. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R441	76450330	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Strong alternating depths establish the 2.1333 d orbital cycle; the historical SAP 1.06665 d branch is the half-period. The deeper parity is fitted for the reported epoch. Next test: Optional precision characterization
R442	95247597	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; S88 has a consecutive 14.376 d same-kind recurrence and S61 adds a long baseline. Earlier S7 windows have only five quality-zero points each, so they are not clean nulls. Current MAST QLP availability includes S7/34/61/88; availability alone is not an eclipse detection. Next test: Optional precision characterization
R443	436762701	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Seven deep raw events across S56 and S83 include consecutive cycles, with consistent depths and a target-centered deficit. Next test: Optional precision characterization
R444	462538667	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Eight deep raw events across S41 and S75 include consecutive cycles, with a well-centered pixel deficit and stable depths. Next test: Optional precision characterization
R445	62433366	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Even historical half-period cycles are modestly but persistently deeper than odd cycles in S76 and S84, supporting the 4.7597 d orbital period. The duration is only a rough trapezoid shape measure. Next test: Optional precision characterization
R446	220213855	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Four consecutive raw eclipses in S87 establish recurrence across 16.43 d. S6 is available but its candidate windows have at most four quality-zero cadences inside half the search duration; no clean null is claimed. The fitted period error is limited by one sector and is not total ephemeris uncertainty. Next test: Optional precision characterization
R450	384076458	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Sixteen measured raw eclipses span sectors 66 and 102 with twelve adjacent-cycle pairs. The only approximately capable Gaia neighbor is 46.6 arcsec away from the target-centered deficit. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R451	27732047	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Deep and shallow eclipses alternate at the 2.58356-d photometric search period. The 5.16713-d orbital ephemeris uses the eleven deeper odd-parity centers; shallow even-parity eclipses independently support the double-period interpretation. Two nominally capable Gaia neighbors lie beyond 54 arcsec; the source image is structured, but its strongest deficit is on the target core. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R452	309253911	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Nine measured eclipses in sectors 1, 38 and 96 include five adjacent-cycle pairs. No Gaia neighbor within 60 arcsec meets the approximate flux budget. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R453	237240609	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Five measured eclipses in sectors 54 and 81 include a sector-81 adjacent pair. No Gaia neighbor within 60 arcsec meets the approximate flux budget. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R455	339368119	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Nine deep measured eclipses in sectors 19, 59 and 86 include five adjacent-cycle pairs. No Gaia neighbor within 60 arcsec meets the approximate flux budget. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R456	270478197	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Four deep measured eclipses include an adjacent sector-70 pair and a sector-71 repeat. No Gaia neighbor within 60 arcsec meets the approximate flux budget. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R457	233095433	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Five deep measured eclipses include adjacent pairs in sectors 14 and 53 and a sector-86 repeat. No Gaia neighbor within 60 arcsec meets the approximate flux budget. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R458	3942765	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Four eclipse-shaped events occur in sector 48; three centers give the ephemeris. Cycle 136 has a positive pre-ingress excursion that biases its fitted baseline and midpoint, so it is retained as morphology and excluded from timing. The remaining adjacent pair tests recurrence; no capable Gaia neighbor lies within 60 arcsec. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R460	258317152	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Eight strong raw eclipses across S10/99/101. The historical S64 deficit peaks on the target, with no capable Gaia neighbor under the approximate flux budget; the image-model residual is large, so its centroid is only descriptive. Next test: Optional precision characterization
R461	102052526	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Deep and shallow events alternate in S9/63/99. The 3.706-day photometric half-period is not the orbital period: the deeper even parity recurs at 7.412 days. The historical S63 aperture depths are strongly diluted and its image-model residual is high, but the deficit core is on the target and no capable Gaia neighbor is found. Next test: Optional precision characterization
R462	261702897	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Five raw eclipses across S12/99/102. Two capable Gaia neighbors at about 45 and 55 arcsec require separate evaluation; the S38 deficit peak is at the target pixel, distinct from the bright southeast neighbor. The poor image-model fit limits centroid precision. Next test: Optional precision characterization
R463	34014384	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Six raw eclipses across S2/69/96, including three consecutive S69 cycles. The S69 deficit peaks at the target and the aperture reductions retain the eclipse. Next test: Optional precision characterization
R464	123267654	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Twelve raw eclipses across S97/104, with consecutive cycles and consistent depths. The S97 cutout has a clean target-centered deficit. The selected earlier S3 QLP file supplied no usable midpoint; it is not treated as a null eclipse. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R465	285232364	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Four broad raw eclipses in S58/85, two adjacent cycles per sector. A capable neighbor about 56 arcsec away is spatially distinct from the S18 target-centered deficit. The image fit is imperfect and aperture depths are diluted. Next test: Optional precision characterization
R467	310996111	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Four shallow S76 raw events recur at the predicted phase, and the historical S49 aperture extraction retains a roughly 3 ppt target-pixel loss. S15/23 are 30-minute cadence: each predicted short event has at most one central sample, so their apparent weak or absent dips are not clean nulls. Capable neighbors about 52 and 55 arcsec away are distinct from the target-centered S49 deficit; the morphology and attribution remain limited by shallow depth. Next test: Optional precision characterization
R468	294082465	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Three broad deep raw events in S8/99, including consecutive S99 cycles. The S99 target-centered deficit is supported by all three apertures, though that historical image stack contains one event only. Next test: Optional precision characterization
R470	24524336	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; S32 cycles 98/100 and S98 cycles 339/341/343/345 have opposite half-period parity. Intervening windows are quality-flagged, not clean nulls. The within-sector 15.099-day spacing together with cross-sector parity excludes a constant doubled period. Next test: Optional precision characterization
R471	141923153	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Consecutive raw eclipses in S7 and S34 support recurrence; the capable 51.6-arcsec neighbor is separated from the target-centered deficit despite substantial image-model residual. Next test: Optional precision characterization
R473	348773879	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; The capable 17.0-arcsec neighbor is about a pixel away; the deficit core falls much closer to the target. Attribution favors the target at TESS resolution, with uncalibrated subpixel precision. Next test: Optional precision characterization
R476	33789753	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Five raw events in S6/S33/S87 agree in timing and shape. No brightness-capable Gaia neighbor appears within 60 arcsec. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R477	330469458	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Five deep raw events in S16/S83 recur at the measured period. A capable neighbor 46.3 arcsec away is distinct from the target-centered deficit. Next test: Optional precision characterization
R478	238126968	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Shallow and deep eclipses alternate at the historical 4.94798-day search interval. Five deep even-cycle midpoints give the 9.89595-day orbital period; shallow odd cycles support the alternating pattern. Next test: Optional precision characterization
R479	220410034	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; S98 contains measured eclipses in both parity classes of the historical period, excluding a simple double-period alias. No brightness-capable Gaia neighbor appears within 60 arcsec. Next test: Optional precision characterization
R480	308609252	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Four raw events include an adjacent sector-58 pair. The sector-18 difference core is centered near the target, while the only capable catalogued Gaia neighbor is 54.4 arcsec away. The 0.94 image-model residual limits centroid precision. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R481	34759144	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Three clean raw eclipses occur one per selected sector. Enumerating neighboring integer cycle counts gives 57 then 37 cycles at 19.33997 d; the next-best pairs disagree by at least 0.0248 d between gaps. This tests nearby aliases, while subharmonics remain bounded by observed coverage. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R482	408189324	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Six consecutive sector-32 events and six sector-98 events support the same 4.39365-day cycle. The deficit is sharply centered on the target and no capable Gaia neighbor is listed within 60 arcsec. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R483	30770290	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Sector-39 and sector-66 measured pairs are separated by two historical cycles, but independent inspection finds a real ~69-ppt usable-core deficit at intervening sector-66 cycle 176. Its partial coverage prevents a reliable midpoint fit; it supports the 8.3804-day recurrence over a simple 16.76-day parity alias. Sector-39 cycle 88 is a gap, not a null. The high image-model residual limits localization precision. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R484	306980055	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Measured sector-39 and sector-93 adjacent pairs span the long baseline. The capable 55.8-arcsec Gaia neighbor is distinct from the target-centered difference core. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R487	144102918	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Thirteen measured eclipses span sectors 4, 31 and 97, including adjacent cycles. Difference imaging is sharply target-centered with no capable Gaia neighbor in the 60-arcsec screen. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R489	189435226	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Eight raw eclipses span sectors 17, 58 and 85, with adjacent cycles in both later sectors. The difference image favors the target and no capable Gaia neighbor is listed within 60 arcsec. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R490	316138676	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Five deep raw eclipses across S18/59/86, including consecutive S59 and S86 cycles. The deficit core is centered on the target; no capable Gaia neighbor is present under the approximate brightness screen. The long fitted duration is geometry dependent. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R491	127267444	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Three consecutive raw eclipses in S92 give a measured 8.299744-day period, close to the Gaia 8.299164-day orbit; the historical 8.29212-day screening estimate was inaccurate. Two capable neighbors at 27.8 and 49.5 arcsec are spatially distinct from the target-centered deficit. The pixel-model residual is substantial. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R492	61260847	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Fifteen deep raw eclipses in S43/70/71 establish recurrence and a measured period. The cutout core favors the target with no capable Gaia neighbor; local image residuals limit precise centroid claims. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R495	63013423	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Twelve raw eclipses in S44/46/72 show stable recurrence. The capable neighbor 49.1 arcsec away is distinct from the target-centered deficit; aperture and model residuals limit exact intrinsic depth. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R496	167314199	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Eight deep raw eclipses in S15/75/83 recur on the Gaia timescale. A capable 35.7-arcsec neighbor requires an extreme intrinsic eclipse and is distinct from the target-favored core; the image-model residual is high. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R497	1643990	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Six deep raw eclipses in S8/61/99 give a long timing baseline. The cutout deficit lies on the target with low model residual and no capable Gaia neighbor under the brightness screen. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R498	313998039	G 12-12.25 spectroscopic-orbit population	CONFIRMED	true	Report; Four deep raw eclipses in S76/83 repeat consistently. Bright capable neighbors at 29.6 and 39.1 arcsec cause severe aperture dilution, but the deficit core favors the target location; high image residual prevents precise localization. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R501	370829832	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; The S63 difference image has a remote single-pixel maximum at [7,14] coincident with a bright field source; its origin is uncertain. The multi-pixel eclipse deficit and centroid lie near the target [7,8]. The remote maximum and large image-model residual limit precision localization. Next test: Optional higher-resolution eclipse imaging and binary characterization.
R502	51564054	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Twenty raw events across S54/55/81 recur at the measured period; the difference core lies on the target and no Gaia neighbor passes the approximate brightness screen. Next test: Optional precision binary characterization.
R503	315929104	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Representative raw eclipses in S18/59/86 and a target-centered S18 difference core support the target; capable neighbors are tens of arcsec away. Next test: Optional precision binary characterization.
R504	400118957	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Raw depths change with sector aperture, while nine measured midpoints retain one recurrence. A capable neighbor 15.3 arcsec away is distinct from the target-centered difference core; crowding limits intrinsic depth claims. Next test: Optional precision binary characterization.
R505	305826287	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Eight raw events across S58/85 have stable depth and a target-centered difference core; the one capable listed neighbor is 44.5 arcsec away. Next test: Optional precision binary characterization.
R507	233199615	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Thirteen representative raw events across S14/53/86 repeat at the same period; the difference core is target-centered and no listed Gaia neighbor passes the approximate brightness screen. Next test: Optional precision binary characterization.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R509	195206508	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Alternating shallow even and deeper odd eclipses across S15/55/82 set a 13.407-day orbital period. The deeper odd cycles alone define the reported epoch and timing fit. The S55 cutout has a central target deficit, but high image-model residual and neighboring field stars limit subpixel localization. Next test: Optional resolved eclipse imaging and refined primary/secondary modeling.
R510	97649150	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; S7/34/87 raw eclipses agree. The closest capable neighbor is G17.24 at 13.7 arcsec and would need roughly 54% intrinsic loss; deficit is within 0.12 pixel of target. Faint close-source ambiguity remains below calibrated pixel precision. Next test: Optional resolved time-series photometry.
R511	210343372	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; S41/82 provide consecutive raw repeats. The closest capable G16.03 source lies 17.8 arcsec away and would require about 22% intrinsic loss; deficit is within 0.20 pixel of target. The Gaia phase offset is descriptive and within its large propagated orbital uncertainty. Next test: Optional resolved time-series photometry.
R512	49901727	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Four same-sector raw eclipses establish recurrence and a photometric period, but the short single-sector baseline leaves a comparatively broad period error. No capable Gaia neighbor within 60 arcsec; target-centered difference image. Next test: Additional-sector timing would refine the period.
R513	424636246	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; S14/55/84 raw repeats and target-centered deficit. Capable catalogued neighbors are at least 50.6 arcsec away and need large intrinsic losses; no nearby source is favored by the deficit image. Next test: Optional precision characterization.
R514	285926385	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Deep S9/63/90 raw eclipses recur at the measured period. Difference image favors the target and lists no capable Gaia neighbor within 60 arcsec. Next test: Optional precision characterization.
R517	392763429	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; S13/94/103 raw eclipses follow the same clock. A brief S103 depth doubling affects consecutive cycles, not stable odd/even alternation; its cause is unresolved. S13 difference image is target-centered with no capable Gaia neighbor within 60 arcsec. Next test: Investigate the S103 depth discontinuity with aperture diagnostics.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R518	1978575696	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Alternating raw depths identify the historical 3.486-day clock as half the orbital eclipse cycle. Deeper even cycles yield a measured 6.97216-day same-kind period, consistent with Gaia; S41 difference image favors the target. A capable G14.97 source at 23.9 arcsec is not favored by the target-centered deficit. Next test: Optional resolved photometry and secondary-eclipse characterization.
R520	351610379	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; S58/86 raw repeats and the S58 deficit favor the target. The nearest capable G15.21 neighbor is 20.3 arcsec away and would require about 42% intrinsic eclipse depth; a brighter capable source at 54.1 arcsec is spatially separate. Pixel localization has no calibrated error. Next test: Optional resolved eclipse photometry.
R521	61462126	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; S13/67/104 raw eclipses repeat. The S27 alternative extraction recovers two events with a target-centered deficit; no Gaia neighbor within 60 arcsec passes the approximate flux-budget test. Next test: Optional precision characterization.
R522	169318948	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; S32/98 raw repeated eclipses and the S98 alternative extraction agree. The difference image favors the target and lists no capable Gaia neighbor within 60 arcsec. Next test: Optional precision characterization.
R523	441257677	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; S13/27/67 raw events recur with no persistent odd/even depth split. The S13 alternative extraction shows four events and a target-centered deficit with no capable Gaia neighbor within 60 arcsec. Next test: Optional precision characterization.
R524	153972241	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Deep S14/41/76 raw eclipses repeat; the S75 extraction recovers six events. Its target-centered difference image lists no capable Gaia neighbor within 60 arcsec. Next test: Optional precision characterization.
R525	384744266	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; S12/39/102 raw repeats agree. The S12 deficit is near the target. Three nominally capable neighbors lie 29–60 arcsec away, each requiring about 78–100% intrinsic loss, and none is favored by the observed deficit. Next test: Optional resolved eclipse photometry.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R526	163946960	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; S10/99/101 raw repeats and the S10 alternative extraction support the eclipse. The deficit lies about 0.26 pixel from the target, away from two capable Gaia neighbors at 15.8 and 29.7 arcsec; those require about 60% and 44% intrinsic losses. Localization is descriptive, without a calibrated uncertainty. Next test: Resolved eclipse photometry would strengthen host attribution.
R527	191634144	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; S57/84 raw shallow eclipses recur. Odd/even fitted depths vary at roughly the local noise level without a stable half-period interpretation. The S57 difference image favors the target over capable neighbors at 23.9, 34.4 and 53.6 arcsec. Next test: Additional-sector timing and resolved photometry could tighten this shallow case.
R528	219399648	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Alternating S32/98 raw depths show that the 4.66571-day SAP search branch is half the orbital eclipse cycle. Deeper even events give the 9.33142-day same-kind period consistent with Gaia. The S98 difference image favors the target with no capable Gaia neighbor within 60 arcsec. Next test: Optional secondary-eclipse characterization.
R530	411063531	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Six raw events span S10, S64 and S100, with adjacent pairs in both later sectors. The crowded S90 deficit centroid is 3.1 arcsec from the target while the light centroid is displaced 26.3 arcsec; four flux-capable catalogued neighbors are 44–47 arcsec from the deficit. The target is favored, but the image-model residual (0.86) limits localization precision. Next test: Resolved eclipse imaging would strengthen attribution in this crowded field.
R531	399293743	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Five raw events span S54 and S81, including three adjacent S81 cycles. The S81 deficit lies near the target, about 17.5 arcsec from the only flux-capable Gaia neighbor; the source is favored at TESS resolution. Next test: Optional resolved photometry and spectroscopy.
R532	44487490	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Adjacent S17 and S84 dips alternate timing around the 3.44695-day photometric search clock. Odd cycles define the same-kind 6.89392-day orbit, matching Gaia; the even events are the other eclipse kind. The S84 image is target-centered, with the sole capable neighbor about 54 arcsec away. Next test: Optional characterization of both eclipse kinds.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R533	426704374	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Raw eclipses recur across S12, S66 and S102, including adjacent S12 and S66 cycles. The S12 deficit is target-centered and no Gaia neighbor in the 60-arcsec screen passes the flux-budget bound. Next test: Optional precision characterization.
R534	316637090	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Ten shallow raw events in S53 and S80 repeat; the S80 aperture extraction recovers six. The target-near deficit is roughly 31 arcsec from the only capable Gaia neighbor. The Gaia conjunction offset is descriptive and is not used as a significance veto. Next test: Additional resolved photometry would strengthen this shallow attribution.
R535	382581210	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; The current MAST QLP availability record lists nine sectors; all nine files were retrieved and checked. Ten measured quality-zero raw eclipses in S2, S3, S29, S68, S69 and S95 now span the long baseline, including adjacent S29 and S68 pairs. The S29 clean intermediate windows reject equal-kind half, third and quarter-period submultiples; flagged windows were not used as nulls. The historical S2 cutout separately stacks two sparse predicted windows into one target-centered deficit, which alone would not establish each window as an eclipse. Its capable Gaia neighbor is 33 arcsec from the target and 32.5 arcsec from the deficit. Next test: Optional later-sector timing and resolved photometry.
R536	277598042	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Fourteen raw events span S13, S92 and S94 with multiple adjacent cycles. The S94 difference core lies near the target and about 20 arcsec from the only capable Gaia neighbor; that neighbor would need an approximately 80% intrinsic eclipse. Next test: Optional resolved photometry.
R537	470627360	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; S59 has two adjacent measured eclipses and S73 supplies a longer-baseline third. The S59 difference core is target-centered, about 35.5 arcsec from its only capable Gaia neighbor; the long event is also recovered by aperture extraction. Next test: Further-sector timing would refine the modest three-event ephemeris.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R538	440168538	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Two adjacent raw eclipses in each of S57 and S84 anchor the recurrence. The S57 deficit favors the target; no Gaia neighbor within 60 arcsec passes the flux-budget screen. The image-model residual (0.77) limits localization precision. Next test: Optional resolved photometry.
R539	446159745	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Seven raw eclipses span S19, S53 and S74, with an adjacent S53 triplet. The S20 alternative extraction is a separate observing occasion with three events; its deficit is sharply target-centered and no capable Gaia neighbor is listed. Next test: Optional precision characterization.
R540	125783923	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Four deep raw eclipses; compact target-centered deficit and no flux-capable Gaia neighbor within 60 arcsec. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.
R541	24928269	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Twelve deep raw eclipses over S3/30/97. Sector-dependent depth is consistent with changing dilution; isolated target-centered deficit. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.
R544	78886343	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Eight deep raw eclipses; compact deficit is near the target, distinct from both capable neighbors (approximately 1.7 and 2.4 pixels away). Bright off-target light displaces the out-of-eclipse centroid; it does not move the deficit peak. Target attribution is favored at TESS resolution. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.
R546	406972595	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Seven repeated raw eclipses with within-sector recurrence in S93; compact target deficit and no capable Gaia neighbor within 60 arcsec. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R547	123220066	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Five actual raw midpoints after inspecting all five currently available QLP sectors (8/35/62/89/99). A bounded 0.8-40 day integer-alias test leaves only the 19.389-day recurrence without clean sampled nulls; the half-period predicts four nulls. Two capable sources at 32 and 45 arcsec are separated from the target-centered deficit. One-event image noise and non-calibrated localization remain limitations. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.
R549	65709882	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Four deep raw eclipses including consecutive S7 events support recurrence. Fresh deficit lies 0.12 pixel from target while all five capable alternatives are at least about one pixel away. The very bright off-target light dominates the full image but not its compact deficit. Attribution favors target at TESS resolution; nearest capable neighbor limits subpixel claims. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.
R550	300339443	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Two near-equal-depth event kinds recur at unequal phase separation. Nearby known EB Gaia DR3 1827899632630791296 is 48.55 arcsec away, G=18.463, has catalogue P=19.923 d, and would require an impossible approximate 32.52 fractional intrinsic loss to produce this depth. Its pixel (9.694,8.783) is away from the target core. This is a contamination check, not an exact target catalogue recovery. Other capable neighbors are spatially separated from the deficit core. Next test: Optional resolved photometry and precision binary characterization.
R551	142502809	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated sharp shallow eclipses across representative sectors. Raw full-sector plot includes additional under-sampled or incompletely bracketed events excluded from free timing. No capable Gaia neighbor within the historical 60-arcsec flux screen. Next test: Optional resolved photometry and precision binary characterization.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R552	246857844	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Only two complete timed eclipses, both in S2 and adjacent with spacing14.17242d, rather than two long-separated events with unresolved integer count. Current MAST query returns QLP sectors2,29,69, all inspected. S29 starts during a partial real eclipse; it is not used as a midpoint. S69 selected-eclipse windows lack clean coverage. Two-center period error propagates timing allowances with zero fit degrees of freedom; it does not measure timing scatter. Half-cycle covered baselines show no comparable eclipse. This short-baseline photometric period remains imprecise; no all-sector long-baseline precision is claimed. Next test: A complete new eclipse for a longer-baseline ephemeris.
R553	206614346	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Six complete selected-kind raw eclipses span S2/68/96. Depth varies with aperture and sector. Narrow sharp eclipses are distinct from broad baseline variability; no capable Gaia neighbor in the saved 60-arcsec screen. Next test: Optional resolved photometry and precision binary characterization.
R554	17033860	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Deep selected eclipse and a much shallower second kind at phase+0.2737 establish unequal event separation. No capable Gaia neighbor under the depth budget. Eclipse shapes remain sharp and recurring across S14/41/75. Next test: Optional resolved photometry and precision binary characterization.
R555	278966364	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Small sharp eclipses repeat in S42/70/92, including adjacent S92 events. Large positive spikes and gaps are visible in the raw sectors; the selected eclipse cores are distinct from those artifacts. The weaker sparse S92 event increases model and timing uncertainty; timing covariance is scatter-inflated. Next test: Optional resolved photometry and precision binary characterization.
R556	338072783	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Twenty complete timing-usable eclipses and17 within-sector consecutive pairs establish recurrence. Two capable neighbors at23.0 and52.7arcsec are separated from the deficit core. Strong out-of-eclipse variability remains visible; local raw eclipse fits do not use detrended QLP flux. Next test: Optional resolved photometry and precision binary characterization.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R557	464680877	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Two recomputed S65 deficits locate the mean0.254px from target versus0.617px from a capable G14.18 neighbor17.29arcsec away (target-neighbor separation0.82px). The maximum core and aperture pattern favor the target. Only two deficit occasions and uncalibrated pixel-response systematics limit subpixel claims, but the neighbor lies above the target-centered core. The repeated shallower kind atphase+0.4834 distinguishes the measured orbital recurrence. Next test: Optional resolved photometry and precision binary characterization.
R559	260765760	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Deep selected eclipse and shallower second kind atphase-0.4677 recur with unequal depths. Generic exact-half-phase baseline windows partly overlap the secondary; their negative depth estimates are not clean eclipse depths. Fresh pixel aperture deficits remain deep and target-centered; no capable Gaia neighbor under the flux budget. Next test: Optional resolved photometry and precision binary characterization.
R564	393761712	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Four deep eclipses repeat across three QLP sectors, with consecutive same-kind S72 events. Compact target deficit and no capable Gaia source in60arcsec support target attribution. Next test: Optional resolved follow-up for physical parameters and orbital eclipse labeling.
R568	349452680	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; All four currently available QLP sectors43/44/59/86 inspected; three clear actual events. Bounded integer aliases0.8-40d reject shorter alternatives through sampled clean nulls;19.34676d and38.69352d survive. Reported19.34673d is observed eclipse recurrence; double orbital-period equal-eclipse labeling remains unresolved. Sole capable neighbor47.1arcsec away is spatially distinct from compact target deficit. Next test: Optional resolved follow-up for physical parameters and orbital eclipse labeling.
R570	308151602	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Four deep events include consecutive pairs. Target-centered deficit is distinct from three capable neighbors at 27.55-42.48 arcsec. Next test: Optional resolved photometry and spectroscopy for improved localization and physical parameters.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R571	206538272	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Twenty-one deep events recur; nineteen shallower secondary fits at phase 0.50006 distinguish eclipse kinds. No capable Gaia neighbor within 60 arcsec. Next test: Optional resolved photometry and spectroscopy for improved localization and physical parameters.
R572	81797376	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Eight deep events repeat across three sectors. The one capable neighbor is 49.17 arcsec away, spatially distinct from the target deficit. Next test: Optional resolved photometry and spectroscopy for improved localization and physical parameters.
R574	28116156	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Nine short eclipses in S30 establish same-kind recurrence; the period/error use this actual within-sector baseline. Coarse S3 does not contribute precise timings. The capable neighbor at 30.94 arcsec is spatially distinct from the compact deficit. Next test: Optional resolved photometry and spectroscopy for improved localization and physical parameters.
R575	322679338	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Four deep events form consecutive pairs in S60/S73. Two shallow secondary candidates in S19 lie at actual phase 0.54415, with other-sector confirmation limited. The capable neighbor at 54.93 arcsec is separate from the target deficit. Next test: Optional resolved photometry and spectroscopy for improved localization and physical parameters.
R576	147278528	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Six selected eclipses repeat on a changing variable baseline. The target deficit is compact and separate from capable neighbors at 32.19 and 51.96 arcsec. Variability limits baseline, timing and depth precision. Next test: Optional resolved photometry and spectroscopy for improved localization and physical parameters.
R577	100756625	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Eleven deep selected events recur; four shallow secondary candidates lie near phase 0.49956. Compact target deficit and no capable neighbor within 60 arcsec. Next test: Optional resolved photometry and spectroscopy for improved localization and physical parameters.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R578	137395412	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Four selected centers form consecutive pairs in S19/S53. No capable neighbor within 60 arcsec. No significant secondary center is claimed; Gaia predicts the other conjunction at actual phase 0.72348, so a half-phase null alone is not an eccentric-secondary test. Next test: Optional resolved photometry and spectroscopy for improved localization and physical parameters.
R579	316913462	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Eight selected events recur in S95/S104, plus five shallow secondary candidates at phase 0.49943, near Gaia prediction 0.49584. Target pixel loss 27.92 e-/s versus 5.85 e-/s at the capable 22.51-arcsec neighbor favors the target at TESS resolution despite overlapping wings. Modern Gaia conjunction offset 0.692 days reflects accumulated period difference: projection near Gaia reference agrees within 0.00132 days; P difference 0.001206 days versus Gaia error 0.000506 days. These descriptive tensions are not a definitive contradiction. Next test: Optional resolved photometry and spectroscopy for improved localization and physical parameters.
R580	153711481	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Five shallow but sharp recurring eclipses, including consecutive cycles in S31 and S98. Fresh S97 four-event deficit is centered on target (0.068pixel). Both capable neighbors, at33.4/54.0arcsec, are spatially distinct from the compact positive core; their pixels show weak target-wing/noise losses rather than distinct matching deficit peaks. A bright remote field source distorts the total light image, explaining high global PSF residual; it does not move the eclipse core. The shallow stack is noisy (target pixel deficit3.13e/s versus0.14e/s at closer capable source); this supports target attribution but no calibrated localization confidence is claimed. Raw per-point scatter is about2ppt, so coherent recurrence and multiple in-event samples, rather than any individual point, establish eclipse morphology. Next test: Optional resolved follow-up and physical binary modeling.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R581	367147644	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Five sharp repeated eclipses and consecutive S59 cycles. Fresh compact target deficit has no flux-capable Gaia neighbor within60arcsec. Strong raw baseline curvature in S17/S86 is not mistaken for a secondary. Gaia conjunction offset0.04136phase is comparable to its0.313d propagated timing allowance, not a hard0.03phase veto. Next test: Optional resolved follow-up and physical binary modeling.
R582	165779351	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Four measured deep flat-bottomed eclipses establish same-kind recurrence. Fresh compact target deficit is distinct from all three capable neighbors at36.8-58.4arcsec. A weak recurring secondary candidate lies near0.72phase, consistent with eccentric Gaia0.7297 +/-0.0483; free-center secondary fits are not secure, so no secondary precision timing is claimed. Next test: Optional resolved follow-up and physical binary modeling.
R583	91614687	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Six deep sharp same-kind eclipses and five measured unequal secondaries near0.50712phase distinguish the11.61516d orbit from its half. Compact target deficit has no capable Gaia neighbor within60arcsec. Next test: Optional resolved follow-up and physical binary modeling.
R584	66503432	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Nine deep selected eclipses and eight measured secondaries establish unequal spacings: secondary phase0.47448, then0.52552 back to selected kind. Half period cannot represent this event sequence. Compact target deficit has no capable Gaia neighbor within60arcsec. Source-specific Gaia predicts other phase0.51951 +/-0.01607 (linear allowance), a0.0450phase tension (~2.8 allowances). Argument-of-periastron and A/B conventions were checked against methods; observed centers, not Gaia predictions, define secondary phase. Primary conjunction agrees within0.0374d versus0.0965d allowance. This modest secondary geometry tension is retained as a limitation, not a calibrated rejection. Approximate minimum-mass central-duration model also remains a plausibility screen. Next test: Refine spectroscopic eccentric geometry and compare measured secondary phases; resolved physical modeling.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R586	152876230	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Four deep selected-kind events in S68/S95 and three measured secondaries near0.71238phase support eccentric12.21853d orbit and reject a half-period interpretation. Fresh target-centered deficit has no capable Gaia neighbor within60arcsec. S1 lacks an accepted selected-center fit but has secondary coverage, not a clean null. Selected depths347ppt in S68 versus300ppt S95 show sector dilution changes; reported median is a pipeline-dependent quantity. Next test: Optional resolved follow-up and physical binary modeling.
R587	47066501	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Eighteen sharp selected eclipses across three sectors establish3.03829d. Brighter G10.68 neighbor34.7arcsec away dominates the baseline image but the compact deficit is on target, spatially displaced from that neighbor. The1x1/3x3/5x5 depth dilution progression and the26.1arcsec light-to-deficit displacement are expected for that blend; source ownership follows actual deficit morphology. No secure secondary is claimed. Next test: Optional resolved follow-up and physical binary modeling.
R588	462346588	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Four flat-bottomed selected eclipses include three consecutive S81 cycles. Compact target deficit is distinct from all three capable sources at47.3-59.0arcsec. Long selected duration8.63h is compatible with approximate central12.73h geometry. No secure secondary is claimed. Next test: Optional resolved follow-up and physical binary modeling.
R591	158072011	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated box eclipses and a secondary near phase 0.397. The deficit favors the target over the capable 36.05 arcsec neighbor; the bright blended out-of-eclipse image displaces the light centroid. Saturation/blend morphology limits precision localization. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.
R592	5903277	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated eclipses with unequal secondary near phase 0.495. Compact target-centered deficit; no flux-capable Gaia neighbor within 60 arcsec under the approximate G-band screen. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R593	314713331	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated box eclipses and weak secondary near phase 0.500; target-centered deficit distinct from two capable neighbors 36.57 and 53.57 arcsec away. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.
R594	356241663	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated narrow eclipses plus weak secondary. Target-centered deficit without a capable neighbor within 60 arcsec. Gaia conjunction offset-0.0457 phase is descriptive and lies within the propagated orbit timing allowance; it is not a fixed phase-cut rejection. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.
R595	392794270	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated sharp box eclipses are distinct from strong changing baseline variability. No secure secondary phase is assigned. The target-centered difference image and no capable Gaia neighbor support ownership. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.
R596	18558131	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Repeated deep eclipses and shallower secondary near phase 0.500; compact target deficit. The nearby known EB Gaia 5634548560328478848 at 48.69 arcsec has period 0.64499 d, unlike this 5.33939 d signal, and is too faint under the G-band loss budget. It is a contamination alert, not a target catalogue recovery. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.
R597	314040568	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Persistent unequal alternating 120/190 ppt dips reject the historical half-period as an orbital period. The quoted 9.180514665 d ephemeris uses seven deeper odd-parity measured eclipses; shallow even-parity events form the phase 0.500 secondary. Fresh pixel stack uses only deeper events and favors target over the 57.49 arcsec capable neighbor. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R598	154652154	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Four QLP primary centers span multiple sectors, with sector 84 primaries two cycles apart. Additional free-center custom 3 x 3 cutout timings in sector 57 include consecutive same-kind 83 ppt eclipses, confirming the 7.47035 d cycle count; see cutout-s 57-timing.json and image. No null is inferred at uncovered QLP cycle 244. Compact deficit has no capable Gaia neighbor. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.
R599	49621708	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Four deep primary eclipses include consecutive pairs in sectors 70 and 92, plus a much shallower secondary near phase 0.496. Compact target deficit and no capable Gaia neighbor within 60 arcsec. Next test: Optional resolved photometry and spectroscopy for precision physical parameters.
R600	287756198	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Representative sectors and simplified trapezoid/linear timing errors; shared TESS photons; approximate Gaia edge-on geometry and G-band neighbor flux budgets. Next test: Additional clean eclipse timing and color-aware neighbor/PRF modeling would refine the bounded result.
R601	301986385	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Representative sectors and simplified trapezoid/linear timing errors; shared TESS photons; approximate Gaia edge-on geometry and G-band neighbor flux budgets. Next test: Additional clean eclipse timing and color-aware neighbor/PRF modeling would refine the bounded result.
R602	164660905	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Representative sectors and simplified trapezoid/linear timing errors; shared TESS photons; approximate Gaia edge-on geometry and G-band neighbor flux budgets. Next test: Additional clean eclipse timing and color-aware neighbor/PRF modeling would refine the bounded result.
R604	376669046	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Representative sectors and simplified trapezoid/linear timing errors; shared TESS photons; approximate Gaia edge-on geometry and G-band neighbor flux budgets. Next test: Additional clean eclipse timing and color-aware neighbor/PRF modeling would refine the bounded result.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R605	338069871	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Representative sectors and simplified trapezoid/linear timing errors; shared TESS photons; approximate Gaia edge-on geometry and G-band neighbor flux budgets. Next test: Additional clean eclipse timing and color-aware neighbor/PRF modeling would refine the bounded result.
R606	21411263	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Representative sectors and simplified trapezoid/linear timing errors; shared TESS photons; approximate Gaia edge-on geometry and G-band neighbor flux budgets. Only two clean timings; flagged S21 windows; local recurrence precision only. Next test: Additional clean eclipse timing and color-aware neighbor/PRF modeling would refine the bounded result.
R607	395396499	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Representative sectors and simplified trapezoid/linear timing errors; shared TESS photons; approximate Gaia edge-on geometry and G-band neighbor flux budgets. Next test: Additional clean eclipse timing and color-aware neighbor/PRF modeling would refine the bounded result.
R608	73023909	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Representative sectors and simplified trapezoid/linear timing errors; shared TESS photons; approximate Gaia edge-on geometry and G-band neighbor flux budgets. Next test: Additional clean eclipse timing and color-aware neighbor/PRF modeling would refine the bounded result.
R611	176048490	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Four deep, flat-bottomed raw eclipses and an unequal secondary near phase0.3683 support eccentric binary morphology. The capable neighbor at56.6 arcsec is separated from the deficit. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R612	76177367	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Six narrow raw eclipses and another eclipse near phase0.5930; two capable neighbors at35 and57 arcsec lie away from the deficit. Very bright field light shifts the full-light centroid and dilutes aperture photometry; the compact lost flux remains near target. No deeper-primary assignment is claimed; baseline/extraction differences limit the depth comparison. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R613	101259338	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Sixteen repeated narrow eclipses and an unequal secondary near0.5004. No capable Gaia neighbor within60 arcsec; compact lost flux favors target. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R614	157367278	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Sixteen deep flat-bottomed eclipses, shallower secondary near0.5000. No capable Gaia neighbor within60 arcsec. Bright displaced field light shifts the light centroid; target-centered deficit remains distinct. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R615	264485594	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Eight repeated shallow narrow raw eclipses. Target-centered compact deficit; no capable Gaia neighbor within60 arcsec. No secure secondary; baseline variability limits null claims. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R616	202442866	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Nine deep repeated raw eclipses plus a broad shallower secondary near0.4988. Target-centered deficit; no capable Gaia neighbor within60 arcsec. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R617	297956126	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Five measured deep eclipses after adding S66 QLP. Consecutive same-kind S66 eclipses resolve the long-baseline cycle-count ambiguity. Unequal eccentric secondary near0.7244; three capable neighbors are spatially separated from compact deficit. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R618	216024097	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Nine repeated flat-bottomed raw eclipses and shallower secondary near0.5051. Three capable neighbors at38-56 arcsec are displaced from target deficit core. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R620	269506784	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Fourteen deep repeated raw eclipses and shallower secondary near phase0.4849. Strong sector97 baseline excursions were kept visible; local eclipse baselines remain stable. No capable Gaia neighbor within60arcsec; compact deficit favors target. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R621	369587471	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Twelve flat-bottomed repeated raw eclipses and a shallower secondary near0.5009. Sector-dependent baseline variability remains visible. No capable Gaia neighbor within60arcsec; target-centered compact deficit. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R622	138080650	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Ten deeper primary timings at6.95225485d. Historical3.47613d SAP branch mixes unequal74/32ppt eclipses; deep even cycles alone fit the primary ephemeris. Actual secondary near0.50006. Both capable44.73/45.92arcsec neighbors are displaced from compact deficit. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R623	165793290	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Sixteen deep V-shaped repeated eclipses plus unequal secondary near0.49975. No capable Gaia neighbor within60arcsec; compact deficit is on target. Gaia conjunction offset0.0644d exceeds the linear0.0196d propagation; its phase0.0267 offset is explicitly retained rather than treated as a Gaussian rejection. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R624	93862657	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Twelve deep repeated raw eclipses and shallower secondary near0.4977. Compact lost flux favors target over capable24.17/48.03arcsec neighbors. Nearby known EB Gaia3041697316497710720 at29.83arcsec has G19.73955 and0.282338d catalog period; its intrinsic-loss proxy74.05 exceeds unity and its period differs from4.80345d. It cannot plausibly make the100ppt target eclipse under this brightness model. Crowding raises whole-field PSF residual; no calibrated localization precision claimed. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R625	334608511	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Twenty-two repeated narrow V-shaped eclipses and unequal eccentric secondary near0.54790. Deficit is near target; capable51.33arcsec neighbor is separated from deficit core. Bright field source shifts overall field light; whole-field PSF residual is descriptive. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R626	138854818	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Eleven consistent shallow raw eclipse timings across three sectors; representative windows show narrow repeatable11ppt dips despite substantial sector98 baseline noise. Compact sector98 deficit favors target over capable33.33/53.79arcsec neighbors. No secure secondary; opposite-phase coverage lacks a comparable primary, without excluding shallow dips. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R627	435209690	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Four deep flat-bottomed measured raw eclipses in sectors56/83, including consecutive same-kind pairs in both sectors. These resolve the long-baseline integer count. Shallower secondary is measured near0.50613, not the Gaia-guided initial search phase. No capable Gaia neighbor within60arcsec; compact target-centered deficit. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R629	186604857	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Twelve repeated narrow flat-bottomed raw eclipses and a shallower secondary near0.49975. The capable26.13arcsec neighbor is displaced from the compact target-centered deficit. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R630	165992260	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R631	31816149	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R632	116849606	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R633	45914154	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R634	295546522	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R639	237092634	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Local sector24 ephemeris only; later primary windows flagged/gapped. Next test: Optional additional clean epochs and resolved photometry.
R640	14163319	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Eight measured narrow repeated eclipses in sectors5/32 with stable local baselines. Sector98 has severe excursions retained in the full phase view but no fitted event; its half-phase artifact is not a secure secondary. No capable Gaia neighbor within60arcsec; compact deficit centered on target. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R641	73248723	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Three deep measured eclipses including consecutive sector47 pair, plus shallower observed secondary near0.633. Consecutive same-kind timings resolve integer counts. No capable Gaia neighbor within60arcsec; compact deficit is on target. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R643	70241645	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Four narrow repeated eclipses in three sectors, including consecutive sector81 same-kind events. No secure secondary. All five capable neighbors at25.25-49.41arcsec are displaced from compact target deficit; crowded field light centroid is descriptive. Custom cutout reduction in sector40 recovers one event, a distinct observing occasion from the retained QLP sectors. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R644	265092213	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Historical3.95216d SAP branch mixed unequal47/76ppt eclipses. Six identifiable deeper odd-cycle timings yield7.90432331d; one odd event has singular timing covariance and is excluded. Actual shallower secondary near0.50015. Both capable42.66/53.06arcsec neighbors are displaced from compact target deficit. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R645	257773097	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Fourteen measured deep repeated eclipses and shallower secondary near0.49823. Sector94 aperture depths are shallower than12/66; differences remain visible. No capable Gaia neighbor within60arcsec; compact deficit favors target. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R646	256464205	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Nine measured narrow repeated eclipses across three sectors. No secure secondary. Both capable Gaia neighbors at28.53/41.16arcsec lie away from compact target deficit. Nearby known EB Gaia6067018781690100608 at43.74arcsec has G18.650408 and0.309493d period versus target7.706608d; intrinsic required-loss proxy exceeds10 and its location is displaced. It cannot plausibly make the target eclipse under the brightness model. Whole-field light offset40.8arcsec reflects crowding, not eclipse localization. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R647	150162108	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Eight deep measured repeated eclipses and unequal secondary near0.500. No capable Gaia neighbor within60arcsec; compact target deficit. Secondary realized depth is used rather than extrapolated model amplitude. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R649	274795477	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Six deep repeated eclipses with shallower eccentric secondary near0.63234. All four capable neighbors at31.82-47.57arcsec lie away from compact target deficit; aperture depth dilution remains explicit. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R650	263479136	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R651	440522293	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R652	123485559	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R653	376539517	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R654	445645985	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R655	429570033	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R656	48879301	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R658	151502866	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R661	191562862	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R662	11547648	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R664	464615512	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R666	260981962	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R667	13882408	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R668	229563422	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R669	54140419	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R671	365899880	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R672	376359531	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R674	138997145	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Next test: Optional additional clean epochs and resolved photometry.
R676	357019754	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	true	Report; Representative three QLP sectors; three s12 center covariances are singular and excluded, while another s12 center remains usable. The shallow secondary is visible but its own ephemeris was not fitted.
R677	187202091	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	true	Report; Sectors 73/86 have curved SAP baselines; their free trapezoid centers are rejected. Adopted local-minimum time widths are heuristic, so period error is conditional. Gaia conjunction phase has weak discrimination after long extrapolation. Six cached pixel events peak on target, but the 17.1-arcsec Gaia alternative is only disfavored by uncalibrated centroid geometry; subpixel companions remain unresolved.
R680	402782627	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	true	Report; Correlated out-of-eclipse variation broadens event scatter; 30-min s18 centers are less precise. TIC radius 6.24 solar radii predicts a longer central event than measured, compatible with grazing geometry or uncertain radius. Cached TESScut is the same TESS photons as QLP.
R682	150465032	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	true	Report; Current six VizieR EB/variable catalogues returned successful empty exact-area results; Final alias and novelty checks were completed in the accepted decision.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R683	331650047	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	true	Report; Prior limitations: VSX lists exact Gaia source as ROT, not an EB; this does not negate the observed eclipse. One Kostov table query remains unavailable; Novelty was reconciled in the accepted decision.; decision: Previously accepted physical review; scoped catalogue clearance, root accepts new claim.
R684	384744604	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	true	Report; Prior limitations: Several catalogue services were unavailable in the initial pass; absence of rows is not a nonmatch. Root must complete novelty reconciliation.; decision: Previously accepted physical review; scoped catalogue clearance, root accepts new claim.
R727	238855958	Gaia spectroscopic-binary cached population; mixed-harvest prioritized dip reviews	CONFIRMED	true	Report; Repeated eclipses, target-favored spatial evidence and Gaia agreement; singular S1 covariance, no formal period uncertainty. S1 timing covariance is singular; reported period is rounded from multiple good later events with no formal uncertainty. Shorter-alias nulls assume similar same-kind ~7-ppt eclipses. Alternative extraction shares TESS photons. Next check: A later resolved eclipse would tighten the ephemeris.

Included probable systems (102)

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R003	298645930	G 10-11.5 spectroscopic-orbit population	PROBABLE	true	Report; Repeated eclipses secure; host attribution/check unresolved. See dated tier reconciliation and unchanged historical report.
R048	279166593	G 10-11.5 spectroscopic-orbit population	PROBABLE	true	Report; Repeated eclipses secure; host attribution/check unresolved. See dated tier reconciliation and unchanged historical report.
R058	236531158	G 10-11.5 spectroscopic-orbit population	PROBABLE	true	Report; Repeated eclipses secure; host attribution/check unresolved. See dated tier reconciliation and unchanged historical report.
R065	250742811	G 10-11.5 spectroscopic-orbit population	PROBABLE	true	Report; Repeated eclipses secure; host attribution/check unresolved. See dated tier reconciliation and unchanged historical report.
R067	351541670	G 10-11.5 spectroscopic-orbit population	PROBABLE	true	Report; Repeated eclipses secure; host attribution/check unresolved. See dated tier reconciliation and unchanged historical report.
R098	186100980	G 10-11.5 spectroscopic-orbit population	PROBABLE	true	Report; Repeated eclipses secure; host attribution/check unresolved. See dated tier reconciliation and unchanged historical report.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R107	39069184	G 10-11.5 spectroscopic-orbit population	PROBABLE	true	Report; Repeated eclipses secure; host attribution/check unresolved. See dated tier reconciliation and unchanged historical report.
R136	101779765	G 10-11.5 spectroscopic-orbit population	PROBABLE	true	Report; Repeated eclipses secure; host attribution/check unresolved. See dated tier reconciliation and unchanged historical report.
R153	47719859	G 10-11.5 spectroscopic-orbit population	PROBABLE	true	Report; Refit the Gaia SB2 radial velocities jointly with the long-baseline eclipse ephemeris to resolve the orbital association.
C23	465263741	fainter SB1 period-fold population	PROBABLE	true	Report; See target report for deciding limitation and next check.
C29	285942031	fainter SB1 period-fold population	PROBABLE	true	Report; Reassess Gaia orbital uncertainties/correlations or obtain an updated radial-velocity orbit to resolve the eclipse-versus-Gaia timing tension.
C31	281393563	fainter SB1 period-fold population	PROBABLE	true	Report; Obtain higher resolution time-series imaging or a validated multi-source PRF difference fit to identify the eclipsing host.
C33	126363604	fainter SB1 period-fold population	PROBABLE	true	Report; Validate the repeated difference-image astrometry with a multi-source PRF or higher resolution event photometry; refine Gaia conjunction uncertainty if needed.
C44	201317466	fainter SB1 period-fold population	PROBABLE	true	Report; Refit the RV orbit or evaluate a nonlinear orbital posterior against the observed eclipse timing to resolve Gaia association.
C50	443106429	fainter SB1 period-fold population	PROBABLE	true	Report; Resolve target versusTIC 443106423 with higher spatial resolution eclipse photometry.
C57	402940895	fainter SB1 period-fold population	PROBABLE	true	Report; Reassessment of historical C57; two capable close neighbors remain unresolved. See updated report and prior archived plausible review.
C67	9396327	fainter SB1 period-fold population	PROBABLE	true	Report; Refine Gaia conjunction predictions with orbital posterior and measure a second clean sector or obtain new RVs to test the orbit association.
R215	167178300	12-14.8 d underreviewed Gaia period-fold pilot	PROBABLE	true	Report; Capable Gaia neighbours remain at 58.583arcsec, 41.62arcsec, 12.984arcsec, 24.468arcsec, 18.174arcsec, 59.428arcsec. Deficit centroid favors target, but close blend is unresolved without a PRF/deblend uncertainty test. Next test: Resolved eclipse photometry or multi-source PRF localization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R216	325573958	12-14.8 d underreviewed Gaia period-fold pilot	PROBABLE	true	Report; Capable Gaia neighbours remain at 57.77arcsec, 7.958arcsec, 20.628arcsec. Deficit centroid favors target, but close blend is unresolved without a PRF/deblend uncertainty test. Next test: Resolved eclipse photometry or multi-source PRF localization
R217	428691463	G 11.5-12 spectroscopic-orbit population	PROBABLE	true	Report; Unresolved attribution: capable neighbor Gaia2514032005529105 92 at6.95arcsec needs only2.3% fractional eclipse. Deficit centroid lies between target and neighbor; PSF residual0.815 and uncalibrated centroid cannot discriminate. Real repeated eclipses remain probable for this target. Next test: Resolved eclipse photometry / multi-source PRF localization
R231	448485777	G 11.5-12 spectroscopic-orbit population	PROBABLE	true	Report; Crowded field: G13.14 neighbor16.26arcsec away requires16.1% eclipse, deficit0.615pixel from neighbor vs0.252pixel target. Another14.44arcsec neighbor requires97.6%. Target is favored by central deficit and deeper1x1 aperture, but blending and uncalibrated localization leave attribution physically unresolved after completed checks. Next test: Resolved eclipse photometry / multi-source PRF localization
R246	128460489	G 11.5-12 spectroscopic-orbit population	PROBABLE	true	Report; Unresolved attribution after completed check: capable Gaia5600259057572857856 neighbor11.02arcsec away requires25.4percent intrinsic eclipse and shares rounded TESS pixel with target. Deficit centroid is0.270pixel from target and0.516pixel from neighbor, favoring target but without calibrated uncertainty sufficient to exclude neighbor. Local blended pixel loss cannot exclude that neighbor. Retain probable for target pending resolved photometry or multi-source PRF localization. Next test: Resolved eclipse photometry / multi-source PRF localization
R247	379054533	G 12-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Crowded field with capable bright neighbor at11.93arcsec requiring only2.1% eclipse; target-centered deficit favors target but uncalibrated localization cannot securely distinguish close blended contributors. Attribution remains uncertain. Next test: Resolved eclipse photometry / multi-source PRF localization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R253	419573851	G 12-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Capable neighbor at4.34arcsec needs49% eclipse. Deficit0.073pixel from target favors target, but close separation and uncalibrated centroid cannot securely exclude neighbor. Completed review retains probable attribution. Next test: Resolved eclipse photometry / multi-source PRF localization
R254	124007093	G 12-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Long eclipses coexist with baseline variability; free local linear fits and conservative timing errors preserve its contribution. Target-centered cutout deficit0.17pixel from target favors target, but nearest capable neighbor18.57arcsec needs only8.3% eclipse, and PSF residual0.75 without calibrated centroid uncertainty leaves attribution unresolved. Independent optimizer timing centers differ by up to0.0152d; model systematics are not total-error calibrated. Next test: Resolved eclipse photometry / multi-source PRF localization
R256	48108063	G 12-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Deep raw eclipses approximately134-143ppt. Independent measured-ephemeris S62 cutout re-extraction gives3x3 core deficit26.5ppt and1x1 32.1ppt; the factor4-5 contrast persists. Dilution and extraction differences are possible, unestablished explanations. Deficit0.036pixel from target vs neighbor0.72pixel favors target, but15arcsec capable neighbor and uncalibrated centroid leave interpretation unresolved. Next test: Resolved eclipse photometry / multi-source PRF localization
R258	88811793	G 12-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Unresolved close-neighbor attribution after completed check. Capable neighbor14.54arcsec away lies0.75pixel from deficit centroid; target is0.087pixel away. Another capable neighbor27.33arcsec away is spatially distinct. Deficit favors target, but no calibrated multi-source PRF localization excludes the closest source. Strong aperture dilution: cutout45/24/15ppt versus QLP approximately60ppt. Pixel fractional loss is blended light, not source-intrinsic required eclipse. Next test: Resolved eclipse photometry / multi-source PRF localization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R272	207813753	G 12-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Completed attribution check remains unresolved:16 capable neighbors, including12.72–18.90arcsec sources. Target-centered deficit favors target but lacks calibrated multistar localization to rule out close sources. Aperture depths differ strongly in crowded field. Real repeating eclipses retained probable. Next test: Resolved eclipse photometry or calibrated multisource PRF localization.
R273	173040554	G 12-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Seventeen selected even-half-cycle times; deeper ordering usually favors even but noisy shapes make primary identity provisional. Thirteen capable neighbors include13.75arcsec G13.70; centroid0.37pixel from target cannot exclude blended sources. Real raw alternating dips and orbital-period fit retained probable. Geometry phase offset-0.040 is discussed with Gaia covariance, not an automatic veto. Next test: Resolved eclipse photometry or calibrated multisource PRF localization.
R281	374589618	G 11.5-12 spectroscopic-orbit population	PROBABLE	true	Report; Repeated shallow alternating eclipses are real; odd half-period cycles define deeper-eclipse timing. Deficit0.099pixel from target favors target, but capable neighbor13.5arcsec away is only0.67pixel from deficit and needs approximately79percent intrinsic eclipse. An undersampled sector54 fit can exaggerate depth (cycle2 fitted15.37ppt versus observed minimum about7-8ppt); that even-parity fit is excluded from the deeper odd-parity ephemeris. Shallow noisy difference image lacks calibrated localization; source attribution remains unresolved after bounded check. Next test: Resolved eclipse photometry / multi-source PRF localization
R324	102716983	G 12-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Repeating eclipses are secure, including consecutive S59 cycles94/95 measured after supplementing the original selection. Attribution is unresolved: two capable neighbours at14.39/14.45arcsec lie within one TESS pixel, and the historical PSF residual is0.920. The third capable neighbour is26.56arcsec away. Source localization requires resolved imaging or multi-source PRF analysis. Next test: Resolved eclipse photometry or multi-source PRF localization.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R336	601646479	G 12-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Unequal alternating eclipses establish full3.535848d orbit from13 deeper even historical half-period cycles, orbital_cycle=cycle/2. Attribution remains unresolved against the capable1.89arcsec neighbor; its deficit distance0.137px is slightly smaller than target0.165px. Image cannot resolve these sources. Depths depend on aperture dilution and local baseline. Pixel centroid uncertainties and timing systematics are not calibrated. Next test: Resolve source with higher-resolution eclipse photometry
R380	343340931	G 11.5-12 spectroscopic-orbit population	PROBABLE	true	Report; Attribution remains unresolved: capable neighbor only2.879arcsec from target, required intrinsic eclipse0.205; deficit distances0.190pixel to target and0.282pixel to neighbor are insufficient to distinguish these sources reliably. TIC mass/radius absent; approximate solar-fallback duration4.79h cannot constrain observed9.82h. Gaia association remains possible; source assignment needs resolved photometry. Next test: Resolved eclipse photometry / multi-source PRF localization
R394	363264601	G 11.5-12 spectroscopic-orbit population	PROBABLE	true	Report; Repeated localized 4-5 ppt depressions have flat surrounding baseline in sectors 100/103 and support genuine eclipses. Attribution remains unresolved after a completed check: a 5.595 arcsec neighbor can supply the signal with 1.85 percent intrinsic loss. Target/neighbor deficit distances 0.093/0.350 pixel favor the target but do not reliably distinguish these close sources without a calibrated localization. This is PROBABLE because the eclipses repeat and attribution is unresolved, not because their morphology is uncertain. Next test: Resolved eclipse photometry or calibrated multi-source PRF localization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R397	119193784	G 11.5-12 spectroscopic-orbit population	PROBABLE	true	Report; Four deep V-shaped/truncated-bottom events across S50/91, all odd historical10.445d cycles. Intermediate cycles0 and106 have zero quality-zero points within0.2d (42 and173 flagged points respectively), so neither is a clean null. Available QLP inventory lists only S50/91. Two20.8906d-separated pairs establish real repetition, but10.4453149d period is conditional on the Gaia-informed cycle count;20.8906d and integer-count aliases are not independently rejected. This missing orbital-alias check retains PROBABLE despite completed bounded inspection. Target deficit0.191pixel versus capable neighbor2.279pixels; dilution limits intrinsic depth. Next test: Clean intermediate-cycle eclipse photometry to test orbital aliases
R419	2055563318	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; Eight repeated raw eclipses S29/95/104 show a real recurring signal, but a capable Gaia neighbor only 1.1 arcsec away shares the central TESS pixel. The image cannot uniquely identify the eclipsing star; geometry agrees with the selected Gaia orbit but does not cure this attribution limit. Next test: Resolved photometry / multi-source PRF localization
R425	409852500	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; Two G~16.3 neighbors at 7.3 and 9.5 arcsec are capable in flux and share a TESS pixel; the deficit centroid favors the target but does not securely resolve them. The sector-104 cycle-294 egress gap excludes that center from timing. Next test: Obtain spatially resolved eclipse photometry, or a stronger difference image in a suitable sector.
R427	251731963	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; A G=12.27 Gaia star 3.69 arcsec away can produce the dip with a ~27% intrinsic eclipse; the TESS pixel centroid cannot resolve the pair. Next test: Resolve the two stars in higher-spatial-resolution time-series photometry at predicted eclipse times.
R432	700593737	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; A raw S19 adjacent-cycle pair confirms repetition. A capable 2.8 arcsec Gaia companion shares the central TESS core; source attribution is unresolved. Next test: Resolved eclipse photometry or multi-source PSF localization of the 2.8 arcsec pair.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R436	66281997	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; Eight usable raw centers lie in six sectors after all eight archived QLP sectors were examined; they include adjacent S56/S57 and S76/S77 cycles. The historical S57 deficit lies 9.1 arcsec from the target, toward a capable G16.8 star 21.8 arcsec away. Noisy pixels and other capable neighbors prevent unique attribution. Next test: Independent high-resolution eclipse photometry or multi-source PSF localization of the crowded deficit core.
R447	265436769	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; Four shallow raw events in S95 and S104 include same-sector consecutive cycles. The S67 difference core favors the target, yet a capable source 22.6 arcsec away and a high image-model residual (0.87) leave attribution unresolved. Early S1 windows are sparse or flagged, not clean nulls. Next test: Resolved source localization
R449	395080968	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; Alternating depths support the 3.6265 d orbital period, but raw fits scatter strongly and residuals reach 75 min. The S56 alternative aperture stack mixes eclipse kinds; its depth is not an intrinsic primary depth. Capable neighbors at 6.3 and 11.9 arcsec share the TESS image core and the image-model residual is 0.982, so source attribution and precision ephemeris remain limited. Next test: Resolved source localization
R459	40076679	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; Seven shallow raw eclipses recur in sectors 39, 66 and 103; three adjacent sector-39 cycles directly test the period. Both sector-39 and newly measured sector-66 difference images have strong structured residuals in a crowded field. Ten Gaia neighbors within 60 arcsec can in principle produce the blended depth. The sector-66 deficit core is near the target, yet the pixel data cannot uniquely assign the signal. Next test: Resolved time-series photometry or multi-source pixel modeling during an eclipse to identify the variable source.
R466	283016810	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; Thirteen raw eclipses across S17/58/85. Three capable Gaia neighbors include one about 19 arcsec away. The S24 deficit peak overlaps the target pixel, but TESS cannot resolve the nearby capable source within roughly one pixel. The image-model residual further limits attribution. Next test: Resolved eclipse photometry or multi-source PRF localization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R469	349749097	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; Sixteen raw eclipses across S13/93/104. Two capable Gaia neighbors about 24 and 33 arcsec away lie in a crowded field. The S13 deficit core overlaps the target pixel, but the high image-model residual and roughly one-pixel separation leave source identity unresolved. Next test: Resolved eclipse photometry or multi-source PRF localization
R472	167656210	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; A 2.9-arcsec capable Gaia neighbor shares the TESS core. The recurring eclipses are real, but these pixels cannot uniquely assign them to the selected Gaia star. Next test: Resolved time-series photometry or multi-source PRF fitting of the 2.9-arcsec pair.
R475	460866286	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; The field has 24 brightness-capable Gaia neighbors within 60 arcsec, including one at 12.3 arcsec. The deficit core is near the target, but the weak signal and 0.97 image-model residual cannot uniquely resolve nearby alternatives. Next test: Higher-resolution eclipse photometry or multi-source PRF fitting in this crowded field.
R485	202634485	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; Sector-55 has three consecutive clean raw eclipses. The sector-82 raw baseline is structured; its same-photon custom-aperture signal is much shallower than QLP SAP, with crowding, aperture and baseline sensitivity likely contributing. A Gaia neighbor 10.1 arcsec away could in principle supply the event if nearly fully eclipsed, and the image-model residual is 0.968. Target attribution remains unresolved. Next test: Resolved eclipse photometry or multi-source pixel-response fitting of the 10.1-arcsec pair.
R488	403410574	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; Sixteen measured shallow but discrete raw eclipses in sectors 99 and 102 establish a repeated 2.11486-day signal. The early sector-39 cadence is too sparse for the local trapezoid timing criterion, although its stacked custom aperture shows the event. A capable Gaia neighbor 12.2 arcsec from the target lies only 10.9 arcsec from the deficit centroid; the residual is 0.896. TESS pixels do not settle attribution. Next test: Resolved eclipse photometry or multi-source pixel-response fitting of the 12.2-arcsec pair.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R493	57047535	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; Seven raw eclipses across S34/35/61/62/88/89, including consecutive cycles 38/39, 77/78/79 and 116/117, resolve the initially sparse timing. Several capable neighbors make this a crowded field. The deficit core favors the target, but the closest capable Gaia source is only about 1.1 TESS pixels away and the image-model residual is large; attribution remains unresolved. Next test: Resolved eclipse photometry or multi-source PRF localization of the crowded field.
R494	17926125	G 12-12.25 spectroscopic-orbit population	PROBABLE	true	Report; Six consecutive raw eclipses in S82/83 support the measured recurrence. The deficit core favors the target, but a capable G16.36 neighbor 15.7 arcsec away shares the central TESS pixel and could in principle produce the loss through an extremely deep intrinsic eclipse. Attribution remains unresolved at this resolution. Next test: Resolved eclipse photometry or multi-source PRF localization of the 15.7-arcsec pair.
R500	391340302	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; The S11 difference image centers on the target pixel, but a Gaia neighbor 6.1 arcsec away lies in that same TESS core and can produce the observed loss. TESS pixels cannot assign the eclipses uniquely. Next test: Resolved eclipse photometry of the 6.1-arcsec pair.
R506	325100028	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Five raw eclipses, including S38 events coincident with the historical TESScut stack, recur at one period. The S38 cutout shows a central deficit but its 3x3 depths of about 23 ppt disagree substantially with QLP raw fits of 164-183 ppt; dilution is plausible but not quantitatively demonstrated. A capable 12.3-arcsec Gaia neighbor remains viable at TESS resolution, so the host is unresolved. Next test: Resolved eclipse imaging and independent aperture/PRF characterization of the crowded field.
R508	87159646	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Three raw eclipses repeat in S59/86, but two capable Gaia neighbors at 3.0 and 4.9 arcsec share the central TESS pixel with the target. The S59 deficit cannot resolve the host. Next test: Resolved eclipse photometry of the close Gaia sources.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R515	376799355	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Raw eclipses repeat in S10/63/90 and the alternative 3x3 aperture also shows a deficit, diluted from roughly 88 ppt QLP to 36 ppt. The field is crowded: capable Gaia sources at 17.4 and 32.0 arcsec remain; the deficit centroid is only approximately target-centered and has no calibrated error. Attribution is unresolved. Next test: Resolved photometry across an eclipse or an independent high-resolution centroid.
R516	234413494	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Raw eclipses repeat in S7/46/72. A separate S44 TESScut extraction recovers four events, but the available cutout is truncated and its peak/residual panel is corrupted. Capable neighbors at 19.3 and 30.4 arcsec remain, so attribution is unresolved. Next test: Acquire a complete cutout or resolved eclipse photometry.
R529	311398753	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; S93/103 raw eclipses repeat and the S13 alternative extraction recovers two. A capable G15.36 source lies only 9.65 arcsec from the target and shares the bright pixel; further capable sources lie at 20.8 and 38.2 arcsec. The target-near deficit cannot distinguish the closest host at uncalibrated pixel precision. Gaia conjunction offset is −0.685 d at the observed epoch versus a 0.335 d linear orbital timing scale; a projection near Gaia2016 gives +0.012 d with only a 0.026 d photometric period allowance, so phase agreement remains descriptive. Next test: Resolved time-series photometry across an eclipse, plus refined Gaia-orbit comparison.
R542	150008480	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Eleven shallow repeated raw eclipses; eleven capable neighbors within 60 arcsec and a noisy deficit leave source attribution unresolved. The Gaia conjunction discrepancy is -0.043 phase, within the descriptive orbit timing uncertainty, and is not evidence of unique target ownership. Next test: Resolved eclipse photometry or multi-source PRF localization; distinguish alternating event kinds.
R543	370553309	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Five long eclipses repeat. A near-equal G=12.65 source at 16.62 arcsec overlaps the deficit core; target ownership is unresolved at this image resolution. Next test: Resolved eclipse photometry or multi-source PRF localization; distinguish alternating event kinds.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R545	1804347444	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Five repeated eclipses; a brighter G=11.54 source only 3.45 arcsec from target is unresolved. Pixel-aperture depth differs from QLP because of strong dilution; this is not a failed eclipse detection. Next test: Resolved eclipse photometry or multi-source PRF localization; distinguish alternating event kinds.
R548	371665307	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Repeated 2.49543-day dips are real. Depth parity is not decisive; the quoted 4.99086-day candidate orbital ephemeris uses even half-period parity only and retains a factor-two ambiguity. The shallow crowded deficit has twenty currently flux-capable neighbors and does not uniquely establish target ownership. Next test: Resolved eclipse photometry or multi-source PRF localization; distinguish alternating event kinds.
R558	62867537	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Real raw eclipses repeat in S15/56/83, with16 usable midpoints and12 consecutive pairs. Nine fresh S56 deficit centers average0.029px from target, so target is favored. A capable G16.03 neighbor8.79arcsec away shares the central TESS resolution element (separation0.42px); a calibrated two-source pixel response/localization uncertainty is unavailable. Deficit event scatter~0.10px and severe aperture dilution do not uniquely exclude that source. Attribution remains unresolved after completed check, so probable rather than confirmed. Next test: Resolved eclipse photometry or calibrated two-source PRF localization of8.8arcsec pair.
R560	272328952	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Deep repeated raw events. Compact difference core lies on target. Capable G12.87 neighbor at24.5arcsec lies1.17pixels away and overlaps the broad core; bright G9.29 neighbor at41.8arcsec dominates baseline but not deficit. Attribution favors target but does not uniquely exclude the closest capable source. Broad off-eclipse variability changes across sectors; S24 custom1x1/3x3/5x5 depths64.9/12.5/4.65ppt differ greatly from QLP144ppt, a major aperture/sector dilution limitation rather than depth agreement. Next test: Resolved photometry through an eclipse to establish ownership; additional timing for remaining period/shape limits.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R561	99569713	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Ten repeated raw eclipses. Capable G15.57 source at5arcsec is unresolved within the target deficit; ownership remains unresolved. A farther source33arcsec away overlaps the wings. Next test: Resolved photometry through an eclipse to establish ownership; additional timing for remaining period/shape limits.
R563	427095568	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Three deep S58 eclipses establish local recurrence; S16/85 small noise fits are explicitly rejected. Full quality-zero light curves lack any comparable deep eclipse anywhere (minima0.994/0.988 of median versus S58 0.746), independently of uncertain extrapolated phase. Including flagged finite cadences gives minima0.886/0.902 in S16/85, still no comparable25percent event; flagged windows are not clean nulls. Fresh S56 target-centered deficit also recovers eclipses. No capable Gaia source in60arcsec. Major cross-sector absence remains unexplained; local S58 timing is not a precise cross-year ephemeris. Next test: Resolved photometry through an eclipse to establish ownership; additional timing for remaining period/shape limits.
R566	24995506	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Six repeated raw eclipses at11.48585d. Compact target-centered deficit overlaps three capable sources19-23arcsec away, so ownership remains unresolved. Nearby known EB Gaia5940721527468980608 at42.94arcsec has G17.062 and period0.72596d: G-band pairwise maximum diluted loss is1.38percent, below observed4.8percent; required intrinsic loss exceeds unity. Its period differs from observed recurrence and its location is distinct from deficit core. This excludes that catalogued EB under approximate passband budget, not every close neighbor. Next test: Resolved photometry through an eclipse to establish ownership; additional timing for remaining period/shape limits.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R567	321533638	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Fresh availability query and actual additional S38/65/101 data increase timing sample to12 across all6 current QLP sectors. Consecutive within-sector recurrences establish7.40605d. Fourteen capable sources include one13.5arcsec away and near-equal source16.5arcsec away; crowded deficit does not uniquely establish target ownership. Next test: Resolved photometry through an eclipse to establish ownership; additional timing for remaining period/shape limits.
R569	2042514759	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Eight deep repeated eclipses. Brighter G11.76 source only2.9arcsec away is unresolved within deficit core; target attribution remains unresolved. Depth in custom aperture is diluted and is not intrinsic binary depth. Next test: Resolved photometry through an eclipse to establish ownership; additional timing for remaining period/shape limits.
R585	400671856	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Five sharp repeated selected eclipses establish10.56733d. Target-centered deficit favors target, but two capable sources at10.24/11.31arcsec overlap its PSF within0.54pixel, so ownership cannot be uniquely established. Two additional capable sources at50.7/60.0arcsec are distinct from the compact core. No secure secondary is claimed. Next test: Resolved eclipse photometry to establish ownership.
R589	298796659	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Six sharp repeated selected eclipses establish10.36267d. Target-centered deficit favors target, but three capable sources at16.2-21.6arcsec overlap the core within roughly onepixel. Two farther capable sources43.7/47.5arcsec are spatially distinct. Unresolved close-source ownership prevents confirmation; large variable raw baselines are kept separate from selected eclipses. Next test: Resolved eclipse photometry to establish ownership.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R590	306936928	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	true	Report; Repeated narrow primary and secondary dips are secure, but five flux-capable Gaia neighbors include a 10.59 arcsec G16.77 source requiring an almost total intrinsic eclipse. The compact blended deficit does not uniquely resolve target ownership; PROBABLE. Next test: Resolved eclipse photometry or multi-source PRF localization; distinguish alternating event kinds.
R603	374606776	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Representative sectors and simplified trapezoid/linear timing errors; shared TESS photons; approximate Gaia edge-on geometry and G-band neighbor flux budgets. Capable close neighbors prevent definitive source attribution. Next test: Higher-resolution time-series localization or a second field orientation resolving capable close sources.
R609	337004236	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Representative sectors and simplified trapezoid/linear timing errors; shared TESS photons; approximate Gaia edge-on geometry and G-band neighbor flux budgets. Capable close neighbors prevent definitive source attribution. Next test: Higher-resolution time-series localization or a second field orientation resolving capable close sources.
R619	380616368	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Six deep repeated raw eclipses and unequal secondary near0.5185 are real. A capable Gaia neighbor only13.06 arcsec away overlaps the TESS footprint; target ownership remains unresolved. Five farther capable neighbors were also inspected. Next test: Resolved eclipse photometry or multi-source PRF localization for target and13.06-arcsec neighbor.
R628	58825156	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Twelve repeated raw eclipses and shallower secondary near0.50704 are real. Capable Gaia2776848815318667392 at12.4345arcsec overlaps the TESS footprint; the0.227pixel deficit offset and broad fractional-loss pattern do not securely separate ownership. Attribution remains unresolved. Next test: Resolved eclipse photometry or multi-source PRF localization for target and12.43-arcsec neighbor.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R635	398837588	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Close capable neighbor leaves attribution unresolved. Next test: Resolved eclipse photometry or multi-source PRF attribution.
R636	94220952	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Close capable neighbor leaves attribution unresolved. Next test: Resolved eclipse photometry or multi-source PRF attribution.
R637	84122354	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Close capable neighbor leaves attribution unresolved. Next test: Resolved eclipse photometry or multi-source PRF attribution.
R638	340860427	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Close capable neighbor leaves attribution unresolved. Next test: Resolved eclipse photometry or multi-source PRF attribution.
R642	352113274	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Five repeated narrow flat-bottomed eclipses and shallower secondary near0.5003. Capable Gaia533640894735716352 at8.83arcsec is inside the unresolved TESS footprint; deficit offset0.248pixel and fractional-loss morphology do not securely separate source ownership. Attribution remains unresolved. Next test: Resolved eclipse photometry or multi-source PRF localization for target and listed close capable neighbor.
R648	441356022	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Five repeated narrow eclipses and a shallower secondary near0.50014. Sector1 broad excursions remain visible in phase review but representative locally normalized primary is consistent. Capable Gaia6793606134390553856 at11.93arcsec overlaps TESS footprint; small centroid offset cannot uniquely establish ownership. Attribution remains unresolved. Next test: Resolved eclipse photometry or multi-source PRF localization for target and listed close capable neighbor.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R657	59955223	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Close capable neighbor leaves attribution unresolved. Next test: Resolved eclipse photometry or multi-source PRF attribution.
R659	404555712	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Close capable neighbor leaves attribution unresolved. Next test: Resolved eclipse photometry or multi-source PRF attribution.
R660	133392449	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Close capable neighbor leaves attribution unresolved. Next test: Resolved eclipse photometry or multi-source PRF attribution.
R663	292916455	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Photometric recurrence1.55009d versus candidate orbital3.10018d remains unresolved; resolve alternating families or secondary. Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Close capable neighbor leaves attribution unresolved. Next test: Resolved eclipse photometry and alternating-family or secondary timing to decide orbital P versus2P.
R665	448983372	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Close capable neighbor leaves attribution unresolved. Next test: Resolved eclipse photometry or multi-source PRF attribution.
R670	396671669	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. P/2 recurrence untested in missing or flagged opposite windows. Next test: Observe clean opposite half-phase windows to test P/2 recurrence.
R673	458606209	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Close capable neighbor leaves attribution unresolved. Next test: Resolved eclipse photometry or multi-source PRF attribution.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R675	214901783	G 12.5-12.75 spectroscopic-orbit population	PROBABLE	true	Report; Approximate Gaia geometry and G-band blend screen; timing errors exclude uncalibrated model systematics. Close capable neighbor leaves attribution unresolved. Next test: Resolved eclipse photometry or multi-source PRF attribution.
R678	54054420	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	true	Report; A G17.60 source only 7.9 arcsec from target is flux-capable and 6.4 arcsec from cached deficit centroid; s24 PSF residual is 0.65 without calibrated centroid covariance. Nearly equal alternating dips leave the photometric half-period alias unresolved by photometry alone. TIC radius 5.07 solar radii gives longer central eclipse than observed; grazing geometry or stellar radius errors could reconcile it.
R685	109560492	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	true	Report; Prior limitations: Initial catalogue services were partly unavailable. The nearby brighter star and its aliases require explicit novelty/attribution checks.; decision: Previously accepted physical review; scoped catalogue clearance, root accepts new claim; crowded attribution limitation remains.
R686	167197264	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	true	Report; Prior limitations: G15.32/G15.35 neighbors at 22-24 arcsec near maximum blended-loss bound; s76 centroid favors target but uncertainty uncalibrated; no calibrated photometric period error; decision: Root accepts physical review and scoped catalogue clearance; crowded attribution and no calibrated photometric period error remain.
R691	342060048	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	true	Report; Decision: Repeated measured eclipses pass bounded review, with unresolved close-source attribution and Gaia-clock differences; accept probable new claim within completed catalogue scope. Four narrow deep events repeat, including consecutive sector 40 cycles. The 17.52-arcsec capable Gaia neighbor lies less than one TESS pixel away; target-centered difference imaging favors the target but does not resolve the blend. Gaia period and conjunction are modestly discrepant at formal error scale.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R692	184398115	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	true	Report; Decision: Repeated deep eclipses and target-centered deficit; material disagreement with Gaia orbit remains. Accept probable new claim, not a confirmed association with the spectroscopic pair. Four very deep repeated eclipses include consecutive sector 97 cycles, and the deficit is target-centered with no capable Gaia neighbor in the 60-arcsec cone. The measured eclipse period and phase are materially inconsistent with the formal Gaia SB2 orbit; a distinct eclipsing binary in a multiple system or a problematic Gaia orbit remains possible.
R706	121277143	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	true	Report; Decision: Root accepts repeated measured eclipses at probable tier; 34.55-arcsec neighbor remains nominally capable under optimistic Gaia-G flux proxy and is not securely excluded. Exact ASAS-SN ROT: counterpart is not a prior EB classification; completed scoped checks support a new EB claim. A custom TESScut aperture of the same TESS photons reproduces the deficit, but it is not an independent observing occasion. 1 Gaia DR3 neighbor(s) within 60 arcsec pass an optimistic G-band/full-throughput capability screen; the target-centered difference image does not conclusively resolve source ownership at TESS pixel scale. Sector 40 lacks egress and sector 53 has broad baseline modulation; both are retained as detections but excluded from center timing.
R725	219345200	Gaia spectroscopic-binary cached population; mixed-harvest prioritized dip reviews	PROBABLE	true	Report; Repeated signal and compatible Gaia geometry; close near-capable neighbor leaves ownership uncertain. Real repeated eclipses and compatible Gaia geometry; close capable or near-capable neighbor prevents firm source ownership. Period is approximate from representative raw centers, not a calibrated ephemeris error. Next check: Obtain higher-resolution eclipse photometry or a multi-pixel PSF fit across separate events.
R726	450652458	Gaia spectroscopic-binary cached population; mixed-harvest prioritized dip reviews	PROBABLE	true	Report; Repeated broad flat-core troughs; crowded aperture leaves ownership uncertain; S65 unusable as null. Crowding, sloping baselines and uncertain broad-event centers limit source ownership and period precision. No formal error is inferred from provisional trapezoid fits. Next check: Obtain a clean fully covered next trough with spatial difference imaging and a second extraction.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R728	458324895	Gaia spectroscopic-binary cached population; mixed-harvest prioritized dip reviews	PROBABLE	true	Report; Four broad recurrent events; aliases unresolved, Gaia-favored clock is not uniquely measured. Real repeated broad eclipses have unresolved integer aliases; no measured orbital period is adopted. S39/S66 incomplete baselines and same-photon alternative extraction limit depth and source precision. Next check: Observe a deep eclipse at a discriminating intermediate cycle and repeat target-centered pixels.
R729	236223193	Gaia spectroscopic-binary cached population; mixed-harvest prioritized dip reviews	PROBABLE	true	Report; Six target-centered events repeat; 35.239 d photometric clock disagrees with Gaia SB2; nearby VSX EB has different clock and insufficient depth. The target owns the recurring photometric dips, but the Gaia SB2 orbital association and source-specific conjunction geometry are unresolved. Local fit covariances understate long-baseline timing scatter; the approximate period has no defensible formal error yet. Next check: Obtain source-specific RV/conjunction timing and independent photometry, then reconcile the 35.239 d eclipse clock with the Gaia 35.3747 d SB2 orbit.
R730	28059397	Gaia spectroscopic-binary cached population; mixed-harvest prioritized dip reviews	PROBABLE	true	Report; Repeated events including partial third occasion; eclipse kind and period conditional; no precise ephemeris. The partial third occasion supports recurrence but its missing center leaves kind and orbital period conditional; no precise period or epoch is adopted. QLP KSPSAP is absent for S56/S83; fresh TESSCut provides a spatial alternative extraction of the same photons, not another observing occasion. Next check: Measure a complete next event in an independent pipeline and compare separate eclipse-kind recurrence with source-specific Gaia conjunction timing.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R731	19385783	Gaia spectroscopic-binary cached population; mixed-harvest prioritized dip reviews	PROBABLE	true	Report; Recurrent 2.065942 d dip cadence; ownership inconclusive and Gaia association unresolved; cadence need not be orbital period. The sector-34 TESSCut difference image is inconclusive for ownership because an unrelated excess biases its centroid. Gaia SB1 P35.0856 d is not the measured dip recurrence and its conjunction is 0.231 phase away at S34; the orbital relationship is unresolved. The physical orbit might be an integer multiple of the observed dip cadence; the reported clock is a photometric recurrence, with no defensible period error. Next check: Obtain a clean difference image or ground-based resolved photometry and test the Gaia orbit versus a separate short-period binary.
R732	51321183	Gaia spectroscopic-binary cached population; mixed-harvest prioritized dip reviews	PROBABLE	true	Report; Recurrent blended-aperture signal; capable G13.015 neighbor 0.36 pixel away; target ownership and Gaia association unresolved. Attribute signal to aperture, not established target EB. The photometric clock is a rounded blended-aperture recurrence, not an established orbit of the Gaia target. Fresh WCS pixels have valid target flux but a capable G13.015 neighbor only 0.36 pixel from the target; local drift moves the difference centroid off both positions. Source ownership is unresolved. Gaia SB1 P38.46865 d is incompatible with the observed 16.89-d clock; no source-specific conjunction tie is claimed. Local formal timing errors omit systematics, so no period error is supplied. Next check: Obtain resolved target and neighbor photometry at another predicted sharp eclipse, then establish source ownership and orbital association.

Known prior eclipsing systems (physical recoveries) (36)

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R043	257800191	G 10-11.5 spectroscopic-orbit population	CONFIRMED	false	Report; Refine diluted two-star dimensions and small-e orbital solution with joint light-curve/RV modelling.
R177	265350422	G 11.5-12 spectroscopic-orbit population	CONFIRMED	false	Report; Three G-band-capable neighbours at 32.85, 42.38 and 51.78 arcsec lie away from the compact deficit on the target. The target-centered dip grows rather than vanishes in the small aperture. No calibrated centroid confidence ellipse is available. Next test: Optional precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R179	253126207	G 11.5-12 spectroscopic-orbit population	PROBABLE	false	Report; Attribution remains unresolved: seven capable neighbours include three within one TESS pixel (15.16, 17.15 and 19.43 arcsec); the historical PSF fractional residual is 0.657. A known Gaia eclipsing binary at 24.77 arcsec with P=0.312563 d is a different source, and its short period does not identify the 17.44 d signal. Real repeating eclipses pass the other checks; source localization is the failed scientific check. Resolved eclipse imaging or defensible multi-source PRF modeling is required. Next test: Resolved eclipse photometry / multi-source PRF localization
R274	67300566	G 12-12.5 spectroscopic-orbit population	PROBABLE	false	Report; Real repeated raw dips; capable G12.84 neighbor18.26arcsec away lies0.84pixel from deficit versus0.05pixel target offset. Uncalibrated close-source localization prevents definitive attribution; completed review retains probable. Next test: Resolved eclipse photometry or calibrated multisource PRF localization.
R276	387275908	G 12-12.5 spectroscopic-orbit population	CONFIRMED	false	Report; Capable neighbors27.34 and36.11arcsec are spatially separated from deficit0.10pixel from target. High total-light PSF residual0.823 reflects crowding and is not used alone as attribution evidence. Gaia phase offset-0.059 has covariance uncertainty0.255d; duration can be noncentral. Next test: Optional precision characterization; no missing bounded admission check.
R358	233462817	G 11.5-12 spectroscopic-orbit population	CONFIRMED	false	Report; Shallow repeated raw eclipses in sectors49/77; no capable listed neighbor. Deficit centroid0.53pixel offset and noisy difference image limit precise localization. Gaia conjunction offset0.041phase is interpreted with full covariance (0.245d), short observed duration allowing noncentral geometry; no arbitrary sigma veto. Next test: Optional precision characterization
R390	399868192	G 11.5-12 spectroscopic-orbit population	CONFIRMED	false	Report; Repeated flat-bottom eclipses across sectors 5/71/98. Sector 98 cycle 398 lacks ingress and has singular center covariance, so it is excluded from timing. Other consecutive events anchor the recurrence. Capable neighbors are at least 1.86 pixels from deficit versus target 0.057 pixel. Next test: Optional resolved photometry or spectroscopy for precision characterization

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R407	348302938	G 11.5-12 spectroscopic-orbit population	CONFIRMED	false	Report; Fourteen shallow narrow V-shaped eclipses repeat in S61/88; individual depths scatter at several ppt noise. Deficit0.270pixel from target versus nearest capable neighbor1.107pixels. Consecutive cycles constrain recurrence; parity depths are consistent within noise. Nearly equal primary/secondary events could leave a doubled orbital alias photometrically ambiguous; Gaia supports the retained branch. Next test: Optional precision characterization
R434	152147232	G 12-12.25 spectroscopic-orbit population	PROBABLE	false	Report; Fifteen raw repeats include many within-sector pairs. A capable 2.7 arcsec Gaia neighbor shares the TESS core and cannot be resolved. Next test: Resolved eclipse photometry or multi-source PSF localization of the 2.7 arcsec pair.
R474	99493790	G 12-12.25 spectroscopic-orbit population	CONFIRMED	false	Report; The target-centered deficit and absence of a brightness-capable Gaia neighbor favor attribution; the image-model residual is high. Next test: Optional precision characterization
R486	457272901	G 12-12.25 spectroscopic-orbit population	CONFIRMED	false	Report; The added sector-65 raw events anchor the 4.649217-day timing baseline; later sector-99/102 troughs repeat. The difference centroid is 2.1 arcsec from the target and the closest capable neighbor is about 17 arcsec from that centroid. Twenty-seven capable Gaia sources lie within 60 arcsec; the crowded image and uncalibrated residual limit precision, but the spatial deficit favors the target. Next test: Optional resolved photometry and spectroscopy for precision binary parameters; no missing bounded admission check.
R499	428703516	G 12-12.25 spectroscopic-orbit population	PROBABLE	false	Report; Four shallow raw eclipses in S19/59/86, including consecutive S86 cycles, support recurrence. The deficit favors the target over the capable 21.5-arcsec neighbor, but a capable Gaia source 11.2 arcsec away is inside one TESS pixel and could in principle produce the 4-ppt loss with a deep intrinsic eclipse. Attribution remains unresolved at this resolution. Next test: Resolved eclipse photometry or multi-source PRF localization of the 11.2-arcsec pair.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R562	386161311	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	false	Report; Repeated eclipse-shaped raw dips with depth variation. Nineteen capable sources include sources9.4/13.4/14arcsec away, unresolved around the weak target-centered deficit. Target ownership remains unresolved; conjunction offset0.0505phase is comparable to propagated timing allowance, rather than a clean veto. Next test: Resolved photometry through an eclipse to establish ownership; additional timing for remaining period/shape limits.
R610	370138999	G 12.5-12.75 spectroscopic-orbit population	CONFIRMED	false	Report; Shallow repeated narrow raw eclipses; compact deficit near target. Three capable neighbors at31-59 arcsec are spatially separated from the deficit core. No secure secondary. Next test: Optional resolved photometry and spectroscopy to refine physical parameters.
R681	24668377	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	false	Report; VSX query was unavailable; the exact IJspeert EB catalogue row already makes the object a known recovery.
R687	155346713	2026-11-03/04 selected spectroscopic-orbit follow-up	PLAUSIBLE	false	Report; Prior limitations: One complete event per kind; sector-7 A ingress truncated; sector-61 B central forecast in Q0 core gap; Gaia period uncertainty projects to days across long baselines; decision: Root accepts plausible review; exact IJspeert prior EB identification; no claim or physical membership.
R689	159060569	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	false	Report; Decision: Accept completed physical review at stated tier; exact published EB identity excludes a new-system claim. Six raw SAP eclipses are visibly repeated across four sectors, though two gap-shaped fits have unidentifiable center covariance and are excluded from the timing fit. Sector 98 is shallower, consistent with sector-dependent aperture dilution.
R690	387268955	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	false	Report; Decision: Accept completed physical review at stated tier; exact published EB identity excludes a new-system claim. Fourteen usable broad same-kind eclipses across many sectors; sector 60 is shallower at an edge window. Two additional fitted windows are excluded for incomplete/unidentifiable timing.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R693	64701575	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	false	Report; Decision: Accept completed physical review at stated tier; exact published EB identity excludes a new-system claim. Four clean rounded eclipses across sectors 33/34/87/88; sector 7 contains a partial predicted window. The timing and target-centered difference image support the Gaia-orbit eclipse identification.
R694	284613090	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Accept completed physical review at stated tier; exact published EB identity excludes a new-system claim. Four deep same-kind eclipses repeat, with two consecutive sector 33 cycles. Deep opposite half-phase events are also measured; unequal depths support the full 23.1095-day photometric branch. The target-centered difference image has no capable Gaia neighbor, but Gaia SB2 period and conjunction differ materially at formal precision.
R695	128720791	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Accept completed physical review at stated tier; exact published EB identity excludes a new-system claim. Three clean narrow same-kind dips in sectors 16/56/83; the earlier sector 56 predicted cycle lies at the sector boundary and is not a clean null. Explicit 15–30 d integer-cycle enumeration found a 16.2798 d alias, but three additional clean predicted windows have near-zero loss and refute that equal-depth branch. The Gaia-near 21.7064 d branch has no contradictory clean window. The signal is reproduced by separate TESScut extraction and a target-centered difference image; Gaia phase/period differences remain a limitation.
R697	56912737	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	false	Report; Decision: Accept completed physical review at stated tier; exact published EB identity excludes a new-system claim. Three compact dips in sectors 21 and 47 give two adjacent sector 47 cycles; measuring the individual centers resolves the apparent phase displacement of the earlier sector 47 window. Timing agrees with the Gaia period within stated errors and the deficit is target-centered.
R698	349978696	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	false	Report; Decision: Root accepts completed physical tier with all stated limitations; exact published IJspeert EB identity excludes a new-system claim. A custom TESScut aperture of the same TESS photons reproduces the deficit, but it is not an independent observing occasion.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R699	230130556	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Root accepts completed physical tier with all stated limitations; exact published IJspeert EB identity excludes a new-system claim. A custom TESScut aperture of the same TESS photons reproduces the deficit, but it is not an independent observing occasion. 1 Gaia DR3 neighbor(s) within 60 arcsec pass an optimistic G-band/full-throughput capability screen; the target-centered difference image does not conclusively resolve source ownership at TESS pixel scale.
R700	174303697	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Root accepts completed physical tier with all stated limitations; exact published IJspeert EB identity excludes a new-system claim. A custom TESScut aperture of the same TESS photons reproduces the deficit, but it is not an independent observing occasion. The measured eclipse is 0.942 d from the closest Gaia conjunction, 6.4 times its propagated Gaia linear error; the published Gaia orbit and eclipse clock may refer to different or inaccurately modeled motion. Sector 16 has a flagged preceding core and a clean later eclipse whose egress is beyond the sector edge; neither supplies a full center fit.
R701	102305337	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	false	Report; Decision: Root accepts completed physical tier with all stated limitations; exact published IJspeert EB identity excludes a new-system claim. A custom TESScut aperture of the same TESS photons reproduces the deficit, but it is not an independent observing occasion. Early and late depths differ by roughly 4 ppt, consistent with extraction or aperture variation; timing remains repeated.
R702	343452418	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Root accepts completed physical tier with all stated limitations; exact published IJspeert EB identity excludes a new-system claim. A custom TESScut aperture of the same TESS photons reproduces the deficit, but it is not an independent observing occasion. 1 Gaia DR3 neighbor(s) within 60 arcsec pass an optimistic G-band/full-throughput capability screen; the target-centered difference image does not conclusively resolve source ownership at TESS pixel scale.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R703	207780183	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Root accepts completed physical tier with all stated limitations; exact published IJspeert EB identity excludes a new-system claim. A custom TESScut aperture of the same TESS photons reproduces the deficit, but it is not an independent observing occasion. 2 Gaia DR3 neighbor(s) within 60 arcsec pass an optimistic G-band/full-throughput capability screen; the target-centered difference image does not conclusively resolve source ownership at TESS pixel scale. Several opposite-half-phase windows have broad, shallower deficits displaced about 0.20 d from exact half phase; these are a possible eccentric-orbit secondary eclipse, not same-kind timing points.
R704	36721228	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	false	Report; Decision: Root accepts completed physical tier with all stated limitations; exact published IJspeert EB identity excludes a new-system claim. A custom TESScut aperture of the same TESS photons reproduces the deficit, but it is not an independent observing occasion.
R705	431996459	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Root accepts completed physical tier with all stated limitations; exact published IJspeert EB identity excludes a new-system claim. A custom TESScut aperture of the same TESS photons reproduces the deficit, but it is not an independent observing occasion. 1 Gaia DR3 neighbor(s) within 60 arcsec pass an optimistic G-band/full-throughput capability screen; the target-centered difference image does not conclusively resolve source ownership at TESS pixel scale.
R707	431589019	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Root accepts completed physical tier with all stated limitations; exact published IJspeert EB identity excludes a new-system claim. A custom TESScut aperture of the same TESS photons reproduces the deficit, but it is not an independent observing occasion. 2 Gaia DR3 neighbor(s) within 60 arcsec pass an optimistic G-band/full-throughput capability screen; the target-centered difference image does not conclusively resolve source ownership at TESS pixel scale. The representative cutout has only four core frames and substantial aperture-dependent dilution amid two bright capable neighbors; target-like difference peak is insufficient for unique attribution.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R708	257992903	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Known IJspeert exact identity; real repeats accepted probable with capable neighbors unresolved. Two Gaia neighbors at 47.26 and 54.64 arcsec remain capable under an optimistic G-band/full-capture screen; the target-like pixel deficit does not calibrate source ownership.
R709	70561926	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	false	Report; Decision: Confirmed recurrent eclipses; four irregular event intervals constrain clock. Root resolves VSX EA association using 0.30 arcsec position, matching clock and no alternative Gaia source within 2 arcsec; known recovery. The Gaia period differs from the eclipse fit by 0.04084 d (1.67 Gaia quoted standard errors), and the closest conjunction is 1.39 d away with 1.48 d source-specific linear covariance uncertainty. The exact VSX counterpart is a known EB.
R710	380395732	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Known IJspeert exact identity; real repeats accepted probable with capable neighbors unresolved. Two Gaia neighbors at 39.52 and 52.68 arcsec remain capable under an optimistic G-band/full-capture screen. The 5×5 pixel loss is diluted relative to the 1×1; source ownership needs higher spatial resolution.
R713	117930511	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Root accepts strongly supported positional prior VSX EA association, despite discrepant prior clock for117930511; no new-system claim. Root required probable because the target-like broad deficit does not exclude the capable overlapping neighbor. Three same-kind raw events include consecutive sector-57 eclipses. The difference image favors the target; the one marginal G-band capable neighbor lies 30.77 arcsec away and would require a roughly 94% intrinsic eclipse under optimistic throughput. The broad TESS PSF overlaps that neighbor at roughly1.5 pixels, so target-like centroid and morphology do not decisively exclude it. Attribution remains unresolved. The measured eclipse clock differs from the prior VSX clock; this does not make the star a new system. TESScut is a separate reduction of the same photons. Gaia G-band flux ratios and full throughput are optimistic neighbor capability proxies, not TESS PRF likelihoods. No exhaustive all-event fitting, precision stellar characterization, or Monte Carlo was performed.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R722	198358825	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	false	Report; Decision: Root accepts strongly supported positional prior VSX EA association, despite discrepant prior clock for117930511; no new-system claim. Five usable representative centers over81 cycles, with numerous intervening raw coverage tests, establish a stable same-kind clock. The gapped sector57 event is detected but excluded from center fitting. The deficit is on target and no capable Gaia neighbor appears in the bounded cone. The prior KELT/VSX EA clock agrees closely, making this a known recovery. TESScut is a separate reduction of the same photons. Gaia G-band flux ratios and full throughput are optimistic neighbor capability proxies, not TESS PRF likelihoods. No exhaustive all-event fitting, precision stellar characterization, or Monte Carlo was performed.

Known recoveries and controls (12)

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R102	432482170	G 10-11.5 spectroscopic-orbit population	CONFIRMED	false	Report; See target report for limitations.
R163	407991939	G 10-11.5 spectroscopic-orbit population	CONFIRMED	false	Report; Joint RV/eclipse modeling and longer-baseline eclipse timing; resolve remaining alias and geometry uncertainty.
R165	407220461	G 10-11.5 spectroscopic-orbit population	CONFIRMED	false	Report; Higher-resolution eclipse photometry or calibrated PRF fitting to resolve the close capable neighbour; joint RV/eclipse timing fit.
C01	195865529	fainter SB1 period-fold population	CONFIRMED	false	Report; Known ASAS-SN EA at 0.16 arcsec (P 3.8310 d); ATLAS DBH P/2. Recovery excluded from new count.
C05	70665829	fainter SB1 period-fold population	CONFIRMED	false	Report; Already objects.csv A4; blind control independently recovered. Archive Gaia-VEB/varisum/SB9 controls failed on later mirror timeout, predecessor empty cones only. No addition.
C18	300903537	fainter SB1 period-fold population	CONFIRMED	false	Report; See target report for deciding limitation and next check.
C58	183994341	fainter SB1 period-fold population	CONFIRMED	false	Report; Resolve ASAS-SN period alias and obtain RV/orbital posterior or a joint timing/geometry analysis to settle the Gaia association.
P024	155303130	G 11.5-12; Gaia spectroscopic orbit 20-60 d	CONFIRMED	false	Report; Pilot characterization; period errors are scatter-only. No exhaustive literature priority claim. Next test: Optional precision characterization; no missing bounded admission check

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R257	377389892	G 12-12.5 spectroscopic-orbit population	CONFIRMED	false	Report; Known VSX EA within0.02arcsec; excluded from new-system count. Catalogue1.147d is a quarter-period alias: raw fold shows deeper primary each4.588d and shallower secondary at phase0.5. Fold at1.147d mixes out-of-eclipse and unequal eclipse points. Primary epoch identifies deeper eclipse; physical Gaia orbit association retained. Next test: Optional precision characterization
R448	238406817	G 12-12.25 spectroscopic-orbit population	CONFIRMED	false	Report; Alternating ~275 and ~116 ppt eclipses identify the 4.7752 d orbital period; the historical 2.38762 d SAP branch is the half-period. The deeper parity defines the epoch. Next test: Optional precision characterization
R519	66072136	G 12.25-12.5 spectroscopic-orbit population	CONFIRMED	false	Report; S16/76/84 raw repeats and the alternative S76 aperture support a real target-centered eclipse. No capable Gaia neighbor within 60 arcsec. Central catalogue reconciliation identifies the close VSX EA V0693 Lac counterpart as the same object, so this is a known eclipsing-binary recovery, not a new-system addition. VSX lists 1.08916 d versus the measured same-kind 4.58920 d; the disagreement does not negate the measured raw eclipses. Next test: Resolve the discordant VSX period for catalogue interpretation; no new-system count.
R573	295771108	G 12.25-12.5 spectroscopic-orbit population	PROBABLE	false	Report; Known ASAS-SN EA recovery, not a new addition. Deep eclipses and shallower phase-0.50052 events repeat, but the capable G=13.79 neighbor at 13.45 arcsec is unresolved: target and neighbor pixels lose 34.85 and 32.18 e-/s. Gaia predicts the other conjunction at phase 0.60848 +/-0.03919 (linear covariance allowance) versus measured 0.50052. This descriptive tension and blend leave geometry/ownership less secure. Next test: Resolved eclipse photometry or multi-source PRF analysis of the 13.45-arcsec blend, and a joint orbit/eclipse fit.

Pending catalogue novelty (8)

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R714	25320385	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Root accepts completed physical tier, but unavailable catalogue scopes prevent any new-system claim. Repeated V-shaped eclipse-like dips are secure, but the raw clock is a harmonic of the screening clock: dips occur near every 3.5906 d and a 7.18125 d branch also fits. Opposite dips have comparable depths with changing widths; primary/secondary identity and a unique orbital factor of two are unestablished. The crowded difference image is offset 0.83 pixel and dilution changes strongly with aperture. The event source cannot be uniquely assigned to this Gaia star, and it cannot be declared a definite Gaia by-catch. Generic orbital ephemeris fields are blank. TESScut is a separate reduction of the same photons. Gaia G-band flux ratios and full throughput are optimistic neighbor capability proxies, not TESS PRF likelihoods. No exhaustive all-event fitting, precision stellar characterization, or Monte Carlo was performed.
R715	122557124	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	false	Report; Decision: Root accepts completed physical tier, but unavailable catalogue scopes prevent any new-system claim. Three full raw eclipses across 110 cycles establish a consistent clock; the other integer counts tested fail actual timing or coverage. Deep, flat-bottomed stellar eclipses survive the separate extraction, with a target-centered deficit and no capable Gaia neighbor in the bounded cone. Gaia conjunction is compatible. TESScut is a separate reduction of the same photons. Gaia G-band flux ratios and full throughput are optimistic neighbor capability proxies, not TESS PRF likelihoods. No exhaustive all-event fitting, precision stellar characterization, or Monte Carlo was performed.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R716	260797021	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Root accepts completed physical tier, but unavailable catalogue scopes prevent any new-system claim. Deep flat-bottomed eclipses repeat with consecutive anchors. A target-centered difference image and separate extraction support an on-target eclipsing source, but Gaia has P=20.3196973 d versus measured about20.29960 d. Both period and conjunction disagree by about five formal Gaia standard deviations. Published-orbit errors or different motion remain possible; this is not a geometry-consistent confirmation of the selecting Gaia companion. The ingress-gapped sector67 fit is excluded from adopted timing. TESScut is a separate reduction of the same photons. Gaia G-band flux ratios and full throughput are optimistic neighbor capability proxies, not TESS PRF likelihoods. No exhaustive all-event fitting, precision stellar characterization, or Monte Carlo was performed.
R717	330664712	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Root accepts completed physical tier, but unavailable catalogue scopes prevent any new-system claim. Repeated short eclipses include consecutive sector76 events and reject tested integer aliases. The independent sector16 extraction has just four core frames, severe aperture dilution, and eight capable neighbors under the optimistic test, including sources near22 arcsec. Its centroid is target-like but does not uniquely resolve ownership at this sampling. Attribution remains unresolved. TESScut is a separate reduction of the same photons. Gaia G-band flux ratios and full throughput are optimistic neighbor capability proxies, not TESS PRF likelihoods. No exhaustive all-event fitting, precision stellar characterization, or Monte Carlo was performed.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R718	436013770	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Root accepts completed physical tier, but unavailable catalogue scopes prevent any new-system claim. Root required probable because the target-like broad deficit does not exclude the capable overlapping neighbor. Two full raw eclipse centers span75 cycles. A third partial raw eclipse in sector98 has actual deep samples nearBTJD4010.12719; it is not used as a precise fitted center. This third occasion rejects the tested nearby integer counts by3.35–10.23 d, while the selected branch agrees to0.039 d. The difference favors the target and 1x1/3x3 losses33.20/33.31 ppt are stable, but the broad TESS PSF overlaps the sole capable neighbor31.7 arcsec away at roughly1.5 pixels. The image does not decisively exclude that source; attribution remains unresolved even though Gaia geometry is compatible. TESScut is a separate reduction of the same photons. Gaia G-band flux ratios and full throughput are optimistic neighbor capability proxies, not TESS PRF likelihoods. No exhaustive all-event fitting, precision stellar characterization, or Monte Carlo was performed.
R720	449012335	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Root accepts completed physical tier, but unavailable catalogue scopes prevent any new-system claim. The screening window clipped a roughly1.6 d flat-bottomed eclipse. New local raw fits require coverage at both contacts; sector11 and101 provide centers, while gapped fits are retained but excluded. A separate TESScut sector100 contact fit gives BTJD4085.18444, agreeing with the raw period prediction to0.00561 d and rejecting nearby integer counts. This reduction shares TESS photons but sector100 is a different observing occasion from sectors11/101. A neighbor10.43 arcsec away and others remain capable, so ownership is unresolved. Long-baseline orbital modulation and aperture drift require local baselines. TESScut is a separate reduction of the same photons. Gaia G-band flux ratios and full throughput are optimistic neighbor capability proxies, not TESS PRF likelihoods. No exhaustive all-event fitting, precision stellar characterization, or Monte Carlo was performed.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R723	241479387	2026-11-03/04 selected spectroscopic-orbit follow-up	PROBABLE	false	Report; Decision: Root accepts recurrent eclipses at probable tier with unresolved8.59arcsec neighbor and disclosed mild Gaia tension. Missing catalogue scopes prevent a new-system claim. The event is real and repeats, but a capable 8.59 arcsec Gaia neighbor prevents definitive target ownership. Aperture dilution varies strongly, and TIC 241479387 has mild roughly 2 sigma Gaia period/conjunction tension. Timing fit errors are conditional and exclude extraction/model systematics. No exhaustive all-sector characterization or Monte Carlo was performed. Novelty remains pending because scoped catalogue coverage is incomplete.
R724	421331604	2026-11-03/04 selected spectroscopic-orbit follow-up	CONFIRMED	false	Report; Decision: Root accepts confirmed repeated deep eclipses and separate recurrent shallow conjunction; alternate reduction, bounded source census and Gaia orbit support physical interpretation. Missing catalogue scopes prevent a new-system claim. The sector 26 egress-only model is excluded from adopted timing and depth. Depths depend on aperture dilution. Pixel centroids and Gaia G-band capability proxies are descriptive rather than calibrated TESS PRF likelihoods. Timing fit errors exclude model/pipeline systematics. No exhaustive all-sector characterization or Monte Carlo was performed. Physical confirmation is separate from pending catalogue novelty.

Plausible signals (28)

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R064	1989681813	G 10-11.5 spectroscopic-orbit population	PLAUSIBLE	false	Report; Obtain spatially resolved time-series photometry across a later narrow deficit and both Gaia conjunctions; use empirical calibrated TESS PRFs in a second sector to distinguish astrophysical deficit from nonstationary spatial contamination.
C16	319185391	fainter SB1 period-fold population	PLAUSIBLE	false	Report; Fold-only ~1 ppt dip, individual events ~1 sigma, dip not pixel-tested, no second pipeline. Report prose says probable-level but batch summary says PLAUSIBLE at best; adopt batch summary outside count. Gaia 434-day astrometric orbit suggests triple.
C49	71866523	fainter SB1 period-fold population	PLAUSIBLE	false	Report; Require a clean repeated negative event before EB classification.
C54	92452368	fainter SB1 period-fold population	PLAUSIBLE	false	Report; Verify a clean recurrent event in independently reduced data.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
C55	94300366	fainter SB1 period-fold population	PLAUSIBLE	false	Report; Obtain detrending resistant multiseason photometry to establish stable discrete repeats.
C59	166781582	fainter SB1 period-fold population	PLAUSIBLE	false	Report; Obtain or inspect another clean sector and require repeated discrete ingress/egress before fitting a binary period.
C60	230270717	fainter SB1 period-fold population	PLAUSIBLE	false	Report; Obtain or inspect another clean sector and require repeated discrete ingress/egress before fitting a binary period.
C61	462845564	fainter SB1 period-fold population	PLAUSIBLE	false	Report; Obtain or inspect another clean sector and require repeated discrete ingress/egress before fitting a binary period.
C63	350661747	fainter SB1 period-fold population	PLAUSIBLE	false	Report; Obtain or inspect another clean sector and require repeated discrete ingress/egress before fitting a binary period.
C64	424565669	fainter SB1 period-fold population	PLAUSIBLE	false	Report; Obtain or inspect another clean sector and require repeated discrete ingress/egress before fitting a binary period.
C68	275022619	fainter SB1 period-fold population	PLAUSIBLE	false	Report; Obtain a well-covered predicted candidate event in a distinct reduction and establish a second coherent detached eclipse before claiming an eclipse period or host.
C72	257575960	fainter SB1 period-fold population	PLAUSIBLE	false	Report; Obtain a well-covered predicted candidate event in a distinct reduction and establish a second coherent detached eclipse before claiming an eclipse period or host.
P020	8853239	G 11-11.5; Gaia spectroscopic orbit 20-60 d	PLAUSIBLE	false	Report; Single secure event; Gaia-only period; KELT null is coverage-limited and not a refutation. Tentative B is not an ephemeris. Next test: Measure a second A eclipse to test the cycle period; a B detection alone would still need repeat timing.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R249	314047448	G 12-12.5 spectroscopic-orbit population	PLAUSIBLE	false	Report; Historical sector24 deficit has large PSF residual and displaced bright-source structure; target attribution remains unresolved after completed independent helper audit. Only two long-separated raw events have usable timings, with no independent adjacent same-kind interval; the period is untested against integer-cycle alternatives, so the physical tier is PLAUSIBLE. Two measured QLP events fourteen historical cycles apart support timing conditional on integer-cycle assignment; alternative cycle aliases are not independently eliminated. The helper S24 cutout center fits failed (bounds/zero depth), not valid timings or nulls. Gaia conjunction offset0.0646phase has propagated uncertainty1.34d, not an arbitrary inconsistency veto. Next test: Consecutive same-kind resolved eclipse timings; source attribution
R326	331861613	G 12-12.5 spectroscopic-orbit population	PLAUSIBLE	false	Report; Three measured events at historical cycles0,2,102 repeat, but no adjacent-cycle pair fixes the integer/subharmonic choice. The18.15346d ephemeris and its conditional fit error assume those cycle assignments; alternative periods remain unresolved. Available archived sectors17,18,58,85 were checked, including added earlierS18. S58 cycle62 has zero quality-zero samples within0.2d, so it is not a null. Twelve capable neighbours include one at13.99arcsec; attribution also remains unresolved. Next test: Resolve cycle-count aliases with an observed adjacent eclipse and resolve source localization.
R335	377129604	G 12-12.5 spectroscopic-orbit population	PLAUSIBLE	false	Report; Only two deep S72 events are established. Their depths50.81/60.59ppt differ in QLP and custom aperture. The18.233234d value is a measured event separation, not an established same-kind orbital period;36.466469d double alias and eclipse-kind ambiguity remain. All available QLP sectors45,46,72 were inspected; earlier weak fits excluded, not used as timings. Earlier absence is not asserted as a clean null. Historical18.266535d period is not adopted. Depths depend on aperture dilution and local baseline. Pixel centroid uncertainties and timing systematics are not calibrated. Next test: Obtain a third deep eclipse and test same-kind/double aliases

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R368	382631802	G 11.5-12 spectroscopic-orbit population	PLAUSIBLE	false	Report; PLAUSIBLE: recurring V-shaped dips ride structured variability. No stable sharp contacts or flat-bottomed occultation signature is resolved across representative windows; free trapezoids alone do not distinguish eclipses from pulsation or tidal minima. Eclipse interpretation itself is insecure. Free trapezoid duration median7.58h exceeds approximate central3.21h using TIC radius1.64solar and main-sequence minimum-mass companion. Difference centroid favors target0.220pixel versus capable neighbours2.38/2.67pixel; PSF fractional residual0.46 limits precision. Peak is measured within the central target5x5 box, excluding invalid peripheral cutout pixels; peripheral invalid pixels do not decide attribution. Model durations vary1.58-14.05h, indicating baseline/shape contamination. Timing reduced chi-square about126 requires scatter inflation. Geometry and eclipse identity are unresolved after completed checks; alternative variability and revised radii require testing. Next test: Joint variability/eclipse model and radii check
R431	210418035	G 12-12.25 spectroscopic-orbit population	PLAUSIBLE	false	Report; Broad, changing 0.96–2.86 ppt depressions repeat near the selected phase, but fitted duration is 27.45 h versus a rough 5.84 h central eclipse duration. S11 includes a narrow-looking dip while S99/S102 are broad. A phase-dependent orbital variable, blend or extraction structure remains possible. The timing fit does not establish stable eclipses; crowded S65 pixels are noisy. Next test: Resolved time-series photometry and a phase-resolved orbital model to distinguish broad variability from eclipses; repeat difference imaging in a cleaner sector.
R440	381076353	G 12-12.25 spectroscopic-orbit population	PLAUSIBLE	false	Report; Recurrent ~3 ppt depressions are comparable to local scatter. Folded S11 and S102 show localized troughs and recovery, but S99 is rounded and broad variability is not decisively excluded; eclipse morphology remains uncertain. Fifty capable Gaia sources within 60 arcsec include several within 8 arcsec; the target-centered core has 0.909 image-model fractional residual, leaving attribution unresolved. Trapezoid durations and timing are shape-sensitive (up to 113 min timing residual). Next test: Test morphology and resolve source attribution

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R454	232533575	G 12-12.25 spectroscopic-orbit population	PLAUSIBLE	false	Report; Two target-centered eclipses in sector 77 are separated by 15.252 d. The Gaia-like 5.084-d period assigns three cycles, but cycles 291 and 292 fall in an observation gap. Sector 23 has no usable repeated center and current MAST lists only sectors 23 and 77. Thus the integer-cycle alias is untested by a third eclipse; the reported period is conditional on the Gaia-supported three-cycle assignment. Next test: A third measured eclipse in future coverage to distinguish 15.252 d divided by three from other integer-cycle aliases.
R565	290749976	G 12.25-12.5 spectroscopic-orbit population	PLAUSIBLE	false	Report; Shallow recurring broad dips have variable baselines and a noisy offset deficit (0.62pixel). Three capable neighbors lie36-49arcsec away. Smooth morphology and variable fitted duration versus approximate central4.85h do not establish eclipse geometry; rotational, tidal or baseline modulation remains possible. No convincing sharp eclipse component is established, so this is PLAUSIBLE, not a demonstrated repeating eclipser. TIC radius2.113 solar radii is already used; broad shape/baseline and attribution remain unresolved. Period describes dip recurrence; trapezoid centers are model-dependent. Next test: Establish sharp eclipse morphology against rotational/tidal/baseline modulation, then resolve source ownership with eclipse photometry.
R679	462088261	2026-11-03/04 selected spectroscopic-orbit follow-up	PLAUSIBLE	false	Report; Large correlated orbital modulation and curved local baselines make the shallow narrow component hard to isolate as an eclipse. Target is an exact VSX RS variable. G15.56 and G18.13 sources 5.2 arcsec away are flux-capable; s37 PSF residual is 0.87 and the target pixel cannot distinguish them. Sparse four usable trapezoid centers do not yield a secure eclipse ephemeris.
R688	412639100	2026-11-03/04 selected spectroscopic-orbit follow-up	PLAUSIBLE	false	Report; Decision: Bounded lead/helper review complete. Broad shallow recurrent trough lacks secure eclipse contacts, measured timings, or localized source; alternative periastron variability unresolved. No physical or new-claim membership.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R696	95623701	2026-11-03/04 selected spectroscopic-orbit follow-up	PLAUSIBLE	false	Report; Decision: Completed review finds contradictory clean null for assumed cycle alias and unresolved source; accept plausible, excluded. The deep sharp sector 12 and 103 events have similar shape at an assumed 112-cycle alias, but sector 39 is much weaker and a fully covered quality-zero sector 65 predicted window is a clean null for that alias. The helper TESScut 3×3 extraction independently finds 93.06, 13.81, 0.124, and 126.81 ppt in sectors 12/39/65/103; aperture size alone does not explain the null. A 16.81-arcsec Gaia neighbor is capable only under a very deep intrinsic eclipse and remains unresolved at the pixel scale. The assumed alias also disagrees strongly with Gaia. The repeated-looking signal has no secure persistent orbital clock or source ownership.
R711	280950412	2026-11-03/04 selected spectroscopic-orbit follow-up	PLAUSIBLE	false	Report; Decision: Completed bounded review remains plausible; sparse or weak events do not resolve integer aliases. No new-system claim. The two putative shallow dips are baseline- and aperture-dependent, with no reliable raw-event timing fit. All tested aliases lack sensitivity to a 1.2-ppt repeat; the screen conjunction is not a measured event. Severe crowding prevents source attribution.
R712	188908998	2026-11-03/04 selected spectroscopic-orbit follow-up	PLAUSIBLE	false	Report; Decision: Completed bounded review remains plausible; sparse or weak events do not resolve integer aliases. No new-system claim. Only two raw same-kind events, 32 putative cycles apart, have usable centers. The neighboring 31-cycle alias has no sensitive intermediate coverage, so the integer period is untested; two capable Gaia neighbors remain.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
R719	349527829	2026-11-03/04 selected spectroscopic-orbit follow-up	PLAUSIBLE	false	Report; Decision: Root accepts complete plausible review; sparse integer-cycle ambiguity prevents physical inclusion. Two similar raw eclipse occasions are separated by about1230 d. Several nearby integer-cycle aliases have no sensitive actual coverage null and remain untested. The sector91 ingress is gapped, making its reconstructed center additionally model dependent. Difference imaging and a separate extraction support the target, but do not supply a third observing occasion. The conditional period and formal fit error in saved numeric artifacts are hypotheses, not an established ephemeris; generic orbital fields are blank. TESScut is a separate reduction of the same photons. Gaia G-band flux ratios and full throughput are optimistic neighbor capability proxies, not TESS PRF likelihoods. No exhaustive all-event fitting, precision stellar characterization, or Monte Carlo was performed.
R721	95836684	2026-11-03/04 selected spectroscopic-orbit follow-up	PLAUSIBLE	false	Report; Decision: Root accepts complete plausible review; sparse integer-cycle ambiguity prevents physical inclusion. Two deep flat-bottomed raw eclipses are target-centered and survive the separate extraction, but the31-cycle assignment is conditional. Nearby integer counts29,30,32,33 remain without sensitive actual coverage nulls. A half-period null does not resolve these distant integer alternatives. A third independent observing occasion is required; generic orbital fields are blank. TESScut is a separate reduction of the same photons. Gaia G-band flux ratios and full throughput are optimistic neighbor capability proxies, not TESS PRF likelihoods. No exhaustive all-event fitting, precision stellar characterization, or Monte Carlo was performed.

Non-EB, neighbouring-source and systematic signals (21)

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
C13	372683900	fainter SB1 period-fold population	ELLIPSOIDAL	false	Report; Smooth double wave, no narrow eclipse; unresolved 1.8 arcsec neighbour; Gaia eccentricity conflicts at 3.3 sigma.
C14	13106408	fainter SB1 period-fold population	ELLIPSOIDAL	false	Report; Smooth double wave; nearby 364-day SR unrelated.
C15	372683622	fainter SB1 period-fold population	ELLIPSOIDAL	false	Report; Double wave plus variable spots; no eclipse.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
C17	62328710	fainter SB1 period-fold population	NEIGHBOUR_SIGNAL	false	Report; Delta Scuti-like signal localized to Gaia DR3 2165710355333626240 at 44 arcsec and independently seen there by ZTF; target flat at Gaia period; suspect SB1 orbit is interpretation.
C27	10790605	fainter SB1 period-fold population	REFUTED	false	Report; See target report for deciding limitation and next check.
C28	376082142	fainter SB1 period-fold population	REFUTED	false	Report; Search higher precision data for a discrete narrow eclipse only if this smooth variable remains of interest.
C34	251357000	fainter SB1 period-fold population	REFUTED	false	Report; Search higher precision data for a discrete narrow eclipse only if this smooth variable remains of interest.
C42	12320469	fainter SB1 period-fold population	REFUTED	false	Report; Higher precision light curves could test for a narrow residual eclipse if further work on this smooth variable is desired.
C43	86908196	fainter SB1 period-fold population	REFUTED	false	Report; Higher precision light curves could test for a narrow residual eclipse if further work on this smooth variable is desired.
C46	411647453	fainter SB1 period-fold population	REFUTED	false	Report; Higher resolution event photometry can establish the contaminant host and measure its orbital period precisely.
C47	450217907	fainter SB1 period-fold population	REFUTED	false	Report; Higher precision light curves could test for a narrow residual eclipse if further work on this smooth variable is desired.
C48	60443793	fainter SB1 period-fold population	REFUTED	false	Report; Review TIC 60443695 separately if desired.
C51	311206604	fainter SB1 period-fold population	REFUTED	false	Report; No further eclipse follow-up required unless new discrete events appear.
C52	428769301	fainter SB1 period-fold population	REFUTED	false	Report; No further eclipse follow-up required unless new discrete events appear.
C53	436947190	fainter SB1 period-fold population	REFUTED	false	Report; No further eclipse follow-up required unless new discrete events appear.
C62	431762229	fainter SB1 period-fold population	REFUTED	false	Report; Higher resolution eclipse photometry can verify the contaminating host.
C65	222498305	fainter SB1 period-fold population	REFUTED	false	Report; Higher resolution eclipse photometry can verify the contaminating host.
C66	90542579	fainter SB1 period-fold population	REFUTED	false	Report; If revisiting, use clean cadences away from downlink boundaries with a validated background and aperture.
C69	297541839	fainter SB1 period-fold population	REFUTED	false	Report; No missing deciding gate for this candidate. Reopen only if new data supply distinct, repeated detached events.

Key	TIC	Population	Physical verdict	Added	Evidence / limitation
C70	396248369	fainter SB1 period-fold population	REFUTED	false	Report; No missing deciding gate for this candidate. Reopen only if new data supply distinct, repeated detached events.
C71	277479128	fainter SB1 period-fold population	REFUTED	false	Report; No missing deciding gate for this candidate. Reopen only if new data supply distinct, repeated detached events.