

*Ph.D. Thesis*

**Probing Electroweakly Interacting  
Massive Particles  
with Drell-Yan Process  
at 100 TeV Colliders**

So Chigusa

*Department of Physics*



THE UNIVERSITY OF TOKYO

December 2019

## Abstract

(♣ To be written ♣) The main part of this thesis is based on our works [1, 2].

# Acknowledgments

I would like to thank my supervisor, Takeo Moroi, who provided stimulating discussions, helpful comments, fruitful suggestions and collaboration and proofread the earlier version of this thesis.

(♣ Appreciate more ♣)

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Overview . . . . .	1
1.2	Organization of this thesis . . . . .	4

# Section 1

## Introduction

### 1.1 Overview

There are many models that extend the standard model (SM) of the particle physics by introducing weakly interacting massive particles (WIMPs), which have interactions with SM particles whose size is comparable to the electroweak gauge coupling. We have many theoretical motivations to introduce WIMPs, some of which are listed below.

- One of the motivations is the existence of the dark matter (DM) in our universe. When we assume a stable WIMP with its interaction strength comparable to the electroweak gauge coupling, its thermal relic abundance agrees with the observation if its mass is of  $\mathcal{O}(\text{TeV})$  or so.
- Such  $\mathcal{O}(\text{TeV})$  WIMPs often appear in well-motivated models beyond the standard model (BSM). For example, the following two models contain such WIMPs: the minimally supersymmetric standard model (MSSM) introduced to solve the hierarchy problem and the minimal dark matter (MDM) model that can explain the existence and stability of the DM in a minimal extension of the SM.

Besides, models with WIMPs are also phenomenologically interesting because

- Such  $\mathcal{O}(\text{TeV})$  WIMPs are likely to be discovered by many kinds of experiments such as the dark matter searches and the particle colliders.

Considering the second point, many of the WIMPs in the BSM models mentioned above have non-zero electroweak charges, which explain its weak interaction with SM particles. In this thesis, we would like to focus on such kind of WIMPs and seek ways to search for them.<sup>[†1](#)</sup>

In the MSSM, it is known that the supersymmetric (SUSY) partner of the electroweak gauge bosons or the Higgs boson can be the lightest supersymmetric particles (LSP) and are natural DM candidate. In particular, there are well motivated scenarios where the so-called Higgsino or Wino play the role of the LSP, which transform as doublet and triplet under

---

<sup>†1</sup>The word “WIMPs” is usually used in a broader sense that includes particles with some unknown weak interactions with SM particles. To distinguish this usage with ours, which only denotes some particles with non-zero electroweak charge, it may be better to call them as “WIMPs”, abbreviation of electroweakly interacting massive particles. However, within this thesis, we will just use “WIMPs” in a narrow sense obeying the widely spread custom.

the weak  $SU(2)_L$  gauge symmetry, respectively; light Higgsino is preferred to reduce the amount of the fine-tuning of the electroweak scale as in the “natural SUSY” set up [3–6], while the so-called “mini-split” spectrum [7–12] with anomaly mediation [13, 14] makes Wino LSP. Another example, the MDM scenario [15–17], introduces a larger  $SU(2)_L$  multiplet, whose stability is automatically ensured by the charge assignment. In particular, a 5-plet Majorana fermion with hypercharge zero is a good DM candidate that escape from the DM search experiments so far.

To search for WIMPs, several different approaches are adopted. One way is to rely on DM search experiments, assuming that the WIMPs are the dominant component of the DM. Firstly, there exist several direct detection experiments that utilize a scattering between the DM and the nucleus [18–20]. Wino is one of the promising targets of these experiments, whose spin independent scattering cross section with a nucleon is almost mass independently given as  $\sigma_p^{\text{SI}} \simeq 2.3 \times 10^{-47} \text{ cm}^2$  [21–25], which is still an order of magnitude below the current experimental limit. The situation for Higgsino highly depends on the size of the mixing between Higgsino and SUSY partners of electroweak gauge bosons (or electroweakinos). However, it is particularly difficult to detect (almost) pure Higgsino DM since its scattering cross section is comparable to or below that of the neutrino background [22]. The detection of MDM may also be difficult [26] since its possibly larger mass of  $\mathcal{O}(10)$  TeV weakens the sensitivity of direct detection experiments.

Secondly, a lot of efforts are devoted to detecting cosmic rays resulting from DM annihilation, namely the DM indirect detection [27–30]. Although the results suffer from some astrophysical uncertainties, they have already excluded, for example, Wino with mass less than 400 GeV and also around 2 TeV [31]. On the other hand, the corresponding Higgsino bound is weaker and it has been probed only up to 350 GeV [32] again due to the smallness of its  $SU(2)_L$  charge. For the MDM, 5-plet fermion is analyzed as an example in [33] and the mass less than 2 TeV and several narrow regions are excluded. Note again that the WIMPs must be the dominant component of the DM for these approaches to be sensitive.

Another way of probing WIMPs is the direct production at the large hadron collider (LHC). The main strategy is to use the disappearing charged track, which indicates a long-lived charged particle, the charged components of the WIMP in the current case. Both ATLAS and CMS collaborations announced a result of this method with the data of  $\sqrt{s} = 13$  TeV LHC [34–36]. The current lower bound on the mass of the pure Higgsino-like (Wino-like) state is 152 (460) GeV at 95 % C.L. We can obtain a similar bound for the MDM using the same method [37]. In this method, however, the bound strongly depends on the lifetime of the charged component, which is sensitive to the mass difference between the charged and the neutral components. In particular, it is often the case in the SUSY model that the Higgsino-like LSP and its charged counterpart possess a non-negligible fraction of electroweakinos, which significantly enhances the mass difference compared to the pure

Higgsino case. In such a case, the lifetime of the charged component is extremely short, making the disappearing track search challenging. There is another option called mono-X search to search for a new physics signal in general. However, the corresponding bound is usually very weak due to the large SM background and no bound is imposed on Higgsino at  $\sqrt{s} = 14$  TeV LHC [38].

To acquire somewhat stronger bounds on the WIMPs, independent of the decay product or the lifetime and whether they are the dominant component of the DM or not, it has been discussed that indirect search of WIMPs at collider experiments is useful [39–43]. It utilizes the WIMP loop effect on various observables. We pursue this possibility and study the effects of WIMPs on the oblique correction to the electroweak gauge bosons. In particular, in this paper, we study the prospect of the indirect search method at future 100 TeV hadron colliders such as FCC-hh at CERN [44–46] or SppC in China [47, 48]. We concentrate on the Drell-Yan process that has two leptons in the final state since it provides a very clean signal without any hadronic jets at least from the final state particles. We will show that it provides a comparable or better experimental reach for Higgsino compared to the direct production search of WIMPs at future colliders [49–52]. This method also provides independent and additional information about Wino and the MDM. Besides, we will reveal that this method is useful to investigate WIMP properties, such as charges, masses, and spins.

## 1.2 Organization of this thesis

This thesis is organized as follows.

In Sec. ??, we briefly review models with WIMPs considered in the thesis. Sec. ?? is devoted to the minimally supersymmetric standard model, while Sec. ?? to the minimal dark matter model. The mass splitting among the multiplet, which is a phenomenologically important property of WIMPs, is described in Sec. ?. The WIMP properties are summarized in Sec. ?.

In Sec. ?, we summarize the WIMP properties as a DM candidate. We show the calculation of thermal relic abundance and derive the required mass for WIMP DMs in Sec. ?. (**♣ More later ♣**)

In Sec. ?, we study the WIMP production at the hadron collider experiments and its detection. Possible production processes and the kinematics of the produced WIMPs are summarized in Sec. ?. Using the production processes described here, the current status and the prospect at the future 100, TeV colliders are reviewed in the following subsections. Sec. ? is devoted to the description of the most promising way for the WIMP search called the disappearing track search. (**♣ More later ♣**)

In Sec. ?, we discuss our own ideas [1, 2] to probe the one-loop effect of WIMPs on the lepton pair production processes through the precise measurement at 100 TeV colliders. Here, we will describe our statistical analysis, show the obtained reach for WIMPs, and see the possibility to determine the WIMP properties after its discovery. Conclusions of the thesis are presented in Sec. ?.

In Appendices, we first summarize the conventions and notations used in the thesis in ?. Then, we briefly review the  $\mathcal{N} = 1$  supersymmetry in Appendix ?, the properties of the so-called transverse mass used in our analysis in Appendix ?, and the statistical analysis method called the profile likelihood method in Sec. ?.

# References

- [1] S. Chigusa, Y. Ema, T. Moroi, Probing electroweakly interacting massive particles with Drell-Yan process at 100 TeV hadron colliders, Phys. Lett. B789 (2019) 106–113. [arXiv:1810.07349](#), [doi:10.1016/j.physletb.2018.12.011](#).
- [2] T. Abe, S. Chigusa, Y. Ema, T. Moroi, Indirect studies of electroweakly interacting particles at 100 TeV hadron colliders, Phys. Rev. D100 (5) (2019) 055018. [arXiv:1904.11162](#), [doi:10.1103/PhysRevD.100.055018](#).



- 
- [3] R. Kitano, Y. Nomura, A Solution to the supersymmetric fine-tuning problem within the MSSM, Phys. Lett. B631 (2005) 58–67. [arXiv:hep-ph/0509039](#), [doi:10.1016/j.physletb.2005.10.003](#).
  - [4] C. Brust, A. Katz, S. Lawrence, R. Sundrum, SUSY, the Third Generation and the LHC, JHEP 03 (2012) 103. [arXiv:1110.6670](#), [doi:10.1007/JHEP03\(2012\)103](#).
  - [5] M. Papucci, J. T. Ruderman, A. Weiler, Natural SUSY Endures, JHEP 09 (2012) 035. [arXiv:1110.6926](#), [doi:10.1007/JHEP09\(2012\)035](#).
  - [6] H. Baer, V. Barger, P. Huang, X. Tata, Natural Supersymmetry: LHC, dark matter and ILC searches, JHEP 05 (2012) 109. [arXiv:1203.5539](#), [doi:10.1007/JHEP05\(2012\)109](#).
  - [7] J. D. Wells, Implications of supersymmetry breaking with a little hierarchy between gauginos and scalars, in: 11th International Conference on Supersymmetry and the Unification of Fundamental Interactions (SUSY 2003) Tucson, Arizona, June 5–10, 2003, 2003. [arXiv:hep-ph/0306127](#).
  - [8] J. D. Wells, PeV-scale supersymmetry, Phys. Rev. D71 (2005) 015013. [arXiv:hep-ph/0411041](#), [doi:10.1103/PhysRevD.71.015013](#).
  - [9] N. Arkani-Hamed, S. Dimopoulos, Supersymmetric unification without low energy supersymmetry and signatures for fine-tuning at the LHC, JHEP 06 (2005) 073. [arXiv:hep-th/0405159](#), [doi:10.1088/1126-6708/2005/06/073](#).
  - [10] G. F. Giudice, A. Romanino, Split supersymmetry, Nucl. Phys. B699 (2004) 65–89, [Erratum: Nucl. Phys.B706,487(2005)]. [arXiv:hep-ph/0406088](#), [doi:10.1016/j.nuclphysb.2004.11.048](#), [doi:10.1016/j.nuclphysb.2004.08.001](#).
  - [11] N. Arkani-Hamed, S. Dimopoulos, G. F. Giudice, A. Romanino, Aspects of split supersymmetry, Nucl. Phys. B709 (2005) 3–46. [arXiv:hep-ph/0409232](#), [doi:10.1016/j.nuclphysb.2004.12.026](#).
  - [12] N. Arkani-Hamed, S. Dimopoulos, S. Kachru, Predictive landscapes and new physics at a TeV (2005). [arXiv:hep-th/0501082](#).
  - [13] L. Randall, R. Sundrum, Out of this world supersymmetry breaking, Nucl. Phys. B557 (1999) 79–118. [arXiv:hep-th/9810155](#), [doi:10.1016/S0550-3213\(99\)00359-4](#).
  - [14] G. F. Giudice, M. A. Luty, H. Murayama, R. Rattazzi, Gaugino mass without singlets, JHEP 12 (1998) 027. [arXiv:hep-ph/9810442](#), [doi:10.1088/1126-6708/1998/12/027](#).

- [15] M. Cirelli, N. Fornengo, A. Strumia, Minimal dark matter, Nucl. Phys. B753 (2006) 178–194. [arXiv:hep-ph/0512090](#), [doi:10.1016/j.nuclphysb.2006.07.012](#).
- [16] M. Cirelli, A. Strumia, M. Tamburini, Cosmology and Astrophysics of Minimal Dark Matter, Nucl. Phys. B787 (2007) 152–175. [arXiv:0706.4071](#), [doi:10.1016/j.nuclphysb.2007.07.023](#).
- [17] M. Cirelli, A. Strumia, Minimal Dark Matter: Model and results, New J. Phys. 11 (2009) 105005. [arXiv:0903.3381](#), [doi:10.1088/1367-2630/11/10/105005](#).
- [18] D. S. Akerib, et al., Results from a search for dark matter in the complete LUX exposure, Phys. Rev. Lett. 118 (2) (2017) 021303. [arXiv:1608.07648](#), [doi:10.1103/PhysRevLett.118.021303](#).
- [19] X. Cui, et al., Dark Matter Results From 54-Ton-Day Exposure of PandaX-II Experiment, Phys. Rev. Lett. 119 (18) (2017) 181302. [arXiv:1708.06917](#), [doi:10.1103/PhysRevLett.119.181302](#).
- [20] E. Aprile, et al., Dark Matter Search Results from a One Tonne×Year Exposure of XENON1T, Phys. Rev. Lett. 121 (11) (2018) 111302. [arXiv:1805.12562](#), [doi:10.1103/PhysRevLett.121.111302](#).
- [21] J. Hisano, K. Ishiwata, N. Nagata, A complete calculation for direct detection of Wino dark matter, Phys. Lett. B690 (2010) 311–315. [arXiv:1004.4090](#), [doi:10.1016/j.physletb.2010.05.047](#).
- [22] J. Hisano, K. Ishiwata, N. Nagata, Direct Search of Dark Matter in High-Scale Supersymmetry, Phys. Rev. D87 (2013) 035020. [arXiv:1210.5985](#), [doi:10.1103/PhysRevD.87.035020](#).
- [23] J. Hisano, K. Ishiwata, N. Nagata, QCD Effects on Direct Detection of Wino Dark Matter, JHEP 06 (2015) 097. [arXiv:1504.00915](#), [doi:10.1007/JHEP06\(2015\)097](#).
- [24] R. J. Hill, M. P. Solon, Universal behavior in the scattering of heavy, weakly interacting dark matter on nuclear targets, Phys. Lett. B707 (2012) 539–545. [arXiv:1111.0016](#), [doi:10.1016/j.physletb.2012.01.013](#).
- [25] R. J. Hill, M. P. Solon, WIMP-nucleon scattering with heavy WIMP effective theory, Phys. Rev. Lett. 112 (2014) 211602. [arXiv:1309.4092](#), [doi:10.1103/PhysRevLett.112.211602](#).

- 
- [26] J. Hisano, K. Ishiwata, N. Nagata, T. Takesako, Direct Detection of Electroweak-Interacting Dark Matter, JHEP 07 (2011) 005. [arXiv:1104.0228](#), [doi:10.1007/JHEP07\(2011\)005](#).
- [27] A. Albert, et al., Searching for Dark Matter Annihilation in Recently Discovered Milky Way Satellites with Fermi-LAT, Astrophys. J. 834 (2) (2017) 110. [arXiv:1611.03184](#), [doi:10.3847/1538-4357/834/2/110](#).
- [28] M. L. Ahnen, et al., Limits to Dark Matter Annihilation Cross-Section from a Combined Analysis of MAGIC and Fermi-LAT Observations of Dwarf Satellite Galaxies, JCAP 1602 (02) (2016) 039. [arXiv:1601.06590](#), [doi:10.1088/1475-7516/2016/02/039](#).
- [29] H. Abdallah, et al., Search for dark matter annihilations towards the inner Galactic halo from 10 years of observations with H.E.S.S., Phys. Rev. Lett. 117 (11) (2016) 111301. [arXiv:1607.08142](#), [doi:10.1103/PhysRevLett.117.111301](#).
- [30] G. A. Gómez-Vargas, M. A. Sánchez-Conde, J.-H. Huh, M. Peiró, F. Prada, A. Morselli, A. Klypin, D. G. Cerdeño, Y. Mambrini, C. Muñoz, Constraints on WIMP annihilation for contracted dark matter in the inner Galaxy with the Fermi-LAT, JCAP 1310 (2013) 029. [arXiv:1308.3515](#), [doi:10.1088/1475-7516/2013/10/029](#).
- [31] B. Bhattacharjee, M. Ibe, K. Ichikawa, S. Matsumoto, K. Nishiyama, Wino Dark Matter and Future dSph Observations, JHEP 07 (2014) 080. [arXiv:1405.4914](#), [doi:10.1007/JHEP07\(2014\)080](#).
- [32] R. Krall, M. Reece, Last Electroweak WIMP Standing: Pseudo-Dirac Higgsino Status and Compact Stars as Future Probes, Chin. Phys. C42 (4) (2018) 043105. [arXiv:1705.04843](#), [doi:10.1088/1674-1137/42/4/043105](#).
- [33] H. Abdalla, et al., Searches for gamma-ray lines and 'pure WIMP' spectra from Dark Matter annihilations in dwarf galaxies with H.E.S.S., JCAP 1811 (11) (2018) 037. [arXiv:1810.00995](#), [doi:10.1088/1475-7516/2018/11/037](#).
- [34] M. Aaboud, et al., Search for long-lived charginos based on a disappearing-track signature in pp collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector, JHEP 06 (2018) 022. [arXiv:1712.02118](#), [doi:10.1007/JHEP06\(2018\)022](#).
- [35] Search for direct pair production of higgsinos by the reinterpretation of the disappearing track analysis with  $36.1 \text{ fb}^{-1}$  of  $\sqrt{s} = 13$  TeV data collected with the ATLAS experiment, Tech. Rep. ATL-PHYS-PUB-2017-019, CERN, Geneva (Dec 2017). URL <http://cds.cern.ch/record/2297480>

- 
- [36] A. M. Sirunyan, et al., Search for disappearing tracks as a signature of new long-lived particles in proton-proton collisions at  $\sqrt{s} = 13$  TeV, JHEP 08 (2018) 016. [arXiv:1804.07321](#), [doi:10.1007/JHEP08\(2018\)016](#).
- [37] B. Ostdiek, Constraining the minimal dark matter fiveplet with LHC searches, Phys. Rev. D92 (2015) 055008. [arXiv:1506.03445](#), [doi:10.1103/PhysRevD.92.055008](#).
- [38] H. Baer, A. Mustafayev, X. Tata, Monojets and mono-photons from light higgsino pair production at LHC14, Phys. Rev. D89 (5) (2014) 055007. [arXiv:1401.1162](#), [doi:10.1103/PhysRevD.89.055007](#).
- [39] D. S. M. Alves, J. Galloway, J. T. Ruderman, J. R. Walsh, Running Electroweak Couplings as a Probe of New Physics, JHEP 02 (2015) 007. [arXiv:1410.6810](#), [doi:10.1007/JHEP02\(2015\)007](#).
- [40] C. Gross, O. Lebedev, J. M. No, Drell-Yan constraints on new electroweak states: LHC as a  $pp \rightarrow l^+l^-$  precision machine, Mod. Phys. Lett. A32 (16) (2017) 1750094. [arXiv:1602.03877](#), [doi:10.1142/S0217732317500948](#).
- [41] M. Farina, G. Panico, D. Pappadopulo, J. T. Ruderman, R. Torre, A. Wulzer, Energy helps accuracy: electroweak precision tests at hadron colliders, Phys. Lett. B772 (2017) 210–215. [arXiv:1609.08157](#), [doi:10.1016/j.physletb.2017.06.043](#).
- [42] K. Harigaya, K. Ichikawa, A. Kundu, S. Matsumoto, S. Shirai, Indirect Probe of Electroweak-Interacting Particles at Future Lepton Colliders, JHEP 09 (2015) 105. [arXiv:1504.03402](#), [doi:10.1007/JHEP09\(2015\)105](#).
- [43] S. Matsumoto, S. Shirai, M. Takeuchi, Indirect Probe of Electroweakly Interacting Particles at the High-Luminosity Large Hadron Collider, JHEP 06 (2018) 049. [arXiv:1711.05449](#), [doi:10.1007/JHEP06\(2018\)049](#).
- [44] M. L. Mangano, et al., Physics at a 100 TeV pp Collider: Standard Model Processes, CERN Yellow Report (3) (2017) 1–254. [arXiv:1607.01831](#), [doi:10.23731/CYRM-2017-003.1](#).
- [45] R. Contino, et al., Physics at a 100 TeV pp collider: Higgs and EW symmetry breaking studies, CERN Yellow Report (3) (2017) 255–440. [arXiv:1606.09408](#), [doi:10.23731/CYRM-2017-003.255](#).
- [46] T. Golling, et al., Physics at a 100 TeV pp collider: beyond the Standard Model phenomena, CERN Yellow Report (3) (2017) 441–634. [arXiv:1606.00947](#), [doi:10.23731/CYRM-2017-003.441](#).

- 
- [47] M. Ahmad, et al., CEPC-SPPC Preliminary Conceptual Design Report. 1. Physics and Detector (2015).
  - [48] C.-S. S. Group, CEPC-SPPC Preliminary Conceptual Design Report. 2. Accelerator (2015).
  - [49] M. Low, L.-T. Wang, Neutralino dark matter at 14 TeV and 100 TeV, JHEP 08 (2014) 161. [arXiv:1404.0682](#), [doi:10.1007/JHEP08\(2014\)161](#).
  - [50] M. Cirelli, F. Sala, M. Taoso, Wino-like Minimal Dark Matter and future colliders, JHEP 10 (2014) 033, [Erratum: JHEP01,041(2015)]. [arXiv:1407.7058](#), [doi:10.1007/JHEP10\(2014\)033](#), [doi:10.1007/JHEP01\(2015\)041](#).
  - [51] T. Han, S. Mukhopadhyay, X. Wang, Electroweak Dark Matter at Future Hadron Colliders, Phys. Rev. D98 (3) (2018) 035026. [arXiv:1805.00015](#), [doi:10.1103/PhysRevD.98.035026](#).
  - [52] R. Mahbubani, P. Schwaller, J. Zurita, Closing the window for compressed Dark Sectors with disappearing charged tracks, JHEP 06 (2017) 119, [Erratum: JHEP10,061(2017)]. [arXiv:1703.05327](#), [doi:10.1007/JHEP06\(2017\)119](#), [doi:10.1007/JHEP10\(2017\)061](#).