

Ph.D. Thesis

**Indirect study of electroweakly
interacting particles
at 100 TeV hadron colliders**

So Chigusa

Department of Physics



THE UNIVERSITY OF TOKYO

December 2019

Abstract

(♣ To be written ♣)

Acknowledgments

(♣ To be written ♣)

Contents

1	Introduction	1
2	Weakly interacting massive particle	1
2.1	WIMPs as a dark matter candidate	1

Section 1

Introduction

(♣ Definition of “SM” ♣)

(♣ Definition of “WIMP” ♣)

Section 2

Weakly interacting massive particle

2.1 WIMPs as a dark matter candidate

One of the most important evidences of the beyond SM is the existence of dark matter (DM) [1]. DM is an unknown object that occupies a non-negligible ratio of the total energy of our Universe, but has not yet been directly observed because of its weak interaction with the SM particles.^{‡1} In spite of its invisibility, the existence of DM is confirmed by several astrophysical observations such as the mass measurement using the gravitational lensing effect caused by galaxies and clusters [2, 3], the flatness of galactic rotation curves further the optical radius [4, 5], the measurement of the power spectrum of the cosmic microwave background (CMB), and so on. In particular, the observation of CMB allows us to determine various cosmological parameters [6, 7] including the density of the non-relativistic matter and baryon, which is currently determined as [8]

$$\Omega_m h^2 = 0.1430 \pm 0.0011, \quad (2.1)$$

$$\Omega_b h^2 = 0.02237 \pm 0.00015, \quad (2.2)$$

where $h \sim \mathcal{O}(1)$ is the Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$. The difference between $\Omega_m h^2$ and $\Omega_b h^2$ implies the existence of DM and its abundance $\Omega_c h^2 \simeq 0.12$.

^{‡1} At worst DM interacts with the SM particles through the gravity, which is considerably weaker than all the other known interactions. (♣ Mention to Ema paper?? ♣)

References

- [1] F. Zwicky, Die Rotverschiebung von extragalaktischen Nebeln, *Helvetica Physica Acta* 6 (1933) 110.
- [2] F. Zwicky, On the Masses of Nebulae and of Clusters of Nebulae, *Astrophysical Journal* 86 (1937) 217.
- [3] V. Trimble, Existence and Nature of Dark Matter in the Universe, *Ann. Rev. Astron. Astrophys.* 25 (1987) 425–472. [doi:10.1146/annurev.aa.25.090187.002233](#).
- [4] H. W. Babcock, The rotation of the Andromeda Nebula, *Lick Observatory Bulletin* 19 (1939) 41–51. [doi:10.5479/ADS/bib/1939LicOB.19.41B](#).
- [5] K. G. Begeman, A. H. Broeils, R. H. Sanders, Extended rotation curves of spiral galaxies: Dark haloes and modified dynamics, *Mon. Not. Roy. Astron. Soc.* 249 (1991) 523.
- [6] G. Jungman, M. Kamionkowski, A. Kosowsky, D. N. Spergel, Weighing the universe with the cosmic microwave background, *Phys. Rev. Lett.* 76 (1996) 1007–1010. [arXiv:astro-ph/9507080](#), [doi:10.1103/PhysRevLett.76.1007](#).
- [7] G. Jungman, M. Kamionkowski, A. Kosowsky, D. N. Spergel, Cosmological parameter determination with microwave background maps, *Phys. Rev. D* 54 (1996) 1332–1344. [arXiv:astro-ph/9512139](#), [doi:10.1103/PhysRevD.54.1332](#).
- [8] N. Aghanim, et al., Planck 2018 results. VI. Cosmological parameters [arXiv:1807.06209](#).