

PAC-Bayes & Variational Inference

Badr-Eddine Chérif-Abdellatif
Chargé de Recherche CNRS

badr.eddine.cherief.abdellatif@gmail.com



Master 2, Sorbonne Université
Paris, Spring 2023

Reminder of the setting

Training dataset : $\mathcal{S} = \{(X_1, Y_1), \dots, (X_n, Y_n)\}$ i.i.d $\sim \mathbb{P}$,
 $X_i \in \mathcal{X} \subset \mathbb{R}^d$, $Y_i \in \mathcal{Y}$.

- $\mathcal{Y} = \{\text{lion, gazelle}\}$: Binary Classification.
- $\mathcal{Y} = \mathbb{R}$: Regression.

Reminder of the setting

Training dataset : $\mathcal{S} = \{(X_1, Y_1), \dots, (X_n, Y_n)\}$ i.i.d $\sim \mathbb{P}$,
 $X_i \in \mathcal{X} \subset \mathbb{R}^d$, $Y_i \in \mathcal{Y}$.

- $\mathcal{Y} = \{\text{lion, gazelle}\}$: Binary Classification.
- $\mathcal{Y} = \mathbb{R}$: Regression.

Loss $\ell(y', y)$ quantifies the price to predict y' instead of y .

- $\mathcal{Y} = \{\text{cat, dog}\}$: $\ell(y', y) = \mathbb{1}(y' \neq y)$.
- $\mathcal{Y} = \mathbb{R}$: $\ell(y', y) = (y' - y)^2$.

Reminder of the setting

Training dataset : $\mathcal{S} = \{(X_1, Y_1), \dots, (X_n, Y_n)\}$ i.i.d $\sim \mathbb{P}$,
 $X_i \in \mathcal{X} \subset \mathbb{R}^d$, $Y_i \in \mathcal{Y}$.

- $\mathcal{Y} = \{\text{lion, gazelle}\}$: Binary Classification.
- $\mathcal{Y} = \mathbb{R}$: Regression.

Loss $\ell(y', y)$ quantifies the price to predict y' instead of y .

- $\mathcal{Y} = \{\text{cat, dog}\}$: $\ell(y', y) = \mathbb{1}(y' \neq y)$.
- $\mathcal{Y} = \mathbb{R}$: $\ell(y', y) = (y' - y)^2$.

We then define the (theoretical) risk of a predictor $f_\theta, \theta \in \Theta$:

$$R(\theta) = \mathbb{E}_{(X, Y) \sim \mathbb{P}} [\ell(f_\theta(X), Y)],$$

and the empirical risk $\hat{R}_\mathcal{S}(\theta) = \frac{1}{n} \sum_{i=1}^n \ell(f_\theta(X_i), Y_i)$.

Reminder of the setting

Training dataset : $\mathcal{S} = \{(X_1, Y_1), \dots, (X_n, Y_n)\}$ i.i.d $\sim \mathbb{P}$,
 $X_i \in \mathcal{X} \subset \mathbb{R}^d$, $Y_i \in \mathcal{Y}$.

- $\mathcal{Y} = \{\text{lion, gazelle}\}$: Binary Classification.
- $\mathcal{Y} = \mathbb{R}$: Regression.

Loss $\ell(y', y)$ quantifies the price to predict y' instead of y .

- $\mathcal{Y} = \{\text{cat, dog}\}$: $\ell(y', y) = \mathbb{1}(y' \neq y)$.
- $\mathcal{Y} = \mathbb{R}$: $\ell(y', y) = (y' - y)^2$.

We then define the (theoretical) risk of a predictor $f_\theta, \theta \in \Theta$:

$$R(\theta) = \mathbb{E}_{(X,Y) \sim \mathbb{P}} [\ell(f_\theta(X), Y)],$$

and the empirical risk $\hat{R}_\mathcal{S}(\theta) = \frac{1}{n} \sum_{i=1}^n \ell(f_\theta(X_i), Y_i)$.

General goal : Learn using the data a predictor $\hat{\theta}$ with small risk

$$R(\hat{\theta}) = \mathbb{E}_{(X,Y) \sim \mathbb{P}} [\ell(f_{\hat{\theta}}(X), Y) | \mathcal{S}].$$

Reminder of the previous lecture

In this course, we want to derive **Probably Approximately Correct** generalization bounds for predictors $\hat{\theta}$ or randomized predictors $\hat{\rho}$.

Reminder of the previous lecture

In this course, we want to derive **Probably Approximately Correct** generalization bounds for predictors $\hat{\theta}$ or randomized predictors $\hat{\rho}$.

Typical PAC bounds : **with high probability**, the **generalization gap** of θ is **at most something we can control & compute**. For any δ ,

$$\mathbb{P}_{\mathcal{S}} \left[\forall \theta \in \Theta, |R(\theta) - \hat{R}_{\mathcal{S}}(\theta)| \lesssim \sqrt{\frac{\text{comp}(\Theta) + \log(\frac{1}{\delta})}{n}} \right] \geq 1 - \delta.$$

Reminder of the previous lecture

In this course, we want to derive **Probably Approximately Correct** generalization bounds for predictors $\hat{\theta}$ or randomized predictors $\hat{\rho}$.

Typical PAC bounds : **with high probability**, **the generalization gap** of θ is **at most something we can control & compute**. For any δ ,

$$\mathbb{P}_{\mathcal{S}} \left[\forall \theta \in \Theta, |R(\theta) - \hat{R}_{\mathcal{S}}(\theta)| \lesssim \sqrt{\frac{\text{comp}(\Theta) + \log(\frac{1}{\delta})}{n}} \right] \geq 1 - \delta.$$

Some limits to this classical statistical learning theory :

Reminder of the previous lecture

In this course, we want to derive **Probably Approximately Correct** generalization bounds for predictors $\hat{\theta}$ or randomized predictors $\hat{\rho}$.

Typical PAC bounds : **with high probability**, the **generalization gap** of θ is **at most something we can control & compute**. For any δ ,

$$\mathbb{P}_{\mathcal{S}} \left[\forall \theta \in \Theta, |R(\theta) - \hat{R}_{\mathcal{S}}(\theta)| \lesssim \sqrt{\frac{\text{comp}(\Theta) + \log(\frac{1}{\delta})}{n}} \right] \geq 1 - \delta.$$

Some limits to this classical statistical learning theory :

- relies on restricting the complexity of Θ ,

Reminder of the previous lecture

In this course, we want to derive **Probably Approximately Correct** generalization bounds for predictors $\hat{\theta}$ or randomized predictors $\hat{\rho}$.

Typical PAC bounds : **with high probability**, the **generalization gap** of θ is **at most something we can control & compute**. For any δ ,

$$\mathbb{P}_{\mathcal{S}} \left[\forall \theta \in \Theta, |R(\theta) - \hat{R}_{\mathcal{S}}(\theta)| \lesssim \sqrt{\frac{\text{comp}(\Theta) + \log(\frac{1}{\delta})}{n}} \right] \geq 1 - \delta.$$

Some limits to this classical statistical learning theory :

- relies on restricting the complexity of Θ ,
- too conservative as Θ is rarely entirely explored by the algorithm $\{\hat{\theta}_t\}_t$,

Reminder of the previous lecture

In this course, we want to derive **Probably Approximately Correct** generalization bounds for predictors $\hat{\theta}$ or randomized predictors $\hat{\rho}$.

Typical PAC bounds : **with high probability**, the **generalization gap** of θ is **at most something we can control & compute**. For any δ ,

$$\mathbb{P}_{\mathcal{S}} \left[\forall \theta \in \Theta, |R(\theta) - \hat{R}_{\mathcal{S}}(\theta)| \lesssim \sqrt{\frac{\text{comp}(\Theta) + \log(\frac{1}{\delta})}{n}} \right] \geq 1 - \delta.$$

Some limits to this classical statistical learning theory :

- relies on restricting the complexity of Θ ,
- too conservative as Θ is rarely entirely explored by the algorithm $\{\hat{\theta}_t\}_t$,
- **ignores the interaction between the dataset \mathcal{S} and the algorithm $\hat{\theta}$.**

An information-theoretic notion of stability

An information-theoretic notion of stability

- Key idea : a (possibly randomized) learning algorithm $\theta \sim \hat{\rho}_{\mathcal{S}}$ is stable if it doesn't reveal too much information about the dataset \mathcal{S} it was trained on.

An information-theoretic notion of stability

- Key idea : a (possibly randomized) learning algorithm $\theta \sim \hat{\rho}_{\mathcal{S}}$ is stable if it doesn't reveal too much information about the dataset \mathcal{S} it was trained on.
- Question : how do we formalize this dependence ?

An information-theoretic notion of stability

- Key idea : a (possibly randomized) learning algorithm $\theta \sim \hat{\rho}_{\mathcal{S}}$ is stable if it doesn't reveal too much information about the dataset \mathcal{S} it was trained on.
- Question : how do we formalize this dependence ?
- Answer : the mutual information !

$$\mathcal{I}(\hat{\rho}_{\mathcal{S}}; \mathcal{S}) = \mathbb{E}_{\mathcal{S}} [\text{KL}(\hat{\rho}_{\mathcal{S}} \| \mathbb{E}_{\mathcal{S}}[\hat{\rho}_{\mathcal{S}}])] = \text{KL}(P_{\theta, \mathcal{S}} \| P_{\theta} \otimes P_{\mathcal{S}}).$$

An information-theoretic notion of stability

- Key idea : a (possibly randomized) learning algorithm $\theta \sim \hat{\rho}_{\mathcal{S}}$ is stable if it doesn't reveal too much information about the dataset \mathcal{S} it was trained on.
- Question : how do we formalize this dependence ?
- Answer : the mutual information !

$$\mathcal{I}(\hat{\rho}_{\mathcal{S}}; \mathcal{S}) = \mathbb{E}_{\mathcal{S}} [\text{KL}(\hat{\rho}_{\mathcal{S}} \| \mathbb{E}_{\mathcal{S}}[\hat{\rho}_{\mathcal{S}}])] = \text{KL}(P_{\theta, \mathcal{S}} \| P_{\theta} \otimes P_{\mathcal{S}}).$$

- MI quantifies the number of bits of information the algorithm leaks about the training data into the parameters it learns.

An information-theoretic notion of stability

- Key idea : a (possibly randomized) learning algorithm $\theta \sim \hat{\rho}_{\mathcal{S}}$ is stable if it doesn't reveal too much information about the dataset \mathcal{S} it was trained on.
- Question : how do we formalize this dependence ?
- Answer : the mutual information !

$$\mathcal{I}(\hat{\rho}_{\mathcal{S}}; \mathcal{S}) = \mathbb{E}_{\mathcal{S}} [\text{KL}(\hat{\rho}_{\mathcal{S}} \| \mathbb{E}_{\mathcal{S}}[\hat{\rho}_{\mathcal{S}}])] = \text{KL}(P_{\theta, \mathcal{S}} \| P_{\theta} \otimes P_{\mathcal{S}}).$$

- MI quantifies the number of bits of information the algorithm leaks about the training data into the parameters it learns.
- The higher the MI, the worse the stability of the algorithm, the worse the generalization ability.

An information-theoretic notion of stability

- Key idea : a (possibly randomized) learning algorithm $\theta \sim \hat{\rho}_{\mathcal{S}}$ is stable if it doesn't reveal too much information about the dataset \mathcal{S} it was trained on.
- Question : how do we formalize this dependence ?
- Answer : the mutual information !

$$\mathcal{I}(\hat{\rho}_{\mathcal{S}}; \mathcal{S}) = \mathbb{E}_{\mathcal{S}} [\text{KL}(\hat{\rho}_{\mathcal{S}} \| \mathbb{E}_{\mathcal{S}}[\hat{\rho}_{\mathcal{S}}])] = \text{KL}(P_{\theta, \mathcal{S}} \| P_{\theta} \otimes P_{\mathcal{S}}).$$

- MI quantifies the number of bits of information the algorithm leaks about the training data into the parameters it learns.
- The higher the MI, the worse the stability of the algorithm, the worse the generalization ability.

Theorem (Russo & Zhou, '16 / Xu & Raginsky, '17 / Catoni, '07) :
if $\ell(\cdot, \cdot) \leq 1$,

$$\mathbb{E}_{\mathcal{S}} \left[R(\hat{\rho}_{\mathcal{S}}) - \hat{R}_{\mathcal{S}}(\hat{\rho}_{\mathcal{S}}) \right] \leq \sqrt{\frac{2 \cdot \mathcal{I}(\hat{\rho}_{\mathcal{S}}; \mathcal{S})}{n}}.$$

Information-theory vs PAC-Bayes

Information theoretic generalization bound ($\ell(\cdot, \cdot) \leq 1$) :

$$\mathbb{E}_{\mathcal{S}} \left[R(\hat{\rho}_{\mathcal{S}}) - \hat{R}_{\mathcal{S}}(\hat{\rho}_{\mathcal{S}}) \right] \leq \sqrt{\frac{2 \cdot \mathbb{E}_{\mathcal{S}} [\text{KL}(\hat{\rho}_{\mathcal{S}} \parallel \mathbb{E}_{\mathcal{S}}[\hat{\rho}_{\mathcal{S}}])]}{n}}.$$

Information-theory vs PAC-Bayes

Information theoretic generalization bound ($\ell(\cdot, \cdot) \leq 1$) :

$$\mathbb{E}_{\mathcal{S}} \left[R(\hat{\rho}_{\mathcal{S}}) - \hat{R}_{\mathcal{S}}(\hat{\rho}_{\mathcal{S}}) \right] \leq \sqrt{\frac{2 \cdot \mathbb{E}_{\mathcal{S}} [\text{KL}(\hat{\rho}_{\mathcal{S}} \| \mathbb{E}_{\mathcal{S}}[\hat{\rho}_{\mathcal{S}}])]}{n}}.$$

vs PAC-Bayes generalization bound ($\ell(\cdot, \cdot) \leq 1$) :

$$\mathbb{E}_{\mathcal{S}} \left[R(\hat{\rho}_{\mathcal{S}}) - \hat{R}_{\mathcal{S}}(\hat{\rho}_{\mathcal{S}}) \right] \leq \sqrt{\frac{2 \cdot \mathbb{E}_{\mathcal{S}} [\text{KL}(\hat{\rho}_{\mathcal{S}} \| \pi)]}{n}}.$$

Information-theory vs PAC-Bayes

Information theoretic generalization bound ($\ell(\cdot, \cdot) \leq 1$) :

$$\mathbb{E}_{\mathcal{S}} \left[R(\hat{\rho}_{\mathcal{S}}) - \hat{R}_{\mathcal{S}}(\hat{\rho}_{\mathcal{S}}) \right] \leq \sqrt{\frac{2 \cdot \mathbb{E}_{\mathcal{S}} [\text{KL}(\hat{\rho}_{\mathcal{S}} \| \mathbb{E}_{\mathcal{S}}[\hat{\rho}_{\mathcal{S}}])]}{n}}.$$

vs PAC-Bayes generalization bound ($\ell(\cdot, \cdot) \leq 1$) :

$$\mathbb{E}_{\mathcal{S}} \left[R(\hat{\rho}_{\mathcal{S}}) - \hat{R}_{\mathcal{S}}(\hat{\rho}_{\mathcal{S}}) \right] \leq \sqrt{\frac{2 \cdot \mathbb{E}_{\mathcal{S}} [\text{KL}(\hat{\rho}_{\mathcal{S}} \| \pi)]}{n}}.$$

The prior minimizing the PAC-Bayes bound is $\pi := \mathbb{E}_{\mathcal{S}}[\hat{\rho}_{\mathcal{S}}]$.

Information-theory vs PAC-Bayes

Information theoretic generalization bound ($\ell(\cdot, \cdot) \leq 1$) :

$$\mathbb{E}_{\mathcal{S}} \left[R(\hat{\rho}_{\mathcal{S}}) - \hat{R}_{\mathcal{S}}(\hat{\rho}_{\mathcal{S}}) \right] \leq \sqrt{\frac{2 \cdot \mathbb{E}_{\mathcal{S}} [\text{KL}(\hat{\rho}_{\mathcal{S}} \| \mathbb{E}_{\mathcal{S}}[\hat{\rho}_{\mathcal{S}}])]}{n}}.$$

vs PAC-Bayes generalization bound ($\ell(\cdot, \cdot) \leq 1$) :

$$\mathbb{E}_{\mathcal{S}} \left[R(\hat{\rho}_{\mathcal{S}}) - \hat{R}_{\mathcal{S}}(\hat{\rho}_{\mathcal{S}}) \right] \leq \sqrt{\frac{2 \cdot \mathbb{E}_{\mathcal{S}} [\text{KL}(\hat{\rho}_{\mathcal{S}} \| \pi)]}{n}}.$$

The prior minimizing the PAC-Bayes bound is $\pi := \mathbb{E}_{\mathcal{S}}[\hat{\rho}_{\mathcal{S}}]$.

Typical PAC-Bayes bound : **with high probability**, the **generalization gap** of ρ is **at most something we can control & compute**. If we assume that $\ell(\cdot, \cdot) \leq 1$, then $\forall \delta \in (0, 1)$,

$$\mathbb{P}_{\mathcal{S}} \left[\forall \rho \in \mathcal{P}(\Theta), \left| R(\rho) - \hat{R}_{\mathcal{S}}(\rho) \right| \lesssim \sqrt{\frac{\text{KL}(\rho \| \pi) + \log(\frac{1}{\delta})}{n}} \right] \geq 1 - \delta.$$

Overview of the course

The course will be divided in 5 lectures :

- Lecture 1 : Introduction & Motivation
- Lecture 2 : Basics of PAC-Bayes Theory
- Lecture 3 : Advances in PAC-Bayes Theory
- Lecture 4 : Basics of Variational Inference
- Lecture 5 : Advances in Variational Inference

Lecture 3 : Advances in PAC-Bayes Theory

Outline of the lecture

- PAC-Bayes bounds robust to heavy-tails.
- PAC-Bayes bounds achieving fast rates.
- Towards tight certificates in Deep Learning.
- Generalization bounds for SGD using information bounds.

PAC-Bayes bounds robust to heavy-tails

A generic PAC-Bayes bound

A generic PAC-Bayes bound

Germain, Lacasse, Laviolette and Marchand [2009]

For any convex function $\mathcal{D} : [0, 1]^2 \rightarrow \mathbb{R}$, with proba $\geq 1 - \delta$:

$$\forall \rho \in \mathcal{P}(\Theta), \quad \mathcal{D}(\hat{R}_S(\rho), R(\rho)) \leq \frac{\text{KL}(\rho \parallel \pi) + \log \left(\frac{\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} [e^{n\mathcal{D}(\hat{R}_S(\theta), R(\theta))}]}{\delta} \right)}{n}.$$

A generic PAC-Bayes bound

Germain, Lacasse, Laviolette and Marchand [2009]

For any convex function $\mathcal{D} : [0, 1]^2 \rightarrow \mathbb{R}$, with proba $\geq 1 - \delta$:

$$\forall \rho \in \mathcal{P}(\Theta), \quad \mathcal{D}(\hat{R}_S(\rho), R(\rho)) \leq \frac{\text{KL}(\rho \parallel \pi) + \log \left(\frac{\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} [e^{n\mathcal{D}(\hat{R}_S(\theta), R(\theta))}]}{\delta} \right)}{n}.$$

$$\begin{aligned} n\mathcal{D}(\mathbb{E}_{\theta \sim \rho} [\hat{R}_S(\theta)], \mathbb{E}_{\theta \sim \rho} [R(\theta)]) &\leq n \cdot \mathbb{E}_{\theta \sim \rho} [\mathcal{D}(\hat{R}_S(\theta), R(\theta))] \\ &\leq \text{KL}(\rho \parallel \pi) + \log \left(\mathbb{E}_{\theta \sim \pi} [e^{n\mathcal{D}(\hat{R}_S(\theta), R(\theta))}] \right) \\ &\leq \text{KL}(\rho \parallel \pi) + \log \left(\frac{\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} [e^{n\mathcal{D}(\hat{R}_S(\theta), R(\theta))}]}{\delta} \right) \end{aligned}$$

A generic PAC-Bayes bound

Germain, Lacasse, Laviolette and Marchand [2009]

For any convex function $\mathcal{D} : [0, 1]^2 \rightarrow \mathbb{R}$, with proba $\geq 1 - \delta$:

$$\forall \rho \in \mathcal{P}(\Theta), \quad \mathcal{D}(\hat{R}_S(\rho), R(\rho)) \leq \frac{\text{KL}(\rho \parallel \pi) + \log \left(\frac{\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} [e^{n\mathcal{D}(\hat{R}_S(\theta), R(\theta))}]}{\delta} \right)}{n}.$$

$$\begin{aligned} n\mathcal{D}(\mathbb{E}_{\theta \sim \rho} [\hat{R}_S(\theta)], \mathbb{E}_{\theta \sim \rho} [R(\theta)]) &\leq n \cdot \mathbb{E}_{\theta \sim \rho} [\mathcal{D}(\hat{R}_S(\theta), R(\theta))] \\ &\leq \text{KL}(\rho \parallel \pi) + \log \left(\mathbb{E}_{\theta \sim \pi} [e^{n\mathcal{D}(\hat{R}_S(\theta), R(\theta))}] \right) \\ &\leq \text{KL}(\rho \parallel \pi) + \log \left(\frac{\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} [e^{n\mathcal{D}(\hat{R}_S(\theta), R(\theta))}]}{\delta} \right) \end{aligned}$$

Germain's bound is a generalization of both McAllester's and Catoni's bounds (and many others) : if $\ell(\cdot, \cdot) \leq 1$,

$$\text{McAllester [1999]} : \quad R(\rho) \leq \hat{R}_S(\rho) + \sqrt{\frac{\text{KL}(\rho \parallel \pi) + \log \left(\frac{2\sqrt{n}}{\delta} \right)}{2n}}.$$

$$\text{Catoni [2003]} : \quad R(\rho) \leq \hat{R}_S(\rho) + \frac{\text{KL}(\rho \parallel \pi)}{\lambda} + \frac{\lambda}{8n} + \frac{\log \left(\frac{1}{\delta} \right)}{\lambda}.$$

Towards a robust PAC-Bayes bound ?

Towards a robust PAC-Bayes bound ?

Catoni's PAC-Bayes bound [2003]

For any λ , with probability $\geq 1 - \delta : \forall \rho \in \mathcal{P}(\Theta)$,

$$\left| \hat{R}_S(\rho) - R(\rho) \right| \leq \frac{\text{KL}(\rho \parallel \pi) + \log \left(\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} [e^{\lambda |\hat{R}_S(\theta) - R(\theta)|}] \right) + \log \left(\frac{1}{\delta} \right)}{\lambda}.$$

Towards a robust PAC-Bayes bound ?

Catoni's PAC-Bayes bound [2003]

For any λ , with probability $\geq 1 - \delta : \forall \rho \in \mathcal{P}(\Theta)$,

$$\left| \hat{R}_S(\rho) - R(\rho) \right| \leq \frac{\text{KL}(\rho \| \pi) + \log \left(\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} [e^{\lambda |\hat{R}_S(\theta) - R(\theta)|}] \right) + \log \left(\frac{1}{\delta} \right)}{\lambda}.$$

What if the exponential moment does not exist ?

Towards a robust PAC-Bayes bound ?

Catoni's PAC-Bayes bound [2003]

For any λ , with probability $\geq 1 - \delta : \forall \rho \in \mathcal{P}(\Theta)$,

$$\left| \hat{R}_S(\rho) - R(\rho) \right| \leq \frac{\text{KL}(\rho \| \pi) + \log \left(\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} [e^{\lambda |\hat{R}_S(\theta) - R(\theta)|}] \right) + \log \left(\frac{1}{\delta} \right)}{\lambda}.$$

What if the exponential moment does not exist ?

We assume instead a much weaker assumption : for some integer q ,

$$\mathcal{M}_q := \mathbb{E}_S \mathbb{E}_{\theta \sim \pi} \left[\left| \hat{R}_S(\theta) - R(\theta) \right|^q \right] < +\infty.$$

Towards a robust PAC-Bayes bound ?

Catoni's PAC-Bayes bound [2003]

For any λ , with probability $\geq 1 - \delta : \forall \rho \in \mathcal{P}(\Theta)$,

$$\left| \hat{R}_S(\rho) - R(\rho) \right| \leq \frac{\text{KL}(\rho \parallel \pi) + \log \left(\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} [e^{\lambda |\hat{R}_S(\theta) - R(\theta)|}] \right) + \log \left(\frac{1}{\delta} \right)}{\lambda}.$$

What if the exponential moment does not exist ?

We assume instead a much weaker assumption : for some integer q ,

$$\mathcal{M}_q := \mathbb{E}_S \mathbb{E}_{\theta \sim \pi} \left[\left| \hat{R}_S(\theta) - R(\theta) \right|^q \right] < +\infty.$$

To get a PAC-Bayes bound, we need to consider **Csiszàr ϕ -divergences** :
let ϕ be a convex function with $\phi(1) = 0$,

$$D_\phi(\rho \parallel \pi) := \mathbb{E}_\pi \left[\phi \left(\frac{d\rho}{d\pi} \right) \right],$$

when $\rho \ll \pi$ and $D_\phi(\rho \parallel \pi) = +\infty$ otherwise.

Towards a robust PAC-Bayes bound ?

Catoni's PAC-Bayes bound [2003]

For any λ , with probability $\geq 1 - \delta : \forall \rho \in \mathcal{P}(\Theta)$,

$$\left| \hat{R}_S(\rho) - R(\rho) \right| \leq \frac{\text{KL}(\rho \| \pi) + \log \left(\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} [e^{\lambda |\hat{R}_S(\theta) - R(\theta)|}] \right) + \log \left(\frac{1}{\delta} \right)}{\lambda}.$$

What if the exponential moment does not exist ?

We assume instead a much weaker assumption : for some integer q ,

$$\mathcal{M}_q := \mathbb{E}_S \mathbb{E}_{\theta \sim \pi} \left[\left| \hat{R}_S(\theta) - R(\theta) \right|^q \right] < +\infty.$$

To get a PAC-Bayes bound, we need to consider **Csiszàr ϕ -divergences** :
let ϕ be a convex function with $\phi(1) = 0$,

$$D_\phi(\rho \| \pi) := \mathbb{E}_\pi \left[\phi \left(\frac{d\rho}{d\pi} \right) \right],$$

when $\rho \ll \pi$ and $D_\phi(\rho \| \pi) = +\infty$ otherwise.

The KL is given by the special case $\text{KL}(\rho \| \pi) = D_{x \log(x)}(\rho \| \pi)$.

A robust PAC-Bayes bound for heavy tails

Fix $p > 1$, $q = p/(p - 1)$, $\delta \in (0, 1)$ and let $\phi_p : x \mapsto x^p$.

A robust PAC-Bayes bound for heavy tails

Fix $p > 1$, $q = p/(p-1)$, $\delta \in (0, 1)$ and let $\phi_p : x \mapsto x^p$.

Alquier & Guedj PAC-Bayes bound [2018]

With probability $\geq 1 - \delta : \forall \rho \in \mathcal{P}(\Theta)$,

$$\left| \hat{R}_S(\rho) - R(\rho) \right| \leq \left(\frac{\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} \left[\left| \hat{R}_S(\theta) - R(\theta) \right|^q \right]}{\delta} \right)^{\frac{1}{q}} \cdot \left(\mathbb{E}_{\pi} \left[\left(\frac{d\rho}{d\pi} \right)^p \right] \right)^{\frac{1}{p}}$$

A robust PAC-Bayes bound for heavy tails

Fix $p > 1$, $q = p/(p-1)$, $\delta \in (0, 1)$ and let $\phi_p : x \mapsto x^p$.

Alquier & Guedj PAC-Bayes bound [2018]

With probability $\geq 1 - \delta : \forall \rho \in \mathcal{P}(\Theta)$,

$$\left| \hat{R}_S(\rho) - R(\rho) \right| \leq \left(\frac{\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} \left[\left| \hat{R}_S(\theta) - R(\theta) \right|^q \right]}{\delta} \right)^{\frac{1}{q}} \cdot \left(\mathbb{E}_{\pi} \left[\left(\frac{d\rho}{d\pi} \right)^p \right] \right)^{\frac{1}{p}}$$

The bound decouples :

- The moment \mathcal{M}_q (depending on the distribution of the data).
- The divergence $D_{\phi_p-1}(\rho \parallel \pi) + 1$ (measure of complexity).

A robust PAC-Bayes bound for heavy tails

Fix $p > 1$, $q = p/(p-1)$, $\delta \in (0, 1)$ and let $\phi_p : x \mapsto x^p$.

Alquier & Guedj PAC-Bayes bound [2018]

With probability $\geq 1 - \delta : \forall \rho \in \mathcal{P}(\Theta)$,

$$\left| \hat{R}_S(\rho) - R(\rho) \right| \leq \left(\frac{\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} \left[\left| \hat{R}_S(\theta) - R(\theta) \right|^q \right]}{\delta} \right)^{\frac{1}{q}} \cdot \left(\mathbb{E}_{\pi} \left[\left(\frac{d\rho}{d\pi} \right)^p \right] \right)^{\frac{1}{p}}$$

The bound decouples :

- The moment \mathcal{M}_q (depending on the distribution of the data).
- The divergence $D_{\phi_p-1}(\rho \parallel \pi) + 1$ (measure of complexity).

Note the weak dependence $\delta^{-1/q}$ vs $\sqrt{\log(1/\delta)}$ (there's no free lunch)...

A robust PAC-Bayes bound for heavy tails

Fix $p > 1$, $q = p/(p-1)$, $\delta \in (0, 1)$ and let $\phi_p : x \mapsto x^p$.

Alquier & Guedj PAC-Bayes bound [2018]

With probability $\geq 1 - \delta : \forall \rho \in \mathcal{P}(\Theta)$,

$$\left| \hat{R}_S(\rho) - R(\rho) \right| \leq \left(\frac{\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} \left[\left| \hat{R}_S(\theta) - R(\theta) \right|^q \right]}{\delta} \right)^{\frac{1}{q}} \cdot \left(\mathbb{E}_{\pi} \left[\left(\frac{d\rho}{d\pi} \right)^p \right] \right)^{\frac{1}{p}}$$

The bound decouples :

- The moment \mathcal{M}_q (depending on the distribution of the data).
- The divergence $D_{\phi_{p-1}}(\rho \parallel \pi) + 1$ (measure of complexity).

Note the weak dependence $\delta^{-1/q}$ vs $\sqrt{\log(1/\delta)}$ (there's no free lunch)...

For $p = q = 2$, for $\mathcal{V} := \mathbb{E}_{\theta \sim \pi} \mathbb{V}_{(X,Y) \sim \mathbb{P}}[\ell(f_{\theta}(x), y)] < +\infty$,

A robust PAC-Bayes bound for heavy tails

Fix $p > 1$, $q = p/(p-1)$, $\delta \in (0, 1)$ and let $\phi_p : x \mapsto x^p$.

Alquier & Guedj PAC-Bayes bound [2018]

With probability $\geq 1 - \delta : \forall \rho \in \mathcal{P}(\Theta)$,

$$\left| \hat{R}_S(\rho) - R(\rho) \right| \leq \left(\frac{\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} \left[\left| \hat{R}_S(\theta) - R(\theta) \right|^q \right]}{\delta} \right)^{\frac{1}{q}} \cdot \left(\mathbb{E}_{\pi} \left[\left(\frac{d\rho}{d\pi} \right)^p \right] \right)^{\frac{1}{p}}$$

The bound decouples :

- The moment \mathcal{M}_q (depending on the distribution of the data).
- The divergence $D_{\phi_{p-1}}(\rho \parallel \pi) + 1$ (measure of complexity).

Note the weak dependence $\delta^{-1/q}$ vs $\sqrt{\log(1/\delta)}$ (there's no free lunch)...

For $p = q = 2$, for $\mathcal{V} := \mathbb{E}_{\theta \sim \pi} \mathbb{V}_{(X,Y) \sim \mathbb{P}}[\ell(f_{\theta}(x), y)] < +\infty$, w.p. $\geq 1 - \delta$,

$$\forall \rho \in \mathcal{P}(\Theta), \quad R(\rho) \leq \hat{R}_S(\rho) + \sqrt{\frac{\mathcal{V}(1 + \chi^2(\rho \parallel \pi))}{n\delta}}.$$

Proof of Alquier & Guedj's bound

Alquier & Guedj PAC-Bayes bound [2018]

With probability $\geq 1 - \delta : \forall \rho \in \mathcal{P}(\Theta)$,

$$\left| \hat{R}_S(\rho) - R(\rho) \right| \leq \left(\frac{\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} \left[\left| \hat{R}_S(\theta) - R(\theta) \right|^q \right]}{\delta} \right)^{\frac{1}{q}} \cdot \left(\mathbb{E}_{\pi} \left[\left(\frac{d\rho}{d\pi} \right)^p \right] \right)^{\frac{1}{p}}.$$

Proof of Alquier & Guedj's bound

Alquier & Guedj PAC-Bayes bound [2018]

With probability $\geq 1 - \delta : \forall \rho \in \mathcal{P}(\Theta)$,

$$\left| \hat{R}_S(\rho) - R(\rho) \right| \leq \left(\frac{\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} \left[\left| \hat{R}_S(\theta) - R(\theta) \right|^q \right]}{\delta} \right)^{\frac{1}{q}} \cdot \left(\mathbb{E}_{\pi} \left[\left(\frac{d\rho}{d\pi} \right)^p \right] \right)^{\frac{1}{p}}.$$

$$\begin{aligned} \left| \mathbb{E}_{\theta \sim \rho} \left[\hat{R}_S(\theta) \right] - \mathbb{E}_{\theta \sim \rho} \left[R(\theta) \right] \right| &\leq \mathbb{E}_{\theta \sim \rho} \left[\left| \hat{R}_S(\theta) - R(\theta) \right| \right] \\ &= \mathbb{E}_{\theta \sim \pi} \left[\left| \hat{R}_S(\theta) - R(\theta) \right| \frac{d\rho}{d\pi}(\theta) \right] \\ &\leq \mathbb{E}_{\theta \sim \pi} \left[\left| \hat{R}_S(\theta) - R(\theta) \right|^q \right]^{\frac{1}{q}} \cdot \mathbb{E}_{\pi} \left[\left(\frac{d\rho}{d\pi} \right)^p \right]^{\frac{1}{p}} \\ &\leq \frac{\mathbb{E}_S \mathbb{E}_{\theta \sim \pi} \left[\left| \hat{R}_S(\theta) - R(\theta) \right|^q \right]^{\frac{1}{q}}}{\delta^{\frac{1}{q}}} \cdot \mathbb{E}_{\pi} \left[\left(\frac{d\rho}{d\pi} \right)^p \right]^{\frac{1}{p}} \end{aligned}$$

PAC-Bayes bounds achieving fast rates

Reminder on Tolstikhin's bound

Reminder on Tolstikhin's bound

Tolstikhin's bound

With proba $\geq 1 - \delta$ ($\ell(\cdot, \cdot) \leq 1$),

$$\forall \rho \in \mathcal{P}(\Theta), \quad R(\rho) \leq \hat{R}_S(\rho) + \sqrt{2\hat{R}_S(\rho) \frac{\text{KL}(\rho \parallel \pi) + \log\left(\frac{2\sqrt{n}}{\delta}\right)}{n}} + 2 \frac{\text{KL}(\rho \parallel \pi) + \log\left(\frac{2\sqrt{n}}{\delta}\right)}{n}.$$

Reminder on Tolstikhin's bound

Tolstikhin's bound

With proba $\geq 1 - \delta$ ($\ell(\cdot, \cdot) \leq 1$),

$$\forall \rho \in \mathcal{P}(\Theta), \quad R(\rho) \leq \hat{R}_S(\rho) + \sqrt{2\hat{R}_S(\rho) \frac{\text{KL}(\rho \parallel \pi) + \log\left(\frac{2\sqrt{n}}{\delta}\right)}{n}} + 2 \frac{\text{KL}(\rho \parallel \pi) + \log\left(\frac{2\sqrt{n}}{\delta}\right)}{n}.$$

We get this bound from Seeger's one via a refinement of Pinsker's inequality $\text{kl}(p \parallel q) \geq (p - q)^2 / 2q$ i.e. $\text{kl}^{-1}(q \parallel b) \leq q + \sqrt{2qb} + 2b$, hence improving over McAllester's bound.

Reminder on Tolstikhin's bound

Tolstikhin's bound

With proba $\geq 1 - \delta$ ($\ell(\cdot, \cdot) \leq 1$),

$$\forall \rho \in \mathcal{P}(\Theta), \quad R(\rho) \leq \hat{R}_S(\rho) + \sqrt{2\hat{R}_S(\rho) \frac{\text{KL}(\rho \parallel \pi) + \log\left(\frac{2\sqrt{n}}{\delta}\right)}{n}} + 2 \frac{\text{KL}(\rho \parallel \pi) + \log\left(\frac{2\sqrt{n}}{\delta}\right)}{n}.$$

We get this bound from Seeger's one via a refinement of Pinsker's inequality $\text{kl}(p \parallel q) \geq (p - q)^2 / 2q$ i.e. $\text{kl}^{-1}(q \parallel b) \leq q + \sqrt{2qb} + 2b$, hence improving over McAllester's bound.

An amazing characteristic of this bound : while its order is in general in $1/\sqrt{n}$, as all the PAC-Bayes bounds seen so far, the dependence drops to $1/n$ in the noiseless case where $\hat{R}_S(\rho) = 0$.

Reminder on Tolstikhin's bound

Tolstikhin's bound

With proba $\geq 1 - \delta$ ($\ell(\cdot, \cdot) \leq 1$),

$$\forall \rho \in \mathcal{P}(\Theta), \quad R(\rho) \leq \hat{R}_S(\rho) + \sqrt{2\hat{R}_S(\rho) \frac{\text{KL}(\rho \parallel \pi) + \log\left(\frac{2\sqrt{n}}{\delta}\right)}{n}} + 2 \frac{\text{KL}(\rho \parallel \pi) + \log\left(\frac{2\sqrt{n}}{\delta}\right)}{n}.$$

We get this bound from Seeger's one via a refinement of Pinsker's inequality $\text{kl}(p \parallel q) \geq (p - q)^2 / 2q$ i.e. $\text{kl}^{-1}(q \parallel b) \leq q + \sqrt{2qb} + 2b$, hence improving over McAllester's bound.

An amazing characteristic of this bound : while its order is in general in $1/\sqrt{n}$, as all the PAC-Bayes bounds seen so far, the dependence drops to $1/n$ in the noiseless case where $\hat{R}_S(\rho) = 0$.

Question : is it possible to achieve fast rates more generally ?

Reminder on Tolstikhin's bound

Tolstikhin's bound

With proba $\geq 1 - \delta$ ($\ell(\cdot, \cdot) \leq 1$),

$$\forall \rho \in \mathcal{P}(\Theta), \quad R(\rho) \leq \hat{R}_S(\rho) + \sqrt{2\hat{R}_S(\rho) \frac{\text{KL}(\rho \parallel \pi) + \log\left(\frac{2\sqrt{n}}{\delta}\right)}{n}} + 2 \frac{\text{KL}(\rho \parallel \pi) + \log\left(\frac{2\sqrt{n}}{\delta}\right)}{n}.$$

We get this bound from Seeger's one via a refinement of Pinsker's inequality $\text{kl}(p \parallel q) \geq (p - q)^2 / 2q$ i.e. $\text{kl}^{-1}(q \parallel b) \leq q + \sqrt{2qb} + 2b$, hence improving over McAllester's bound.

An amazing characteristic of this bound : while its order is in general in $1/\sqrt{n}$, as all the PAC-Bayes bounds seen so far, the dependence drops to $1/n$ in the noiseless case where $\hat{R}_S(\rho) = 0$.

Question : is it possible to achieve fast rates more generally? Yes!
Under some specific required assumptions.

Oracle inequalities : controlling the excess risk

Let's start in a deterministic setting where we want to quantify the performance of $f_{\hat{\theta}}$ via *excess risk bounds/oracle inequalities*.

Oracle inequalities : controlling the excess risk

Let's start in a deterministic setting where we want to quantify the performance of $f_{\hat{\theta}}$ via *excess risk bounds/oracle inequalities*.

Given \mathcal{S} , how well does $f_{\hat{\theta}}$ predict on unseen data ?

$$R(\hat{\theta}) = \mathbb{E} [\ell(f_{\hat{\theta}}(X), Y) | \mathcal{S}] .$$

Oracle inequalities : controlling the excess risk

Let's start in a deterministic setting where we want to quantify the performance of $f_{\hat{\theta}}$ via *excess risk bounds/oracle inequalities*.

Given \mathcal{S} , how well does $f_{\hat{\theta}}$ predict on unseen data ?

$$R(\hat{\theta}) = \mathbb{E} [\ell(f_{\hat{\theta}}(X), Y) | \mathcal{S}].$$

What is the best predictor I could (should?) have chosen ?

$$R(\theta^*) = \inf_{\theta \in \Theta} \mathbb{E} [\ell(f_{\theta}(X), Y)].$$

Oracle inequalities : controlling the excess risk

Let's start in a deterministic setting where we want to quantify the performance of $f_{\hat{\theta}}$ via *excess risk bounds/oracle inequalities*.

Given \mathcal{S} , how well does $f_{\hat{\theta}}$ predict on unseen data ?

$$R(\hat{\theta}) = \mathbb{E} [\ell(f_{\hat{\theta}}(X), Y) | \mathcal{S}].$$

What is the best predictor I could (should?) have chosen ?

$$R(\theta^*) = \inf_{\theta \in \Theta} \mathbb{E} [\ell(f_{\theta}(X), Y)].$$

With high probability, the excess risk of $f_{\hat{\theta}}$ within Θ is expected to be (for bounded losses), for any δ , of order,

$$\mathbb{P}_{\mathcal{S}} \left[R(\hat{\theta}) - \inf_{\theta \in \Theta} R(\theta) \lesssim \sqrt{\frac{\text{comp}(\Theta)}{n}} \times \sqrt{\log \left(\frac{1}{\delta} \right)} \right] \geq 1 - \delta.$$

Oracle inequalities : controlling the excess risk

Let's start in a deterministic setting where we want to quantify the performance of $f_{\hat{\theta}}$ via *excess risk bounds/oracle inequalities*.

Given \mathcal{S} , how well does $f_{\hat{\theta}}$ predict on unseen data ?

$$R(\hat{\theta}) = \mathbb{E} [\ell(f_{\hat{\theta}}(X), Y) | \mathcal{S}].$$

What is the best predictor I could (should?) have chosen ?

$$R(\theta^*) = \inf_{\theta \in \Theta} \mathbb{E} [\ell(f_{\theta}(X), Y)].$$

With high probability, the excess risk of $f_{\hat{\theta}}$ within Θ is expected to be (for bounded losses), for any δ , of order,

$$\mathbb{P}_{\mathcal{S}} \left[R(\hat{\theta}) - \inf_{\theta \in \Theta} R(\theta) \lesssim \sqrt{\frac{\text{comp}(\Theta)}{n}} \times \sqrt{\log \left(\frac{1}{\delta} \right)} \right] \geq 1 - \delta.$$

Is it possible to achieve faster rates for bounded losses ?

Oracle inequalities : controlling the excess risk

Let's start in a deterministic setting where we want to quantify the performance of $f_{\hat{\theta}}$ via *excess risk bounds/oracle inequalities*.

Given \mathcal{S} , how well does $f_{\hat{\theta}}$ predict on unseen data ?

$$R(\hat{\theta}) = \mathbb{E} [\ell(f_{\hat{\theta}}(X), Y) | \mathcal{S}].$$

What is the best predictor I could (should?) have chosen ?

$$R(\theta^*) = \inf_{\theta \in \Theta} \mathbb{E} [\ell(f_{\theta}(X), Y)].$$

With high probability, the excess risk of $f_{\hat{\theta}}$ within Θ is expected to be (for bounded losses), for any δ , of order,

$$\mathbb{P}_{\mathcal{S}} \left[R(\hat{\theta}) - \inf_{\theta \in \Theta} R(\theta) \lesssim \sqrt{\frac{\text{comp}(\Theta)}{n}} \times \sqrt{\log \left(\frac{1}{\delta} \right)} \right] \geq 1 - \delta.$$

Is it possible to achieve faster rates for bounded losses? Yes!
Under further assumptions on the "easiness" of the problem.

Faster rates of convergence

Actually, the optimal excess risk of a rule $f_{\hat{\theta}}$ is usually of order

$$R(\hat{\theta}) - \inf_{\theta \in \Theta} R(\theta) \lesssim \left(\frac{\text{comp}(\Theta)}{n} \right)^{\gamma}$$

where $\gamma \in [\frac{1}{2}, 1]$ reflects the easiness of the problem $(\mathbb{P}, \ell, \Theta)$.

Faster rates of convergence

Actually, the optimal excess risk of a rule $f_{\hat{\theta}}$ is usually of order

$$R(\hat{\theta}) - \inf_{\theta \in \Theta} R(\theta) \lesssim \left(\frac{\text{comp}(\Theta)}{n} \right)^{\gamma}$$

where $\gamma \in [\frac{1}{2}, 1]$ reflects the easiness of the problem $(\mathbb{P}, \ell, \Theta)$.

- If the problem is "hard" (e.g. no specific assumption), $\gamma = 1/2$ and :

$$R(\hat{\theta}) - \inf_{\theta \in \Theta} R(\theta) \lesssim \sqrt{\frac{\text{comp}(\Theta)}{n}}.$$

Faster rates of convergence

Actually, the optimal excess risk of a rule $f_{\hat{\theta}}$ is usually of order

$$R(\hat{\theta}) - \inf_{\theta \in \Theta} R(\theta) \lesssim \left(\frac{\text{comp}(\Theta)}{n} \right)^{\gamma}$$

where $\gamma \in [\frac{1}{2}, 1]$ reflects the easiness of the problem $(\mathbb{P}, \ell, \Theta)$.

- If the problem is "hard" (e.g. no specific assumption), $\gamma = 1/2$ and :

$$R(\hat{\theta}) - \inf_{\theta \in \Theta} R(\theta) \lesssim \sqrt{\frac{\text{comp}(\Theta)}{n}}.$$

- If the problem is "easy" (e.g. noiseless/low-noise settings), $\gamma = 1$ and :

$$R(\hat{\theta}) - \inf_{\theta \in \Theta} R(\theta) \lesssim \frac{\text{comp}(\Theta)}{n}.$$

Fast rates in classification

Consider the following binary classification problem $(\mathbb{P}, \ell, \Theta)$:

Fast rates in classification

Consider the following binary classification problem $(\mathbb{P}, \ell, \Theta)$:

- $\ell(y', y) = \mathbb{1}(y' \neq y)$ so that $R(\theta) = \mathbb{P}(f_\theta(X) \neq Y)$

Fast rates in classification

Consider the following binary classification problem $(\mathbb{P}, \ell, \Theta)$:

- $\ell(y', y) = \mathbb{1}(y' \neq y)$ so that $R(\theta) = \mathbb{P}(f_\theta(X) \neq Y)$
- $f^*(x) = \arg \max_y \mathbb{P}(Y = y | X = x) \in \{f_\theta; \theta \in \Theta\}$
(assumption)

Fast rates in classification

Consider the following binary classification problem $(\mathbb{P}, \ell, \Theta)$:

- $\ell(y', y) = \mathbb{1}(y' \neq y)$ so that $R(\theta) = \mathbb{P}(f_\theta(X) \neq Y)$
- $f^*(x) = \arg \max_y \mathbb{P}(Y = y | X = x) \in \{f_\theta; \theta \in \Theta\}$
(assumption)

Question : when is the problem difficult ?

Fast rates in classification

Consider the following binary classification problem $(\mathbb{P}, \ell, \Theta)$:

- $\ell(y', y) = \mathbb{1}(y' \neq y)$ so that $R(\theta) = \mathbb{P}(f_\theta(X) \neq Y)$
- $f^*(x) = \arg \max_y \mathbb{P}(Y = y|X = x) \in \{f_\theta; \theta \in \Theta\}$
(assumption)

Question : when is the problem difficult ?

- Answer : when $\mathbb{P}(Y = 1|X)$ very close to $1/2$! But then learning is (almost) useless...

Fast rates in classification

Consider the following binary classification problem $(\mathbb{P}, \ell, \Theta)$:

- $\ell(y', y) = \mathbb{1}(y' \neq y)$ so that $R(\theta) = \mathbb{P}(f_\theta(X) \neq Y)$
- $f^*(x) = \arg \max_y \mathbb{P}(Y = y|X = x) \in \{f_\theta; \theta \in \Theta\}$
(assumption)

Question : when is the problem difficult ?

- Answer : when $\mathbb{P}(Y = 1|X)$ very close to $1/2$! But then learning is (almost) useless...
- Solution : assume that $\mathbb{P}(Y = 1|X)$ not too close to $1/2$!

Fast rates in classification

Consider the following binary classification problem $(\mathbb{P}, \ell, \Theta)$:

- $\ell(y', y) = \mathbb{1}(y' \neq y)$ so that $R(\theta) = \mathbb{P}(f_\theta(X) \neq Y)$
- $f^*(x) = \arg \max_y \mathbb{P}(Y = y|X = x) \in \{f_\theta; \theta \in \Theta\}$
(assumption)

Question : when is the problem difficult ?

- Answer : when $\mathbb{P}(Y = 1|X)$ very close to $1/2$! But then learning is (almost) useless...
- Solution : assume that $\mathbb{P}(Y = 1|X)$ not too close to $1/2$!

w.h.p. over X , $\left| \mathbb{P}(Y = 1|X) - \frac{1}{2} \right|$ is large.

The margin condition in classification


Tsybakov's α -margin condition [Tsybakov, AoS 2004]

$$\mathbb{P}_X \left[\left| \mathbb{P}(Y = 1|X) - \frac{1}{2} \right| \leq t \right] \leq ct^\alpha.$$

The margin condition in classification


Tsybakov's α -margin condition [Tsybakov, AoS 2004]

$$\mathbb{P}_X \left[\left| \mathbb{P}(Y = 1|X) - \frac{1}{2} \right| \leq t \right] \leq ct^\alpha.$$




easy

$\alpha = \infty$



moderate

$\alpha = 1$




hard

$\alpha = 0$


The margin condition in classification

Tsybakov's α -margin condition [Tsybakov, AoS 2004]


$$\mathbb{P}_X \left[\left| \mathbb{P}(Y = 1|X) - \frac{1}{2} \right| \leq t \right] \leq ct^\alpha.$$



easy
 $\alpha = \infty$



moderate
 $\alpha = 1$



hard
 $\alpha = 0$

$$R(\hat{\theta}) - \inf_{\theta \in \Theta} R(\theta) \lesssim \left(\frac{\text{comp}(\Theta)}{n} \right)^{\frac{1+\alpha}{2+\alpha}}.$$

The general setting : Bernstein condition

Bernstein's condition [Bartlett & Mendelson, PTFR 2006]

For some $\beta \in [0, 1]$ and $B > 0$, with the notation $\ell_\theta = \ell(f_\theta(X), Y)$,

$$\mathbb{E} \left[(\ell_\theta - \ell_{\theta^*})^2 \right] \leq B \mathbb{E} [\ell_\theta - \ell_{\theta^*}]^\beta.$$

The general setting : Bernstein condition

Bernstein's condition [Bartlett & Mendelson, PTFR 2006]

For some $\beta \in [0, 1]$ and $B > 0$, with the notation $\ell_\theta = \ell(f_\theta(X), Y)$,

$$\mathbb{E} \left[(\ell_\theta - \ell_{\theta^*})^2 \right] \leq B \mathbb{E} [\ell_\theta - \ell_{\theta^*}]^\beta.$$

Under β -Bernstein condition,

$$R(\hat{\theta}) - \inf_{\theta \in \Theta} R(\theta) \lesssim \left(\frac{\text{comp}(\Theta)}{n} \right)^{\frac{1}{2-\beta}}.$$

The general setting : Bernstein condition

Bernstein's condition [Bartlett & Mendelson, PTFR 2006]

For some $\beta \in [0, 1]$ and $B > 0$, with the notation $\ell_\theta = \ell(f_\theta(X), Y)$,

$$\mathbb{E} \left[(\ell_\theta - \ell_{\theta^*})^2 \right] \leq B \mathbb{E} [\ell_\theta - \ell_{\theta^*}]^\beta.$$


Under β -Bernstein condition,

$$R(\hat{\theta}) - \inf_{\theta \in \Theta} R(\theta) \lesssim \left(\frac{\text{comp}(\Theta)}{n} \right)^{\frac{1}{2-\beta}}.$$

Bernstein condition is satisfied in the following settings :

- In noiseless classification $R(\theta^*) = 0$, with $\beta = 1$.
- Under Tsybakov's α -margin assumption, with $\beta = \frac{\alpha}{1+\alpha}$.
- For Lipschitz and strongly-convex loss functions, with $\beta = 1$.

Welcome to the zoo [Van Erven et al., JMLR 2015]



How about fast PAC-Bayes bounds ?

Reminder on the Gibbs posterior :

$$\hat{\rho}_{\lambda}(d\theta) := \arg \min_{\rho \in \mathcal{P}(\Theta)} \left\{ \hat{R}_S(\rho) + \frac{\text{KL}(\rho \parallel \pi)}{\lambda} \right\} \propto \exp \left(-\lambda \sum_{i=1}^n \ell(f_{\theta}(X_i), Y_i) \right) \pi(d\theta).$$

How about fast PAC-Bayes bounds ?

Reminder on the Gibbs posterior :

$$\hat{\rho}_\lambda(d\theta) := \arg \min_{\rho \in \mathcal{P}(\Theta)} \left\{ \hat{R}_S(\rho) + \frac{\text{KL}(\rho \parallel \pi)}{\lambda} \right\} \propto \exp \left(-\lambda \sum_{i=1}^n \ell(f_\theta(X_i), Y_i) \right) \pi(d\theta).$$

We denote $\mathcal{R}(\theta) = R(\theta) - \inf_{\Theta} R$ the excess risk, and assume $\ell(\cdot, \cdot) \leq 1$.

How about fast PAC-Bayes bounds ?

Reminder on the Gibbs posterior :

$$\hat{\rho}_{\lambda}(d\theta) := \arg \min_{\rho \in \mathcal{P}(\Theta)} \left\{ \hat{R}_S(\rho) + \frac{\text{KL}(\rho \parallel \pi)}{\lambda} \right\} \propto \exp \left(-\lambda \sum_{i=1}^n \ell(f_{\theta}(X_i), Y_i) \right) \pi(d\theta).$$

We denote $\mathcal{R}(\theta) = R(\theta) - \inf_{\Theta} R$ the excess risk, and assume $\ell(\cdot, \cdot) \leq 1$.

Slow excess risk bound (in expectation) for the Gibbs posterior :

$$\mathbb{E}_S \mathbb{E}_{\theta \sim \hat{\rho}_{\lambda}} [\mathcal{R}(\theta)] \leq \mathbb{E}_S \left[\inf_{\rho \in \mathcal{P}(\Theta)} \left\{ \mathbb{E}_{\theta \sim \rho} [\mathcal{R}(\theta)] + \frac{\text{KL}(\rho \parallel \pi)}{\lambda} \right\} \right] + \frac{\lambda}{8n}.$$

How about fast PAC-Bayes bounds ?

Reminder on the Gibbs posterior :

$$\hat{\rho}_\lambda(d\theta) := \arg \min_{\rho \in \mathcal{P}(\Theta)} \left\{ \hat{R}_S(\rho) + \frac{\text{KL}(\rho \parallel \pi)}{\lambda} \right\} \propto \exp \left(-\lambda \sum_{i=1}^n \ell(f_\theta(X_i), Y_i) \right) \pi(d\theta).$$

We denote $\mathcal{R}(\theta) = R(\theta) - \inf_{\Theta} R$ the excess risk, and assume $\ell(\cdot, \cdot) \leq 1$.

Slow excess risk bound (in expectation) for the Gibbs posterior :

$$\mathbb{E}_S \mathbb{E}_{\theta \sim \hat{\rho}_\lambda} [\mathcal{R}(\theta)] \leq \mathbb{E}_S \left[\inf_{\rho \in \mathcal{P}(\Theta)} \left\{ \mathbb{E}_{\theta \sim \rho} [\mathcal{R}(\theta)] + \frac{\text{KL}(\rho \parallel \pi)}{\lambda} \right\} \right] + \frac{\lambda}{8n}.$$

Fast excess risk bound (in expectation) for the Gibbs posterior under Bernstein's condition ($\beta = 1$) :

$$\mathbb{E}_S \mathbb{E}_{\theta \sim \hat{\rho}_\lambda} [\mathcal{R}(\theta)] \leq \frac{1}{1 - \frac{B\lambda/n}{2(1-\lambda/n)}} \mathbb{E}_S \left[\inf_{\rho \in \mathcal{P}(\Theta)} \left\{ \mathbb{E}_{\theta \sim \rho} [\mathcal{R}(\theta)] + \frac{\text{KL}(\rho \parallel \pi)}{\lambda} \right\} \right].$$

How about fast PAC-Bayes bounds?

Reminder on the Gibbs posterior :

$$\hat{\rho}_\lambda(d\theta) := \arg \min_{\rho \in \mathcal{P}(\Theta)} \left\{ \hat{R}_S(\rho) + \frac{\text{KL}(\rho \parallel \pi)}{\lambda} \right\} \propto \exp \left(-\lambda \sum_{i=1}^n \ell(f_\theta(X_i), Y_i) \right) \pi(d\theta).$$

We denote $\mathcal{R}(\theta) = R(\theta) - \inf_{\Theta} R$ the excess risk, and assume $\ell(\cdot, \cdot) \leq 1$.

Slow excess risk bound (in expectation) for the Gibbs posterior :

$$\mathbb{E}_S \mathbb{E}_{\theta \sim \hat{\rho}_\lambda} [\mathcal{R}(\theta)] \leq \mathbb{E}_S \left[\inf_{\rho \in \mathcal{P}(\Theta)} \left\{ \mathbb{E}_{\theta \sim \rho} [\mathcal{R}(\theta)] + \frac{\text{KL}(\rho \parallel \pi)}{\lambda} \right\} \right] + \frac{\lambda}{8n}.$$

Fast excess risk bound (in expectation) for the Gibbs posterior under Bernstein's condition ($\beta = 1$) :


$$\mathbb{E}_S \mathbb{E}_{\theta \sim \hat{\rho}_\lambda} [\mathcal{R}(\theta)] \leq \frac{1}{1 - \frac{B\lambda/n}{2(1-\lambda/n)}} \mathbb{E}_S \left[\inf_{\rho \in \mathcal{P}(\Theta)} \left\{ \mathbb{E}_{\theta \sim \rho} [\mathcal{R}(\theta)] + \frac{\text{KL}(\rho \parallel \pi)}{\lambda} \right\} \right].$$

For $\lambda = n/(1+B)$, under Bernstein's condition ($\beta = 1$) :

$$\mathbb{E}_S \mathbb{E}_{\theta \sim \hat{\rho}_\lambda} [\mathcal{R}(\theta)] \leq 2 \cdot \mathbb{E}_S \left[\inf_{\rho \in \mathcal{P}(\Theta)} \left\{ \mathbb{E}_{\theta \sim \rho} [\mathcal{R}(\theta)] + \frac{(1+B)\text{KL}(\rho \parallel \pi)}{n} \right\} \right].$$


Towards Tight Certificates in Deep Learning

Rethinking generalization with DL



- Many parameters !
- NNs trained with SGD achieve 0 training error.
- NNs can overfit but in practice don't : **why** ?
- **Hypothesis** : complexity \ll number of parameters.

Rethinking generalization with DL



(b)

A breakthrough : [Dziugaite and Roy, 2017]

[Dziugaite and Roy, 2017] achieved non-vacuous bounds on binary MNIST using PAC-Bayes bounds (≈ 0.2 vs 0.03 test error).

A breakthrough : [Dziugaite and Roy, 2017]

[Dziugaite and Roy, 2017] achieved non-vacuous bounds on binary MNIST using PAC-Bayes bounds (≈ 0.2 vs 0.03 test error).

- Choose a Gaussian posterior $\rho_{w,s^2} = \mathcal{N}(w, s^2 I_p)$ and minimize McAllester's PAC-Bayes bound wrt (w, s^2) .
- Upper bound the 0-1 loss by a convex, Lipschitz upper bound in order to make the bound easier to minimize
$$\mathbb{1}(f_\theta(x) \neq y) \leq \log(1 + e^{-y f_\theta(x)}) / \log(2).$$
- Use SGD to solve the optimization problem (importance of achieving flat minima).
- Important : use a data-dependent prior ! Optimize the prior variance (union bound argument), mean equal to 0 or randomly chosen.
- Do not use $\text{kl}(\hat{R}_S(\rho), R(\rho)) \geq 2 \left(R(\rho) - \hat{R}_S(\rho)\right)^2$ but (right) invert the kl directly \rightarrow evaluate the subsequent bound at $\hat{\rho}_S$.

Two different bounds

Langford & Seeger's PAC-Bayes bound

With probability $\geq 1 - \delta$,

$$\forall \rho \in \mathcal{P}(\mathcal{F}), \quad R(\rho) \leq \text{kl}^{-1} \left(\hat{R}_S(\rho) \left\| \frac{\text{KL}(\rho \parallel \pi) + \log \left(\frac{2\sqrt{n}}{\delta} \right)}{n} \right\| \right).$$

Two different bounds

Langford & Seeger's PAC-Bayes bound

With probability $\geq 1 - \delta$,

$$\forall \rho \in \mathcal{P}(\mathcal{F}), \quad R(\rho) \leq \text{kl}^{-1} \left(\hat{R}_S(\rho) \parallel \frac{\text{KL}(\rho \parallel \pi) + \log \left(\frac{2\sqrt{n}}{\delta} \right)}{n} \right).$$

This leads to two different PAC-Bayes bounds :

- A bound for the training stage (not tight) : wp $\geq 1 - \delta$ over data samples, uniformly over $\rho \in \mathcal{P}(\mathcal{F})$,

$$R^x(\rho) \leq \hat{R}_S^x(\rho) + \sqrt{\frac{\text{KL}(\rho \parallel \pi) + \log \left(\frac{2\sqrt{n}}{\delta} \right)}{2n}}.$$

- A bound for the evaluation stage (not practical) : wp $\geq 1 - \delta - \delta'$ over data + MC samples, uniformly over $\rho \in \mathcal{P}(\mathcal{F})$,

$$R^{01}(\rho) \leq \text{kl}^{-1} \left(\text{kl}^{-1} \left(\hat{R}_S^{01}(\tilde{\rho}_m), \frac{\log \left(\frac{2\sqrt{m}}{\delta'} \right)}{m} \right), \frac{\text{KL}(\rho \parallel \pi) + \log \left(\frac{2\sqrt{n}}{\delta} \right)}{2n} \right).$$

Algorithm 1 PAC-Bayes with Backprop (PBB)

Input:

μ_0 \triangleright Prior center parameters (random init.)
 ρ_0 \triangleright Prior scale hyper-parameter
 $Z_{1:n}$ \triangleright Training examples (inputs + labels)
 $\delta \in (0, 1)$ \triangleright Confidence parameter
 $\alpha \in (0, 1), T$ \triangleright Learning rate; # of iterations

Output: Optimal μ, ρ \triangleright Centers, scales

```
1: procedure PB_QUAD_GAUSS
2:    $\mu \leftarrow \mu_0$   $\triangleright$  Set posterior centers to init. of prior
3:    $\rho \leftarrow \rho_0$   $\triangleright$  Set posterior scale to  $\rho_0$  hyperparam.
4:   for  $t \leftarrow 1 : T$  do  $\triangleright$  Run SGD for T iterations.
5:     Sample  $V \sim \mathcal{N}(0, I)$ 
6:      $W = \mu + \log(1 + \exp(\rho)) \odot V$ 
7:      $f(\mu, \rho) = f_{\text{quad}}(Z_{1:n}, W, \mu, \rho, \mu_0, \rho_0, \delta)$ 
8:     SGD gradient step using  $\begin{bmatrix} \nabla_{\mu} f \\ \nabla_{\rho} f \end{bmatrix}$ 
9:   return  $\mu, \rho$ 
```

PAC-Bayes workflow [Pérez-Ortiz et al., 2021]





- Split the dataset in two separate subsets $\mathcal{S} = \mathcal{S}_{\text{prior}} \cup \mathcal{S}_{\text{eval}}$.
- Learn the prior using $\mathcal{S}_{\text{prior}}$ by ERM with dropout.
- Learn the posterior using the whole dataset \mathcal{S} ,

$$\min_{\rho} \left\{ \hat{R}_{\mathcal{S}}^x(\rho) + \sqrt{\frac{\text{KL}(\rho \parallel \pi) + \log\left(\frac{2\sqrt{n}}{\delta}\right)}{2n}} \right\}.$$





- Evaluate the bound at the learned posterior ρ using $\mathcal{S}_{\text{eval}}$,

$$\text{kl}^{-1} \left(\text{kl}^{-1} \left(\hat{R}_{\mathcal{S}}^{01}(\tilde{\rho}_m), \frac{\log\left(\frac{2}{\delta'}\right)}{m} \right), \frac{\text{KL}(\rho \parallel \pi) + \log\left(\frac{2\sqrt{|\mathcal{S}_{\text{eval}}|}}{\delta}\right)}{2|\mathcal{S}_{\text{eval}}|} \right).$$

MNIST experiments [Pérez-Ortiz et al., 2021]



CIFAR 10 experiments [Pérez-Ortiz et al., 2021]



Conclusions of [Pérez-Ortiz et al., 2021]

- Model selection feasible without data splitting.
- Non-vacuous and tight bounds achievable.
- Choosing a prior centered at the ERM is key.
- Different trade-offs between test error and risk certificate.
- Extensive experiments for FCNs and CNNs.
- How about specific models ?
- How about different learning strategies ?

Generalization bounds for SGD using information bounds

Stochastic Gradient Descent

SGD algorithm :

$$\theta_{t+1} = \theta_t - \eta_t g(\theta_t; X_{l_t}, X_{l_t})$$

where η_t is the learning rate, l_t is the index set of minibatch of datapoints (ind. of \mathcal{S}) at step t , and $g(\theta; x, y) = \nabla_{\theta} \ell(f_{\theta}(x), y)$ is the gradient (averaged over the minibatch). The stepsize and sampling rule are fixed but arbitrary.

Stochastic Gradient Descent

SGD algorithm :

$$\theta_{t+1} = \theta_t - \eta_t g(\theta_t; X_{l_t}, X_{l_t})$$

where η_t is the learning rate, l_t is the index set of minibatch of datapoints (ind. of \mathcal{S}) at step t , and $g(\theta; x, y) = \nabla_{\theta} \ell(f_{\theta}(x), y)$ is the gradient (averaged over the minibatch). The stepsize and sampling rule are fixed but arbitrary.

(Expected) generalization gap of SGD

$$\mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \leq ?$$

Stochastic Gradient Descent

SGD algorithm :

$$\theta_{t+1} = \theta_t - \eta_t g(\theta_t; X_{l_t}, X_{l_t})$$

where η_t is the learning rate, l_t is the index set of minibatch of datapoints (ind. of \mathcal{S}) at step t , and $g(\theta; x, y) = \nabla_{\theta} \ell(f_{\theta}(x), y)$ is the gradient (averaged over the minibatch). The stepsize and sampling rule are fixed but arbitrary.

(Expected) generalization gap of SGD

$$\mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \leq ?$$


Question : when does SGD generalize ?

Attempt by Neu, Dziugaite, Haghifam & Roy (COLT 2021)
via Information bounds !

When does a predictor generalize?


When does a predictor generalize?

- Flatness (Hochreiter & Schmidhuber, Neural computation 1997, Keskar et al., ICLR 2017)
 - belief that algorithms that find wide optima of the loss landscape generalize well to test data
 - some flaws (difficult to define, parameterization,...)



When does a predictor generalize?

- Flatness (Hochreiter & Schmidhuber, Neural computation 1997, Keskar et al., ICLR 2017)
 - belief that algorithms that find wide optima of the loss landscape generalize well to test data
 - some flaws (difficult to define, parameterization,...)



- Stability (Hardt, Recht & Singer, ICML 2016)
 - SGD has strong stability conditions
 - stability improves as assumptions get stronger

Fixing the mutual information bounds

The problem with mutual information bounds?

Fixing the mutual information bounds

The problem with mutual information bounds? The MI between deterministic quantities is equal to $+\infty$!

Fixing the mutual information bounds

The problem with mutual information bounds? The MI between deterministic quantities is equal to $+\infty$!

Idea : *artificially* perturb the iterates!

Fixing the mutual information bounds

The problem with mutual information bounds? The MI between deterministic quantities is equal to $+\infty$!

Idea : *artificially* perturb the iterates! We can define a randomly perturbed version of SGD :

$$\tilde{\theta}_{t+1} = \theta_t + \zeta_t \quad \text{with} \quad \zeta_t = \sum_{k=1}^{t-1} \varepsilon_k \sim \mathcal{N}(0, \sigma_{1:t}^2),$$

$$\tilde{\theta}_{t+1} = \tilde{\theta}_t - \eta_t g(\theta_t; X_{l_t}, Y_{l_t}) + \varepsilon_t \quad \text{where} \quad \varepsilon_t \sim \mathcal{N}(0, \sigma_t^2 I).$$

Fixing the mutual information bounds

The problem with mutual information bounds? The MI between deterministic quantities is equal to $+\infty$!

Idea : *artificially* perturb the iterates! We can define a randomly perturbed version of SGD :

$$\tilde{\theta}_{t+1} = \theta_t + \zeta_t \quad \text{with} \quad \zeta_t = \sum_{k=1}^{t-1} \varepsilon_k \sim \mathcal{N}(0, \sigma_{1:t}^2),$$

$$\tilde{\theta}_{t+1} = \tilde{\theta}_t - \eta_t g(\theta_t; X_{l_t}, Y_{l_t}) + \varepsilon_t \quad \text{where} \quad \varepsilon_t \sim \mathcal{N}(0, \sigma_t^2 I).$$

We get :

$$\begin{aligned} \mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] &= \mathbb{E}_{\zeta_T, \mathcal{S}} \left[R(\tilde{\theta}_T) - \hat{R}_{\mathcal{S}}(\tilde{\theta}_T) \right] \\ &+ \mathbb{E}_{\zeta_T, \mathcal{S}, \mathcal{S}'} \left[\hat{R}_{\mathcal{S}'}(\theta_T) - \hat{R}_{\mathcal{S}'}(\tilde{\theta}_T) \right] + \mathbb{E}_{\zeta_T, \mathcal{S}, \mathcal{S}'} \left[\hat{R}_{\mathcal{S}}(\tilde{\theta}_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \\ &\leq \sqrt{\frac{2 \cdot \mathcal{I}(\tilde{\theta}_T; \mathcal{S})}{n}} + \mathbb{E}_{\mathcal{S}, \mathcal{S}'} [\Delta_{\sigma_{1:T}}(\theta_T, \mathcal{S}') - \Delta_{\sigma_{1:T}}(\theta_T, \mathcal{S})]. \end{aligned}$$

Main result

A generalization bound depending on the :

Main result

A generalization bound depending on the :

- variance of the gradients along the SGD path,

Main result

A generalization bound depending on the :

- variance of the gradients along the SGD path,
- perturbation-sensitivity of the gradients along the SGD path,

Main result

A generalization bound depending on the :

- variance of the gradients along the SGD path,
- perturbation-sensitivity of the gradients along the SGD path,
- perturbation-sensitivity of the loss at the final output.

Main result

A generalization bound depending on the :

- variance of the gradients along the SGD path,
- perturbation-sensitivity of the gradients along the SGD path,
- perturbation-sensitivity of the loss at the final output.

Thm [Neu, Dziugaite, Haghifam & Roy, COLT 2021] : assume that $\ell(\cdot, \cdot) \leq 1$, then for any $(\sigma_1, \dots, \sigma_T)$, $\sigma_{1:T} = \sqrt{\sum_{k=1}^{T-1} \sigma_k^2}$,

$$\left| \mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \right| \leq \sqrt{\frac{4}{n} \sum_{t=1}^T \frac{\eta_t^2}{\sigma_t^2} \mathbb{E}_{\mathcal{S} \sim P_{\mathcal{S}}} [\Gamma_{\sigma_{1:t}}(\theta_t) + V_t(\theta_t)]} + \left| \mathbb{E}_{\mathcal{S}, \mathcal{S}'} [\Delta_{\sigma_{1:T}}(\theta_T, \mathcal{S}') - \Delta_{\sigma_{1:T}}(\theta_T, \mathcal{S})] \right|.$$

Variance of the gradients

The gradient variance V_t measures the variability of the gradients with respect to the randomness of the data :

$$V_t(\theta) = \mathbb{E}_{\mathcal{S}} \left[\left\| g(\theta; X_{I_t}, Y_{I_t}) - \bar{g}(\theta) \right\|_2^2 \middle| \theta_t = \theta \right]$$

where $\bar{g}(\theta) = \mathbb{E}_{(X,Y) \sim \mathbb{P}}[g(\theta; X, Y)]$.



Sensitivity of the gradients

The gradient sensitivity Γ_σ measures the variability of the gradients to small perturbations in the parameter space.


$$\Gamma_\sigma(\theta) = \mathbb{E}_{(X,Y) \sim \mathbb{P}, \zeta \sim \mathcal{N}(0, \sigma^2 I)} \left[\|g(\theta, Z) - g(\theta + \zeta, Z)\|_2^2 \right].$$



Value sensitivity

The value sensitivity Δ_σ measures the variability of the loss function to small perturbations in the parameter space.

$$\Delta_\sigma(\theta, s) = \mathbb{E}_{\zeta \sim \mathcal{N}(0, \sigma^2 I)} \left[\|\hat{R}_s(\theta) - \hat{R}_s(\theta + \zeta)\|_2^2 \right].$$



Summary of the quantities

Thm : for any $(\sigma_1, \dots, \sigma_T)$, $\sigma_{1:T} = \sqrt{\sum_{k=1}^{T-1} \sigma_k^2}$, for losses ≤ 1 ,

$$\begin{aligned} \left| \mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \right| &\leq \sqrt{\frac{4}{n} \sum_{t=1}^T \frac{\eta_t^2}{\sigma_t^2} \mathbb{E}_{\mathcal{S}} [\Gamma_{\sigma_{1:t}}(\theta_t) + V_t(\theta_t)]} \\ &\quad + |\mathbb{E}_{\mathcal{S}, \mathcal{S}'} [\Delta_{\sigma_{1:T}}(\theta_T, \mathcal{S}') - \Delta_{\sigma_{1:T}}(\theta_T, \mathcal{S})]|, \end{aligned}$$

with the variance of the gradients along the SGD path

$$V_t(\theta) = \mathbb{E}_{\mathcal{S}} \left[\|g(\theta; X_t, Y_t) - \bar{g}(\theta)\|_2^2 \middle| \theta_t = \theta \right],$$

the sensitivity of the gradients along the SGD path

$$\Gamma_{\sigma_{1:t}}(\theta) = \mathbb{E}_{(X, Y) \sim \mathbb{P}, \zeta \sim \mathcal{N}(0, \sigma_{1:t}^2 I)} \left[\|g(\theta; X, Y) - g(\theta + \zeta; X, Y)\|_2^2 \right],$$

and the sensitivity of the loss at the final output :

$$\Delta_{\sigma_{1:t}}(\theta, \mathcal{S}) = \mathbb{E}_{\zeta \sim \mathcal{N}(0, \sigma_{1:t}^2 I)} \left[\|\hat{R}_{\mathcal{S}}(\theta) - \hat{R}_{\mathcal{S}}(\theta + \zeta)\|_2^2 \right].$$

Result for smooth functions

Assume that :

- $\eta_t = \eta$ and minibatches of size b ,
- for each $i = 1, \dots, n$, there is exactly one index t such that $i \in I_t$,
- $\mathbb{E}_{(X,Y) \sim \mathbb{P}}[\|g(\theta; X, Y) - \bar{g}(\theta)\|_2^2] \leq \nu$ for all θ ,
- ℓ is globally μ -smooth i.e.

$$\|g(\theta; x, y) - g(\theta + u; x, y)\|_2 \leq \mu \|u\|_2$$

for all θ, u and all x, y .

Result for smooth functions

Assume that :

- $\eta_t = \eta$ and minibatches of size b ,
- for each $i = 1, \dots, n$, there is exactly one index t such that $i \in I_t$,
- $\mathbb{E}_{(X,Y) \sim \mathbb{P}}[\|g(\theta; X, Y) - \bar{g}(\theta)\|_2^2] \leq \nu$ for all θ ,
- ℓ is globally μ -smooth i.e.

$$\|g(\theta; x, y) - g(\theta + u; x, y)\|_2 \leq \mu \|u\|_2$$

for all θ, u and all x, y .

Proposition : for any σ ,

Result for smooth functions

Assume that :

- $\eta_t = \eta$ and minibatches of size b ,
- for each $i = 1, \dots, n$, there is exactly one index t such that $i \in I_t$,
- $\mathbb{E}_{(X,Y) \sim \mathbb{P}}[\|g(\theta; X, Y) - \bar{g}(\theta)\|_2^2] \leq \nu$ for all θ ,
- ℓ is globally μ -smooth i.e.

$$\|g(\theta; x, y) - g(\theta + u; x, y)\|_2 \leq \mu \|u\|_2$$

for all θ, u and all x, y .

Proposition : for any σ ,

$$\left| \mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \right| = \mathcal{O} \left(\sqrt{\frac{R^2 \eta^2 T}{\sigma^2 n} \left(\mu^2 \sigma^2 dT + \frac{\nu}{b} \right)} + \mu \sigma^2 dT \right)$$

Different rates and regimes

Proposition : for any σ ,

$$\left| \mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \right| = \mathcal{O} \left(\sqrt{\frac{R^2 \eta^2 T}{\sigma^2 n} \left(\mu^2 \sigma^2 dT + \frac{\nu}{b} \right)} + \mu \sigma^2 dT \right)$$

Different rates and regimes

Proposition : for any σ ,

$$\left| \mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \right| = \mathcal{O} \left(\sqrt{\frac{R^2 \eta^2 T}{\sigma^2 n} \left(\mu^2 \sigma^2 dT + \frac{v}{b} \right)} + \mu \sigma^2 dT \right)$$

- Small-batch SGD : $T = \mathcal{O}(n)$ and $b = \mathcal{O}(1)$.

Different rates and regimes

Proposition : for any σ ,

$$\left| \mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \right| = \mathcal{O} \left(\sqrt{\frac{R^2 \eta^2 T}{\sigma^2 n} \left(\mu^2 \sigma^2 dT + \frac{v}{b} \right)} + \mu \sigma^2 dT \right)$$

- Small-batch SGD : $T = \mathcal{O}(n)$ and $b = \mathcal{O}(1)$. With $\eta = \mathcal{O}(1/n)$ & $\sigma = \Theta(n^{-4/3})$:

$$\left| \mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \right| = n^{-1/3}.$$

Different rates and regimes

Proposition : for any σ ,

$$\left| \mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \right| = \mathcal{O} \left(\sqrt{\frac{R^2 \eta^2 T}{\sigma^2 n} \left(\mu^2 \sigma^2 d T + \frac{\nu}{b} \right)} + \mu \sigma^2 d T \right)$$

- Small-batch SGD : $T = \mathcal{O}(n)$ and $b = \mathcal{O}(1)$. With $\eta = \mathcal{O}(1/n)$ & $\sigma = \Theta(n^{-4/3})$:

$$\left| \mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \right| = n^{-1/3}.$$

- Large-batch SGD : $T = \mathcal{O}(\sqrt{n})$ & $b = \Omega(\sqrt{n})$.

Different rates and regimes

Proposition : for any σ ,

$$\left| \mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \right| = \mathcal{O} \left(\sqrt{\frac{R^2 \eta^2 T}{\sigma^2 n} \left(\mu^2 \sigma^2 d T + \frac{v}{b} \right)} + \mu \sigma^2 d T \right)$$

- Small-batch SGD : $T = \mathcal{O}(n)$ and $b = \mathcal{O}(1)$. With $\eta = \mathcal{O}(1/n)$ & $\sigma = \Theta(n^{-4/3})$:

$$\left| \mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \right| = n^{-1/3}.$$

- Large-batch SGD : $T = \mathcal{O}(\sqrt{n})$ & $b = \Omega(\sqrt{n})$. With $\eta = \mathcal{O}(1/T) = \mathcal{O}(1/\sqrt{n})$ and $\sigma = \Theta(1/\sqrt{n})$:

$$\left| \mathbb{E}_{\mathcal{S}} \left[R(\theta_T) - \hat{R}_{\mathcal{S}}(\theta_T) \right] \right| = 1/\sqrt{n}.$$

Many points to discuss

- Guarantees obtained for non-randomized predictors
- Small values of Γ , V and Δ imply good generalization
- How do we measure them ?
- Why would SGD make them small ?
- How do we adjust SGD so that they become smaller ?
- Is it necessary ?
- Limitations of the geometry
- Choice of the surrogate in the proof
- How about the subGaussian assumption ?

Next lecture : Variational inference !