

# The genus *Gelsemium*: An update

V. Dutt, S. Thakur<sup>1</sup>, V. J. Dhar<sup>2</sup>, A. Sharma<sup>3</sup>

Guru Gobind Singh College of Pharmacy, Yamuna Nagar, Haryana, <sup>1</sup>S.D. College of Pharmacy, Barnala, <sup>2</sup>Swift School of Pharmacy, Rajpura, Punjab, <sup>3</sup>University Institute of Pharmaceutical Science, Punjab University, Chandigarh, India

Submitted: 10-03-10

## ABSTRACT

The review includes 103 references on the genus *Gelsemium*, and comprises ethnopharmacology, morphology, phytoconstituents, pharmacological reports, clinical studies and toxicology of the prominent species of *Gelsemium*. Alkaloids and iridoids constitute major classes of phytoconstituents of the genus. Most popular species of the genus are the Asian *G. elegans* and the two North American related species, *G. sempervirens* and *G. rankinii*. *Gelsemium* species are categorized under medicinal as well as poisonous plants. Amongst various species, *G. elegans* and *G. sempervirens* possess medicinal value, and have been traditionally used as nervous system relaxant. These plants have been explored exhaustively for their anticancer activity. In the concluding part, the future scope of *Gelsemium* species has been emphasized with a view to establish their multifarious biological activities and mode of actions

**Key words:** Alkaloids, anticancer, *gelsemium*, Iridoids

## INTRODUCTION

The literature review on *Gelsemium* has been compiled using references from major databases such as Chemical Abstracts, Medicinal and Aromatic Plants Abstracts, PubMed, King's American Dispensatory, Raintree Nutrition Incorporation, Henriette's Herbal Homepage, National Agricultural Library (AGRICOLA), Duke's Phytochemical and Ethnobotany database, UK Cropnet Ethnobotany database, Archives of American Folk Medicine and USPTO Patent Full Text and Image database.

The available information on *Gelsemium* has been divided into six sections, that is, ethnopharmacology, morphology, phytoconstituents, pharmacological studies, clinical studies, toxicology, covering prominent species of *Gelsemium*. The ethnopharmacological section has been further subdivided into two sections, that is, traditional uses, and alternative and complementary medicinal uses. The reports, in which *Gelsemium* species have been used as domestic remedy by common men without any prescription for the treatment of various ailments, have been discussed under traditional uses. The subhead 'alternative and complementary medicinal uses' highlights *Gelsemium* species as medicine prescribed by medical practitioners for the treatment of various ailments. It also mentions uses for which *Gelsemium* species or their preparations are available in

the market. Under every section, *Gelsemium* species have been arranged in an alphabetical order.

## THE GENUS *GELSEMIUM*

### Taxonomic classification

Kingdom	: Plantae (plants)
Superkingdom	: Tracheobionta (Vascular plants)
Superdivision	: Spermatophyta (Seed plants)
Phylum	: Embryophyta (Higher plants)
Class	: Magnoliopsida (Dicotyledons)
Order	: Gentianales
Family	: Loganiaceae
Genus	: <i>Gelsemium</i>
Species	: <i>elegans</i> , <i>rankinii</i> , <i>sempervirens</i>

The genus *Gelsemium* belongs to family Loganiaceae, and comprises about five species that are widely distributed throughout Central America. *Gelsemium elegans* known as Lemuan is an extremely poisonous plant that is indigenous to the South East Asian countries and found predominantly in Malaysia and Sarawak.<sup>[1-3]</sup> Most popular species of the genus are the Asian *G. elegans* and the two North American related species, *G. sempervirens* and *G. rankinii*.<sup>[4]</sup> *Gelsemium* is a climbing plant with dark evergreen leaves and all parts are poisonous.<sup>[5,6]</sup>

## ETHNOPHARMACOLOGY

### Traditional uses

*Gelsemium elegans* has been traditionally used as a nervous system relaxant to treat various types of pain including headache and

### Address for correspondence:

Dr. Vandana Dutt,  
E-mail: Vandudatt2006@rediffmail.com

DOI: 10.4103/0973-7847.70916

pain associated with inflammatory conditions.<sup>[5,6]</sup> *Gelsemium* is one of homeopathy's important remedy for influenza.<sup>[7]</sup> *Gelsemium sempervirens* Ait. Syn. *G. nitidum*, commonly known as Yellow jasmine, has been used in the treatment of restlessness, mental irritability, insomnia, headache, irritation of the urinary tract, hyperemia and convulsions.<sup>[8]</sup> The roots of *G. sempervirens* have been used in the treatment of migraine, neuralgia, rheumatism, and in ovarian and uterine pain.<sup>[9]</sup> In the US, it has been extensively used as an arterial sedative and febrifuge in various fevers. It has been used in treatment of spasmodic disorders such as asthma and whooping cough. The plant has been used in hysteria, dysmenorrhea, chorea, pneumonia and bronchitis.<sup>[10]</sup>

### Alternative and complimentary medicinal uses

An herbal preparation containing *G. sempervirens* has been used for evening cold and influenza.<sup>[11]</sup> Pharmaceutical formulations containing *G. sempervirens* as one of the constituents have been used for the treatment of psoriasis and neurodermatitis.<sup>[12]</sup> Nebera *et al.* reported that homoeopathic preparations, containing *G. sempervirens* in the form of sugar granules, are used in treating myopia of I, II or III degree.<sup>[13]</sup> Fluid extract of the plant has been used for its spasmolytic actions.<sup>[14]</sup>

## MORPHOLOGY

*Gelsemium* includes three species of shrubs and climbers from United States and Central America and one species from China, South East Asia and Indo-Malaysia. *Gelsemium elegans* Benth. is a large, woody, evergreen climber with corky bark; leaves ovate or ovate-lanceolate; flowers golden yellow; sepals ovate, margin minutely ciliate; corolla funnel shaped, lobes imbricate in bud; ovules numerous in each cell; style filiform, at the apex with four short stigmatic branches.<sup>[15,16]</sup> *Gelsemium sempervirens* is a climbing shrub indigenous to the Southern United States from Virginia to Florida and Texas.<sup>[17]</sup> The plant is supplied commercially as segments of the cylindrical rhizome with attached wiry roots;<sup>[18]</sup> rhizome horizontal, the segments 3 to 20 cm in length and 3 to 30 mm in diameter, externally moderate brown to dark yellowish-orange, frequently spirally twisted, longitudinally wrinkled, with purplish-brown longitudinal lines and transverse fissures, upper surface with few stem scars, the under and lateral portions with several roots and root scars, fracture of rhizome tough, splintery, internally exhibiting a narrow purplish-brown bark, a broad, pale yellowish orange to light yellow, finely radiate and eccentric wood, a minute disintegrated pith; roots up to 20 cm in length and 2 to 8 mm thick, light brown, nearly smooth and wiry; fracture one-half transverse, the other oblique and splintery, fractured surface showing a broad, radiate, yellow wood and a thin bark.<sup>[15]</sup>

## PHYTOCONSTITUENTS

A survey of literature reveals that alkaloid constitutes the major class of phytoconstituents in *Gelsemium* species. Table 1 summarizes phytoconstituents reported from various species of *Gelsemium*.

## PHARMACOLOGICAL STUDIES

It has been reported that alkaloidal fraction isolated from *G. elegans* exhibits analgesic and anti-inflammatory activities.<sup>[72]</sup> A growth stimulant for pig has been prepared from gouwen (*G. elegans*) that served to promote nutrient absorption of organism, to increase immunity, and raise lean-to-fat pork ratio.<sup>[73]</sup> Cueilleron *et al.* reported that *Gelsemium* tincture at higher concentration inhibits dopamine, noradrenaline and serotonin uptake into synaptosomal preparations from different parts of the rat brain, whereas lower concentrations enhances noradrenaline and serotonin uptake into mesencephale preparation.<sup>[74]</sup> Methanol extract of *G. elegans* leaves exhibited high cytotoxicity against the human ovarian cancer cell lines CaOV-3 with an IC50 value of 5 µg/ml after 96 h of incubation while less toxicity against the human breast cancer cells MDA-MB-231 suggesting its selectivity towards CaOV-3 cells.<sup>[75]</sup> Uncarinic acid E, isolated from *G. elegans*, reported to exhibit antitumor effects due to its growth inhibitory activity for HepG2 cells in dose-dependent manner.<sup>[76]</sup> Gelsedine type alkaloids of *G. elegans* exhibited potent cytotoxic activity in an A431 human epidermoid carcinoma cell line.<sup>[52-77]</sup> Koumine (50 mmol/L) induced apoptosis of human colon adenocarcinoma LoVo cells in a time-dependent manner, and inhibited DNA synthesis in LoVo cells *in vitro*.<sup>[78]</sup> Koumine (20–320 µg/ml) dose dependently inhibited concanavalin A or phytohemagglutinin-induced proliferation of murine lymphocytes determined by MTT colorimetry.<sup>[79]</sup> It also decreased IL-2 level in the cell culture supernatant measured by enzyme-linked immunosorbent assay. Koumine showed remarkable inhibitory effect on mouse vaginal epithelial cell mitosis and promoted the formation of epidermal glandular layer in the scales at the mouse tail.<sup>[80]</sup> It also decreased serum IL-2 level in mice at concentrations 6, 30, 150 mg/kg. These activities infer that the therapeutic effect of Koumine against psoriasis is related to the inhibition of epidermal cell proliferation, promoting the formation of glandular cells and decreasing the serum level of IL-2. Koumine injection significantly decreased mouse spontaneous activity in moderate and high doses but did not produce any effect on the respiratory and cardiovascular system of dogs.<sup>[81]</sup>

Low doses of *G. sempervirens* and *Atropa belladonna* showed significant neurotropic and protective effects on behavioral and gastric alterations induced by foot shock stress in mice.<sup>[82]</sup> Ethanolic extract of *G. sempervirens* has been reported to increase the resistance of rabbits to pneumococcus toxin.<sup>[83]</sup> *G. sempervirens* alkaloidal fraction possessed anticancer activity as evidenced by significant inhibition of hepatic carcinoma HepG<sub>2</sub> cells *in vitro* using crystal violet dyeing method.<sup>[84]</sup> Sempervirine nitrate, isolated from *G. sempervirens*, exhibited antimitotic activity in mice bearing several types of tumors.<sup>[85]</sup> Sempervirine has also been reported to possess vasoconstrictor action in the perfused isolated rabbit ear.<sup>[86]</sup> In an experiment performed by Hinsdale on isolated rat intestine, *G. sempervirens* produced an immediate relaxation of the intestine tissue.<sup>[87]</sup> *G. sempervirens* and *Datura stramonium* in appropriate concentration reported to prevent development of spontaneous seizures induced in rats by lithium or pilocarpine.<sup>[88]</sup>

## CLINICAL STUDIES

Gouwen injection, prepared by using alkaloids extracted from *G. elegans* roots, has been reported to exhibit antitumor effect against esophagus cancer in humans.<sup>[89]</sup> An injection containing *G. sempervirens* extract as one of the ingredients when given to patients suffering from neurodermatitis, resulted in subsidence of itching associated with neurodermatitis after 1 week, disappearance of large erythema after 3–4 weeks and elimination of disease after 6 weeks.<sup>[12]</sup>

## TOXICOLOGY

*Gelsemium* acts in the similar manner as nicotine and coniine. Sometimes small doses of *G. sempervirens* can produce toxic symptoms.<sup>[9]</sup> A drachm of fluid extract of the plant can cause death, and 30 minims are dangerous. An alkaloid gelsemicine from *G. sempervirens* is toxic, and symptoms of toxicity are depressed respiration, tremors, paralysis of extremities, convulsions, urination, defecation, retchings and salivation.<sup>[90]</sup> Gelsemicine, in small doses, stimulates respiration and paralyzes the respiratory centers in larger doses.<sup>[91]</sup> The minimum lethal dose is 0.02 to 0.03 mg/g, s.c. for frogs, 0.00010 to 0.00012 mg/g, s.c. for rats, 0.00005 to 0.00006 mg/g, i.v. for rabbits and 0.0005 to 0.001 mg/g, i.v. for dogs. E-Bay website rated extracts of *Gelsemium* 'extremely toxic'.<sup>[92]</sup> The roots of *G. sempervirens* contain a resin, which is poisonous in very small doses. A tincture prepared by digesting it in undiluted alcohol is fatal. Small doses of *G. sempervirens* relax the muscles but larger doses cause dropping of the lower jaws and difficulty in managing the eyelids. The continued administration of it affects





the brain spinal centers and medulla, causing marked feebleness of muscular movements, confusion of vision and vertigo. Large doses paralyze the spinal cord and cause almost complete loss of muscular power. These effects are due to its action upon the spinal marrow. The characteristic toxic symptoms are peripheral relaxation, disturbance of the ocular muscles, drooping of the lower jaw and profound prostration and muscular relaxation. When applied locally to the eye, it dilates the pupils. Overdosage of the plant may cause death due to asphyxia. Alkaloid fraction, isolated from *G. elegans* leaves, at a lethal dose produced violent clonic convulsions that led to respiratory failure. Authors suggested that alkaloids act centrally against GABA as evidenced by prevention of convulsions by pentobarbital or diazepam and potentiation by reserpine.<sup>[8]</sup>

Persons are reported to have been poisoned by eating honey gathered by the bees from *G. sempervirens* flowers. *Gelsemium* when administered to rabbits and guinea pigs produced a marked generalized congestion of all organs, depressive action on heart and respiration and severe toxic action on liver, kidney and testes.<sup>[93]</sup> Histological studies showed neurological signs characterized by marked progressive weakness and convulsions culminating in death in three goats over a 24-h period after ingestion of *G. sempervirens* leaves.<sup>[94]</sup> Aqueous extract of the plant greatly depressed the activity of the isolated frog heart muscles.<sup>[95]</sup> It has been reported that single intravenous injection of gelsemine (0.2 mg/kg), isolated from *G. sempervirens*, in chloralosed dog produced a marked and prolonged decrease in blood pressure.<sup>[96]</sup> Alkaloids gelsemine, sempervirine and gelsemicine, isolated from *G. sempervirens*, increased the hypertensive action of adrenaline<sup>[97]</sup> and inhibited the cholinesterases of nervous tissue and serum.<sup>[97]</sup> The drug is used as poison as its effects vary rapidly.<sup>[98,99]</sup>




**Table 1: Phytoconstituents of various species of *Gelsemium***

Species	Phytoconstituents
<i>G. elegans</i>	Alkaloids koumine, <sup>[1]</sup> kounidine, kouminine, kouminidine, koumicine, kouminicine, <sup>[19-28]</sup> 19-(R)- and 19-(S)-hydroxydihydrokoumine, <sup>[2,3,29]</sup> 19-(R)- and 9-(S)-koumino, <sup>[4,5,30]</sup> humanenine, <sup>[6]</sup> 11-hydroxyhumanenine, <sup>[7]</sup> 15-hydroxyhumanenine, 11-methoxy-humanenine, <sup>[8]</sup> N-desmethoxyhumanenine, <sup>[9]</sup> rankinidine, <sup>[10]</sup> 11-hydroxyrankinidine, <sup>[11]</sup> N-desmethoxy-rankinidine, <sup>[12]</sup> 20-hydroxydihydrorankinidine, humanenirine, <sup>[13]</sup> humantendine, <sup>[14]</sup> gelsemoxonine, <sup>[15,31-36]</sup> gelsamydine, <sup>[37]</sup> 19 $\alpha$ -hydroxygelsamydine, <sup>[16]</sup> 14 $\alpha$ -hydroxygelsamydine, <sup>[17]</sup> 14 -hydroxyelegansamine, <sup>[18]</sup> Gelsebanine, <sup>[19,38,39]</sup> koumidine, <sup>[20]</sup> 19-(Z)-akuaminidine, <sup>[21]</sup> 16-epi-voacarpine, <sup>[22]</sup> 19-hydroxy-dihydrogelsevirine (40), gelsemine, <sup>[20-22,23]</sup> gelsevirine, <sup>[24]</sup> 19-(R)-acetyldihydrogelsevirine, 19-(R)-hydroxydihydrogelsemine, <sup>[41]</sup> gelsemicine, <sup>[25]</sup> gelsedine, <sup>[26,42]</sup> elegansamine, <sup>[27,43]</sup> N-methoxy-anhydrobasinediol, <sup>[28,44]</sup> gelsemamide, <sup>[29]</sup> 11-methoxygelsemamide, <sup>[45]</sup> 11-methoxy-19-(R)-hydroxygelselegine, <sup>[29,30]</sup> sempervirine, <sup>[46-50]</sup> Gelsedilam, <sup>[31]</sup> 14-acetoxygelsedilam, <sup>[32]</sup> gelsefuranidine, <sup>[33]</sup> gelseinidone, <sup>[34,51]</sup> 14 acetoxygelsenicine, <sup>[35]</sup> 14 acetoxo-15-hydroxygelsenicine, <sup>[36]</sup> 14-acetoxo-19-oxogelsenicine, <sup>[37]</sup> 14-acetoxygelselegine, <sup>[38,52]</sup> humanenine N(4)-oxide, <sup>[100]</sup> Iridoids (53) gelsemide, <sup>[39]</sup> GEIR-1, <sup>[40]</sup> GRIR-1, <sup>[41]</sup> GEIR-2, <sup>[42]</sup> GEIR-3, <sup>[43]</sup> Di(2-ethylhexyl) phthalate, 3 $\beta$ -hydroxyl-27-p-(Z)-coumaroyloxy ursan-12-en-28-oic acid, 3 $\beta$ -hydroxyl-27-p-(E)-coumaroyloxy ursan-12-en-28-oic acid, uncarinic acid (Wei et al., 2007), $\beta$ -sitostreol, stigmasterol, daucosterol, stigmasterol, $\beta$ -D-glucopyranoside, ursolic acid, gallic acid, ferullic acid, protocate chulic acid. <sup>[101]</sup> Glycosides; Gelsemiunside A and B. <sup>[102]</sup>
<i>G. rankini</i> Small	Alkaloids 21-oxogelsevirine, 21-oxogelsimine, gelsivirine, <sup>[24]</sup> gelsemine, <sup>[24,54]</sup> rankinidine, <sup>[10]</sup> humanenirine, <sup>[14]</sup> humanenine, <sup>[6,55]</sup> Iridoids gelsemide, <sup>[39]</sup> GEIR-1, <sup>[40]</sup> GRIR-1, <sup>[41]</sup> GEIR-2, <sup>[42]</sup> GEIR-3. <sup>[43,53]</sup>
<i>G. sempervirens</i>	Alkaloids gelsemine, <sup>[23]</sup> gelseminine, gelsemide, gelsemicine, <sup>[25,56-62]</sup> 21-oxogelsemine, <sup>[63]</sup> gelsevirine, <sup>[24,64]</sup> gelsedine, <sup>[24,48]</sup> 14-hydroxygelsemicine, <sup>[65,66]</sup> 14 $\beta$ -hydroxygelsedine, <sup>[67,68]</sup> sempervirine, <sup>[46,47,60]</sup> gelsedine type oxindole alkaloids, <sup>[69]</sup> gelsemic acid, <sup>[70]</sup> Iridoids gelsemide, <sup>[39]</sup> gelsemide-7-glucoside, gelsemiol, <sup>[44]</sup> gelsemiol 1- and 3-glucoside, 9-hydroxysemperoside, <sup>[71]</sup> 2008 Steroids pregna-4, 16-diene-3,20-dione, 12 $\beta$ -hydroxy-5 $\alpha$ -pregna-16-ene-3,20-dione, 12 $\beta$ -hydroxypregna-4, 16-diene-3,20-dione, <sup>[68]</sup> Scopoletin, 7-O-beta-D-glucopyranosylscopoletin, 7-O-beta-D-apiofuranosyl-(1->6) $\beta$ -lucopyranosylscopoletin, Uvaol, 2-(4-hydroxyphenyl) ethylheptadecanoate. <sup>[103]</sup>

## APPENDIX





**1****2** 19-(R)**3** 19-(S)**R****4** CHOHCH<sub>3</sub> (R)**5** CHOHCH<sub>3</sub> (S)**R<sub>1</sub>****R<sub>2</sub>****R<sub>3</sub>**

<b>6</b>	OCH <sub>3</sub>	CH <sub>3</sub>	H
<b>7</b>	OCH <sub>3</sub>	CH <sub>3</sub>	OH
<b>8</b>	OCH <sub>3</sub>	CH <sub>3</sub>	OCH <sub>3</sub>
<b>9</b>	H	CH <sub>3</sub>	H
<b>10</b>	OCH <sub>3</sub>	H	H
<b>11</b>	OCH <sub>3</sub>	H	OH
<b>12</b>	H	H	H
<b>13</b>	OCH <sub>3</sub>	H	OCH <sub>3</sub>


**14****15****16**

Appendix (contd...)


## Appendix (contd...)




	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>
<b>20</b>	CH <sub>2</sub> OH	H	H	H
<b>21</b>	CO <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> OH	H	H
<b>22</b>	CH <sub>2</sub> OH	CO <sub>2</sub> CH <sub>3</sub>	OH	H



<b>23</b>	H	H <sub>2</sub>
<b>24</b>	OCH <sub>3</sub>	H <sub>2</sub>




<b>25</b>	H	H	OCH <sub>3</sub>
<b>26</b>	H	H	H




Appendix (contd...)


Appendix (contd...)




28



29




30




31 R = H


32 R = OAc



33




34



35 R = H

36 R = OH









37

Appendix (contd...)



## Appendix (contd...)



## CONCLUSION

About 05 species of the genus *Gelsemium* have been reported in various floras. An exhaustive survey of literature revealed that information is available on 03 species. Among these 05 species, most of ethnopharmacological reports are available on *G. elegans*

and *G. sempervirens*. Further, these plants [Table 1] have been investigated for their phytoconstituents.

*Gelsemium sempervirens* has been included in a number of herbal and homoeopathic formulations, which are in clinical use for the treatment of various ailments. Mother tinctures of the

plant are available in Indian market, and are frequently used for the treatment of CNS disorders, but no pharmacological work supports its efficacy in CNS disorders. Keeping in view the traditional, alternative and complimentary medicinal uses, and frequency of use in homoeopathic formulations, *G. sempervirens* seems to hold great potential for in-depth investigation on various biological activities, especially its effect on the central nervous systems.

A close scrutiny of literature on *Gelsemium* reveals that two species have been investigated pharmacologically. Pharmacological studies infer that *G. elegans* exhibits analgesic, antiinflammatory and cytotoxic properties; *G. sempervirens* exhibits neurotropic and antitumor activities. Koumine, gelsedine type alkaloids and uncarinic acid E have been considered bioactive constituents of *G. elegans*. Toxicological studies have confirmed high toxicity of *Gelsemium* species at higher doses. These plants have narrow therapeutic index, that is, margin between therapeutic efficacy and toxic value is very less. Although the plants of the genus *Gelsemium* hold great potential to be developed as antitumor drugs, their toxicity could not be ruled out.

## ACKNOWLEDGMENTS

The authors are grateful to Mr. Bhupinder Singh Jauhar, Chairman, Guru Nanak Khalsa Group of Institutions, Yamuna Nagar for encouragement, financial assistance and providing laboratory facilities for carrying out this research work and to Dr. R.P. Kapoor and Dr. P. Das for their insight on the work.

## REFERENCES

- Latiff A, Ibrahim AZ, Mat-Salleh K. An account and checklist of the flowering plants at Kelabit Highlands. Bario Sarawak: ASEAN Review of Biodiversity and Environmental Conservation; 1999. p. 15.
- Ridley HN. The Flora of the Malay Peninsular. Ashford, London: Reeve and Co.; 1925.
- Perry LM. Medicinal plant of East and Southeast Asia: Attributed properties and uses. Cambridge, MA: MIT Press; 1980.
- Burkhill JA. A Dictionary of the economic products of the Malay Peninsular. Kuala Lumpur: Ministry of Agriculture and Cooperatives; 1966.
- Ponglux D, Wongseripipatana S, Subhadhirasakul S. Studies on the indole alkaloids of *Gelsemium elegans* (Thailand): Structure elucidation and proposal of biogenetic route. *Tetrahedron* 1988;44:5075-4.
- Watson L, Dallwitz MJ. Families of Flowering Plants: Descriptions, Illustrations, Identification, and Information Retrieval; 2002.
- Blackwell WH. Poisonous and Medicinal plants. Prentice Hall, New Jersey: Englewoods and Cliff; 1990. p. 329.
- Felter HW, Lloyd JU. *Gelsemium* (U.S.P.). King's American Dispensatory; 1898.
- Chopra RN. Poisonous plants of India. India: Council of Agricultural Research; 1940. p. 692.
- Grieve M. A Modern Herbal (first Published in 1931). London: Tiger Books International; 1994. p. 249.
- Voorhees J, Nachman L. inventors: Herbal composition for treating symptoms of influenza. US patent 6455070; 2002. P. 3.
- Calarasu C, inventor; Pharmaceuticals containing diphenhydramine and Echinacea and Eupatorium and Gelsemium and Lachesis extracts for the treatment of Psoriasis and neurodermatitis. German patent 3641220; 1988. P. 4.
- Nebera SA, Nebera OA, Pakhomova NN. inventors: Homoeopathic preparation for treating myopia. RU patent 2203674; 2003.
- Uhlenbroock K, Schweer M, Maschmann L. Spasmolytic effect of the ingredients of *Gelsemium* fluid extract. *Arzneimittelforschung* 1959;9:419-22.
- The Wealth of India. A Dictionary of Indian Raw Materials and Industrial Product. New Delhi: Council of Scientific and Industrial Research; 1956. p. 123.
- Brandis D. Indian Trees. In: Singh B, Singh MP, editors. Dehradun, India; 1987. p. 476.
- Wallis T. Textbook of Pharmacognosy. 5th ed. New Delhi: CBS Publishers and Distributors; 1999. p. 387, 558-62.
- Gahalot A, Patra A, Sharma A. Pharmacognostical studies on *Gelsemium sempervirens* Ait. *Roots P' Cog Mag*; 2008.
- Chou TQ. The alkaloids of *Gelsemium elegans* Benth. III. *Zhangguo Shenglixue Zazhi* 1931;5:345-52.
- Chou TQ, Wang CH, Cheng WC. The alkaloids of Chinese gelsemium. *Ta-ch'a-yeh Zhangguo Shenglixue Zazhi* 1936; 10: 79-84.
- Chi YF, Kao YS, Hueng YT. Alkaloids of Chinese gelsemium, Kou-wen. *J Am Chem Soc* 1938;60:1723-4.
- Liu CT, Loh JY, Liu CC, Lu JY, Chu TC, Wang CH. *Gelsemium* alkaloids. I. Reinvestigation of the alkaloids of *Gelsemium elegans* and the constitution of koumine. *Huaxue xuebao* 1961;27:47-58.
- Khuong-Huu F, Chiaroni A, Riche C. Structure of koumine, an alkaloid from *Gelsemium elegans* Benth. *Tetrahedron Lett* 1981;22:733-4.
- Jin H, Xu RS. Studies on the alkaloids of *Gelsemium elegans* Benth. The structure of koumidine. *Acta Chim Sin* 1982;40: 1129-35.
- Lounasmaa M, Koskinen A. A plausible biogenetic proposal for koumine. *Planta Med* 1982;44:120-1.
- Yang J, Chen Y. Chemical studies on the alkaloids from Hu Man Teng (*Gelsemium elegans* Benth). II. structure of humantenidine. *Yaoxue Xuebao* 1984;19:437-40.
- Zhang L, Lin J, Wu Z. Advances in the study on chemical constituents and pharmacology of *Gelsemium elegans* (Gardn. et Champ.) Benth. *Zhong Yao Cai* 2003;26:451-3.
- Zhang LL, Wang ZR, Huang CQ, Zhang ZY, Lin JM. Extraction and separation of koumine from *Gelsemium* alkaloids. *Di Yi Jun Yi Da Xue Xue Bao* 2004;24:1006-8.
- Lin LZ, Cordell GA, Ni CZ, Clardy J. Two oxindole alkaloids from *Gelsemium elegans*. *Phytochem* 1990;29:3013-7.
- Sun F, Xing QY, Liang XT. Structure of (19R)-kouminol and (19S)-kouminol from *Gelsemium elegans*. *J Nat Prod* 1989;52:1180-2.
- Yang J, Chen Y. Chemical study of the alkaloids of *Gelsemium elegans*. *Yaoxue Tongbao* 1982;17:119-20.
- Yang J, Chen Y. Chemical studies on the alkaloids from Hu Man Teng (*Gelsemium elegans* Benth.) II. structure of humantendine. *Yaoxue Xuebao* 1982;17:633-4.
- Yang JS, Chen YW. Chemical studies on the alkaloids of Hu-



- Man-Teng (*Gelsemium elegans* Benth.). I. Isolation of the alkaloids and structure of humantenmine. Yao Xue Xue Bao 1983;18:104-12.
34. Lin LZ, Cordell GA. New humantenine-type alkaloids from *Gelsemium elegans*. J Nat Prod 1989;52:588-94.
  35. Lin LZ, Cordell GA, Ni CZ, Clardy J. Oxindole alkaloids from *Gelsemium elegans*. Phytochemistry 1991;30:1311-5.
  36. Kitajima M, Kogure N, Yamaguchi K, Takayama H, Aimi N. Structure reinvestigation of gelsemoxonine, a constituent of *Gelsemium elegans*, reveals a novel azetidine containing indole alkaloid. Org Lett 2003;5:2075-8.
  37. Lin LZ, Cordell GA, Ni CZ, Clardy J. Gelsamydine, an indole alkaloid from *Gelsemium elegans* with two monoterpene units. J Org Chem 1989;54:3199-202.
  38. Lin LZ, Hu SF, Cordell GA. 19  $\alpha$ -Hydroxygelsamydine from *Gelsemium elegans*. Phytochemistry 1996;43:723-6.
  39. Xu YK, Yang SP, Liao SG. Alkaloids from *Gelsemium elegans*. J Nat Prod 2006;69:1347-50.
  40. Sakai S, Wongseripipatana S, Ponglux D, Yokota K, Ogata K, Takayama M, *et al.* Indole alkaloids isolated from *Gelsemium elegans* (Thailand): 19-(z)-aluammidine, 16-epivocarpine, 19-hydroxydihydro gelsevirine, and the revised structure of koumidine. Chem Pharm Bull 1987;35:4668-71.
  41. Lin LZ, Cordell GA, Ni CZ, Clardy J. Three oxindole Alkaloids from *Gelsemium* species. Phytochemistry 1991;30:679-83.
  42. Du X, Dai Y, Zhang C, Lu S, Liu Z. Studies on the *Gelsemium* alkaloids. I. The structure of gelsenicine. Huaxue Xuebao 1982;40:1137-41.
  43. Ponglux D, Wangseripipatana S, Takayama H, Ogata K, Aimi N, Sakai S. A new class of indole alkaloid, elegansamine, constructed from a monoterpene indole alkaloid and an iridoid. Tetrahedron Lett 1988;29:5395-6.
  44. Lin LZ, Cordell GA, Ni CZ, Clardy J. N-methoxyanhydrovobasinediol from *Gelsemium elegans*. Phytochemistry 1989;28:2827-31.
  45. Lin LZ, Cordell GA, Ni CZ, Clardy J. Gelsemamide and 11-methoxy gelsemamide, two novel secoindole alkaloids from *Gelsemium elegans*. Tetrahedron Lett 1989;30:1177-80.
  46. Stevenson AE, Sayre LE. Third alkaloid of *Gelsemium*. J Am Pharm Assoc 1915;4:60-2.
  47. Stevenson AR, Sayre LE. Sempervirine from *Gelsemium* root. J Am Pharm Assoc 1915;4:1458-63.
  48. Sayre LE. Final report on the alkaloids of *Gelsemium*. J Am Pharm Assoc 1919;8:708-11.
  49. Hasenfratz V. The presence of an oxygen-free alkaloid in *Gelsemium sempervirens*. Comptes Rendus des Seances de la Societe de Biologie et de ses Filiales 1933;196:1530-2.
  50. Janot MM, Goutarel R, Perezamador Y, Barron MC. Isolation of sempervirine from *Gelsemium elegans*. Ann Pharm Fr 1953;11:602-8.
  51. Kogure N, Ishii N, Kitajima M, Wongseripipatana S, Takayama H. Four novel gelsenicine-related oxindole alkaloids from the leaves of *Gelsemium elegans* Benth. Org Lett 2006;8:3085-8.
  52. Kitajima M, Nakamura T, Kogure N, Ogawa M, Mitsuno Y, Ono K, *et al.* Isolation of gelsedine-type indole alkaloids from *Gelsemium elegans* and evaluation of the cytotoxic activity of gelsemium alkaloids for A431 epidermoid carcinoma cells. J Nat Prod 2006;69:715-8.
  53. Kogure N, Ishii N, Kobayashi H, Kitajima M, Wongseripipatana S, Takayama H. New iridoids from *Gelsemium* species. Chem Pharm Bull 2008;56:870-2.
  54. Chun Y, Cordell GA, Garland M. 21-oxogelsevirine, a new alkaloid from *Gelsemium rankinii*. J Nat Prod 1986;49:483-7.
  55. Chun Y, Cordell GA. Rankinidine, a new indole alkaloid from *Gelsemium rankinii*. J Nat Prod 1986;49:806-8.
  56. Sayre LE. Gelseminine and other constituents of *Gelsemium*. Pharm J 1911;86:242-3.
  57. Sayre LE. A study of the more toxic alkaloids of *Gelsemium*. Midland Druggist and Pharmaceutical Review 1912;45:439-41.
  58. Moore CW. Constituents of *Gelsemium*. J Chem Soc Abs 1911;97:2223.
  59. Chou TA. The alkaloids of *Gelsemium*. I. Gelsemine and gelsemicine. Zhongguo shenglixue Zazhi 1931;5:131-9.
  60. Ferrerio M. The alkaloids of *Gelsemium sempervirens* Ait. (Loganiaceae). Anal Sci Nat Bot 1945;6:148.
  61. Janot MM, Gautarol R, Friedrich W. Gelsemicine. Ann Pharm Fr 1951;9:305-7.
  62. Schwarz H, Marion L. Three minor alkaloids of *Gelsemium sempervirens*. Can J Chem 1953;31:958-75.
  63. Nikiforov A, Latzel J, Varmuza K, Wichtl M. *Gelsemium* alkaloids.3. structure elucidation of 21-oxogelsemine by mass spectrometry. Monatsh Chem 1974;105:1292-8.
  64. Wenkert E, Chang CJ, Cochran DW, Pollicciari R. Carbon-13 nuclear magnetic resonance spectroscopy of naturally occurring substances. VIII. Structure of gelsevirine. Experientia 1972;28:377-9.
  65. Wichtl M, Nikiforov A, Schulz G, Sponer S, Jentzsch K. Structural elucidation of 21-oxogelsemine by NMR and mass spectrometry. Monatsh Chem 1973;104:99-104.
  66. Wichtl M, Nikiforov A, Sponer S, Jentzsch K. Structure of *Gelsemium* alkaloid A. mass spectrometry of *Gelsemium* alkaloids. Monatsh Chem 1973;104:87-98.
  67. Chun Y, Cordell GA. 14  $\beta$ -Hydroxygelsedine, a new oxindole alkaloid from *Gelsemium sempervirens*. J Nat Prod 1985;48:788-91.
  68. Chun Y, Cordell GA. Cytotoxic steroids of *Gelsemium sempervirens*. J Nat Prod 1987;50:195-8.
  69. Kitajima M, Urano A, Kogure N, Takayama H, Aimi N. New oxindole alkaloids and iridoid from Carolina jasmine (*Gelsemium sempervirens* Ait.). Chem Pharm Bull 2003;51:1211-4.
  70. Chou TQ. The alkaloids of *Gelsemium* II. Sempervine and an amorphous alkaloid. Zhongguo Shenglixue Zazhi 1931;5:295-300.
  71. Jensen SR, Kirk O, Nielsen BJ, Norrestam R. 9-hydroxy substituted iridoids from *Gelsemium sempervirens*. Phytochemistry 1987;26:1725-31.
  72. Rujjanawate C, Kanjanapothi D, Panthong A. Pharmacological effect and toxicity of alkaloids from *Gelsemium elegans* Benth. J Ethnopharmacol 2003;89:91-5.
  73. Hu X. inventor: Growth Stimulant for Swine. Chinese patent; 1997.
  74. Cueilleron P, Fonlupt P, Pacheco H. Effect of *Gelsemium* tincture on dopamine, noradrenaline and serotonin uptake in synaptosomal preparations of rat brain. Comptes Rendus des Seances de l'Academie des Sciences, Serie 2 1982;294:853-5.
  75. Abdul Wahab K, Ahmad FB, Din LB, Swee Hung C, Shiueh Lian M. *In vitro* A study of the cytotoxic activity of *Gelsemium elegans* using human ovarian and breast cancer cell lines. Trop Biomed 2004;21:139-44.
  76. Zhao M, Guo T, Wang M, Minwei W, Zhao Q, Liu Y, *et al.* The course of uncarinic acid E-induced apoptosis of hep G2 cells from damage to DNA and p53 activation to mitochondrial

- release of cytochrome C. *Biol Pharm Bull* 2006;29:1639-44.
77. Kitajima M. Chemical studies on monoterpenoid indole alkaloids from medicinal plant resources *Gelsemium* and *Ophiorrhiza*. *J Nat Med* 2006;61:14-23.
  78. Chi DB, Lei LS, Jin H, Pang JX, Jiang YP. *In vitro* Study of koumine-induced apoptosis of human colon adenocarcinoma LoVo cells. *Di Yi Jun Yi Da Xue Xue Bao* 2003;23:911-3.
  79. Wang ZR, Huang CQ, Zhang ZY, Zhang LL, Lin JM. *In vitro* Effect of koumine on proliferation of murine CD4+ T cells purified by magnetic-activated cell sorting. *Di Yi Jun Yi Da Xue Xue Bao* 2005;25:562-4.
  80. Zhang LL, Huang CQ, Zhang ZY, Wang ZR, Lin JM. Therapeutic effects of koumine on psoriasis: an experimental study in mice. *Di Yi Jun Yi Da Xue Xue Bao* 2005;25:547-9.
  81. Chi DB, Lei LS, Yang HX, Sun LS. General pharmacology of koumine parenteral solution. *Di Yi Jun Yi Da Xue Xue Bao* 2004;24:32-4.
  82. Bousta D, Soulimani R, Jarmouni I, Belon P, Falla J, Froment N, *et al.* Neurotropic, immunological and gastric effects of low doses of *Atropa belladonna* L., *Gelsemium sempervirens* L. and *Poumon histamine* in stressed mice. *J Ethnopharmacol* 2001;74:205-15.
  83. Baker WF. *Gelsemium sempervirens* and *Bryonia alba* in influenza. *J Am Inst Homeopath* 1920;12:695-8.
  84. Wang Y, Fang Y, Lin W, Cheng M, Jiang Y, Yin M. *In vitro* Inhibitory effect of *Gelsemium* alkaloids extract on hepatic carcinoma HepG2 cells. *Zhong Yao Cai* 2001;24:579-81.
  85. Bassleer R, Clerment D, Marnette JM, Caprasse M, Tits M, Angenot L. Effect of dihydroflavopereirine and sempervirine ( $\beta$ -carbolinium alkaloid) on cancer cells in culture. *Ann Pharm Fr* 1985;43:83-8.
  86. Rothlin E, Raymond H. Vascular action of sempervirine, an oxygen-free alkaloid of *Gelsemium sempervirens*. *Comptes Rendus des Seances de la Societe de Biologie et de ses Filiales* 1934;117:754-6.
  87. Hinsdale AE. Action of *Gelsemium* upon intestinal movement. *J Am Inst Homeopath* 1918;10:969-70.
  88. Peredery O, Persinger MA. Herbal treatment following post-seizure induction in rat by lithium pilocarpine: *Scutellaria lateriflora* (Skullcap), *Gelsemium sempervirens* (*Gelsemium*) and *Datura stramonium* (Jimson Weed) may prevent development of spontaneous seizures. *Phytother Res* 2004;18:700-5.
  89. Dai R. Preparation of Gouwen injection. *Zhong Yao* 1993;24:471-2.
  90. Hou HC. The pharmacological action of gelsemicine, an alkaloid from *Gelsemium*. I. Acute toxicity. *Zhangguo Shenglixue Zazhi* 1931;5:181-90.
  91. Raymond H. Gelsemicine. *Comptes Rendus des Seances de la Societe de Biologie et de ses Filiales* 1937;126:1151-4.
  92. Cantrell FL. Look what I found. Poison hunting on e-Bay. *Clin Toxicol (Phila)* 2005;43:375-9.
  93. Mitchell T. *Gelsemium sempervirens*. *J Am Inst Homeopath* 1926;19:707-13.
  94. Thompson LJ, Frazier K, Stiver S, Styer E. Multiple animal intoxications associated with Carolina jasmine (*Gelsemium sempervirens*) ingestions. *Vet Hum Toxicol* 2002;44:272-3.
  95. Espanes D, Moisset E. Action of extract of *Gelsemium sempervirens* on the isolated frog heart. *Comptes Rendus des Seances de la Societe de Biologie et de ses Filiales* 1938;127:1088-90.
  96. Espanes D, Moisset E. Cardiovascular action of gelsemine. Action on the carotid pressure of the dog. *Comptes Rendus des Seances de la Societe de Biologie et de ses Filiales* 1938;127:1002-4.
  97. Vincent D, Lagreu R. Action of the alkaloids of *Gelsemium sempervirens* on cholinesterases. *Comptes Rendus des Seances de la Societe de Biologie et de ses Filiales* 1951;145:348-50.
  98. Kitajima M, Arai Y, Takayama H, Aimi N. A chemical study on 'Yakatsu' stored in Shosoin repository: Isolation and characterization of four indole alkaloids from a 1250 year-old sample of the Chinese toxic medicine. *Proc Japan Acad* 1998;74: 159-63.
  99. Blaw ME, Adkisson MA, Levin D, Garriott JC, Tindall RS. Poisoning with Carolina jessamine (*Gelsemium sempervirens* [L.] Ait.). *J Pediatr* 1979;94:998-1001.
  100. Yin S, He XF, Wu Y, Yue JM. Monoterpenoid indole alkaloids bearing an N4-Iridoid from *Gelsemium elegans*. *Chem Asian J* 2008;3:1824-9.
  101. Qing-chun Z, Yan-Hui F, Tao G, Wei H, Li-jun W. Isolation and identification of non alkaloids constituents from *G. elegans* Benth. *Yaoke Daxue Xuebao* 2007;24:619-22.
  102. Hua W, Zhao QC, Yang J, Shi GB, Wu LJ, Guo T. Two new benzofuran from lignan Glycosides. *Chinese Chem Lett* 2008;19:1327-9.
  103. Zhang Z, Wang P, Yuan W, Li S. Steroids, alkaloids and coumarins from *G. sempervirens*. *Planta Med* 2008;74: 1818-22.

**Source of Support:** Guru Nanak Khalsa Group of Institutions,  
**Conflict of Interest:** None declared

Copyright of Pharmacognosy Reviews is the property of Medknow Publications & Media Pvt. Ltd. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.